Published in final edited form as:

Neurosci Lett. 2013 August 26; 548: 27–32. doi:10.1016/j.neulet.2013.05.027.

Intrathecal neurosteroids and a neurosteroid antagonist: Effects on inflammation-evoked thermal hyperalgesia and tactile allodynia

Elin Svensson*, Josefin Persson@, Bethany Fitzsimmons, and Tony L. Yaksh

Department of Anesthesiology, University of California, San Diego.

Abstract

Neurosteroids regulate neuronal excitability through binding sites associated with the ionotropic γ -aminobutyric acid (GABA_A) receptor. We sought to characterize the spinal analgesic actions in rats of two 5 α -reduced neurosteroids, allopregnanolone and alphaxalone, on nociceptive processing and to determine whether a putative neurosteroid antagonist attenuates this effect: (3 α , 5 α)-17-phenylandroster-16-en-3-ol (17PA). Intrathecal (IT) injection of allopregnanolone (1–30 μ g/10 μ L in 20% cyclodextrin) delivered through lumbar catheters produced a dose-dependent analgesia in rats as measured by thermal thresholds in the ipsilateral (inflamed by intraplantar carrageenan) and in the contralateral (un-inflamed paws). Similar observations were made with alphaxalone (30–60 μ g in 20% cyclodextrin). Effective doses were not associated with suppressive effects on pinnae, blink or placing and stepping reflex. Effects of allopregnanolone (30 μ g) on the normal and hyperalgesic paw were completely prevented by IT 17PA (30 μ g). Reversal by IT 17PA of an equi-analgesic dose of alphaxalone occurred only at higher antagonist dosing. These results suggest that a spinal neurosteroid-binding site with which 17 PA interacts may regulate spinal nociceptive processing in normal and inflamed tissue. (NIDA DA02110) (171 words)

Keywords

Neurosteroid; allopregnanolone; alphaxalone; hyperalgesia; Tactile allodynia; GABA_A; intrathecal

Introduction

Neurosteroids have analgesic, anesthetic and anticonvulsant effects.^{14, 26} Spinal delivery of 5 α -reduced neurosteroids such as allopregnanolone (3 α , 5 α -tetrahydroprogesterone or 3 α , 5 α -THP) reduces the facilitated nociceptive response produced by repetitive nerve stimulation^{4, 20, 23} or visceral inflammation³¹. Antagonism of these effects by IT bicuculline⁹ suggests that these agents facilitate activation by γ -amino butyric acid (GABA) of the GABA_A chloride ionophore and at higher concentrations by directly activating the ionophore,¹ promoting the channel open state.² Molecular studies have

Corresponding author: Tony L. Yaksh, Department of Anesthesiology, University of California, San Diego, 9500 Gilman drive, La Jolla Ca 92103. Tel: 619-543-3597. tyaksh@ucsd.edu.

*Present address: Svartbäcksgatan 83B753 33 Uppsala Sweden

@Present address: St Johannesgatan 32B, 75233 Uppsala Sweden

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

The first two authors contributed equally to this work.

shown two neurosteroid binding sites that mediate the potentiating and direct channel activating effect of neurosteroids.¹² Ample evidence supports the ability of neurosteroids to amplify dorsal horn GABA_A receptor activity regulating components relevant to dorsal horn pain processing.^{6,24}

Evidence for the action of neurosteroids at a specific site is further provided by the observation that neurosteroid activity is antagonized by agents such as (3 α ,5 α)-17-phenylandroster-16-en-3-ol (17PA). 17PA has little effect on baseline GABA-mediated Cl flux but antagonizes gating and conductance augmentation produced by 5 α -reduced steroids, such as allopregnanolone.^{15,18} We note that two 5 α -reduced steroids, allopregnanolone and alphaxalone ((3 α ,5 α)-3-hydroxypregnane-11,20-dione), both produce sedation but only the effects of allopregnanolone are reversed by 17PA when administered systemically *in vivo*. This differential effect has been taken to support different neurosteroid binding sites.¹⁵ Whether this differential effect is observed *viz* the analgesic effects of these two neurosteroids is not known. Thus, we examined: i) spinal actions of allopregnanolone and alphaxalone in regulating inflammation evoked hyperalgesia; ii) dose dependency of this spinal action; and, iii) whether IT 17PA would antagonize spinal actions of IT allopregnanolone and/or alphaxalone, supporting an interaction with a spinal neurosteroid binding site.

Materials and Methods

Animals

Adult male rats (Holtzman, 200-250g) (12/12 day-night cycle, *ad libitum* access to food/water) were used in accordance with protocols approved by the Institutional Animal Care and Use Committee at the University of California, San Diego. To allow IT delivery, rats were prepared with chronic lumbar IT catheters placed 8.5 cm from the cisterna magna and externalized on the top of the head.^{17, 35}

Drugs

The 5 α -reduced neurosteroid agonists, allopregnanolone and alphaxalone, and the neurosteroid antagonist (3 α ,5 α)-17-phenylandroster-16-en-3-ol (17PA) (Tocris Bioscience, Ellisville, MO, USA) were delivered in β -cyclodextrin (20% w/v) (Sigma-Aldrich, St Louis, MO, USA) and 0.9% NaCl with the final doses delivered in 10 μ L followed by 10 μ l of saline. Vehicle controls were the cyclodextrin formulation. Solutions were made and injected by two different individuals and testing was without knowledge of the solution.

Behavioral testing

IT drugs were delivered 30 min prior to unilateral intraplantar delivery of carrageenan (2% in 100 μ L) at time = 0. For thermal threshold assessments, a Hargreaves system (UCSD University Anesthesia Research and Development Group, La Jolla, Ca)⁸ was employed. After 60 min adaptation, a radiant heat stimulus was delivered to each individual paw and the latency to withdraw noted.⁸ To prevent tissue damage, a cut-off time of 20 sec was used. In a limited series of experiments, we assessed tactile thresholds using von Frey hairs as previously described using the “up-down” method, with the maximum pressure being 15 grams.⁵ The effects of treatment on behavior and function were assessed before and during the subsequent 4 hr post injection test interval during which antinociception was observed by noting the presence or absence of i) hindpaw placing / stepping response, ii) pinnae response, iii) blink (corneal) reflex, iv) righting reflex and v) normal ambulation

Statistical analysis

Data are presented as mean and SEM of thermal escape latency (seconds) or tactile threshold (grams). Drug response data were converted to a Hyperalgesic Index (area under the curve for % Hyperalgesia: e.g. [Premeasurement – Postmeasurement / Premeasurement] $\times$ 100). In each case, hyperalgesia is indicated by a larger value whereas reversal of hyperalgesia or analgesia is indicated by a smaller or negative number. Time-dependent response data were analyzed by two-way analysis of variance-repeated measures across time. Hyperalgesic indices between treatment groups were compared using 1-way ANOVA, with a Bonferroni multiple comparison analysis for *post hoc* comparisons. Differences reaching a $p < 0.05$ level of significance were considered to be significant. Statistics were carried out using Prism (v4.2c) (LaJolla, Ca).

RESULTS

Agonist effects upon thermal hyperalgesia

Baseline thermal escape latencies prior to drug treatment was similar for all treatment groups (approximately 9-11 sec). In vehicle treated animals, intraplantar carrageenan reduced the thermal escape latency of the inflamed (ipsilateral) paw, which persisted undiminished through the 4 hr time point (approximately 3-5 sec, see Figure 1), with no change in the normal (contralateral) paw withdrawal latency. IT cyclodextrin (20%/10 μ L) had no effect upon the thermal withdrawal latency of the ipsilateral vs. the contralateral paw, as compared to IT saline (data not shown). IT allopregnanolone (3-30 μ g/10 μ L) resulted in a significant dose-dependent reversal of the hyperalgesic paw latency (as measured by the latency and the hyperalgesic index) and an increase in latency above baseline for the uninjured paw (Fig. 1). Intraperitoneal delivery of the maximum IT allopregnanolone dose (30 μ g) had no effect upon thermal withdrawal latencies of either the ipsilateral or contralateral paw ($N = 4$; $p > 0.05$), indicating that the IT drug effect did not represent a simple systemic redistribution (data not shown). The effects of IT alphaxalone (30 and 60 μ g/10 μ L) were examined on thermal escape. As shown (Fig. 2), a significant increase in the thermal escape latency was noted with both the ipsilateral (hyperalgesic) and contralateral paw as

Agonist effects upon tactile allodynia

After carrageenan, rats displayed a tactile allodynia in the inflamed (ipsilateral) paw with no change in the contralateral paw. Rats pre-treated with IT allopregnanolone or alphaxalone (but not 20% cyclodextrin) displayed a reduction in the tactile allodynia otherwise observed after intraplantar carrageenan as compared to vehicle: Post IT Vehicle threshold at 3 hr. = 3.1 ± 0.6 grams vs. IT allopregnanolone, 30 μ g/10 μ L: 11.6 ± 1.9 grams; ($p < 0.05$ vs. vehicle) or alphaxalone (60 μ g/10 μ L): 13.5 ± 2.1 grams ($p < 0.05$ vs. vehicle) (data not shown).

Agonist effects upon behavior

Over the intrathecal doses examined for alphaxalone (30-60 μ g) and allopregnanolone (3-30 μ g), there was no change in the ability to evoke a pinnae or blink response. There was no loss of hind paw placing/stepping, or the righting response secondary to being placed recumbent. In the uninflamed (contralateral) paw, there was typically a modest increase in the thermal escape latency, but most animals even at the highest doses continued to display withdrawal. Though not systematically examined, in the absence of stimulation, high dose alphaxalone animals appeared to show less spontaneous activity in the test cage, than adjacent animals run in parallel receiving vehicle only, though they were not immobile and showed the previously noted evoked responses.

Intrathecal Antagonist

IT 17 PA (30 μ g/10 μ L) alone had no effect upon either the hyperalgesic thermal escape latency (ipsilateral paw) or the normal escape latency (contralateral paw). To examine antagonism by 17PA of the agonist-evoked antihyperalgesia, IT 17 PA (30 μ g) was given 15 min prior to the agonist. At -20 min, the just maximally effective IT doses of allopregnanolone (30 μ g/10 μ L) or alphaxalone (60 μ g/10 μ L) were given, followed at T = 0 by intraplantar carrageenan. IT 17 PA pretreatment (Fig. 3) resulted in a near complete reversal of the antinociceptive effects of IT allopregnanolone as measured by the latency and the hyperalgesic index on both the ipsilateral and contralateral paw (e.g. no statistical difference between Allopregnanolone + 17PA vs. vehicle).

In contrast, as shown in Figure 4, IT 17PA (30 μ g) produced only a modest none significant reversal of the antihyperalgesic actions of IT alphaxalone in the ipsilateral and contralateral paws. Increasing the 17PA dose to 60 μ g resulted in a reduction in the antihyperalgesic effects in the ipsilateral paw and contralateral paw.

In separate experiments, we sought to determine if the post-treatment of 17PA at 120 min post-carrageenan would reverse the effects of IT allopregnanolone (30 μ g/10 μ L). In these studies (data not shown), there was a modest but statistically significant reversal after the IT 17PA injection: Post-Vehicle latency = 3.4 ± 0.8 sec vs. Post 17PA 6.9 ± 1.8 sec ($p < 0.05$).

DISCUSSION

Intrathecal delivery of 5 α -reduced neurosteroid agonists allopregnanolone and alphaxalone in a cyclodextrin-water-based vehicle reverse, in a dose-dependent fashion, thermal and mechanical hyperalgesia. These effects occurred at doses that are not systemically active and which have no effect upon general function or motor function. These results are in accord with previous work showing that IT delivery of several neurosteroid-like molecules produces a significant effect upon facilitated pain states.^{9, 20, 31} Aside from their antihyperalgesic action, these agents also modestly increased the response latency of the un-inflamed paw, suggesting that their action was not limited to a facilitated state. This observation is comparable to work showing the effects of IT neurosteroids on tail flick response latency.⁹

Spinal pharmacology of neurosteroid agonist action

Neurosteroids interact with an allosterically coupled binding complex of the GABA_A chloride ionophore. This coupling augments the GABA initiated opening of the channel. At higher concentrations, neurosteroids can activate the channel. The likelihood that the neurosteroids are exerting their spinal actions through a GABA_A channel is supported by reports that the antihyperalgesic actions are prevented by GABA_A block with bicuculline.⁹

Both allopregnanolone and alphaxalone display potent anesthetic effects. Although previous systemic work with alphaxalone reported no specific antinociceptive effects²⁰, both agents comparably activate the GABA_A ionophore⁷ and prevent thermal hyperalgesia. To address the likelihood that these agents were interacting with a neurosteroid binding site after IT delivery, we examined the ability of the putative neurosteroid antagonist 17PA to reduce the observed antihyperalgesic action. This agent behaves as a competitive antagonist at a GABA_A-associated binding site.¹⁸ IT-17PA abolished the antihyperalgesic and analgesic actions of IT-allopregnanolone. The antihyperalgesic effects of an equianalgesic IT dose of alphaxalone were also reversed, albeit only at a higher 17PA dose. Given the assertion that 17PA is a competitive antagonist, then the apparent difference in antagonist potency of 17PA against equianalgesic doses of the two agonists suggests distinguishable binding sites for these two agonists, e.g. a binding site interaction with an alphaxalone site at which 17PA

had a lower affinity than the binding site for allopregnanolone. Whether or not this hypothesis is correct, the observations showing a different antagonist potency are comparable to previous work in which it was shown that systemic 17PA reversed the sedative effects of allopregnanolone, but had little effect upon sedation after alphaxalone.¹⁵

Cyclodextrin as a vehicle

Cyclodextrin was employed as a vehicle compatible with IT delivery. It had no effect when given alone as compared to IT saline. IT cyclodextrin is a vehicle for solubilizing some lipophilic compounds, such as capsaicin, small peptides and neurosteroids.³⁴ Cyclodextrin the permitted use of a water-based vehicle that is generally preferable to other lipid solubilizing or detergent-like vehicles for neuraxial delivery. There are several *caveats* to the use of IT cyclodextrin. First, previous work with gamma-cyclodextrin suggested that the complex may result in a reduction in the apparent potency of steroids. Nevertheless, the use of IT β -cyclodextrin revealed a monotonic dose-effect relationship with drug effects of several hours duration. Accordingly, to the degree that cyclodextrin sequesters neurosteroids, the observed actions are compatible with a dose-proportional delivery of drug to the active site after IT delivery. Second, β -cyclodextrins can exert direct potentiating effects on GABA_A receptors.²⁵ We did not see evidence of this action with the present study, as the IT cyclodextrin alone had no effect upon either normal or hyperalgesic paw latencies. Whether other response systems might be affected by this action remains to be determined. Third, cyclodextrin binds cholesterol and extracts cholesterol from cell membranes and disrupts caveolae formation.^{10, 21, 36}

Mechanisms of neurosteroid associated antinociception

The interaction of neurosteroids with the GABA ionophore involves specific subunits, notably those of the alpha and beta forming binding pockets with which the neurosteroids may interact.^{11,12} In the spinal cord, alpha subunits are expressed in the superficial dorsal horn³ on primary afferents, which, because of the superficial laminar distribution, likely represent nociceptive afferents, and on second order neurons. The presynaptic location is believed to represent mechanisms underlying mediation by GABA of preterminal inhibition (e.g. primary afferent depolarization)²⁷ while the postsynaptic effect is considered to be associated with a hyperpolarization resulting from the increased chloride conductance.³¹ Knockout of the $\alpha 2$ - GABA_A subunit revealed that the antihyperalgesic effects of spinal benzodiazepines were prevented.³² We note that the reduced effect of IT-17PA on the antihyperalgesic effects of alphaxalone in contrast to the allopregnanolone may also be interpreted as suggesting multiple potential targets other than the GABA_A receptor. Interactions of neurosteroids with NMDA ionophores^{16, 13, 29} and T type calcium channels²² have been described. Neurosteroids have also been shown to interact with glycine receptor mediated chloride currents.^{19, 30} The role of these interactions at the spinal level is not excluded in the present study and requires further investigation. We are however aware of no data considering an interaction of 17 PA with such binding.

Conclusions

We demonstrated that two structurally similar 5 α -reduced neurosteroids produce a potent antinociception and antihyperalgesic action. A putative neurosteroid receptor antagonist had distinguishable potency in reversing the effects of the two agents, consistent with the hypothesis of possible multiple targets for spinal neurosteroids.

Acknowledgments

Funding was provided by NIDA 02110. We would like to thank Damon McCumber for performing some of the studies shown here.

References

1. Akk G, Covey DF, Evers AS, Steinbach JH, Zorumski CF, Mennerick S. Mechanisms of neurosteroid interactions with GABA_A receptors. *Pharmacology and Therapeutics*. 2007; 116:35–57. [PubMed: 17524487]
2. Belelli D, Lambert JJ. Neurosteroids: endogenous regulators of the GABA_A receptor. *Nature reviews*. 2005; 6:565–574.
3. Bohlhalter S, Weinmann O, Mohler H, Fritschy JM. Laminar compartmentalization of GABA_A-receptor subtypes in the spinal cord: an immunohistochemical study. *J Neurosci*. 1996; 16:283–297. [PubMed: 8613794]
4. Chang JL, Peng HY, Wu HC, Lu HT, Pan SF, Chen MJ, Lin TB. Acute neurosteroids inhibit the spinal reflex potentiation via GABAergic neurotransmission. *Am J Physiol Renal Physiol*. 2010; 299:F43–8. [PubMed: 20357028]
5. Chaplan SR, Bach FW, Pogrel JW, Chung JM, Yaksh TL. Quantitative assessment of tactile allodynia in the rat paw. *J Neurosci Methods*. 1994; 53:55–63. [PubMed: 7990513]
6. Chéry N, De Koninck Y. Junctional versus extrajunctional glycine and GABA_A receptor-mediated IPSCs in identified lamina I neurons of the adult rat spinal cord. *J Neurosci*. 1999; 19:7342–7355. [PubMed: 10460241]
7. Cottrell GA, Lambert JJ, Peters JA. Modulation of GABA_A receptor activity by alloxalone. *Br J Pharmacol*. 1987; 90:491–500. [PubMed: 3032320]
8. Dirig DM, Salami A, Rathbun ML, Ozaki GT, Yaksh TL. Characterization of variables defining hindpaw withdrawal latency evoked by radiant thermal stimuli. *J Neurosci Methods*. 1997; 76:183–91. [PubMed: 9350970]
9. Goodchild CS, Guo Z, Nadeson R. Antinociceptive properties of neurosteroids I. Spinally-mediated antinociceptive effects of water-soluble aminosteroids. *Pain*. 2000; 88:23–2. [PubMed: 11098096]
10. Horikawa YT, Patel HH, Tsutsumi YM, Jennings MM, Kidd MW, Hagiwara Y, Ishikawa Y, Insel PA, Roth DM. Caveolin-3 expression and caveolae are required for isoflurane-induced cardiac protection from hypoxia and ischemia/reperfusion injury. *J Mol Cell Cardiol*. 2008; 44:123–30. [PubMed: 18054955]
11. Hosie AM, Clarke L, da Silva H, Smart TG. Conserved site for neurosteroid modulation of GABA_A receptors. *Neuropharmacology*. Jan. 2009; 56(1):149–54. [PubMed: 18762201]
12. Hosie AM, Wilkins ME, da Silva HM, Smart TG. Endogenous neurosteroids regulate GABA_A receptors through two discrete transmembrane sites. *Nature*. 2006; 444:486–9. [PubMed: 17108970]
13. Jang MK, Mierke DF, Russek SJ, Farb DG. A steroid modulatory domain on NR2B controls N-methyl-D-aspartate receptor proton sensitivity. *Proc. Natl. Acad. Sci. USA*. 2004; 101:8198–203. [PubMed: 15150412]
14. Jevtovic-Todorovic V, Covey DF, Todorovic SM. Are neuroactive steroids promising therapeutic agents in the management of acute and chronic pain? *Psychoneuroendocrinology*. 2009; 34:178–185.
15. Kelley SP, Alan JK, O'Buckley TK, Mennerick S, Krishnan K, Covey DF, Leslie Morrow A. Antagonism of neurosteroid modulation of native gamma-aminobutyric acid receptors by (3 α , 5 α)-17-phenylandrosterone-16-en-3-ol. *Eur J Pharmacol*. 2007; 572:94–101. [PubMed: 17658511]
16. Majewska MD. Neurosteroids: endogenous bimodal modulators of the GABA_A receptor. Mechanism of action and physiological significance. *Prog. Neurobiol*. 1992; 38:379–395. [PubMed: 1349441]
17. Malkmus SA, Yaksh TL. Intrathecal Catheterization and Drug Delivery in rat. *Pain Research, methods and protocols* edited by Z. David Luo, MD, PhD. Humana Press Inc. 2004:109–121.

18. Mennerick S, He Y, Jiang X, Manion BD, Wang M, Shute A, Benz A, Evers AS, Covey DF, Zorumski CF. Selective antagonism of 5 α -reduced neurosteroid effects at GABA(A) receptors. *Mol Pharmacol*. 2004; 65:1191–7. [PubMed: 15102947]
19. Mitchell EA, Gentet LJ, Dempster J, Belelli D. GABAA and glycine receptor-mediated transmission in rat lamina II neurones: relevance to the analgesic actions of neuroactive steroids. *J Physiol*. 2007; 583(Pt 3):1021–40. [PubMed: 17656439]
20. Nadeson R, Goodchild CS. Antinociceptive properties of neurosteroids II. Experiments with Saffan and its components alphaxalone and alphadolone to reveal separation of anaesthetic and antinociceptive effects and the involvement of spinal cord GABA_A receptors. *Pain*. 2000; 88:31–39. [PubMed: 11098097]
21. Ostrom RS, Liu X. Detergent and detergent-free methods to define lipid rafts and caveolae. *Methods Mol Biol*. 2007; 2(400):459–68. [PubMed: 17951752]
22. Pathirathna S, Brimelow BC, Jagodic MM, Krishnan K, Jiang X, Zorumski CF, Mennerick S, Covey DF, Todorovic SM, Jevtovic-Todorovic V. New evidence that both T-type calcium channels and GABAA channels are responsible for the potent peripheral analgesic effects of 5 α -reduced neuroactive steroids. *Pain*. 2005; 114:429–43. [PubMed: 15777868]
23. Peng HY, Chen GD, Lee SD, Lai CY, Chiu CH, Cheng CL, Chang YS, Hsieh MC, Tung KC, Lin TB. Neuroactive steroids inhibit spinal reflex potentiation by selectively enhancing specific spinal GABA(A) receptor subtypes. *Pain*. 2009; 143:12–20. [PubMed: 19250751]
24. Poisbeau P, Patte-Mensah C, Keller AF, Barrot M, Breton JD, Luis-Delgado OE, Freund-Mercier MJ, Mensah-Nyagan AG, Schlichter R. Inflammatory pain upregulates spinal inhibition via endogenous neurosteroid production. *J Neurosci*. 2005; 25:1768–76.
25. Pytel M, Mercik K, Mozrzymas JW. Interaction between cyclodextrin and neuronal membrane results in modulation of GABA_A receptor conformational transitions. *Br J Pharmacol*. 2006; 148:413–42. [PubMed: 16702996]
26. Reddy DS. Neurosteroids: endogenous role in the human brain and therapeutic potentials. *Prog Brain Res*. 2010; 186:113–37. [PubMed: 21094889]
27. Rudomin P. In search of lost presynaptic inhibition. *Exp Brain Res*. 2009; 196:139–51. [PubMed: 19322562]
28. Shu HJ, Zeng CM, Wang C, Covey DF, Zorumski CF, Mennerick S. Cyclodextrins sequester neuroactive steroids and differentiate mechanisms that rate limit steroid actions. *Lipids*. 2004; 39:811–9. [PubMed: 15638252]
29. Spivak V, Lin A, Beebe P, Stoll L, Gentile L. Identification of a neurosteroid binding site contained within the GluR2-S1S2 domain. *Br J Pharmacol*. 2007; 150:164–75. [PubMed: 17160009]
30. Weir CJ, Ling AT, Belelli D, Wildsmith JA, Peters JA, Lambert JJ. The interaction of anaesthetic steroids with recombinant glycine and GABAA receptors. *Br J Anaesth*. 2004; 92:704–11. [PubMed: 15033889]
31. Winfree CJ, Coombs DW, DeLeo JA, Colburn RW. Analgesic effects of intrathecally-administered 3 α -hydroxy-5 α -pregnan-20-one in a rat mechanical visceral pain model. *Life Sci*. 1992; 5:1007–12. [PubMed: 1552820]
32. Witschi R, Punnakal P, Paul J, Walczak JS, Cervero F, Fritschy JM, Kuner R, Keist R, Rudolph U, Zeilhofer HU. Presynaptic α 2-GABAA receptors in primary afferent depolarization and spinal pain control. *J Neurosci*. 2011; 31:8134–42. [PubMed: 21632935]
33. Wright R, Raimondo JV, Akerman CJ. Spatial and Temporal Dynamics in the Ionic Driving Force for GABA(A) Receptors. *Neural Plast*. 2011; 72:83–95.
34. Yaksh TL, Jang JD, Nishiuchi Y, Braun KP, Ro SG, Goodman M. The utility of 2-hydroxypropyl-beta-cyclodextrin as a vehicle for the intracerebral and intrathecal administration of drugs. *Life Sci*. 1991; 48:623–33. [PubMed: 1703620]
35. Yaksh TL, Rudy TA. Chronic catheterization of the spinal subarachnoid space. *Physiol Behav*. 1976; 17:1031–6. [PubMed: 14677603]
36. Zidovetzki R, Levitan I. Use of cyclodextrins to manipulate plasma membrane cholesterol content: evidence, misconceptions and control strategies. *Biochim Biophys Acta*. 2007; 1768:1311–24. [PubMed: 17493580]

High Lights

- Spinal allopregnanolone and alphaxalone increase thermal threshold in normal and inflamed paw.
- Effects of allopregnanolone were prevented by neurosteroid antagonist 17PA
- Reversal of an equi-analgesic dose of alphaxalone occurred only at higher antagonist dosing.
- Spinal neurosteroid binding site(s) acted upon by 17 PA regulates spinal pain processing.

Fig 1.

TOP: Effect of IT allopregnanolone or cyclodextrin (20%) vehicle on escape latency of ipsilateral (A) and contralateral (B) hindpaw as a function of time after ipsilateral intraplantar carrageenan given at T=0. BOTTOM: Hyperalgesic index for ipsilateral (C) and contralateral (D) hindpaw shows that IT alloprenanolone resulted in dose dependent reversal of thermal hyperalgesia of inflamed paw and a significant increase above baseline (analgesia) in contralateral, un-inflamed paw. Positive scores indicate hyperalgesia; negative scores indicate analgesia. Each plot presents means $\pm$ SEM, 5-9 animals. Comparisons: A,B: Two Way repeated measures ANOVA with Bonferroni *post hoc* comparison: *: $p < 0.05$ vs baseline; C,D: Bars under each AUC graph link treatments that do not differ from each other (Bonferroni multiple comparisons, $p < 0.05$)

Fig 2.

TOP: Effect of IT alphaxalone or cyclodextrin (20%) vehicle on escape latency of ipsilateral (A) and contralateral (B) hindpaw as a function of time after ipsilateral intraplantar carrageenan given at T=0. BOTTOM: Hyperalgesic index for ipsilateral (C) and contralateral (D) hindpaw shows that IT alloprenanolone resulted in dose dependent reversal of thermal hyperalgesia of the inflamed paw and a significant increase above baseline (analgesia) in contralateral, un-inflamed paw. Positive scores indicate hyperalgesia; negative scores indicate analgesia. Each plot presents means $\pm$ SEM, 5-9 animals. Comparisons: A,B: Two Way repeated measures ANOVA with Bonferroni *post hoc* comparison: *: $p < 0.05$ vs baseline; C,D: Bars under each AUC graph link treatments that do not differ from each other (Bonferroni multiple comparisons, $p < 0.05$)

Fig 3.

TOP: Effect on escape latency of IT neurosteroid antagonist, (17PA) on escape latency in the ipsilateral -carrageenan injected-(A) and the contralateral (B) hindpaw when given alone or in the presence of allopregnanolone (30µg/10µL). IT 17PA (30 µg/10µL) was given 15 min (arrow 1) prior to IT allopregnanolone or vehicle (arrow 2). Carrageenan was injected into the ipsilateral hind paw at t=0 (arrow 3). BOTTOM: Hyperalgesic index for ipsilateral (C) and contralateral (D) hindpaw shows that IT 17PA had no effect upon the carrageenan evoked thermal hyperalgesia, but reversed effects of IT allopregnanolone. Each plot presents means $\pm$ SEM, 5-9 animals. Comparisons: A,B: Two Way repeated measures ANOVA with Bonferroni *post hoc* comparison: *: $p < 0.05$ vs baseline; C,D: Bars under each AUC graph link treatments that do not differ from each other (Bonferroni multiple comparisons, $p < 0.05$)

Fig 4.

TOP: Effect on escape latency of IT neurosteroid antagonist, (17PA) on escape latency in the ipsilateral-carrageenan injected-(A) and contralateral (B) hindpaw when given in the presence of alphaxalone (60µg/10µL). IT 17PA (30 or 60µg/10µL) was given 15 min (arrow 1) prior to IT alphaxalone or vehicle (arrow 2). Carrageenan was injected into ipsilateral hind paw at t=0 (arrow 3). BOTTOM: Hyperalgesic index for ipsilateral (C) and contralateral (D) hindpaw shows that IT 17PA, 60, but not 30 µg/10µL resulted in reversal of the antinociceptive effects of IT alphaxalone. Each plot presents means±SEM, 5-9 animals. Comparisons: A,B: Two Way repeated measures ANOVA with Bonferroni *post hoc* comparison: *: p<0.05 vs baseline; C,D: Bars under each AUC graph link treatments that do not differ from each other (Bonferroni multiple comparisons, p<0.05)