Published in final edited form as:

Neurobiol Aging. 2013 November ; 34(11): 2525–2537. doi:10.1016/j.neurobiolaging.2013.04.021.

Amyloid β precursor protein as a molecular target for amyloid β -induced neuronal degeneration in Alzheimer's disease

Elena Anahi Bignante^a, Florencia Heredia^a, Gerardo Morfini^b, and Alfredo Lorenzo^{a,c,*}

^aLaboratory of Experimental Neuropathology, Instituto de Investigación Médica “Mercedes y Martín Ferreyra,” INIMEC-CONICET, Universidad Nacional de Córdoba, 5016 Córdoba, Argentina

^bDepartment of Anatomy and Cell Biology, University of Illinois at Chicago, Chicago, IL, USA

^cDepartamento de Farmacología, Facultad de Ciencias Químicas, Universidad Nacional de Córdoba, Haya de la Torre y Medina Allende, Ciudad Universitaria, 5000 Córdoba, Argentina

Abstract

A role of amyloid β (A β) peptide aggregation and deposition in Alzheimer's disease (AD) pathogenesis is widely accepted. Significantly, abnormalities induced by aggregated A β have been linked to synaptic and neuritic degeneration, consistent with the “dying-back” pattern of degeneration that characterizes neurons affected in AD. However, molecular mechanisms underlying the toxic effect of aggregated A β remain elusive. In the last 2 decades, a variety of aggregated A β species have been identified and their toxic properties demonstrated in diverse experimental systems. Concurrently, specific A β assemblies have been shown to interact and misregulate a growing number of molecular effectors with diverse physiological functions. Such pleiotropic effects of aggregated A β posit a mayor challenge for the identification of the most cardinal A β effectors relevant to AD pathology. In this review, we discuss recent experimental evidence implicating amyloid β precursor protein (APP) as a molecular target for toxic A β assemblies. Based on a significant body of pathologic observations and experimental evidence, we propose a novel pathologic feed-forward mechanism linking A β aggregation to abnormalities in APP processing and function, which in turn would trigger the progressive loss of neuronal connectivity observed early in AD.

Keywords

APP; Alzheimer's disease; A β ; Neuritic dystrophy; Synaptic degeneration; Dying-back degeneration

1. Introduction

Alzheimer's disease (AD) is a highly prevalent cause of dementia in elderly people. It is estimated that >30 million people worldwide suffer this disease, and epidemiologic studies predict that this number will exponentially grow in the next decades because of the sustained aging of human population. AD is characterized by progressive mental deterioration

© 2013 Elsevier Inc. All rights reserved.

*Corresponding author at: Instituto de Investigación Médica “Mercedes y Martín Ferreyra,” Casilla de Correo 389, 5000 Córdoba, Argentina. Tel.: +54 351 468 1466; fax: +54 351 469 5163. alorenzo@immf.uncor.edu (A. Lorenzo).

Disclosure statementThe authors of this work disclose that there are no conflicts of interest of any type that could inappropriately influence the work.

associated with degeneration and loss of neurons located in brain regions relevant to superior cognitive functions. The confirmatory AD diagnosis is based on 2 major histopathologic hallmarks: senile plaques, which are extracellular deposits of amyloid β ($A\beta$) peptide, and neurofibrillary tangles, which are somatic inclusions of the microtubule-associated protein tau. Early in the course of disease, AD is characterized by loss of synaptic function and neuritic dystrophy, characteristic features of neurons undergoing a “dying-back” pattern of degeneration (Kanaan et al., 2012). Increasing pathologic evidence suggests that the cognitive decline observed in the early stages of AD results from deficits in neuronal connectivity, whereas neuronal loss becomes more relevant in the later stages of the disease (Kanaan et al., 2012).

2. $A\beta$ and neurodegeneration in AD

The most accepted mechanistic hypothesis for AD posits that abnormal aggregation and accumulation of $A\beta$ in the brain trigger a cascade of pathologic events leading to progressive neuronal dysfunction (Hardy, 2009; Hardy and Selkoe, 2002; Huang and Mucke, 2012; Karran et al., 2011). Although the physiological role of monomeric, soluble $A\beta$ remains a matter of debate, several lines of evidence indicate that this form of $A\beta$ modulates synaptic function (Abramov et al., 2009; Hsieh et al., 2006; Puzzo et al., 2008; Russell et al., 2012). Therefore, an enormous effort has been dedicated to elucidate mechanisms underlying $A\beta$ aggregation and neurotoxicity. Consistent with the natural propensity of $A\beta$ to self-aggregate, diverse $A\beta$ assemblies displaying various toxic properties have been generated from synthetic peptides (Lambert et al., 1998; Lorenzo and Yankner, 1994; Pike et al., 1991) and/or isolated from human AD brain (Walsh et al., 2002). $A\beta$ assemblies identified to date include insoluble fibrils found in neuritic plaques, soluble protofibrils, oligomers, and even trimers and dimers, all of which may exist in a dynamic equilibrium in AD brains (Roychaudhuri et al., 2009).

The mechanisms by which $A\beta$ assemblies cause neuronal toxicity remain elusive. However, the identification of several molecular mediators of $A\beta$ toxicity (Table 1) suggests that $A\beta$ elicits a unique mechanism of toxicity, where the property of specific assemblies to recruit multiple molecular effectors would interfere with the normal functionality of various metabolic pathways and intracellular signaling cascades. Accordingly, diverse cellular and systemic functions are affected by $A\beta$ aggregates including ionic plasmalemma conductance, Ca^{2+} homeostasis, mitochondrial function, energy metabolism, cytoskeletal organization, intracellular transport, kinase activation, synaptic transmission, vascular permeability, and inflammatory processes, all of which may ultimately contribute to neuronal degeneration in AD (reviewed by Chen and Yan, 2010; Götz et al., 2011; Mattson, 2004; Reddy and Beal, 2008; Yankner and Lu, 2009). Results from quantitative proteomic experiments using artificial exogenous proteins with intrinsic propensity to form amyloid-like aggregates support this notion. Specifically, it was found that the toxic effects of these artificial proteins correlated with abnormal protein interactions in their aggregated form (Olzscha et al., 2011). Aberrant interactions with these proteins would then trigger alterations in multiple cellular functions (Olzscha et al., 2011). Such pleiotropic mechanism of toxicity poses a challenge for the design of therapeutic interventions for AD and calls for the identification of cardinal $A\beta$ effectors relevant to AD pathogenesis. In this review, we focus on amyloid precursor protein (APP), a protein unequivocally linked to AD pathogenesis. APP is the unique source of $A\beta$, and genetic alterations in *App*, including gene duplication or missense mutations, lead to early-onset familial forms of AD (FAD) (see Alzheimer's Disease and Frontotemporal Dementia Mutation Database: <http://www.molgen.vib-ua.be/ADMutations>). Because pathogenic *App* mutations cluster around or within the $A\beta$ sequence and typically predispose to $A\beta$ aggregation and/or deposition (Tanzi, 2012), it is generally accepted that the pathogenic role of APP in AD is mainly

restricted to be the source of A β . However, recent data from various independent groups suggest that A β aggregates interact with APP, triggering abnormalities in its function and processing that would promote neuronal dysfunction. These observations represent a major focus of this review. First, we briefly summarize various physiological functions proposed for APP and the protein fragments derived from its metabolic processing. Then, we discuss various published reports providing evidence that APP mediates the toxic properties of A β assemblies. Finally, we propose a mechanistic model where abnormal APP/A β interactions play an important role in the development of early and critical pathogenic events of AD, including synaptic dysfunction and neuritic degeneration.

3. APP: protein structure and metabolic processing

APP was discovered in 1987 as the only source of A β (Kang et al., 1987), and the gene encoding this protein (*App*) was mapped to chromosome 21 (Goldgaber et al., 1987; Tanzi et al., 1987). The *App* gene encodes a transmembrane type I protein belonging to a family of conserved proteins that in mammals also include APLP1 and APLP2 (Wasco et al., 1993). Whereas all 3 APP family members are predominantly expressed in the brain, APP and APLP2 are also ubiquitously expressed in other tissues. APP family members display a high degree of homology in the extracellular E1 and E2 domains and in the intracellular C-terminal portion (Fig. 1). Reduced homology exists in the transmembrane and juxtamembrane domains where only APP, but not APLP1 or APLP2, contains the sequence encoding the A β peptide. This observation, together with the discovery that >20 mutations in *App* gene are linked to early-onset FAD (see Alzheimer's Disease and Fronto-temporal Dementia Mutation Database: <http://www.molgen.vibua.be/ADMutations>), motivated interest of the AD scientific community for understanding the basic biology of APP and its relation to AD pathogenesis (Tanzi, 2012).

Several APP isoforms resulting from alternative splicing have been described, the most abundant being a 695-amino acid-long isoform mainly found in neurons (Mattson, 1997). Also, 770- and 751-amino acid-long isoforms containing a Kunitz protease inhibitor domain are expressed in non-neuronal cells, including glia and endothelial cells (Mattson, 1997). However, the functional relevance of cell type-specific expression of APP splicing variants remains poorly understood.

APP is subject to a highly dynamic distribution and intracellular trafficking within cells (Allinquant et al., 1994; Simons et al., 1995; Yamazaki et al., 1995; reviewed by Brunholz et al., 2011). Initially, immature APP traffics through the endoplasmic reticulum and the Golgi apparatus where it undergoes a series of post-translational modifications, including *N*- and *O*-glycosylation and phosphorylation. The modified, mature APP protein is then transported in secretory vesicles to the plasma membrane via axonal transport mechanisms involving conventional kinesin, a major microtubule-dependent molecular motor (Brunholz et al., 2011; Szodorai et al., 2009). After insertion in the plasma membrane, APP can be reinternalized to endosomes from where it can be either recycled back to the cell surface, targeted to lysosomes for degradation, or redistributed to distant cellular compartments through transcytosis (Back et al., 2007; Golde et al., 1992; Haass et al., 1992; Tienari et al., 1996; reviewed by Haass et al., 2012). In addition, APP and its proteolytic fragments (including A β) have been observed in mitochondria (Devi et al., 2006; Lustbader et al., 2004; Manczak et al., 2006; Yamaguchi et al., 1992). In neurons, most APP is initially targeted from the trans-Golgi network to the axonal membrane and then to endosomes, from where it can also be redirected to the dendritic compartment via transcytosis (Tienari et al., 1996; Yamazaki et al., 1995). Most APP is found within intracellular compartments, and only 10%–20% is present in the plasma membrane (Thinakaran and Koo, 2008). During development, APP is enriched at the growth cones of developing neurites. In more mature

neurons, APP localizes to focal adhesion sites and within pre- and postsynaptic structures of the central and peripheral nervous tissue, suggesting a functional role in neuritic growth and synaptic plasticity (Ashley et al., 2005; Sabo et al., 2003; Yamazaki et al., 1997).

The dynamic distribution of APP at different subcellular compartments is accompanied by a complex metabolic processing that includes 2 basic metabolic routes: one precluding and other promoting the generation of A β peptides. Accordingly, these routes are designated as the nonamyloidogenic and amyloidogenic pathways, respectively (Fig. 2). A detailed analysis of APP processing is beyond the scope of this review but has been extensively revised elsewhere (Brunholz et al., 2011; Haass et al., 2012). Within the context of this review, suffices to say that in the nonamyloidogenic pathway, APP is first cleaved by α -secretase within the A β sequence to generate 2 proteolytic fragments: sAPP α (a secreted protein that encompasses most of the APP ectodomain) and C83 (a protein stub that remains tethered to the plasma membrane for further proteolytic processing). In the amyloidogenic pathway, APP is cleaved by β -secretase at the N-terminus of the A β sequence, thus generating a secreted protein (sAPP β) and a membrane-associated fragment known as C99 comprising the entire A β sequence. Both C99 and C83 stubs can be cleaved by γ -secretase within the transmembrane domain, resulting in the release of A β or p3 peptides, respectively, and the intracytoplasmic domain of APP (AICD). Because cleavage by γ -secretase is heterogeneous, A β peptides from various lengths can be generated, with A β 1–40 being the most abundant and soluble and the A β 1–42 the most prone to aggregation (Jarret et al., 1993; McGowan et al., 2006).

It is noteworthy that APP represents only 1 of several secretase substrates. Indeed, several cell surface receptors and proteins including Notch 1, ERBB4, Robo, N-cadherins, neuregulins, P75NTR, LRP1, APLP1, and APLP2, among others, are also cleaved by secretases. Secretase-mediated cleavage of these proteins results in the production of extracellular and intracellular fragments with unique physiological functions (De Strooper et al., 1999; Ni et al., 2001; Reiss et al., 2005; Scheinfeld et al., 2002; reviewed by Bai and Pfaff, 2011). The similarity of APP structure and processing with many of these receptors suggests that APP might also act as a cell surface receptor with specialized functions; however, natural physiological ligands for APP remain elusive.

4. Physiological functions of APP

Various cellular functions have been proposed for APP, but the actual physiological role(s) of this protein remains a matter of debate. *App*-knockout mice are viable and fertile, indicating that loss of APP is not essential for embryonic development and/or the control of vital functions (Li et al., 1996; Zheng et al., 1995). Nevertheless, APP^{-/-} mice display some discrete phenotypic abnormalities including reduced body weight, decreased locomotor activity and forelimb grip strength, defective long-term potentiation, and impairments in learning and memory (Ring et al., 2007; Zheng et al., 1995). Increased reactive gliosis was also reported in these animals, but no obvious evidence of neuronal degeneration was observed (Zheng et al., 1995). Reports of early lethality in APP^{-/-}/APLP1^{-/-}/APLP2^{-/-} triple knockout, APP^{-/-}/APLP2^{-/-}, or APLP1^{-/-}/APLP2^{-/-} double knockout mice raised the possibility that compensatory mechanisms may mask the functional relevance of APP in APP^{-/-} mice (Heber et al., 2000). Surprisingly, APP^{-/-}/APLP1^{-/-} animals are viable, suggesting that APLP2 plays some particular function that cannot be compensated by other APP family members (Heber et al., 2000). Additionally, no apparent abnormalities were reported in APLP2^{-/-} mice (von Koch et al., 1997), further suggesting a functional redundancy among APP family members (for a comprehensive review, see Müller and Zheng, 2012). These in vivo observations strongly suggest that AD-like pathology does not

result from loss of APP function alone. A brief description of experimental data suggesting discrete functional roles for APP and APP metabolites is provided subsequently.

5. Roles of APP on cell adhesion, neuritogenesis, and synaptic plasticity

APP family members have been proposed to play a role in cell-to-cell and cell-substratum interactions. For example, membrane-associated forms of APP, APLP1, and APLP2 promote cell-cell adhesion through homo- and heterodimerization (Baumkötter et al., 2012; Soba et al., 2005). Also, Förster resonance energy transfer experiments using quantum dots also revealed a ligand/receptor interaction between the proteolytic APP fragment sAPP and holo-APP in the plasma membrane (Gralle et al., 2009), suggesting a link between the functional activities of sAPP and holo-APP. In addition, holo-APP can form antiparallel dimers through interactions involving the E2 domain (Wang and Ha, 2004), and both the E1 domain and the GAIIG motif within the transmembrane region also play a role in APP dimerization (Kaden et al., 2008; Munter et al., 2007). Moreover, binding of heparin to E1–E2 domains also promotes APP dimerization (Dahms et al., 2010; Gralle et al., 2006). Providing a physiological consequence for these interactions, APP dimerization was shown to affect its processing by secretases (Kaden et al., 2008; Lefort et al., 2012; Munter et al., 2007; Scheuermann et al., 2001).

Because the extracellular portion of APP contains various structural domains, it is not surprising that several putative physiological ligands have been identified for APP. Interestingly, many of these ligands are extracellular matrix components and cell adhesion–related proteins including heparan sulfate proteoglycans, laminin, collagen type I, F-spondin, Nogo-66 receptor, reelin, netrin, and integrin $\alpha\beta 1$ (Behr et al., 1996; Clarris et al., 1997; Ho and Südhof, 2004; Hoe et al., 2009; Kibbey et al., 1993; Park et al., 2006). Importantly, binding of APP to each of these molecules differentially affects its cellular distribution and/or metabolic processing by secretases, indicating that these interactions are functionally relevant. For example, binding of F-spondin to the central E2 domain of APP prevents its cleavage by secretases (Ho and Südhof, 2004). Also, reelin binding to the E1 domain enhances the nonamyloidogenic processing of APP (Hoe et al., 2006, 2009). Finally, netrin reportedly interacts with APP within the A β domain, and this interaction results in reduced A β generation (Loureço et al., 2009).

The adhesive properties of APP have also been linked to a tetrapeptide sequence RHDS within the N-terminus of the A β peptide, which promotes cell adhesion in an integrin-like manner (Ghisso et al., 1992). In fact, APP colocalizes and interacts with integrin $\alpha\beta 1$ at focal adhesion sites and within synapses. Additionally, reelin binds to the E1 extracellular domain of APP, increasing the levels of holo-APP at the cell surface, the secretion of sAPP, and the elongation of neurites by a mechanism dependent on $\alpha\beta 1$ integrin (Hoe et al., 2009). Results from experiments using APP messenger RNA (mRNA) antisense and APP^{−/−} mice also support a role of cellular APP in the control of neuritic outgrowth and further demonstrate that the growth-promoting effect of sAPP requires both membrane-associated holo-APP and integrins (Allinquant et al., 1995; Perez et al., 1997). Additionally, in utero electroporation experiments revealed that APP is required for proper migration of neuronal precursors to the cortical plate, a process dependent on cell-cell and cell-substratum adhesions (Young-Pearse et al., 2007). Significantly, these migration defects were fully rescued by human APP, APLP1, or APLP2, suggesting functional compensation among APP family members (Young-Pearse et al., 2008).

The role of APP and its proteolytic fragments on the regulation of synaptic plasticity has been extensively documented. For example, initial reports found increased number of synapses in cortical areas of transgenic mice overexpressing wild-type human APP (Mucke

et al., 1994). More recently, it was shown that APP regulates the presynaptic expression and activity of the high-affinity choline transporter at neuromuscular junctions (Wang et al., 2007). Using APLP2^{-/-} mice that also allow the conditional depletion of APP in either motoneurons or muscle, it was determined that the proper formation and function of neuromuscular synapses require APP expression in both pre- and postsynaptic membranes (Wang et al., 2009). Moreover, expression of APP in HEK cells cocultured with primary neurons induced hemisynapse formation (Wang et al., 2009). Based on these observations, APP was proposed to represent a novel class of synaptic adhesion molecule with biochemical properties similar to those reported for neurexins/neuroligins, SynCAMs, and leucine-rich repeat transmembrane neuronal proteins (Baumkötter et al., 2012; Müller and Zheng, 2012; Wang et al., 2009). Collectively, these observations mentioned previously suggest that the neuritogenic and synaptogenic activities of APP involve the modulation of cell-cell and cell-substratum interactions and also the homo- and heterodimerization properties of this molecule. By modulating APP dimerization and processing by secretases, diverse extracellular components and membrane proteins would further regulate APP function.

6. Role of APP on trophic support

Initial experiments in cultured fibroblasts showed that sAPP modulates cell division through the RERMS sequence located in the middle portion of the APP ectodomain (Ninomiya et al., 1993; Saitoh et al., 1989). Additionally, trophic effects of sAPP were widely documented in neuronal cells. For example, application of sAPP to cultured primary neurons promoted long-term survival, enhanced neurite outgrowth, increased synaptic formation, and protected cultured neurons against various toxic stimuli (Mattson, 1997). Extending these observations, intracranial infusion of sAPP-specific antibodies impaired memory, whereas application of sAPP has a memory-enhancing effect (Doyle et al., 1990; Meziane et al., 1998). Together, these findings indicated that sAPP modulates cellular activities involved in cognitive processes, including the functionality and maintenance of synapses and neurites. Interestingly, sAPP binds to and regulates holo-APP dimerization at the plasma membrane, an event required for the neuroprotective effect of sAPP against starvation-induced cell death (Gralle et al., 2009). Whereas sAPP appears to have a beneficial effect to neurons, in vitro over-expression of C99 (a membrane-tethered APP fragment generated after β -secretase cleavage) promotes neuronal degeneration (Yankner et al., 1989; Yoshikawa et al., 1992). The contrasting effect of these APP fragments suggests that neuronal cells depend on an appropriate balance of membrane-tethered APP and sAPP.

It is noteworthy that cultured human Down syndrome (DS) neurons display poor survival rates and defective APP secretion, despite having increased APP expression levels because of chromosome 21 trisomy (Busciglio and Yankner, 1995). The reduced in vitro survival of DS neurons likely results from pathogenic effects originated from multigenic imbalance. Nevertheless, APP appears to play a cardinal role because addition of exogenous sAPP at physiological levels restores long-term viability of cultured DS neurons (Busciglio and Yankner, 1995; Busciglio et al., 2002). Because reduced levels of sAPP have been observed in both DS and AD (Busciglio et al., 2002; Lannfelt et al., 1995; Van Nostrand et al., 1992), it is reasonable to speculate that alterations in APP metabolism and/or function play a role in AD pathogenesis beyond its participation as a source for A β . Abnormal App dosage also appears to play a significant role in neuronal dysfunction in TS65Dn mice, an animal DS model bearing 3 copies of the *App* gene. For example, a reduction in the number of cholinergic neurons was observed in the entorhinal cortex of TS65Dn mice, but removal of the extra *App* copy was sufficient to prevent this defect (Salehi et al., 2006). Also, altered intracellular Ca²⁺ homeostasis and defective cholinergic function were both described in neuronal cell lines derived from Ts16 mice, but normalizing APP expression through APP-

specific mRNA antisense abrogated these deficits (Opazo et al., 2006; Rojas et al., 2008). It remains unclear how alterations in APP levels relate to the AD-like pathology seen in individuals with genetic overexpression of APP, such as DS or FAD because of *App* gene locus duplication (Rovelet-Lecrux et al., 2006) or mutations in the *App* gene-promoter region (Theuns et al., 2006). However, the collective evidence suggests that a delicate balance between APP expression and secretion modulates important neuronal activities including trophic support, synaptic and neuritic plasticity, and survival.

7. Regulation of intracellular signaling pathways by APP

The highest degree of homology among APP family members is found within their intracellular domain, suggesting that this region plays a critical functional role. Indeed, the cytoplasmic domain of all APP family members contain a GYENPTY motif, which mediates their interaction with a number of adaptor proteins including Fe65, Mint/X11-family proteins, Dab1, c-Jun N-terminal kinase, Shc, and Grd2, among others (Matsuda et al., 2001; McLoughlin and Miller, 2008; Russo et al., 2005; Saito et al., 2011). Moreover, the interaction of APP with these proteins modulates the generation of the AICD fragment by secretases. The significance of this modulation relies on reports that AICD, Fe65, and Tip60 can form a tripartite transcriptional complex that controls the expression of several genes, including *KAI1*, *neprilysin*, *LRP1*, and *EGF* receptor (Baek et al., 2002; Cao and Südhof, 2004; Liu et al., 2007; Pardossi-Piquard et al., 2005; Zhang et al., 2007). The similarity of APP and Notch 1 processing by secretases and the regulation of gene transcription by proteolytic fragments derived from these proteins appear consistent with a role of APP as a cell surface receptor.

It is generally assumed that receptors coupled to heterotrimeric G proteins, including Go and Gs, must have a 7 transmembrane domain structure. However, increasing evidence indicates that single transmembrane domain proteins including epidermal growth factor receptor, insulin receptor, insulin-like growth factor receptor, fibroblast growth factor receptor, and natriuretic peptide receptor C-type can also interact and signal through G proteins (Patel, 2004). Significantly, a consensus sequence for Go protein binding was identified in the intracellular juxtamembrane domain of APP (Nishimoto et al., 1993). In vitro experiments confirmed this interaction, further revealing histidines at positions 657e658 (APP695 numbering) as critical residues for this interaction (Brouillet et al., 1999; Nishimoto et al., 1993; Okamoto et al., 1995). More recently, it was also observed that a RHLSK motif within the intracellular domain of APP mediates an interaction with Gs (Deyts et al., 2012). Extending these observations, it was observed that APPL, the *Drosophila* homolog of the mammalian APP, colocalizes with Go in migrating neurons (Swanson et al., 2005) and regulates synaptic structure and number in a manner that requires both the putative Go protein binding site and the GYENPTY motif (Torroja et al., 1999). Furthermore, APP was shown to promote axonal and dendritic outgrowth in mammalian neurons through a mechanism involving Gs-dependent activation (Deyts et al., 2012). Therefore, through interactions between its intracellular domain and various binding partners, including G proteins, membrane-associated APP modulates the activation of various intracellular signaling cascades.

8. Role of APP on AD pathogenesis

The involvement of APP in AD pathology is indisputable because A β is only generated from APP metabolic processing, and genetic alterations in *App* suffice to cause early-onset FAD. In addition, a growing body of evidence indicates that APP plays a role in neuronal degeneration in both genetic and sporadic AD. Specifically, APP appears to act as a molecular target for pathologic Ab assemblies. Such interaction might promote neuronal

degeneration through various mechanisms involving both loss and gain of APP function. Experimental evidence supporting this possibility is provided subsequently.

9. APP as a molecular target for toxic, AD-related A β assemblies

A direct interaction between A β fibrils and the ectodomain of APP has been demonstrated using a variety of experimental conditions (Lorenzo et al., 2000; Melchor et al., 2000; Wagner et al., 2000). In contrast with fibrillar Ab, monomeric A β peptide does not bind APP (Lorenzo et al., 2000; Melchor and Van Nostrand, 2000; Wagner et al., 2000), suggesting that conversion of monomeric Ab into toxic amyloid fibrils is required for binding to APP. The potential pathologic relevance of these observations is suggested by reports showing accumulation of APP at the proximity of fibrillar neuritic Ab plaques in AD brains and in con-gophilic blood vessels of brains affected by hereditary cerebral hemorrhage with amyloidosis Dutch type, an AD-related pathology (Rijal Upadhyaya et al., 2012; Rozemuller et al., 1993). However, experimental work is required to evaluate whether an interaction with A β promotes the pathologic accumulation of APP in degenerating neurons in vivo.

Two sequences have been identified on APP that mediate its binding to A β fibrils. One is located within the N-terminus of APP and encompasses amino acids 18e119, with His110, Val112, and Ile113 being critical for this binding (Van Nostrand et al., 2002). This 18e119 sequence appears to be relevant for binding of A β fibrils to sAPP, rather than holo-APP (Kedikian et al., 2010; White et al., 2003). Consistent with this possibility, it was found that sAPP is sequestered by Ab fibrils from the media of cultured cells (Heredia et al., 2004), an event that might interfere with the normal trophic functions of sAPP and/or its modulatory role in holo-APP dimerization. A second sequence was mapped to the extracellular juxtamembrane A β domain (Shaked et al., 2006; Sola Vigo et al., 2009), which mainly mediates binding of membrane-anchored holo-APP to A β fibrils (Kedikian et al., 2010). Interestingly, no sequence homology exists between the 2 A β -binding sequences of APP mentioned previously. Interestingly, synthetic peptides corresponding to these sequences aggregate and form fibrils in vitro (Hilbich et al., 1993; Rochet and Lansbury, 2000), suggesting that the binding of APP to A β fibrils is conformation dependent and not sequence dependent. Further supporting this notion, it was found that both holo-APP and sAPP interact with fibrils formed by amyloidogenic peptides other than A β , including amylin and prion peptides (Lorenzo et al., 2000; White et al., 2003). Therefore, it is likely that toxic A β assemblies other than fibrils could also bind APP including dimers, oligomers, or protofibrils. Additional experiments are required to elucidate the structural mechanisms underlying the interaction of APP with toxic A β assemblies.

10. A pathogenic feedback loop triggered by APP-A β interactions

The first evidence suggesting a pathogenic loop between aggregated A β and APP came from the observation that treatment of cultured human leptomeningeal smooth muscle cells with synthetic A β fibrils promotes cellular APP accumulation and enhanced A β secretion (Davis-Salinas et al., 1995). Additionally, it was also found that exogenously added fibrillar A β 42 promotes APP accumulation and the formation of insoluble newly synthesized Ab42 in HEK cells (Yang et al., 1999). More recently, it was shown that A β oligomers enhance the amyloidogenic processing of APP in primary neurons (Marsden et al., 2011). Interestingly, it was recently shown that cross-linking of APP with anti-APP antibody 22C11 (which recognizes an epitope at the N-terminus of APP) is sufficient to raise the levels of A β in viable neurons with a concomitant increase in the levels of the b-secretase BACE1 (Lefort et al., 2012). The mechanism appears to involve a sorting defect that stems from the caspase-3-mediated inactivation of a key sorting adaptor protein, namely GGA3, which

prevents the lysosomal degradation of BACE1 (Lefort et al., 2012). Likely, the aforementioned mechanism is triggered by A β fibrils that also promote APP multimerization (Heredia et al., 2004; Lu et al., 2003; Melchor and Van Nostrand, 2000) and raise BACE1 levels (Zhao et al., 2007). Collectively, these observations suggest a pathologic, APP-dependent feed-forward process for A β deposition (Fig. 3). Considering that A β is mainly produced at synapses (Brody et al., 2008; Cirrito et al., 2005; Kamenetz et al., 2003; Lazarov et al., 2002) and its accumulation first takes place at this compartment (Capetillo-Zarate et al., 2011), the potential of APP to bind A β assemblies and to undergo transcytosis might help to explain the transsynaptic spreading of A β deposition in transgenic mice and in AD (Harris et al., 2010a; Lazarov et al., 2002; Sheng et al., 2002; Thal et al., 2002). An understanding of molecular mechanisms underlying the feed-forward process between aggregated A β and APP will prove essential for the development of rational therapeutic interventions aimed to halt the spreading of A β pathology in AD.

11. Pathogenic pathways triggered by Ab/APP interactions

The interaction between A β fibrils and APP might have patho-logic relevance for neuronal degeneration. Fibrillar, but not monomeric A β , promotes accumulation and multimerization of holo-APP at the cell surface of diverse cell types including neurons, astrocytes, and human cerebrovascular smooth muscle cells (Lu et al., 2003; Melchor et al., 2000; Tsuruma et al., 2012; White et al., 2003). Interestingly, toxic A β -induced APP multimerization has been linked to pathogenic events consistent with the early loss of neuronal connectivity characteristic of AD. For example, in cultured hippocampal neurons, A β fibrils induced holo-APP accumulation at aberrant focal adhesion sites of dystrophic neurites (Grace and Busciglio, 2003; Grace et al., 2002; Heredia et al., 2004; Pike et al., 1992). This observation was extended to human AD brains, and brains of transgenic AD mouse models, with APP reportedly accumulating in dystrophic neurites close to fibrillar A β plaques (Borchelt et al., 1997; Cras et al., 1991; Ohgami et al., 1992; Phinney et al., 1999). Cultured neurons from APP-knockout mice were partially resistant to A β -induced toxicity (Lorenzo et al., 2000). Also, APP overexpression enhanced A β toxicity in cultured hippocampal neurons (Shaked et al., 2006; Sola Vigo et al., 2009), and deletion of the A β -binding sequences on APP abrogated both A β -induced APP multimerization and toxicity (Kedikian et al., 2010; Shaked et al., 2006; Sola Vigo et al., 2009). These observations strongly suggest that A β -induced APP multimerization promotes neuronal degeneration, consistent with the dying-back pattern of neuronal degeneration observed early in the course of AD (Kanaan et al., 2012).

Activation of caspases and neuronal apoptosis has both been proposed to play a role in AD pathology (reviewed by Hyman and Yuan, 2012). Supporting an involvement of aggregated A β deposition in this process, it was found that A β promotes caspase activation and apoptotic death of cultured neurons in vitro (Harada and Sugimoto, 1999; Ivins et al., 1999; Loo et al., 1993; Nakagawa et al., 2000; Troy et al., 2000). A β also promotes caspase-dependent cleavage of APP, and event resulting in the release of C31, an intracellular APP fragment with cytotoxic properties (Galvan et al., 2002; Lu et al., 2000). The toxic mechanism of C31 in cultured cells was initially linked to alterations in the transcriptional activity of AICD (Kinoshita et al., 2002). More recently, it was observed that C31 toxicity requires holo-APP expression and depends on the ability of the peptide to promote holo-APP multimerization by a yet undefined mechanism (Park et al., 2009). The relevance of caspase-mediated APP cleavage in vivo has been addressed using genetically modified mice that express a mutant version of APP (D664A) that precludes caspase-dependent generation of C31 (Galvan et al., 2006). Significantly, neither A β pathology nor accumulation of A β plaques was affected in transgenic mice expressing D664A APP along with the Swedish and Indiana FAD-APP mutations. However, several pathologic events found in the Swedish and Indiana FAD-APP double-mutant mice were prevented in these triple-mutant mice,

including astrogliosis, synapse loss, defective long-term potentiation, and cognitive deficits (Galvan et al., 2006, 2008; Saganich et al., 2006; Zhang et al., 2009). These observations suggest that pathogenic changes triggered by A β aggregation and/or accumulation require caspase-dependent generation of an APP fragment comprising the intracellular cytoplasmic domain. However, a recent work challenged these observations (Harris et al., 2010b), and thus, the toxic effect of C31 in vivo remains on debate (Bredesen et al., 2010).

Several observations suggest that APP modulates G protein-dependent intracellular signaling. Overexpression of several FAD-APP mutations (i.e., Val642) promoted constitutively activation of heterotrimeric G proteins (Okamoto et al., 1996), ultimately inducing apoptotic cell death in neuroblastoma cell lines (Giambarella et al., 1997; Yamatsuji et al., 1996a, 1996b). Experiments in cultured hippocampal neurons extended these results, showing that binding of A β fibrils to holo-APP promotes APP multimerization and trigger toxicity through a mechanism involving G activation (Heredia et al., 2004; Sola Vigo et al., 2009). Together, these evidences suggest that APP-mediated alterations in G signaling contribute to the toxic effects elicited by A β assemblies. Also, it was found that molecules that bind APP and promote its multimerization such as transforming growth factor β 2 and APP-specific antibodies (i.e., 22C11) induce toxicity by a G-dependent mechanism (Hashimoto et al., 2003, 2005; Okamoto et al., 1995; Rohn et al., 2000; Sudo et al., 2000, 2001). Additional evidence suggests that multimerization of holo-APP at the plasma membrane induces additional alterations in intracellular signaling. For example, overexpression of FAD-linked APP in primary neurons promoted apoptosis by a mechanism involving activation of G, c-Jun N-terminal kinase (Niikura et al., 2004), and p21-activated kinase (PAK) (McPhie et al., 2003). Interestingly, in both AD brains and in FAD-APP transgenic mice, affected neurons display increased accumulation of active PAK in the surroundings of A β plaques (Ma et al., 2008; Zhao et al., 2006).

12. Concluding remarks and future directions

A β deposition is an early event in AD that precedes neuronal degeneration and cognitive decline by several years or even decades (Bateman et al., 2012; Ingelsson et al., 2004; Thal et al., 2002). Significantly, studies from various independent groups established APP as a molecular target for pathologic A β assemblies and a critical mediator of A β -induced toxicity. Based on the pathologic and experimental evidence discussed previously, we propose a model of AD pathogenesis where aggregated A β -induced multimerization of APP at the cell surface represents a primary pathogenic event that triggers dying-back degeneration of neurons (Fig. 3). In this model, the normal physiological role of APP as a modulator of cell-cell and cell-substratum adhesions in neurites and synapses would be compromised by its interaction with toxic A β aggregates. By inducing aberrant APP multimerization, A β aggregates would trigger aberrant intracellular signaling cascades promoting synaptic and neuritic abnormalities (Grace and Busciglio, 2003; Heredia et al., 2004; Tsai et al., 2004), cytoskeletal alterations (Busciglio et al., 1995; Heredia et al., 2006), and axonal transport defects (Brunholz et al., 2011; Pigino et al., 2009). Additionally, binding of aggregated A β to APP would also promote increased metabolic processing of APP through the amyloidogenic pathway, further contributing to A β deposition, neuritic degeneration, and synapse loss in AD.

An understanding of the mechanisms underlying APP interactions with specific A β assemblies might lead to the development of novel therapeutic strategies for AD. Such strategies would aim to prevent the selective APP interactions with toxic A β assemblies. Several important questions await experimental testing. Among these, it remains unclear whether all toxic A β assemblies interact with APP or induce APP multimerization. Structural studies will also be required to determine whether a conformational change in the

ectodomain of APP involves binding to aggregated A β . This knowledge would be potentially helpful for rational therapeutic drug design. Equally important will be to address how toxic A β assemblies enhance the amyloidogenic processing of APP. Finally, mechanisms linking Ab-induced APP multimerization to neuritic dystrophy, synaptic loss, and tau pathology remain to be defined (Brunholz et al., 2011). Addressing these questions might shed light into the normal physiological function of APP and its potential as a therapeutic target for AD.

Acknowledgments

This work was supported by grants from CONICET, SECyT-UNC, and ANPCyT (PICT 2010-1895) to AL and National Institutes of Health NS066942A, ALS/CVS Therapy Alliance, and Brain Research Foundation grants to GM. AL is career members of CONICET.

References

- Abramov E, Dolev I, Fogel H, Ciccotosto GD, Ruff E, Slutsky I. Amyloid-beta as a positive endogenous regulator of release probability at hippocampal synapses. *Nat Neurosci.* 2009; 12:1567–1576. [PubMed: 19935655]
- Allinquant B, Hantraye P, Mailleux P, Moya K, Bouillot C, Prochiantz A. Downregulation of amyloid precursor protein inhibits neurite outgrowth in vitro. *J Cell Biol.* 1995; 128:919–927. [PubMed: 7876315]
- Allinquant B, Moya KL, Bouillot C, Prochiantz A. Amyloid precursor protein in cortical neurons: coexistence of two pools differentially distributed in axons and dendrites and association with cytoskeleton. *J Neurosci.* 1994; 14:6842–6854. [PubMed: 7965082]
- Allinson TM, Parkin ET, Turner AJ, Hooper NM. ADAMs family members as amyloid precursor protein alpha-secretases. *J Neurosci Res.* 2003; 74:342–352. [PubMed: 14598310]
- Alvarez A, Alarcón R, Opazo C, Campos EO, Muñoz FJ, Calderón FH, Dajas F, Gentry MK, Doctor BP, De Mello FG, Inestrosa NC. Stable complexes involving acetylcholinesterase and amyloid-beta peptide change the biochemical properties of the enzyme and increase the neurotoxicity of Alzheimer's fibrils. *J Neurosci.* 1998; 18:3213–3223. [PubMed: 9547230]
- Alvarez A, Opazo C, Alarcón R, Garrido J, Inestrosa NC. Acetylcholinesterase promotes the aggregation of amyloid-beta-peptide fragments by forming a complex with the growing fibrils. *J Mol Biol.* 1997; 272:348–361. [PubMed: 9325095]
- Ashley J, Packard M, Ataman B, Budnik V. Fasciclin II signals new synapse formation through amyloid precursor protein and the scaffolding protein dX11/Mint. *J Neurosci.* 2005; 25:5943–5955. [PubMed: 15976083]
- Back S, Haas P, Tschape JA, Gruebl T, Kirsch J, Muller U, Beyreuther K, Kins S. Beta-amyloid precursor protein can be transported independent of any sorting signal to the axonal and dendritic compartment. *J Neurosci Res.* 2007; 85:2580–2590. [PubMed: 17335075]
- Baek SH, Ohgi KA, Rose DW, Koo EH, Glass CK, Rosenfeld MG. Exchange of N-CoR corepressor and Tip60 coactivator complexes links gene expression by NF-kappaB and beta-amyloid precursor protein. *Cell.* 2002; 110:55–67. [PubMed: 12150997]
- Bai G, Pfaff SL. Protease regulation: the Yin and Yang of neural development and disease. *Neuron.* 2011; 72:9–21. [PubMed: 21982365]
- Bamberger ME, Harris ME, McDonald DR, Husemann J, Landreth GE. A cell surface receptor complex for fibrillar beta-amyloid mediates microglial activation. *J Neurosci.* 2003; 23:2665–2674. [PubMed: 12684452]
- Barry AE, Klyubin I, McDonald JM, Mably AJ, Farrell MA, Scott M, Walsh DM, Rowan MJ. Alzheimer's disease brain-derived amyloid- β -mediated inhibition of LTP in vivo is prevented by immunotargeting cellular prion protein. *J Neurosci.* 2011; 31:7259–7263. [PubMed: 21593310]
- Bate C, Williams A. Amyloid- β -induced synapse damage is mediated via cross-linkage of cellular prion proteins. *J Biol Chem.* 2011; 286:37955–37963. [PubMed: 21900234]

- Bateman RJ, Xiong C, Benzinger TL, Fagan AM, Goate A, Fox NC, Marcus DS, Cairns NJ, Xie X, Blazey TM, Holtzman DM, Santacruz A, Buckles V, Oliver A, Moulder K, Aisen PS, Ghetti B, Klunk WE, McDade E, Martins RN, Masters CL, Mayeux R, Ringman JM, Rossor MN, Schofield PR, Sperling RA, Salloway S, Morris JC. Dominantly Inherited Alzheimer Network. Clinical and biomarker changes in dominantly inherited Alzheimer's disease. *N Engl J Med*. 2012; 367:795–804. [PubMed: 22784036]
- Baumkötter F, Wagner K, Eggert S, Wild K, Kins S. Structural aspects and physiological consequences of APP/APLP trans-dimerization. *Exp Brain Res*. 2012; 217:389–395. [PubMed: 21952790]
- Behr D, Hesse L, Masters CL, Multhaup G. Regulation of amyloid protein precursor (APP) binding to collagen and mapping of the binding sites on APP and collagen type I. *J Biol Chem*. 1996; 271:1613–1620. [PubMed: 8576160]
- Borchelt DR, Ratovitski T, van Lare J, Lee MK, Gonzales V, Jenkins NA, Copeland NG, Price DL, Sisodia SS. Accelerated amyloid deposition in the brains of transgenic mice co-expressing mutant presenilin 1 and amyloid precursor proteins. *Neuron*. 1997; 19:939–945. [PubMed: 9354339]
- Bredesen DE, John V, Galvan V. Importance of the caspase cleavage site in amyloid- β protein precursor. *J Alzheimers Dis*. 2010; 22:57–63. [PubMed: 20847422]
- Brody DL, Magnoni S, Schwetye KE, Spinner ML, Esparza TJ, Stocchetti N, Zipfel GJ, Holtzman DM. Amyloid-beta dynamics correlate with neurological status in the injured human brain. *Science*. 2008; 321:1221–1224. [PubMed: 18755980]
- Brouillet E, Trembleau A, Galanaud D, Volovitch M, Bouillot C, Valenza C, Prochiantz A, Allinquant B. The amyloid precursor protein interacts with Go heterotrimeric protein within a cell compartment specialized in signal transduction. *J Neurosci*. 1999; 19:1717–1727. [PubMed: 10024358]
- Brunholz S, Sisodia S, Lorenzo A, Deyts C, Kins S, Morfini G. Axonal transport of APP and the spatial regulation of APP cleavage and function in neuronal cells. *Exp Brain Res*. 2011; 217:353–364. [PubMed: 21960299]
- Busciglio J, Lorenzo A, Yeh J, Yankner BA. Beta amyloid fibrils induce tau phosphorylation and loss of microtubule binding. *Neuron*. 1995; 14:879–888. [PubMed: 7718249]
- Busciglio J, Pelsman A, Wong C, Pigino G, Yuan M, Mori H, Yankner BA. Altered metabolism of the amyloid beta precursor protein is associated with mitochondrial dysfunction in Down's syndrome. *Neuron*. 2002; 33:677–688. [PubMed: 11879646]
- Busciglio J, Yankner BA. Apoptosis and increased generation of reactive oxygen species in Down's syndrome neurons in vitro. *Nature*. 1995; 378:776–779. [PubMed: 8524410]
- Cao X, Südhof TC. Dissection of amyloid-beta precursor protein-dependent transcriptional transactivation. *J Biol Chem*. 2004; 279:24601–24611. [PubMed: 15044485]
- Capetillo-Zarate E, Gracia L, Yu F, Banfelder JR, Lin MT, Tampellini D, Gouras GK. High-resolution 3D reconstruction reveals intra-synaptic amyloid fibrils. *Am J Pathol*. 2011; 79:2551–2588. [PubMed: 21925470]
- Chen JX, Yan SS. Role of mitochondrial amyloid- β in Alzheimer's disease. *J Alzheimers Dis*. 2010; 20(suppl 2):569–578. [PubMed: 20164556]
- Choo-Smith LP, Garzon-Rodriguez W, Glabe CG, Surewicz WK. Acceleration of amyloid fibril formation by specific binding of Abeta-(1-40) peptide to ganglioside-containing membrane vesicles. *J Biol Chem*. 1997; 272:22987–22990. [PubMed: 9287293]
- Cirrito JR, Yamada KA, Finn MB, Sloviter RS, Bales KR, May PC, Schoepp DD, Paul SM, Mennerick S, Holtzman DM. Synaptic activity regulates interstitial fluid amyloid-beta levels in vivo. *Neuron*. 2005; 48:913–922. [PubMed: 16364896]
- Cissé M, Halabisky B, Harris J, Devidze N, Dubal DB, Sun B, Orr A, Lotz G, Kim DH, Hamto P, Ho K, Yu GQ, Mucke L. Reversing EphB2 depletion rescues cognitive functions in Alzheimer model. *Nature*. 2011; 469:47–52. [PubMed: 21113149]
- Clariss HJ, Cappai R, Heffernan D, Beyreuther K, Masters CL, Small DH. Identification of heparin-binding domains in the amyloid precursor protein of Alzheimer's disease by deletion mutagenesis and peptide mapping. *J Neurochem*. 1997; 68:1164–1172. [PubMed: 9048763]

- Coraci IS, Husemann J, Berman JW, Hulette C, Dufour JH, Campanella GK, Luster AD, Silverstein SC, El-Khoury JB. CD36, a class B scavenger receptor, is expressed on microglia in Alzheimer's disease brains and can mediate production of reactive oxygen species in response to beta-amyloid fibrils. *Am J Pathol.* 2002; 160:101–112. [PubMed: 11786404]
- Cras P, Kawai M, Lowery D, Gonzalez-DeWhitt P, Greenberg B, Perry G. Senile plaque neurites in Alzheimer disease accumulate amyloid precursor protein. *Proc Natl Acad Sci USA.* 1991; 88:7552–7556. [PubMed: 1652752]
- Dahms SO, Hoefgen S, Roeser D, Schlott B, Gührs KH, Than ME. Structure and biochemical analysis of the heparin-induced E1 dimer of the amyloid precursor protein. *Proc Natl Acad Sci USA.* 2010; 107:5381–5386. [PubMed: 20212142]
- Davis-Salinas J, Saporito-Irwin SM, Cotman CW, Van Nostrand WE. Amyloid beta-protein induces its own production in cultured degenerating cerebrovascular smooth muscle cells. *J Neurochem.* 1995; 65:931–934. [PubMed: 7616257]
- De Felice FG, Velasco PT, Lambert MP, Viola K, Fernandez SJ, Ferreira ST, Klein WL. Abeta oligomers induce neuronal oxidative stress through an N-methyl-D-aspartate receptor-dependent mechanism that is blocked by the Alzheimer drug memantine. *J Biol Chem.* 2007; 282:11590–11601. [PubMed: 17308309]
- De Felice FG, Vieira MN, Bomfim TR, Decker H, Velasco PT, Lambert MP, Viola KL, Zhao WQ, Ferreira ST, Klein WL. Protection of synapses against Alzheimer's-linked toxins: insulin signaling prevents the pathogenic binding of Abeta oligomers. *Proc Natl Acad Sci USA.* 2009; 106:1971–1976. [PubMed: 19188609]
- De Strooper B, Annaert W, Cupers P, Saftig P, Craessaerts K, Mumm JS, Schroeter EH, Schrijvers V, Wolfe MS, Ray WJ, Goate A, Kopan R. A presenilin-1-dependent gamma-secretase-like protease mediates release of Notch intracellular domain. *Nature.* 1999; 398:518–522. [PubMed: 10206645]
- Devi L, Prabhu BM, Galati DF, Avadhani NG, Anandatheerthavarada HK. Accumulation of amyloid precursor protein in the mitochondrial import channels of human Alzheimer's disease brain is associated with mitochondrial dysfunction. *J Neurosci.* 2006; 26:9057–9068. [PubMed: 16943564]
- Deyts C, Vetrivel KS, Das S, Shepherd YM, Dupré DJ, Thinakaran G, Parent AT. Novel Gα_s-protein signaling associated with membrane-tethered amyloid precursor protein intracellular domain. *J Neurosci.* 2012; 32:1714–1729. [PubMed: 22302812]
- Dinamarca MC, Weinstein D, Monasterio O, Inestrosa NC. The synaptic protein neuroligin-1 interacts with the amyloid β-peptide. Is there a role in Alzheimer's disease? *Biochemistry.* 2011; 50:8127–8137. [PubMed: 21838267]
- Dineley KT, Westerman M, Bui D, Bell K, Ashe KH, Sweatt JD. Beta-amyloid activates the mitogen-activated protein kinase cascade via hippocampal α7 nicotinic acetylcholine receptors: in vitro and in vivo mechanisms related to Alzheimer's disease. *J Neurosci.* 2001; 21:4125–4133. [PubMed: 11404397]
- Doyle E, Bruce MT, Breen KC, Smith DC, Anderton B, Regan CM. Intra-ventricular infusions of antibodies to amyloid-beta-protein precursor impair the acquisition of a passive avoidance response in the rat. *Neurosci Lett.* 1990; 115:97–102. [PubMed: 2120637]
- Du H, Guo L, Fang F, Chen D, Sosunov AA, McKhann GM, Yan Y, Wang C, Zhang H, Molkentin JD, Gunn-Moore FJ, Vonsattel JP, Arancio O, Chen JX, Yan SD. Cyclophilin D deficiency attenuates mitochondrial and neuronal perturbation and ameliorates learning and memory in Alzheimer's disease. *Nat Med.* 2008; 14:1097–1105. [PubMed: 18806802]
- El Khoury J, Hickman SE, Thomas CA, Cao L, Silverstein SC, Loike JD. Scavenger receptor-mediated adhesion of microglia to beta-amyloid fibrils. *Nature.* 1996; 382:716–719. [PubMed: 8751442]
- Fang F, Lue LF, Yan S, Xu H, Luddy JS, Chen D, Walker DG, Stern DM, Yan S, Schmidt AM, Chen JX, Yan SS. RAGE-dependent signaling in microglia contributes to neuroinflammation, Abeta accumulation, and impaired learning/memory in a mouse model of Alzheimer's disease. *FASEB J.* 2010; 24:1043–1055. [PubMed: 19906677]
- Fodero LR, Mok SS, Losic D, Martin LL, Aguilar MI, Barrow CJ, Livett BG, Small DH. Alpha7-nicotinic acetylcholine receptors mediate an Abeta(1-42)-induced increase in the level of acetylcholinesterase in primary cortical neurones. *J Neurochem.* 2004; 88:1186–1193. [PubMed: 15009674]

- Galvan V, Chen S, Lu D, Logvinova A, Goldsmith P, Koo EH, Bredesen DE. Caspase cleavage of members of the amyloid precursor family of proteins. *J Neurochem.* 2002; 82:283–294. [PubMed: 12124429]
- Galvan V, Gorostiza OF, Banwait S, Ataie M, Lonvinova AV, Sitaraman S, Carlson E, Sagi SA, Chevallier N, Greenberg DA, Bredesen DE. Reversal of Alzheimer's-like pathology and behavior in human APP transgenic mice by mutation of Asp664. *Proc Natl Acad Sci USA.* 2006; 103:7130–7135. [PubMed: 16641106]
- Galvan V, Zhanga J, Gorostiza OF, Banwait S, Huang W, Ataie M, Tanga H, Bredesen DE. Long-term prevention of Alzheimer's disease-like behavioral deficits in PDAPP mice carrying a mutation in Asp664. *Behav Brain Res.* 2008; 191:246–255. [PubMed: 18485495]
- Ghisso J, Rostagno A, Gardella JE, Liem L, Gorevic PD, Frangione B. A 109-amino-acid C-terminal fragment of Alzheimer's-disease amyloid precursor protein contains a sequence, -RHDS-, that promotes cell adhesion. *Biochem J.* 1992; 288:1053–1059. [PubMed: 1281980]
- Giambarella U, Yamatsuji T, Okamoto T, Matsui T, Ikezu T, Murayama Y, Levine MA, Katz A, Gautam N, Nishimoto I. G-protein betagamma complex-mediated apoptosis by familial Alzheimer's disease mutant of APP. *EMBO J.* 1997; 16:4897–4907. [PubMed: 9305632]
- Gimbel DA, Nygaard HB, Coffey EE, Gunther EC, Laurén J, Gimbel ZA, Strittmatter SM. Memory impairment in transgenic Alzheimer mice requires cellular prion protein. *J Neurosci.* 2010; 30:6367–6374. [PubMed: 20445063]
- Golde TE, Estus S, Younkin LH, Selkoe DJ, Younkin SG. Processing of the amyloid protein precursor to potentially amyloidogenic derivatives. *Science.* 1992; 255:728–730. [PubMed: 1738847]
- Goldgaber D, Lerman MI, McBride OW, Saffiotti U, Gajdusek DC. Characterization and chromosomal localization of a cDNA encoding brain amyloid of Alzheimer's disease. *Science.* 1987; 235:877–880. [PubMed: 3810169]
- Götz J, Eckert A, Matamalas M, Ittner LM, Liu X. Modes of A β toxicity in Alzheimer's disease. *Cell Mol Life Sci.* 2011; 68:3359–3375. [PubMed: 21706148]
- Grace E, Busciglio J. Aberrant activation of focal adhesion proteins mediates fibrillar amyloid h-induced neuronal dystrophy. *J Neurosci.* 2003; 23:493–502. [PubMed: 12533609]
- Grace EA, Rabiner AC, Busciglio JA. Characterization of neuronal dystrophy induced by fibrillar amyloid: implications for Alzheimer's disease. *Neuroscience.* 2002; 114:265–273. [PubMed: 12207971]
- Gralle M, Botelho MG, Wouters FS. Neuroprotective secreted amyloid precursor protein acts by disrupting amyloid precursor protein dimers. *J Biol Chem.* 2009; 284:15016–15025. [PubMed: 19336403]
- Gralle M, Oliveira CL, Guerreiro LH, McKinstry WJ, Galatis D, Masters CL, Cappai R, Parker MW, Ramos CH, Torriani I, Ferreira ST. Solution conformation and heparin-induced dimerization of the full-length extracellular domain of the human amyloid precursor protein. *J Mol Biol.* 2006; 357:493–508. [PubMed: 16436282]
- Gupta-Bansal R, Frederickson RC, Brunden KR. Proteoglycan-mediated inhibition of A beta proteolysis. A potential cause of senile plaque accumulation. *J Biol Chem.* 1995; 270:18666–18671. [PubMed: 7629198]
- Haass C, Kaether C, Thinakaran G, Sisodia S. Trafficking and proteolytic processing of APP. *Cold Spring Harb Perspect Med.* 2012; 2:a006270. [PubMed: 22553493]
- Haass C, Koo EH, Mellon A, Hung AY, Selkoe DJ. Targeting of cell-surface beta-amyloid precursor protein to lysosomes: alternative processing into amyloid-bearing fragments. *Nature.* 1992; 357:500–503. [PubMed: 1608449]
- Harada J, Sugimoto M. Activation of caspase-3 in amyloid beta induced apoptosis of cultured rat cortical neurons. *Brain Res.* 1999; 842:311–323. [PubMed: 10526127]
- Hardy J. The amyloid hypothesis for Alzheimer's disease: a critical reappraisal. *J Neurochem.* 2009; 110:1129–1134. [PubMed: 19457065]
- Hardy J, Selkoe DJ. The amyloid hypothesis of Alzheimer's disease: progress and problems on the road to therapeutics. *Science.* 2002; 297:353–356. [PubMed: 12130773]
- Harris JA, Devidze N, Halabisky B, Lo I, Thwin MT, Yu GQ, Bredesen DE, Masliah E, Mucke L. Many neuronal and behavioral impairments in transgenic mouse models of Alzheimer's disease

- are independent of caspase cleavage of the amyloid precursor protein. *J Neurosci.* 2010b; 30:372–381. [PubMed: 20053918]
- Harris JA, Devidze N, Verret L, Ho K, Halabisky B, Thwin MT, Kim D, Hamto P, Lo I, Yu GQ, Palop JJ, Masliah E, Mucke L. Transsynaptic progression of amyloid- β -induced neuronal dysfunction within the entorhinalhippocampal network. *Neuron.* 2010a; 68:428–441. [PubMed: 21040845]
- Hashimoto Y, Chiba T, Yamada M, Nawa M, Kanekura K, Suzuki H, Terashita K, Aiso S, Nishimoto I, Matsuoka M. Transforming growth factor beta2 is a neuronal death-inducing ligand for amyloid-beta precursor protein. *Mol Cell Biol.* 2005; 25:9304–9317. [PubMed: 16227582]
- Hashimoto Y, Tsuji O, Niikura T, Yamagishi Y, Ishizaka M, Kawasumi M, Chiba T, Kanekura K, Yamada M, Tsukamoto E, Kouyama K, Terashita K, Aiso S, Lin A, Nishimoto I. Involvement of c-Jun N-terminal kinase in amyloid precursor protein-mediated neuronal cell death. *J Neurochem.* 2003; 84:864–877. [PubMed: 12562529]
- Heber S, Herms J, Gajic V, Hainfellner J, Aguzzi A, Rülcke T, von Kretschmar H, von Koch C, Sisodia S, Tremml P, Lipp HP, Wolfner DP, Müller U. Mice with combined gene knock-outs reveal essential and partially redundant functions of amyloid precursor protein family members. *J Neurosci.* 2000; 20:7951–7963. [PubMed: 11050115]
- Heredia L, Helguera P, de Olmos S, Kedikian G, Solá Vigo F, LaFerla F, Staufenbiel M, de Olmos J, Busciglio J, Cáceres A, Lorenzo A. Phosphorylation of ADF/Cofilin by LIM-kinase mediates amyloid beta-induced degeneration: a potential mechanism of neuronal dystrophy in Alzheimer's disease. *J Neurosci.* 2006; 26:6533–6542. [PubMed: 16775141]
- Heredia L, Lin R, Vigo FS, Kedikian G, Busciglio J, Lorenzo A. Deposition of amyloid fibrils promotes cell-surface accumulation of amyloid beta precursor protein. *Neurobiol Dis.* 2004; 16:617–629. [PubMed: 15262274]
- Hilbich C, Monning U, Grund C, Masters CL, Beyreuther K. Amyloid-like properties of peptides flanking the epitope of amyloid precursor protein-specific monoclonal antibody 22C11. *J Biol Chem.* 1993; 268:26571–26577. [PubMed: 7504673]
- Ho A, Südhof TC. Binding of F-spondin to amyloid-beta precursor protein: a candidate amyloid-beta precursor protein ligand that modulates amyloid-beta precursor protein cleavage. *Proc Natl Acad Sci USA.* 2004; 101:2548–2553. [PubMed: 14983046]
- Hoe HS, Lee KJ, Carney RS, Lee J, Markova A, Lee JY, Howell BW, Hyman BT, Pak DT, Bu G, Rebeck GW. Interaction of Reelin with amyloid precursor protein promotes neurite outgrowth. *J Neurosci.* 2009; 29:7459–7473. [PubMed: 19515914]
- Hoe HS, Tran TS, Matsuoka Y, Howell BW, Rebeck GW. DAB1 and Reelin effects on amyloid precursor protein and ApoE receptor 2 trafficking and processing. *J Biol Chem.* 2006; 281:35176–35185. [PubMed: 16951405]
- Hsieh H, Boehm J, Sato C, Iwatsubo T, Tomita T, Sisodia S, Malinow R. AMPAR removal underlies Abeta-induced synaptic depression and dendritic spine loss. *Neuron.* 2006; 52:831–843. [PubMed: 17145504]
- Huang Y, Mucke L. Alzheimer mechanisms and therapeutic strategies. *Cell.* 2012; 148:1204–1222. [PubMed: 22424230]
- Hyman BT, Yuan J. Apoptotic and non-apoptotic roles of caspases in neuronal physiology and pathophysiology. *Nat Rev Neurosci.* 2012; 13:395–406. [PubMed: 22595785]
- Inestrosa NC, Alvarez A, Pérez CA, Moreno RD, Vicente M, Linker C, Casanueva OI, Soto C, Garrido J. Acetylcholinesterase accelerates assembly of amyloid-beta-peptides into Alzheimer's fibrils: possible role of the peripheral site of the enzyme. *Neuron.* 1996; 16:881–891. [PubMed: 8608006]
- Ingelsson M, Fukumoto H, Newell KL, Growdon JH, Hedley-Whyte ET, Frosch MP, Albert MS, Hyman BT, Irazarry MC. Early A β accumulation and progressive synaptic loss, gliosis, and tangle formation in AD brain. *Neurology.* 2004; 62:925–931. [PubMed: 15037694]
- Ivins KJ, Thornton PL, Rohn TT, Cotman CW. Neuronal apoptosis induced by amyloid beta is mediated by caspase-8. *Neurobiol Dis.* 1999; 6:440–449. [PubMed: 10527810]
- Jarrett JT, Berger EP, Lansbury PT Jr. The carboxy terminus of the beta amyloid protein is critical for the seeding of amyloid formation: implications for the pathogenesis of Alzheimer's disease. *Biochemistry.* 1993; 32:4693–4697. [PubMed: 8490014]

- Kaden D, Munter LM, Joshi M, Treiber C, Weise C, Bethge T, Voigt P, Schaefer M, Beyermann M, Reif B, Multhaup G. Homophilic interactions of the amyloid precursor protein (APP) ectodomain are regulated by the loop region and affect beta-secretase cleavage of APP. *J Biol Chem.* 2008; 283:7271–7279. [PubMed: 18182389]
- Kakio A, Nishimoto SI, Yanagisawa K, Kozutsumi Y, Matsuzaki K. Cholesterol-dependent formation of GM1 ganglioside-bound amyloid beta-protein, an endogenous seed for Alzheimer amyloid. *J Biol Chem.* 2001; 276:24985–24990. [PubMed: 11342534]
- Kamenetz F, Tomita T, Hsieh H, Seabrook G, Borchelt D, Iwatsubo T, Sisodia S, Malinow R. APP processing and synaptic function. *Neuron.* 2003; 37:925–937. [PubMed: 12670422]
- Kanaan NM, Pigino GF, Brady ST, Lazarov O, Binder LI, Morfini GA. Axonal degeneration in Alzheimer's disease: when signaling abnormalities meet the axonal transport system. *Exp Neurol.* 2012 [Epub ahead of print].
- Kandalepas PC, Vassar R. Identification and biology of β -secretase. *J Neurochem.* 2012; 120(suppl 1): 55–61. [PubMed: 22122287]
- Kanekiyo T, Zhang J, Liu Q, Liu CC, Zhang L, Bu G. Heparan sulphate proteoglycan and the low-density lipoprotein receptor-related protein 1 constitute major pathways for neuronal amyloid-beta uptake. *J Neurosci.* 2011; 31:1644–1651. [PubMed: 21289173]
- Kang J, Lemaire HG, Unterbeck A, Salbaum JM, Masters CL, Grzeschik KH, Multhaup G, Beyreuther K, Müller-Hill B. The precursor of Alzheimer's disease amyloid A4 protein resembles a cell-surface receptor. *Nature.* 1987; 325:733–736. [PubMed: 2881207]
- Karran E, Mercken M, De Strooper B. The amyloid cascade hypothesis for Alzheimer's disease: an appraisal for the development of therapeutics. *Nat Rev Drug Discov.* 2011; 10:698–712. [PubMed: 21852788]
- Kedikian G, Heredia F, Salvador VR, Raimunda D, Isoardi N, Heredia L, Lorenzo A. Secreted amyloid precursor protein and holo-APP bind amyloid beta through distinct domains eliciting different toxic responses on hippocampal neurons. *J Neurosci Res.* 2010; 88:1795–1803. [PubMed: 20155808]
- Kessels HW, Nguyen LN, Nabavi S, Malinow R. The prion protein as a receptor for amyloid-beta. *Nature.* 2010; 466:E3–E4. [PubMed: 20703260]
- Kibbey MC, Jucker M, Weeks BS, Neve RL, Van Nostrand WE, Kleinman HK. Beta-amyloid precursor protein binds to the neurite-promoting IKVAV site of laminin. *Proc Natl Acad Sci USA.* 1993; 90:10150–10153. [PubMed: 8234269]
- Kinoshita A, Whelan CM, Berezovska O, Hyman BT. The gamma secretase-generated carboxyl terminal domain of the amyloid precursor protein induces apoptosis via Tip60 in H4 cells. *J Biol Chem.* 2002; 277:28530–28536. [PubMed: 12032152]
- Knowles JK, Rajadas J, Nguyen TV, Yang T, LeMieux MC, Vander Griend L, Ishikawa C, Massa SM, Wyss-Coray T, Longo FM. The p75 neurotrophin receptor promotes amyloid-beta(1–42)-induced neuritic dystrophy in vitro and in vivo. *J Neurosci.* 2009; 29:10627–10637. [PubMed: 19710315]
- Koenigsnecht J, Landreth G. Microglial phagocytosis of fibrillar beta-amyloid through a beta1 integrin-dependent mechanism. *J Neurosci.* 2004; 24:9838–9846. [PubMed: 15525768]
- Lacor PN, Buniel MC, Chang L, Fernandez SJ, Gong Y, Viola KL, Lambert MP, Velasco PT, Bigio EH, Finch CE, Krafft GA, Klein WL. Synaptic targeting by Alzheimer's-related amyloid beta oligomers. *J Neurosci.* 2004; 24:10191–10200. [PubMed: 15537891]
- Lacor PN, Buniel MC, Furlow PW, Clemente AS, Velasco PT, Wood M, Viola KL, Klein WL. Abeta oligomer-induced aberrations in synapse composition, shape, and density provide a molecular basis for loss of connectivity in Alzheimer's disease. *J Neurosci.* 2007; 27:796–807. [PubMed: 17251419]
- Lambert MP, Barlow AK, Chromy BA, Edwards C, Freed R, Liosatos M, Morgan TE, Rozovsky I, Trommer B, Viola KL, Wals P, Zhang C, Finch CE, Krafft GA, Klein WL. Diffusible, nonfibrillar ligands derived from Abeta1–42 are potent central nervous system neurotoxins. *Proc Natl Acad Sci USA.* 1998; 95:6448–6453. [PubMed: 9600986]
- Lannfelt L, Basun H, Wahlund LO, Rowe BA, Wagner SL. Decreased alpha-secretase-cleaved amyloid precursor protein as a diagnostic marker for Alzheimer's disease. *Nat Med.* 1995; 1:829–832. [PubMed: 7585189]

- Laurén J, Gimbel DA, Nygaard HB, Gilbert JW, Strittmatter SM. Cellular prion protein mediates impairment of synaptic plasticity by amyloid-beta oligomers. *Nature*. 2009; 457:1128–1132. [PubMed: 19242475]
- Lazarov O, Lee M, Peterson DA, Sisodia SS. Evidence that synaptically released beta-amyloid accumulates as extracellular deposits in the hippocampus of transgenic mice. *J Neurosci*. 2002; 22:9785–9793. [PubMed: 12427834]
- Lefort R, Pozueta J, Shelanski M. Cross-linking of cell surface amyloid precursor protein leads to increased b-Amyloid peptide production in hippocampal neurons: implications for Alzheimer's disease. *J Neurosci*. 2012; 32:10674–10685. [PubMed: 22855816]
- Li ZW, Stark G, Gotz J, Rulicke T, Gschwind M, Huber G, Müller U, Weissmann C. Generation of mice with a 200-kb amyloid precursor protein gene deletion by Cre recombinase-mediated site-specific recombination in embryonic stem cells. *Proc Natl Acad Sci USA*. 1996; 93:6158–6162. [PubMed: 8650236]
- Liu Q, Zerbinatti CV, Zhang J, Hoe HS, Wang B, Cole SL, Herz J, Muglia L, Bu G. Amyloid precursor protein regulates brain apolipoprotein E and cholesterol metabolism through lipoprotein receptor LRP1. *Neuron*. 2007; 56:66–78. [PubMed: 17920016]
- Loo DT, Copani A, Pike CJ, Whittemore ER, Walencewicz AJ, Cotman CW. Apoptosis is induced by beta-amyloid in cultured central nervous system neurons. *Proc Natl Acad Sci USA*. 1993; 90:7951–7955. [PubMed: 8367446]
- Lorenzo A, Yankner BA. Beta amyloid neurotoxicity requires fibril formation and is inhibited by Congo red. *Proc Natl Acad Sci USA*. 1994; 91:12243–12247. [PubMed: 7991613]
- Lorenzo A, Yuan M, Zhang Z, Paganetti PA, Sturchler-Pierrat C, Staufenbiel M, Mautino J, Vigo FS, Sommer B, Yankner BA. Amyloid beta interacts with the amyloid precursor protein: a potential toxic mechanism in Alzheimer's disease. *Nat Neurosci*. 2000; 3:460–464. [PubMed: 10769385]
- Loureño FC, Galvan V, Fombonne J, Corset V, Llambi F, Müller U, Bredesen DE, Mehlen P. Netrin-1 interacts with amyloid precursor protein and regulates amyloid-beta production. *Cell Death Differ*. 2009; 16:655–663. [PubMed: 19148186]
- Lu DC, Rabizadeh S, Chandra S, Shayya RF, Ellerby LM, Ye X, Salvesen GS, Koo EH, Bredesen DE. A second cytotoxic proteolytic peptide derived from amyloid beta-protein precursor. *Nat Med*. 2000; 6:397–404. [PubMed: 10742146]
- Lu DC, Shaked GM, Masliah E, Bredesen DE, Koo EH. Amyloid β protein toxicity mediated by the formation of amyloid- β protein precursor complexes. *Ann Neurol*. 2003; 54:781–789. [PubMed: 14681887]
- Lustbader JW, Cirilli M, Lin C, Xu HW, Takuma K, Wang N, Caspersen C, Chen X, Pollak S, Chaney M, Trinchese F, Liu S, Gunn-Moore F, Lue LF, Walker DG, Kuppusamy P, Zewier ZL, Arancio O, Stern D, Yan SS, Wu H. A β AD directly links A β to mitochondrial toxicity in Alzheimer's disease. *Science*. 2004; 304:448–452. [PubMed: 15087549]
- Ma QL, Yang F, Calon F, Ubeda OJ, Hansen JE, Weisbart RH, Beech W, Frautschy SA, Cole GM. p21-activated kinase-aberrant activation and translocation in Alzheimer disease pathogenesis. *J Biol Chem*. 2008; 283:14132–14143. [PubMed: 18347024]
- Magdesian MH, Carvalho MM, Mendes FA, Saraiva LM, Juliano MA, Juliano L, Garcia-Abreu J, Ferreira ST. Amyloid-beta binds to the extracellular cysteine-rich domain of Frizzled and inhibits Wnt/beta-catenin signaling. *J Biol Chem*. 2008; 283:9359–9368. [PubMed: 18234671]
- Manczak M, Anekonda TS, Henson E, Park BS, Quinn J, Reddy PH. Mitochondria are a direct site of A β accumulation in Alzheimer's disease neurons: implications for free radical generation and oxidative damage in disease progression. *Hum Mol Genet*. 2006; 15:1437–1449. [PubMed: 16551656]
- Manczak M, Reddy PH. Abnormal interaction of VDAC1 with amyloid beta and phosphorylated tau causes mitochondrial dysfunction in Alzheimer's disease. *Hum Mol Genet*. 2012; 21:5131–5146. [PubMed: 22926141]
- Marsden IT, Minamide LS, Bamberg JR. Amyloid- β -induced amyloid- β secretion: a possible feed-forward mechanism in Alzheimer's disease. *J Alzheimers Dis*. 2011; 24:1–11.
- Matsuda S, Yasukawa T, Homma Y, Ito Y, Niikura T, Hiraki T, Hirai S, Ohno S, Kita Y, Kawasumi M, Kouyama K, Yamamoto T, Kyriakis JM, Nishimoto I. c-Jun N-terminal kinase (JNK)-

- interacting protein-1b/islet-brain-1 scaffolds Alzheimer's amyloid precursor protein with JNK. *J Neurosci.* 2001; 21:6597–6607. [PubMed: 11517249]
- Mattson MP. Cellular actions of beta-amyloid precursor protein and its soluble and fibrillogenic derivatives. *Physiol Rev.* 1997; 77:1081–1132. [PubMed: 9354812]
- Mattson MP. Pathways towards and away from Alzheimer's disease. *Nature.* 2004; 430:631–639. [PubMed: 15295589]
- McGowan E, Pickford F, Kim J, Onstead L, Eriksen J, Yu C, Skipper L, Murphy MP, Beard J, Das P, Jansen K, DeLucia M, Lin WL, Dolios G, Wang R, Eckman CB, Dickson DW, Hutton M, Hardy J, Golde T. Abeta42 is essential for parenchymal and vascular amyloid deposition in mice. *Neuron.* 2006; 47:191–199. [PubMed: 16039562]
- McLaurin J, Chakrabarty A. Membrane disruption by Alzheimer beta-amyloid peptides mediated through specific binding to either phospholipids or gangliosides. Implications for neurotoxicity. *J Biol Chem.* 1996; 271:26482–26489. [PubMed: 8900116]
- McLaurin J, Franklin T, Kuhns WJ, Fraser PE. A sulfated proteoglycan aggregation factor mediates amyloid-beta peptide fibril formation and neuro-toxicity. *Amyloid.* 1999; 6:233–243. [PubMed: 10611943]
- McLoughlin DM, Miller CC. The FE65 proteins and Alzheimer's disease. *J Neurosci Res.* 2008; 86:744–754. [PubMed: 17828772]
- McPhie DL, Coopersmith R, Hines-Peralta A, Chen Y, Ivins KJ, Manly SP, Kozlowski MR, Neve KA, Neve RL. DNA synthesis and neuronal apoptosis caused by familial Alzheimer disease mutants of the amyloid precursor protein are mediated by the p21-activated kinase PAK3. *J Neurosci.* 2003; 23:6914–6927. [PubMed: 12890786]
- Melchor JP, Van Nostrand WE. Fibrillar amyloid beta-protein mediates the pathologic accumulation of its secreted precursor in human cerebrovascular smooth muscle cells. *J Biol Chem.* 2000; 275:9782–9791. [PubMed: 10734132]
- Meziane H, Dodart JC, Mathis C, Little S, Clemens J, Paul SM, Ungerer A. Memory-enhancing effects of secreted forms of the β -amyloid precursor protein in normal and amnesic mice. *Proc Natl Acad Sci USA.* 1998; 95:12683–12688. [PubMed: 9770546]
- Moore KJ, El Khoury J, Medeiros LA, Terada K, Geula C, Luster AD, Freeman MW. A CD36-initiated signaling cascade mediates inflammatory effects of beta-amyloid. *J Biol Chem.* 2002; 277:47373–47379. [PubMed: 12239221]
- Mucke L, Masliah E, Johnson W, Ruppe M, Alford M, Rockenstein E, Forss-Petter S, Pietropaolo M, Mallory M, Abraham C. Synaptotrophic effects of human amyloid protein precursors in the cortex of transgenic mice. *Brain Res.* 1994; 666:151–167. [PubMed: 7882025]
- Müller UC, Zheng H. Physiological functions of APP family proteins. *Cold Spring Harb Perspect Med.* 2012; 4:a006288. [PubMed: 22355794]
- Munter LM, Voigt P, Harmeier A, Kaden D, Gottschalk KE, Weise C, Pipkorn R, Schaefer M, Langosch D, Multhaup G. GxxxG motifs within the amyloid precursor protein transmembrane sequence are critical for the etiology of Abeta42. *EMBO J.* 2007; 26:1702–1712. [PubMed: 17332749]
- Nagele RG, D'Andrea MR, Anderson WJ, Wang HY. Intracellular accumulation of beta-amyloid(1-42) in neurons is facilitated by the alpha 7 nicotinic acetylcholine receptor in Alzheimer's disease. *Neuroscience.* 2002; 110:199–211. [PubMed: 11958863]
- Nakagawa T, Zhu H, Morishima N, Li E, Xu J, Yankner BA, Yuan J. Caspase-12 mediates endoplasmic-reticulum-specific apoptosis and cytotoxicity by amyloid beta. *Nature.* 2000; 403:98–103. [PubMed: 10638761]
- Narindrasorasak S, Lowery D, Gonzalez-DeWhitt P, Poorman RA, Greenberg B, Kisilevsky R. High affinity interactions between the Alzheimer's beta-amyloid precursor proteins and the basement membrane form of heparan sulfate proteoglycan. *J Biol Chem.* 1991; 266:12878–12883. [PubMed: 1906461]
- Ni CY, Murphy MP, Golde TE, Carpenter G. Gamma-secretase cleavage and nuclear localization of ErbB-4 receptor tyrosine kinase. *Science.* 2001; 294:2179–2181. [PubMed: 11679632]

- Niikura T, Yamada M, Chiba T, Aiso S, Matsuoka M, Nishimoto I. Characterization of V642I-Aβ-induced cytotoxicity in primary neurons. *J Neurosci Res*. 2004; 77:54–62. [PubMed: 15197738]
- Ninomiya H, Roch JM, Sundsmo MP, Otero DA, Saitoh T. Amino acid sequence RERMS represents the active domain of amyloid beta/A4 protein precursor that promotes fibroblast growth. *J Cell Biol*. 1993; 121:879–886. [PubMed: 8491779]
- Nishimoto I, Okamoto T, Matsuura Y, Takahashi S, Okamoto T, Murayama Y, Ogata E. Alzheimer amyloid protein precursor complexes with brain GTP-binding protein G(o). *Nature*. 1993; 362:75–79. [PubMed: 8446172]
- Ohgami T, Kitamoto T, Tateishi J. Alzheimer's amyloid precursor protein accumulates within axonal swellings in human brain lesions. *Neurosci Lett*. 1992; 136:75–78. [PubMed: 1635670]
- Okamoto T, Takeda S, Giambarella U, Murayama Y, Matsui T, Katada T, Matsuura Y, Nishimoto I. Intrinsic signaling function of APP as a novel target of three V642 mutations linked to familial Alzheimer's disease. *EMBO J*. 1996; 15:3769–3777. [PubMed: 8670881]
- Okamoto T, Takeda S, Murayama Y, Ogata E, Nishimoto I. Ligand-dependent G protein coupling function of amyloid transmembrane precursor. *J Biol Chem*. 1995; 270:4205–4208. [PubMed: 7876177]
- Olzscha H, Schermann SM, Woerner AC, Pinkert S, Hecht MH, Tartaglia GG, Vendruscolo M, Hayer-Hartl M, Hartl FU, Vabulas RM. Amyloid-like aggregates sequester numerous metastable proteins with essential cellular functions. *Cell*. 2011; 144:67–78. [PubMed: 21215370]
- Opazo P, Saud K, de Saint Pierre M, Cárdenas AM, Allen DD, Segura-Aguilar J, Caviedes R, Caviedes P. Knockdown of amyloid precursor protein normalizes cholinergic function in a cell line derived from the cerebral cortex of a trisomy 16 mouse: an animal model of Down syndrome. *J Neurosci Res*. 2006; 84:1303–1310. [PubMed: 16941497]
- Origlia N, Bonadonna C, Rosellini A, Leznik E, Arancio O, Yan SS, Domenici L. Microglial receptor for advanced glycation end product-dependent signal pathway drives beta-amyloid-induced synaptic depression and long-term depression impairment in entorhinal cortex. *J Neurosci*. 2010; 30:11414–11425. [PubMed: 20739563]
- Pardossi-Piquard R, Petit A, Kawai T, Sunyach C, Alves da Costa C, Vincent B, Ring S, D'Adamio L, Shen J, Müller U, St George Hyslop P, Checler F. Presenilin-dependent transcriptional control of the Aβ-degrading enzyme neprilysin by intracellular domains of βAPP and APLP. *Neuron*. 2005; 46:541–554. [PubMed: 15944124]
- Paresce DM, Chung H, Maxfield FR. Slow degradation of aggregates of the Alzheimer's disease amyloid beta-protein by microglial cells. *J Biol Chem*. 1997; 272:29390–29397. [PubMed: 9361021]
- Paresce DM, Ghosh RN, Maxfield FR. Microglial cells internalize aggregates of the Alzheimer's disease amyloid beta-protein via a scavenger receptor. *Neuron*. 1996; 17:553–565. [PubMed: 8816718]
- Park JH, Gimbel DA, GrandPre T, Lee JK, Kim JE, Li W, Lee DH, Strittmatter SM. Alzheimer precursor protein interaction with the Nogo-66 receptor reduces amyloid-beta plaque deposition. *J Neurosci*. 2006; 26:1386–1395. [PubMed: 16452662]
- Park SA, Shaked GM, Bredesen DE, Koo EH. Mechanism of cytotoxicity mediated by the C31 fragment of the amyloid precursor protein. *Biochem Biophys Res Commun*. 2009; 388:450–455. [PubMed: 19679105]
- Patel TB. Single transmembrane spanning heterotrimeric G protein-coupled receptors and their signaling cascades. *Pharmacol Rev*. 2004; 56:371–385. [PubMed: 15317909]
- Perez RG, Zheng H, Van der Ploeg LH, Koo EH. The beta-amyloid precursor protein of Alzheimer's disease enhances neuron viability and modulates neuronal polarity. *J Neurosci*. 1997; 17:9407–9414. [PubMed: 9390996]
- Perini G, Della-Bianca V, Politi V, Della Valle G, Dal-Pra I, Rossi F, Armato U. Role of p75 neurotrophin receptor in the neurotoxicity by beta-amyloid peptides and synergistic effect of inflammatory cytokines. *J Exp Med*. 2002; 195:907–918. [PubMed: 11927634]

- Phinney AL, Deller T, Stalder M, Calhoun ME, Frotscher M, Sommer B, Staufenbiel M, Jucker M. Cerebral amyloid induces aberrant axonal sprouting and ectopic terminal formation in amyloid precursor protein transgenic mice. *J Neurosci*. 1999; 19:8552–8559. [PubMed: 10493755]
- Pigino G, Morfini G, Atagi Y, Deshpande A, Yu C, Jungbauer L, LaDu M, Busciglio J, Brady S. Disruption of fast axonal transport is a pathogenic mechanism for intraneuronal amyloid beta. *Proc Natl Acad Sci USA*. 2009; 106:5907–5912. [PubMed: 19321417]
- Pike CJ, Cummings BJ, Cotman CW. Beta-amyloid induces neuritic dystrophy in vitro: similarities with Alzheimer's disease. *Neuroreport*. 1992; 3:769–772. [PubMed: 1421135]
- Pike CJ, Walencewicz AJ, Glabe CG, Cotman CW. In vitro aging of b-amyloid protein causes peptide aggregation and neurotoxicity. *Brain Res*. 1991; 563:311–314. [PubMed: 1786545]
- Puzzo D, Privitera L, Leznik E, Fà M, Staniszewski A, Palmeri A, Arancio O. Picomolar amyloid-beta positively modulates synaptic plasticity and memory in hippocampus. *J Neurosci*. 2008; 28:14537–14545. [PubMed: 19118188]
- Reddy PH, Beal MF. Amyloid beta, mitochondrial dysfunction and synaptic damage: implications for cognitive decline in aging and Alzheimer's disease. *Trends Mol Med*. 2008; 14:45–53. [PubMed: 18218341]
- Reiss K, Maretzky T, Ludwig A, Tousseyn T, de Strooper B, Hartmann D, Saftig P. ADAM10 cleavage of N-cadherin and regulation of cell-cell adhesion and beta-catenin nuclear signalling. *EMBO J*. 2005; 24:742–752. Erratum in: *EMBO J*. 24:1762. [PubMed: 15692570]
- Renner M, Lacor PN, Velasco PT, Xu J, Contractor A, Klein WL, Triller A. Deleterious effects of amyloid beta oligomers acting as an extracellular scaffold for mGluR5. *Neuron*. 2010; 66:739–754. [PubMed: 20547131]
- Rijal Upadhaya A, Capetillo-Zarate E, Kosterin I, Abramowski D, Kumar S, Yamaguchi H, Walter J, Fändrich M, Staufenbiel M, Thal DR. Dispersible amyloid β -protein oligomers, protofibrils, and fibrils represent diffusible but not soluble aggregates: their role in neurodegeneration in amyloid precursor protein (APP) transgenic mice. *Neurobiol Aging*. 2012; 33:2641–2660. [PubMed: 22305478]
- Ring S, Weyer SW, Kilian SB, Waldron E, Pietrzik CU, Filippov MA, Herms J, Buchholz C, Eckman CB, Korte M, Wolfer DP, Müller UC. The secreted beta-amyloid precursor protein ectodomain APPs alpha is sufficient to rescue the anatomical behavioral, and electrophysiological abnormalities of APP-deficient mice. *J Neurosci*. 2007; 27:7817–7826. [PubMed: 17634375]
- Rochet JC, Lansbury PT Jr. Amyloid fibrillogenesis: themes and variations. *Curr Opin Struct Biol*. 2000; 10:60–68. [PubMed: 10679462]
- Rohn TT, Ivins KJ, Bahr BA, Cotman CW, Cribbs DH. A monoclonal antibody to amyloid precursor protein induces neuronal apoptosis. *J Neurochem*. 2000; 74:2331–2342. [PubMed: 10820193]
- Rojas G, Cárdenas AM, Fernández-Olivares P, Shimahara T, Segura-Aguilar J, Caviedes R, Caviedes P. Effect of the knockdown of amyloid precursor protein on intracellular calcium increases in a neuronal cell line derived from the cerebral cortex of a trisomy 16 mouse. *Exp Neurol*. 2008; 209:234–242. [PubMed: 17976585]
- Rovelet-Lecrux A, Hannequin D, Raux G, Le Meur N, Laquerrière A, Vital A, Dumanchin C, Feuillet S, Brice A, Vercelletto M, Dubas F, Frebourg T, Campion D. APP locus duplication causes autosomal dominant early-onset Alzheimer disease with cerebral amyloid angiopathy. *Nat Genet*. 2006; 38:24–26. [PubMed: 16369530]
- Roychoudhuri R, Yang M, Hoshi MM, Teplow DB. Amyloid beta-protein assembly and Alzheimer disease. *J Biol Chem*. 2009; 284:4749–4753. [PubMed: 18845536]
- Rozemuller AJ, Roos RA, Bots GT, Kamphorst W, Eikelenboom P, Van Nostrand WE. Distribution of beta/A4 protein and amyloid precursor protein in hereditary cerebral hemorrhage with amyloidosis-Dutch type and Alzheimer's disease. *Am J Pathol*. 1993; 142:1449–1457. [PubMed: 7684195]
- Russell CL, Semerdjieva S, Empson RM, Austen BM, Beesley PW, Alifragis P. Amyloid-b acts as a regulator of neurotransmitter release disrupting the interaction between synaptophysin and VAMP2. *PLoS One*. 2012; 7:e43201. [PubMed: 22905234]

- Russo C, Venezia V, Repetto E, Nizzari M, Violani E, Carlo P, Schettini G. The amyloid precursor protein and its network of interacting proteins: physiological and pathological implications. *Brain Res Brain Res Rev.* 2005; 48:257–264. [PubMed: 15850665]
- Sabo SL, Ikin AF, Buxbaum JD, Greengard P. The amyloid precursor protein and its regulatory protein, FE65, in growth cones and synapses in vitro and in vivo. *J Neurosci.* 2003; 23:5407–5415. [PubMed: 12843239]
- Saganich MJ, Schroeder BE, Galvan V, Bredesen DE, Koo EH, Heinemann SF. Deficits in synaptic transmission and learning in amyloid precursor protein (APP) transgenic mice require C-terminal cleavage of APP. *J Neurosci.* 2006; 26:13428–13436. [PubMed: 17192425]
- Saito Y, Akiyama M, Araki Y, Sumioka A, Shiono M, Taru H, Nakaya T, Yamamoto T, Suzuki T. Intracellular trafficking of the amyloid b-protein precursor (APP) regulated by novel function of X11-like. *PLoS One.* 2011; 6:e22108. [PubMed: 21818298]
- Saitoh T, Sundsmo M, Roch JM, Kimura N, Cole G, Schubert D, Oltersdorf T, Schenk DB. Secreted form of amyloid beta protein precursor is involved in the growth regulation of fibroblasts. *Cell.* 1989; 58:615–622. [PubMed: 2475254]
- Salehi A, Delcroix JD, Belichenko PV, Zhan K, Wu C, Valletta JS, Takimoto-Kimura R, Kleschevnikov AM, Sambamurti K, Chung PP, Xia W, Villar A, Campbell WA, Kulnane LS, Nixon RA, Lamb BT, Epstein CJ, Stokin GB, Goldstein LS, Mobley WC. Increased App expression in a mouse model of Down's syndrome disrupts NGF transport and causes cholinergic neuron degeneration. *Neuron.* 2006; 51:29–42. [PubMed: 16815330]
- Scheinfeld MH, Ghersi E, Laky K, Fowlkes BJ, D'Adamio L. Processing of beta-amyloid precursor-like protein-1 and -2 by gamma-secretase regulates transcription. *J Biol Chem.* 2002; 277:44195–44201. [PubMed: 12228233]
- Scheuermann S, Hamsch B, Hesse L, Stumm J, Schmidt C, Behr D, Bayer TA, Beyreuther K, Multhaup G. Homodimerization of amyloid precursor protein and its implication in the amyloidogenic pathway of Alzheimer's disease. *J Biol Chem.* 2001; 276:33923–33929. [PubMed: 11438549]
- Shaked GM, Kummer MP, Lu DC, Galvan V, Bredesen DE, Koo EH. Abeta induces cell death by direct interaction with its cognate extracellular domain on APP (APP 597-624). *FASEB J.* 2006; 20:1254–1256. [PubMed: 16636103]
- Sheng JG, Price DL, Koliatsos VE. Disruption of corticocortical connections ameliorates amyloid burden in terminal fields in a transgenic model of Abeta amyloidosis. *J Neurosci.* 2002; 22:9794–9799. [PubMed: 12427835]
- Simons M, Ikonen E, Tienari PJ, Cid-Arregui A, Mönning U, Beyreuther K, Dotti CG. Intracellular routing of human amyloid proteinprecursor: axonal delivery followed by transport to the dendrites. *J Neurosci Res.* 1995; 41:121–128. [PubMed: 7674373]
- Snyder EM, Nong Y, Almeida CG, Paul S, Moran T, Choi EY, Nairn AC, Salter MW, Lombroso PJ, Gouras GK, Greengard P. Regulation of NMDA receptor trafficking by amyloid-beta. *Nat Neurosci.* 2005; 8:1051–1058. [PubMed: 16025111]
- Soba P, Eggert S, Wagner K, Zentgraf H, Siehl K, Kreger S, Löwer A, Langer A, Merdes G, Paro R, Masters CL, Müller U, Kins S, Beyreuther K. Homo- and heterodimerization of APP family members promotes intercellular adhesion. *EMBO J.* 2005; 24:3624–3634. [PubMed: 16193067]
- Sola Vigo F, Kedikian G, Heredia L, Heredia F, Añel AD, Rosa AL, Lorenzo A. Amyloid-beta precursor protein mediates neuronal toxicity of amyloid beta through Go protein activation. *Neurobiol Aging.* 2009; 30:1379–1392. [PubMed: 18187234]
- Sothibundhu A, Sykes AM, Fox B, Underwood CK, Thangnipon W, Coulson EJ. Beta-amyloid(1-42) induces neuronal death through the p75 neurotrophin receptor. *J Neurosci.* 2008; 28:3941–3946. [PubMed: 18400893]
- Sudo H, Hashimoto Y, Niikura T, Shao Z, Yasukawa T, Ito Y, Yamada M, Hata M, Hiraki T, Kawasumi M, Kouyama K, Nishimoto I. Secreted Abeta does not mediate neurotoxicity by antibody-stimulated amyloid precursor protein. *Biochem Biophys Res Commun.* 2001; 282:548–556. [PubMed: 11401495]

- Sudo H, Jiang H, Yasukawa T, Hashimoto Y, Niikura T, Kawasumi M, Matsuoka M, Murayama Y, Nishimoto I. Antibody-regulated neurotoxic function of cell-surface beta amyloid precursor protein. *Mol Cell Neurosci.* 2000; 16:708–723. [PubMed: 11124892]
- Swanson TL, Knittel LM, Coate TM, Farley SM, Snyder MA, Copenhaver PF. The insect homologue of the amyloid precursor protein interacts with the heterotrimeric G protein Go alpha in an identified population of migratory neurons. *Dev Biol.* 2005; 288:160–178. [PubMed: 16229831]
- Szodrai A, Kuan YH, Hunzelmann S, Engel U, Sakane A, Sasaki T, Takai Y, Kirsch J, Müller U, Beyreuther K, Brady S, Morfini G, Kins S. APP anterograde transport requires Rab3A GTPase activity for assembly of the transport vesicle. *J Neurosci.* 2009; 29:14534–14544. [PubMed: 19923287]
- Tanzi R. The genetics of Alzheimer disease. *Cold Spring Harb Perspect Med.* 2012; 2:a006296. [PubMed: 23028126]
- Tanzi RE, Gusella JF, Watkins PC, Bruns GA, St George-Hyslop P, Van Keuren ML, Patterson D, Pagan S, Kurnit DM, Neve RL. Amyloid beta protein gene: cDNA, mRNA distribution, and genetic linkage near the Alzheimer locus. *Science.* 1987; 235:880–884. [PubMed: 2949367]
- Thal DR, Rüb U, Orantes M, Braak H. Phases of A beta-deposition in the human brain and its relevance for the development of AD. *Neurology.* 2002; 58:1791–1800. [PubMed: 12084879]
- Theuns J, Brouwers N, Engelborghs S, Sleegers K, Bogaerts V, Corsmit E, De Pooter T, van Duijn CM, De Deyn PP, Van Broeckhoven C. Promoter mutations that increase amyloid precursor-protein expression are associated with Alzheimer disease. *Am J Hum Genet.* 2006; 78:936–946. [PubMed: 16685645]
- Thinakaran G, Koo EH. Amyloid precursor protein trafficking, processing, and function. *J Biol Chem.* 2008; 283:29615–29619. [PubMed: 18650430]
- Tienari PJ, De Strooper B, Ikonen E, Simons M, Weidemann A, Czech C, Hartmann T, Ida N, Multhaup G, Masters CL, Van Leuven F, Beyreuther K, Dotti CG. The beta-amyloid domain is essential for axonal sorting of amyloid precursor protein. *EMBO J.* 1996; 5:5218–5229. [PubMed: 8895567]
- Torroja L, Packard M, Gorczyca M, White K, Budnik V. The Drosophila beta-amyloid precursor protein homolog promotes synapse differentiation at the neuromuscular junction. *J Neurosci.* 1999; 19:7793–7803. [PubMed: 10479682]
- Troy CM, Rabacchi SA, Friedman WJ, Frappier TF, Brown K, Shelanski ML. Caspase-2 mediates neuronal cell death induced by amyloid beta. *J Neurosci.* 2000; 20:1386–1392. [PubMed: 10662829]
- Tsai J, Grutzendler J, Duff K, Gan WB. Fibrillar amyloid deposition leads to local synaptic abnormalities and breakage of neuronal branches. *Nat Neurosci.* 2004; 7:1181–1183. [PubMed: 15475950]
- Tsuruma K, Tanaka Y, Shimazawa M, Hara H. Induction of amyloid precursor protein by the neurotoxic peptide, amyloid-beta 25-35, causes retinal ganglion cell death. *J Neurochem.* 2012; 113:1545–1554. [PubMed: 20374419]
- Um JW, Nygaard HB, Heiss JK, Kostylev MA, Stagi M, Vortmeyer A, Wisniewski T, Gunther EC, Strittmatter SM. Alzheimer amyloid- β oligomer bound to postsynaptic prion protein activates Fyn to impair neurons. *Nat Neurosci.* 2012; 15:1227–1235. [PubMed: 22820466]
- Van Nostrand WE, Melchor JP, Keane DM, Saporito-Irwin SM, Romanov G, Davis J, Xu F. Localization of a fibrillar amyloid beta-protein binding domain on its precursor. *J Biol Chem.* 2002; 277:36392–36398. [PubMed: 12107175]
- Van Nostrand WE, Wagner SL, Shankle WR, Farrow JS, Dick M, Rozemuller JM, Kuiper MA, Wolters EC, Zimmerman J, Cotman CW, et al. Decreased levels of soluble amyloid beta-protein precursor in cerebro-spinal fluid of live Alzheimer disease patients. *Proc Natl Acad Sci USA.* 1992; 89:2551–2555. [PubMed: 1557359]
- von Koch CS, Zheng H, Chen H, Trumbauer M, Thinakaran G, van der Ploeg LH, Price DL, Sisodia SS. Generation of APLP2 KO mice and early postnatal lethality in APLP2/APP double KO mice. *Neurobiol Aging.* 1997; 18:661–669. [PubMed: 9461064]

- Wagner MR, Keane DM, Melchor JP, Auspaker KR, Van Nostrand WE. Fibrillar amyloid beta-protein binds protease nexin-2/amyloid beta-protein precursor: stimulation of its inhibition of coagulation factor XIa. *Biochemistry*. 2000; 39:7420–7427. [PubMed: 10858290]
- Walsh DM, Klyubin I, Fadeeva JV, Cullen WK, Anwyl R, Wolfe MS, Rowan MJ, Selkoe DJ. Naturally secreted oligomers of amyloid beta protein potently inhibit hippocampal long-term potentiation in vivo. *Nature*. 2002; 416:535–539. [PubMed: 11932745]
- Wang B, Yang L, Wang Z, Zheng H. Amyloid precursor protein mediates presynaptic localization and activity of the high-affinity choline transporter. *Proc Natl Acad Sci USA*. 2007; 104:14140–14145. [PubMed: 17709753]
- Wang HY, Lee DH, D'Andrea MR, Peterson PA, Shank RP, Reitz AB. Beta-amyloid(1–42) binds to alpha7 nicotinic acetylcholine receptor with high affinity. Implications for Alzheimer's disease pathology. *J Biol Chem*. 2000; 275:5626–5632. [PubMed: 10681545]
- Wang Q, Klyubin I, Wright S, Griswold-Prenner I, Rowan MJ, Anwyl R. Alpha v integrins mediate beta-amyloid induced inhibition of long-term potentiation. *Neurobiol Aging*. 2008; 29:1485–1493. [PubMed: 17442458]
- Wang Y, Ha Y. The X-ray structure of an antiparallel dimer of the human amyloid precursor protein E2 domain. *Mol Cell*. 2004; 15:343–353. [PubMed: 15304215]
- Wang Z, Wang B, Yang L, Guo Q, Aithmitti N, Songyang Z, Zheng H. Presynaptic and postsynaptic interaction of the amyloid precursor protein promotes peripheral and central synaptogenesis. *J Neurosci*. 2009; 29:10788–10801. [PubMed: 19726636]
- Wasco W, Gurubhagavatula S, Paradis MD, Romano DM, Sisodia SS, Hyman BT, Neve RL, Tanzi RE. Isolation and characterization of APLP2 encoding a homologue of the Alzheimer's associated amyloid beta protein precursor. *Nat Genet*. 1993; 5:95–100. [PubMed: 8220435]
- White A, Maher F, Brazier M, Jobling M, Thyer J, Stewart L, Thompson A, Gibson R, Masters C, Multhaup G, Beyreuther K, Barrow C, Collins S, Cappai R. Diverse fibrillar peptides directly bind the Alzheimer's amyloid precursor protein and amyloid-like protein 2 resulting in cellular accumulation. *Brain Res*. 2003; 966:231–244. [PubMed: 12618346]
- Williams TL, Johnson BR, Urbanc B, Jenkins AT, Connell SD, Serpell LC. Aβ42 oligomers, but not fibrils, simultaneously bind to and cause damage to ganglioside-containing lipid membranes. *Biochem J*. 2011; 439:67–77. [PubMed: 21702743]
- Wolfe MS. γ-Secretase as a target for Alzheimer's disease. *Adv Pharmacol*. 2012; 64:127–153. [PubMed: 22840746]
- Wright S, Malinin NL, Powell KA, Yednock T, Rydel RE, Griswold-Prenner I. Alpha2beta1 and alphaVbeta1 integrin signaling pathways mediate amyloid-beta-induced neurotoxicity. *Neurobiol Aging*. 2007; 28:226–237. [PubMed: 16448724]
- Yaar M, Zhai S, Pilch PF, Doyle SM, Eisenhauer PB, Fine RE, Gilchrist BA. Binding of beta-amyloid to the p75 neurotrophin receptor induces apoptosis. A possible mechanism for Alzheimer's disease. *J Clin Invest*. 1997; 100:2333–2340. [PubMed: 9410912]
- Yamaguchi H, Yamazaki T, Ishiguro K, Shoji M, Nakazato Y, Hirai S. Ultrastructural localization of Alzheimer amyloid beta/A4 protein precursor in the cytoplasm of neurons and senile plaque-associated astrocytes. *Acta Neuropathol*. 1992; 85:15–22. [PubMed: 1363016]
- Yamatsuji T, Matsui T, Okamoto T, Komatsuzaki K, Takeda S, Fukumoto H, Iwatsubo T, Suzuki N, Asami-Odaka A, Ireland S, Kinane TB, Giambarella U, Nishimoto I. G Protein-mediated neuronal DNA fragmentation induced by familial Alzheimer's disease-associated mutants of APP. *Science*. 1996a; 272:1349–1352. [PubMed: 8650548]
- Yamatsuji T, Okamoto T, Takeda S, Murayama Y, Tanaka N, Nishimoto I. Expression of V642 APP mutant causes cellular apoptosis as Alzheimer trait-linked phenotype. *EMBO J*. 1996b; 15:498–509. [PubMed: 8599933]
- Yamazaki T, Koo EH, Selkoe DJ. Cell surface amyloid beta-protein precursor colocalizes with beta 1 integrins at substrate contact sites in neural cells. *J Neurosci*. 1997; 17:1004–1010. [PubMed: 8994055]
- Yamazaki T, Selkoe DJ, Koo EH. Trafficking of cell surface beta-amyloid precursor protein: retrograde and transcytotic transport in cultured neurons. *J Cell Biol*. 1995; 129:431–442. [PubMed: 7721945]

- Yan SD, Chen X, Chen M, Zhu H, Roher A, Slattery T, Zhao L, Nagashima M, Morser J, Migheli A, Nawroth P, Stern DM, Schmidt AM. RAGE and Abeta peptide neurotoxicity in Alzheimer's disease. *Nature*. 1996; 382:685–691. [PubMed: 8751438]
- Yan SD, Zhu H, Fu J, Yan SF, Roher A, Tourtellotte WW, Rajavashisth T, Chen X, Godman GC, Stern D, Schmidt AM. Amyloid-beta peptide-receptor for advanced glycation endproduct interaction elicits neuronal expression of macrophage-colony stimulating factor: a proinflammatory pathway in Alzheimer disease. *Proc Natl Acad Sci USA*. 1997; 94:5296–5301. [PubMed: 9144231]
- Yang AJ, Chandswangbhuvana D, Shu T, Henschen A, Glabe CG. Intracellular accumulation of insoluble, newly synthesized abeta-42 in amyloid precursor protein-transfected cells that have been treated with Abeta1-42. *J Biol Chem*. 1999; 274:20650–20656. [PubMed: 10400697]
- Yankner BA, Dawes LR, Fisher S, Villa-Komaroff L, Oster-Granite ML, Neve RL. Neurotoxicity of a fragment of the amyloid precursor associated with Alzheimer's disease. *Science*. 1989; 245:417–420. [PubMed: 2474201]
- Yankner BA, Lu T. Amyloid beta-protein toxicity and the pathogenesis of Alzheimer disease. *J Biol Chem*. 2009; 284:4755–4759. [PubMed: 18957434]
- Yoshikawa K, Aizawa T, Hayashi Y. Degeneration in vitro of post-mitotic neurons overexpressing the Alzheimer amyloid protein precursor. *Nature*. 1992; 359:64–67. [PubMed: 1301020]
- Young-Pearse TL, Bai J, Chang R, Zheng JB, LoTurco JJ, Selkoe DJ. A critical function for beta-amyloid precursor protein in neuronal migration revealed by in utero RNA interference. *J Neurosci*. 2007; 27:14459–14469. [PubMed: 18160654]
- Young-Pearse TL, Chen AC, Chang R, Marquez C, Selkoe DJ. Secreted APP regulates the function of full-length APP in neurite outgrowth through interaction with integrin beta1. *Neural Dev*. 2008; 3:15. [PubMed: 18573216]
- Zhang J, Gorostiza O, Tang H, Bredesen DE, Galvan V. Reversal of learning deficits in hAPP transgenic mice carrying a mutation at Asp664: a role for early experience. *Behav Brain Res*. 2009; 206:202–207. [PubMed: 19751769]
- Zhang YW, Wang R, Liu Q, Zhang H, Liao FF, Xu H. Presenilin/gamma-secretase-dependent processing of beta-amyloid precursor protein regulates EGF receptor expression. *Proc Natl Acad Sci USA*. 2007; 104:10613–10618. [PubMed: 17556541]
- Zhao J, Fu Y, Yasvoina M, Shao P, Hitt B, O'Connor T, Logan S, Maus E, Citron M, Berry R, Binder L, Vassar R. Beta-site amyloid precursor protein cleaving enzyme 1 levels become elevated in neurons around amyloid plaques: implications for Alzheimer's disease pathogenesis. *J Neurosci*. 2007; 27:3639–3649. [PubMed: 17409228]
- Zhao L, Ma QL, Calon F, Harris-White ME, Yang F, Lim GP, Morihara T, Ubeda OJ, Ambegaokar S, Hansen JE, Weisbart RH, Teter B, Frautschy SA, Cole GM. Role of p21-activated kinase pathway defects in the cognitive deficits of Alzheimer disease. *Nat Neurosci*. 2006; 9:234–242. [PubMed: 16415866]
- Zhao WQ, De Felice FG, Fernandez S, Chen H, Lambert MP, Quon MJ, Krafft GA, Klein WL. Amyloid beta oligomers induce impairment of neuronal insulin receptors. *FASEB J*. 2008; 22:246–260. [PubMed: 17720802]
- Zheng H, Jiang M, Trumbauer ME, Sirinathsinghi DJ, Hopkins R, Smith DW, Heavens RP, Dawson GR, Boyce S, Conner MW, Stevens KA, Slunt HH, Sisoda SS, Chen HY, Van der Ploeg LH. Beta-amyloid precursor protein deficient mice show reactive gliosis and decreased locomotor activity. *Cell*. 1995; 81:525–531. [PubMed: 7758106]

Fig. 1.

Scheme depicting the domain structure of amyloid β precursor protein (APP). The E1 domain consisting of the growth factor like domain (GFLD) and the copper-binding domain (CuBD). The acidic domain (AcD) and the Kunitz-type inhibitory domain (Kunitz protease inhibitor [KPI], not present in neurons) bridge the E1 and E2 domains. The E2 domain consists of 2 coiled-coil substructures connected through a continuous helix. The amino acid sequence of amyloid β (A β)/transmembrane/intracellular (intracytoplasmic domain of APP [AICD]) domains is shown in the lower part of the figure. Asterisks denote amino acids substituted in familiar forms of AD and APP variants. Arrows indicate cleavage sites for both secretases and caspases. Binding motifs for heterotrimeric Go/s proteins (Go/s-BM) and scaffolding proteins (SP-BM; e.g., Fe65, Mint/X11-family proteins, Dab1, c-Jun N-terminal kinase) are underlined. The A β sequence is also underlined.

Fig. 2.

Pathways for metabolic processing of amyloid β precursor protein (APP). The 2 most prominent metabolic routes and APP proteolytic derivatives are illustrated. APP is substrate for either α - or β -secretase. Therefore, the nonamyloidogenic and the amyloidogenic pathways are mutually exclusive. Whereas several metalloproteinases display α -secretase activity (reviewed by Allison et al., 2003), β -site APP-cleaving enzyme 1 (BACE1) is the only known aspartylprotease with β -secretase activity (reviewed by Kandalepas and Vassar, 2012). Membrane-tethered APP stubs, C83 and C99 (also known as α - and β -carboxy-terminal fragment), are substrates for γ -secretase, a protein complex that includes at least 4 different proteins: anterior pharynx defective 1, nicastrin, presenilin 1 or presenilin 2, and presenilin enhancer-2 (reviewed by Wolfe, 2012).

Fig. 3.

Scheme representing physiological and maladaptive regulation of amyloid β precursor protein (APP) signaling and processing. In physiological conditions, the formation of *cis* and *trans* homo- and heterodimers of APP is modulated by interactions with transmembrane proteins (e.g., integrins, APLP1), extracellular secreted proteins (e.g., sAPP), or matrix components (e.g., heparan sulfate proteoglycans). These events regulate metabolic processing of APP (preferentially through the nonamyloidogenic pathway) and trigger signaling cascades that functionally modulate synaptic and neuritic plasticity. Binding of amyloid β (A β) fibrils to APP abrogates the interaction of APP with physiological ligands and promotes aberrant APP multimerization. These events shift the metabolic processing of APP to the amyloidogenic pathway, spreading A β deposition, and trigger dysfunctional signaling cascades leading to synaptic and neuritic degeneration.

Table 1

Molecular targets reported for various A β assemblies

Molecular target	A β assembly	Pathologic effect	Reference
p75NTR	Oligomers	Neuronal degeneration Neuritic dystrophy	Yaar et al. (1997) Perini et al. (2002) and Sothibundhu et al. (2008) Knowles et al. (2009)
NR1 (NMDA receptor)	Oligomers	Synaptic toxicity	Lacor et al. (2004)
		Oxidative stress	De Felice et al. (2007)
NR1/NR2B (NMDA receptor)	Oligomers	Synaptic toxicity	Lacor et al. (2007)
mGluR5 (metabotropic glutamate receptor)	Oligomers	Synaptic toxicity	Renner et al. (2010)
		Altered Ca ²⁺ signaling	
EphB2	Oligomers	Synaptic toxicity	Cissé et al. (2011)
$\alpha 7$ nAChRs (acetylcholine receptor)	Oligomers	Synaptic toxicity	Wang et al. (2000)
			Dineley et al. (2001)
			Nagele et al. (2002)
			Fodero et al. (2004)
			Snyder et al. (2005)
			Yan et al. (1996)
RAGE	Fibrils	Oxidative stress	Yan et al. (1997)
	A β from AD brain	Activation of microglia and astrocytes	
	Oligomers	Inflammation	Origlia et al. (2010)
		Synaptic toxicity	Fang et al. (2010)
		Oxidative stress	El Khoury et al. (1996)
Class A scavenger receptors	Fibrils	Activation of microglia	Paresce et al. (1996)
			Paresce et al. (1997)
CD36 (class B scavenger receptor) and CD47	Fibrils	Oxidative stress	Bamberger et al. (2003)
		Inflammation	Coraci et al. (2002)
		Activation of microglia and monocytes	Moore et al. (2002)
Heparan sulfate proteoglycans	Fibrils	Nucleation and A β aggregation Inhibition of A β degradation	Narindrasarak et al. (1991) and McLaurin et al. (1999)
		A β uptake in neurons	Gupta-Bansal et al. (1995)
Gangliosides	Fibrils and oligomers	Nucleation and A β aggregation	Kanekiyo et al. (2011)
		Membrane disruption and neurotoxicity	McLaurin and Chakrabarty (1996) Williams et al. (2011)

Molecular target	A β assembly	Pathologic effect	Reference
PrPc	Oligomers	Synaptic toxicity	Choo-Smith et al. (1997)
			Kakio et al. (2001)
			Laurén et al. (2009)
			Gimbel et al. (2010)
			Barry et al. (2011)
			Kessels et al. (2010)
Neurotrophin-1	Oligomers	Nucleation and A β aggregation Synaptic toxicity	Cissé et al. (2011)
			Bate and Williams (2011)
			Um et al. (2012)
			Dinamarca et al. (2011)
			Inestrosa et al. (1996)
			Alvarez et al. (1997)
Acetylcholinesterase	Fibrils	Nucleation and A β aggregation Synaptic toxicity	Alvarez et al. (1998)
			Koenigsnecht and Landreth (2004)
			Wright et al. (2007)
			Wang et al. (2008)
			Lorenzo et al. (2000)
			Van Nostrand et al. (2002)
α 6 β 1, α 2 β 1, and α V β 1 integrins	Fibrils and oligomers	A β uptake in microglia Neuronal toxicity Synaptic toxicity Neurotoxicity	Shaked et al. (2006)
			Galvan et al. (2006)
			Sola Vigo et al. (2009)
			Kedikian et al. (2010)
			Magdesian et al. (2008)
			Zhao et al. (2008)
APP	Fibrils	Inhibition of Wnt signaling pathway Synaptic toxicity	De Felice et al. (2009)
			Yan et al. (1997); Lustbader et al. (2004)
			Du et al. (2008)
			Maniczak and Reddy (2012)
Frizzled (Fz5-CRD)	Oligomers	Inhibition of Wnt signaling pathway	
Insulin receptor (IR)	Oligomers	Synaptic toxicity	
ABAD	Intracellular A β	Mitochondrial damage	
Cyclophilin D	Intracellular A β	Mitochondrial damage	
VDAC1	Oligomers	Mitochondrial damage	

Key: A β , amyloid β ; AD, Alzheimer's disease; APP, amyloid β precursor protein; NMDA, N-methyl-D-aspartate.