Published in final edited form as:

Pestic Biochem Physiol. 2013 July 1; 106(3): 93–100. doi:10.1016/j.pestbp.2013.02.007.

Diversity and Convergence of Sodium Channel Mutations Involved in Resistance to Pyrethroids

Frank D. Rinkevich, Yuzhe Du, and Ke Dong

Department of Entomology, Genetics and Neuroscience Programs, Michigan State University, East Lansing, MI 48824-1115 USA

Abstract

Pyrethroid insecticides target voltage-gated sodium channels, which are critical for electrical signaling in the nervous system. The intensive use of pyrethroids in controlling arthropod pests and disease vectors has led to many instances of pyrethroid resistance around the globe. In the past two decades, studies have identified a large number of sodium channel mutations that are associated with resistance to pyrethroids. The purpose of this review is to summarize both common and unique sodium channel mutations that have been identified in arthropod pests of importance to agriculture or human health. Identification of these mutations provides valuable molecular markers for resistance monitoring in the field and helped the discovery of the elusive pyrethroid receptor site(s) on the sodium channel.

INTRODUCTION

Pyrethroids are a major class of synthetic insecticide widely used for controlling arthropod pests and disease vectors because of their fast acting, high insecticidal activities and low mammalian toxicity. However, the intensive use of pyrethroids has led to many instances of pest resistance around the globe. The primary target sites of pyrethroids are voltage-gated sodium channels [1; 2; 3]. One of major mechanisms of pyrethroid resistance, which reduces neuronal sensitivity to this class of insecticides, is known as knockdown resistance (*kdr*) [4]. The *kdr* trait was first documented in the house fly and was eventually mapped to a sodium channel locus in house flies where a single nucleotide polymorphism resulted in a substitution of leucine for phenylalanine at position 1014 (L1014F) [5; 6; 7; 8; 9]. A second mutation of M918T in conjunction with L1014F (M918T+ L1014F) is the genotype that leads to higher levels of resistance to pyrethroids, the so called *super-kdr* phenotype [7; 8]. The *kdr* mutation has been documented globally in many major arthropod pests and disease vectors and it is well-established now that mutations in the sodium channel are responsible for *kdr* [10; 11; 12]. Identification of *kdr* mutations has already led to successful development of rapid and accurate molecular methods to detect *kdr*-based pyrethroid resistance in field populations [13; 14; 15; 16]. Several comprehensive reviews on insect sodium channels and *kdr* have summarized pyrethroid resistance-associated sodium channel mutations identified in pyrethroid-resistant populations of pest species [10; 11; 12; 17]. However, since these reviews published, more new mutations associated with pyrethroid

© 2013 Elsevier Inc. All rights reserved.

Address Correspondence to: Dr. Ke Dong, Department of Entomology, 438 Giltner Hall, Michigan State University, East Lansing, MI 48824-1115, USA, Phone: (517) 432-2034, Fax: (517) 353-4354, dongk@msu.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

resistance have been identified in sodium channels from various arthropod pests, particularly in disease vectors, highlighting the complexity of the interactions between pyrethroids and insect sodium channels at the molecular level. The intention of this review is to update the inventory of mutations associated with pyrethroid resistance and discuss both their diversity and convergence among diverse arthropod pest species. In addition, we will examine the relationship of these *kdr* mutations to the pyrethroid receptor(s) on insect sodium channels.

VOLTAGE-GATED SODIUM CHANNELS

Voltage-gated sodium channels are essential for the initiation and propagation of action potentials in the nervous system and other excitable cells. Our current knowledge of voltage-gated sodium channels originates mainly from molecular and functional analyses of mammalian sodium channels [18]. Mammalian sodium channels are composed of one pore-forming α -subunit of about 260 kDa and up to four much smaller auxiliary β -subunits of about 30–40 kDa. The α -subunit contains four homologous repeats (I–IV), each having six transmembrane segments (S1–S6). The S1–S4 segments in each repeat function as the voltage-sensing domains, whereas the S5 and S6 segment, and the re-entrant loops (called the P-region) connecting the S5 and S6 segments compose the pore-forming domains. The voltage-sensing domain is linked to the pore-forming domain by a small intracellular linker connecting the S4 and S5 segments.

The opening and closing of sodium channels are voltage-gated. In response to membrane depolarization, the S4 segments (the voltage sensors; rich in positively charged residues) move outward, initiating the voltage-dependent activation, which results in the opening of the activation gate (presumably formed by the intracellular end of the S6 segments). A few milliseconds after the channel opening, the sodium channel is inactivated (i.e., closed). This inactivation process is mediated by an inactivation particle formed by residues (IFM in mammals, MFM in insects) in the linker between domains III and IV, which blocks the inner pore of the sodium channel. Upon repolarization, the S4 voltage sensors move backward causing the closing of the activation gate, which is known as channel deactivation.

Unlike mammals which have at least nine sodium channel genes [19], insects have only one functional sodium channel gene [20]. However, insect sodium channel transcripts undergo extensive alternative splicing and RNA editing to produce functionally and pharmacologically distinct sodium channel variants [11; 20]. The first sodium channel gene, *para*, was identified in *Drosophila melanogaster* [21]. Due to the involvement of sodium channels in pyrethroid resistance in various arthropod pest populations, *para*-orthologs have been identified from many insect and arachnid pest species [22]. To date, sodium channels from *Drosophila melanogaster*, *Musca domestica*, and *Blattella germanica*, [22], as well as one arachnid sodium channel from *Varroa destructor* [23], have been functionally expressed in *Xenopus* oocytes. Successful expression of these sodium channels *in vitro* allows us to functionally examine pyrethroid resistance-associated mutations.

MODE OF ACTION OF PYRETHROIDS

As discussed above, sodium channels undergo activation (opening) followed by inactivation and deactivation (closing). The gating transitions between the closed and open states are intricately linked to the generation and propagation of electrical impulses (i.e., action potentials). Pyrethroids modify the gating transitions by inhibiting deactivation and inactivation, resulting in prolonged channel opening [1; 2; 3]. At the cellular level, pyrethroids disrupt nerve function causing repetitive discharges, membrane depolarization, and synaptic disturbances [1; 2; 3]. Studies of insect sodium channels expressed in *Xenopus* oocytes show that pyrethroids, particularly type II pyrethroids, preferably bind to the activated (open) state of insect sodium channels and cause the prolonged opening of sodium

channels, evident as a pyrethroid-induced tail current associated with repolarization in voltage-clamp experiments [24; 25; 26; 27] provided the initial electrophysiological evidence for pyrethroid trapping sodium channels in the open state in crayfish giant axons. However, very little is known on how pyrethroids trap sodium channels in the open state at the molecular level.

Earlier electrophysiological and pharmacological studies suggest that pyrethroids have a distinct receptor site on the sodium channel [28; 29; 30; 31; 32]. Attempts to detect specific pyrethroid binding to the sodium channels in insect nerve tissues failed, mainly because the high lipophilicity of pyrethroids resulted in extremely high levels of nonspecific binding to membranes and filters [33; 34; 35]. A high-affinity pyrethroid-binding site has been reported on rat brain sodium channel preparations [36], however, the molecular determinants of the pyrethroid-binding site on the sodium channel have remained elusive until recently with the identification of *kdr* mutations and computer modeling of insect sodium channels (see below).

COMMON AND UNIQUE RESISTANCE-ASSOCIATED MUTATIONS

Pyrethroid resistance-associated mutations were identified by comparing partial or complete coding sequences of the *para*-orthologous sodium channel genes from pyrethroid-sensitive and pyrethroid-resistant strains of various arthropod species. With the increased use of pyrethroids in controlling arthropod pests and disease vectors, more pyrethroid-resistant populations have been documented [37]. This trend, coupled with the recent increase in the accessibility and affordability of molecular tools has had a tremendous influence on the identification of a wide array of mutations associated with pyrethroid resistance in various species. These unique amino acid substitutions are found throughout the sodium channel protein (Fig. 1 and 2).

More than 30 unique resistance-associated mutations or combinations of mutations have been detected in more than one species (Table 1 and Fig. 1). In contrast, Table 2 and Figure 2 summarize resistance-associated mutations that have been detected in only a single species. Most of these mutations in Table 1 have been shown to reduce the pyrethroid sensitivity of house fly, cockroach and/or *Drosophila* sodium channels expressed in *Xenopus* oocytes confirming their role in *kdr* (Table 3). However, the mutations in Table 2 and Figure 2 are mostly functionally uncharacterized.

Intriguingly, the majority of functionally confirmed mutations are found in IIS5, IIS6, and IIIS6 segments (Fig. 1 and Table 3). Computer modeling (see below) predicts that IIS5, IIIS6 and the linker connecting S4 and S5 in domain II compose a pyrethroid binding site and most of the *kdr* mutations in these regions (such as M918T and L925I) likely confer resistance by reducing binding of pyrethroids to sodium channels [38]. Subsequent studies from systematic site-directed mutagenesis of residues in the linker connecting S4 and S5 in domain II, IIS5 and IIIS6 uncovered more pyrethroid-sensing residues in these regions, supporting this model [39; 40]. However, according to this model, many other *kdr* mutations including the L1014F mutation in IIS6 that has been detected in many species is not close to this receptor site (further discussion below).

The functional impact of mutations identified in insect species other than the house fly, cockroach or *Drosophila* has been assessed by introducing those mutations into one of the three sodium channels by site-directed mutagenesis and then characterizing the mutated channels in *Xenopus* oocytes. For example, the *H. virescens* V410M mutation in IS6 reduced the pyrethroid sensitivity by 10-fold when introduced into *Drosophila*, house fly and cockroach sodium channels [27; 41; 42], the T929I mutation in IIS5, identified in diamondback moth and other pest species, drastically reduced the Para channel sensitivity to

deltamethrin [43], and the F1538I mutation in IIS6 completely abolishes the pyrethroid sensitivity of the cockroach sodium channel to structurally diverse pyrethroids [44].

The L1014F mutation in IIS6 was the first resistance-associated mutation that was detected and confirmed to be the cause of *kdr* [8; 25; 26; 45]. Since the first reports from the house fly and the German cockroach [7; 8; 9; 46], pyrethroid resistance has been attributed to substitution of F, C, H, S, or W at this position in other insect species across evolutionarily divergent insect groups (Table 1). Variability (e.g. L1014F/C/H/S/W) in substitution at a single site resulting in pyrethroid resistance is not unique for L1014. Divergent substitutions have been shown at other sites, including V410 (M/A/G/L) in IS6, M918 (T/L/V) in the linker connecting S4 and S5 in domain II, and T929 (I/C/N/V) in IIS5 (in each of these three cases only the first amino acid substitution has been functionally confirmed to cause a reduction in pyrethroid sensitivity). This massively parallel and divergent evolution of resistance demonstrates not only are these sites critical for pyrethroid action, but also raises the possibility that the substituting amino acid at these sites can potentially vary based on the type of pyrethroid used to select the resistant population. For example, sodium channels with the L1014F, L1014H, and L1014S mutations provide variable levels of protection to Type I or Type II pyrethroids or DDT [47]. Similarly, M918T in the linker connecting S4 and S5 in domain II provides extremely high levels of protection against permethrin and deltamethrin [26], but does not provide protection from DDT [48], and F1534C confers channel resistance to type I, but not type II pyrethroids [49]. It will be of great interest to determine if compound-specificity is evident with other sites of divergent substitution that have not yet been functionally characterized.

Co-occurrence of more than one resistance-associated mutation often more drastically reduces the channel sensitivity to pyrethroids than individual mutations alone. For example, the L1014F mutation or the M918T mutation alone caused about 5–10 fold reduction in the sensitivity of the Para channel to deltamethrin, but the double mutations almost abolished the sensitivity of the Para channel to deltamethrin [26; 50]. Similarly, the T929I mutation either alone or in combination with M827I and L932F completely eliminated permethrin sensitivity in Vssc1 channels [51]. Two mutations, E435K and C785R, in the linker connecting domains I and II were found to co-exist with the L1014F only in the German cockroach. Each mutation alone did not reduce the sensitivity of the cockroach sodium channel to deltamethrin [25]. However, when either the E435K or C785R mutation was combined with the L1014F mutation, the channel sensitivity was reduced by 100-fold [25]. Concomitant presence of all three mutations reduced channel sensitivity to deltamethrin by 500-fold [25]. Similarly, E435K and C785R mutations also further reduced the sensitivity of the V410M channel to pyrethroids [41]. These two mutations are considered as enhancers of the V410M and L1014F mutations [25; 41].

DO KDR MUTATIONS DEFINE THE PYRETHROID RECEPTOR SITE(S)?

Relying on information of key *kdr* mutations, O'Reilly et al. [38] used the X-ray structure of the K_v1.2 potassium channel as a template to predict the open conformation of the house fly sodium channel. This interesting model predicts that the pyrethroid receptor site is located in a hydrophobic cavity delimited by the IIS4–S5 linker and IIS5 and IIS6 helices. Subsequent systematic site-directed mutagenesis of these regions uncovered more pyrethroid-sensing residues in these segments, providing further experimental support for this model [39; 40]. Electrophysiological studies also showed that *kdr* mutations in the IIS4–S5 linker and IIS5 and IIS6 helices likely reduce pyrethroid binding to the receptor site on the sodium channel [26; 43; 44; 52]. How this mutation reduces sodium channel sensitivity to pyrethroids remains elusive. The L1014F mutation has been shown to increase close-state inactivation which could reduce the availability of a *Drosophila* sodium channel in the open state for

pyrethroid action [52; 53]. In the same study, Hill plot analysis showed that the house fly M918T mutation in the linker connecting IIS4 and IIS5 (part of the first receptor site) reduces the number of pyrethroid-binding sites per channel from two to one in the *Drosophila* sodium channel [52]. This led to another possibility that L1014 in IIS6 could be at a second pyrethroid receptor site; and the L1014F mutation confers resistance by reducing pyrethroid binding. This hypothesis is supported by results from specialized analysis of pyrethroid binding utilizing the competitive binding of active and inactive isomers of permethrin, called Schild analysis. Schild analysis shows that the L1014F mutation (i.e., L993F in the cockroach sodium channel) reduces pyrethroid binding to the cockroach sodium channel [44]. Future mutational analysis coupled with computer modeling is necessary to identify any possible new pyrethroid receptor site on insect sodium channels.

CONCLUSIONS

Investigations into the molecular mechanism of pyrethroid resistance due to mutations in the voltage-gated sodium channel has yielded a tremendous amount of information that has far-reaching implications in both basic and applied aspects of research. Identification of mutations associated with pyrethroid resistance provides precise molecular markers for rapidly assessing the frequency of resistance alleles in field populations. The diversity in amino acid positions that harbor mutations and the divergent mutations found in those positions likely reflect the evolutionary plasticity of pyrethroid resistance manifested in the field. This plasticity is reinforced by the observation from functional analyses that these mutations provide varying levels of protection against different pyrethroids, implying that the diversity of resistance mutations may also be driven in part by the specific pyrethroid used to select for resistance. In addition to deepening our understanding of the pyrethroid resistance mechanism, information accumulated from the functional characterization of pyrethroid resistance mutations in *Xenopus* oocytes and the elucidation of the binding and action of pyrethroids at the molecular and atomic levels has also contributed significantly to the general knowledge of the gating mechanisms and toxin pharmacology of the sodium channel.

While research on *kdr* mutations has yielded many answers, many important questions still remain. Will new mutations emerge as pyrethroids continue to be a major component of many pest control programs? What are the physiological and evolutionary reasons that restrict the number and frequency of resistance alleles in the field? While it is easy to understand why *kdr* mutations that are located in the predicted pyrethroid receptor site would confer pyrethroid resistance, how do the mutations that are not located at the receptor site confer pyrethroid resistance? Are there additional pyrethroid receptor sites on the sodium channel? Can altered channel gating (e.g., activation, deactivation and/or inactivation) by *kdr* mutations confer pyrethroid resistance by counteracting the action of pyrethroids? One can only hope that the next decade of research in this field will be as productive as it has been for the past two decades and provide answers to these remaining questions.

Acknowledgments

The authors thank Dr. Kris Silver for critical review of this manuscript; and thank past and current lab members for their contributions to the research in the authors' laboratory. The research is supported by NIH (GM57440 and GM80255), NSF (IBN 9696092 and IBN 9808156), USDA-NRI (35607-14866) and BARD (IS-3480-03)

LITERATURE CITED

1. Narahashi T. Neuronal ion channels as the target sites of insecticides. *Pharmacol Toxicol.* 1996; 78:1–14. [PubMed: 8821391]

2. Narahashi T. Nerve membrane ion channels as the target site of insecticides. *Mini Rev Medic Chem*. 2002; 2:419–432.
3. Soderlund DM. Molecular mechanisms of pyrethroid insecticide neurotoxicity: recent advances. *Arch Toxicol*. 2012; 86:165–181. [PubMed: 21710279]
4. Soderlund, DM.; Bloomquist, JR. Molecular Mechanisms of Insecticide Resistance. In: Roush, RT.; Tabashnik, BE., editors. *Pesticide Resistance in Arthropods*. Chapman and Hall; New York and London: 1990. p. 58
5. Williamson MS, Denholm I, Bell CA, Devonshire AL. Knockdown resistance (*kdr*) to DDT and pyrethroid insecticides maps to a sodium channel gene locus in the housefly (*Musca domestica*). *Molec Gen Genet*. 1993; 240:17–22. [PubMed: 8101963]
6. Knipple DC, Doyle KE, Marsella-Herrick PA, Soderlund DM. Tight genetic linkage between the *kdr* insecticide resistance trait and a voltage-sensitive sodium channel gene in the house fly. *Proc Natl Acad Sci*. 1994; 91:2483–2487. [PubMed: 8146143]
7. Ingles PJ, Adams PM, Knipple DC, Soderlund DM. Characterization of voltage-sensitive sodium channel gene coding sequences from insecticide-susceptible and knockdown-resistant house fly strains. *Insect Biochem Molec Biol*. 1996; 26:319–326. [PubMed: 8814780]
8. Williamson M, Martinez-Torres D, Hick C, Devonshire A. Identification of mutations in the housefly *para*-type sodium channel gene associated with knockdown resistance (*kdr*) to pyrethroid insecticides. *Molec Gen Genet*. 1996; 252:51–60. [PubMed: 8804403]
9. Miyazaki M, Ohyama K, Dunlap D, Matsumura F. Cloning and sequencing of the *para*-type sodium channel gene from susceptible and *kdr*-resistant German cockroaches (*Blattella germanica*) and house fly (*Musca domestica*). *Mol Gen Genet*. 1996; 252:61–68. [PubMed: 8804404]
10. Soderlund, DM. Sodium channels. In: Gilbert, LI.; Iatrou, K.; Gill, SS., editors. *Comprehensive molecular insect science*. Elsevier; Amsterdam: 2005. p. 1-24.
11. Dong K. Insect sodium channels and insecticide resistance. *Invert Neurosci*. 2007; 7:17–30. [PubMed: 17206406]
12. Davies TGE, Field LM, Usherwood PNR, Williamson MS. DDT, pyrethrins, pyrethroids and insect sodium channels. *IUBMB Life*. 2007; 59:151–162. [PubMed: 17487686]
13. Hodgdon HE, Yoon KS, Previte D, Kim HJ, Aboelghar GE, Lee SH, Clark JM. Determination of knockdown resistance allele frequencies in global human head louse populations using the serial invasive signal amplification reaction. *Pest Manag Sci*. 2010; 66:1031–1040. [PubMed: 20564731]
14. Nyoni BN, Gorman K, Mzilahowa T, Williamson MS, Navajas M, Field LM, Bass C. Pyrethroid resistance in the tomato red spider mite, *Tetranychus evansi*, is associated with mutation of the *para*-type sodium channel. *Pest Manag Sci*. 2011; 67:891–897. [PubMed: 21432985]
15. Yanola J, Somboon P, Walton C, Nachaiwieng W, Somwang P, Prapanthadara L. High-throughput assays for detection of the F1534C mutation in the voltage-gated sodium channel gene in permethrin-resistant *Aedes aegypti* and the distribution of this mutation throughout Thailand. *Tropical Medicine & International Health*. 2011; 16:501–509. [PubMed: 21342372]
16. Kulkarni MA, Rowland M, Alifrangis M, Mosha FW, Matowo J, Malima R, Peter J, Kweka E, Lyimo I, Magesa S. Occurrence of the leucine-to-phenylalanine knockdown resistance (*kdr*) mutation in *Anopheles arabiensis* populations in Tanzania, detected by a simplified high-throughput SSOP-ELISA method. *Malaria Journal*. 2006; 5:56. [PubMed: 16820067]
17. Soderlund DM, Knipple DC. The molecular biology of knockdown resistance to pyrethroid insecticides. *Insect Biochem Molec Biology*. 2003; 33:563–577.
18. Catterall WA. From ionic currents to molecular mechanisms: the structure and function of voltage-gated sodium channels. *Neuron*. 2000; 26:13–25. [PubMed: 10798388]
19. Goldin AL. Evolution of voltage-gated Na⁺ channels. *Journal of experimental biology*. 2002; 205:575–584. [PubMed: 11907047]
20. Dong, K. Progress in insect sodium channel research. In: Gilbert, LI.; Gill, SS., editors. *Insect Pharmacology, Channels, Receptors, Toxins and Enzymes*. Academic Press; 2010. p. 25-27.
21. Loughney K, Kreber R, Ganetzky B. Molecular analysis of the *para* locus, a sodium channel gene in *Drosophila* Cell. 1989; 58:1143–1154.
22. Soderlund DM. Pyrethroids, knockdown resistance and sodium channels. *Pest Man Sci*. 2008; 64:610–616.

23. Du Y, Nomura Y, Liu Z, Huang Z, Dong K. Functional expression of an arachnid sodium channel reveals residues responsible for tetrodotoxin resistance in invertebrate sodium channels. *J Biol Chem*. 2009; 284:33869–33875. [PubMed: 19828457]
24. Smith M, Goldin AL. Interaction between the sodium channel inactivation linker and domain III S4–S5. *Biophys J*. 1997; 73:1885–95. [PubMed: 9336184]
25. Tan J, Liu Z, Tsai TD, Valles SM, Goldin AL, Dong K. Novel sodium channel gene mutations in *Blattella germanica* reduce the sensitivity of expressed channels to deltamethrin. *Insect Biochem Molec Biol*. 2002; 32:445–454. [PubMed: 11886779]
26. Vais H, Williamson MS, Goodson SJ, Devonshire AL, Warmke JW, Usherwood PNR, Cohen CJ. Activation of *Drosophila* sodium channels promotes modification by deltamethrin: Reductions in affinity caused by knock-down resistance mutations. *Journal of General Physiology*. 2000; 115:305–318. [PubMed: 10694259]
27. Zhao Y, Park Y, Adams ME. Functional and evolutionary consequences of pyrethroid resistance mutations in S6 transmembrane segments of a voltage-gated sodium channel. *Biochem Biophys Res Comm*. 2000; 278:516–521. [PubMed: 11095943]
28. Bloomquist JR, Soderlund DM. Pyrethroid insecticides and DDT modify alkaloid-dependent sodium channel activation and its enhancement by sea anemone toxin. *Molecular pharmacology*. 1988; 33:543–550. [PubMed: 2452970]
29. Brown GB, Gaupp JE, Olsen RW. Pyrethroid insecticides: stereospecific allosteric interaction with the batrachotoxinin-A benzoate binding site of mammalian voltage-sensitive sodium channels. *Molecular pharmacology*. 1988; 34:54–59. [PubMed: 2455860]
30. Jacques Y, Romey G, Cavey MT, Kartalovski B, Lazdunski M. Interaction of pyrethroids with the Na⁺ channel in mammalian neuronal cells in culture. *Biochimica et Biophysica Acta (BBA)-Biomembranes*. 1980; 600:882–897.
31. Lombet A, Mourre C, Lazdunski M. Interaction of insecticides of the pyrethroid family with specific binding sites on the voltage-dependent sodium channel from mammalian brain. *Brain Res*. 1988; 459:44–53. [PubMed: 2844361]
32. Takeda K, Narahashi T. Chemical modification of sodium channel inactivation: separate sites for the action of grayanotoxin and tetramethrin. *Brain Res*. 1988; 448:308–312. [PubMed: 2454144]
33. Rossignol DP. Reduction in number of nerve membrane sodium channels in pyrethroid resistant house flies. *Pestic Biochem Physiol*. 1988; 32:146–152.
34. Pauron D, Barhanin J, Amichot M, Pralavorio M, Berge JB, Lazdunski M. Pyrethroid Receptor in the Insect Na⁺ Channel: Alteration of Its Properties in Pyrethroid-Resistant Flies. *Biochem*. 1989; 28:1673–1377.
35. Dong K, Scott JG. Linkage of *kdr*-type resistance and the *para*-homologous sodium channel in German cockroaches (*Blattella germanica*). *Insect Biochem Molec Biol*. 1994; 24:647–654. [PubMed: 7915170]
36. Trainer VL, McPhee JC, Boutelet-Bochan H, Baker C, Scheuer T, Babin D, DeMoute JP, Guedin D, Catterall WA. High affinity binding of pyrethroids to the α subunit of brain sodium channels. *Molecular Pharmacology*. 1997; 51:651–657. [PubMed: 9106631]
37. Whalon, M.; Mota-Sanchez, D.; Hollingsworth, R. Global pesticide resistance in arthropods. CABI; Cambridge, MA: 2008.
38. O'Reilly AO, Khambay BPS, Williamson MS, Field LA, Wallace BA, Davies TGE. Modelling insecticide-binding sites in the voltage-gated sodium channel. *Biochem J*. 2006; 396:255–263. [PubMed: 16475981]
39. Usherwood PNR, Davies TGE, Mellor IR, O'Reilly AO, Peng F, Vais H, Khambay BPS, Field LM, Williamson MS. Mutations in DIIS5 and the DIIS4–S5 linker of *Drosophila melanogaster* sodium channel define binding domains for pyrethroids and DDT. *FEBS Letters*. 2007; 581:5485–5492. [PubMed: 17991435]
40. Du Y, Lee JE, Nomura Y, Zhang T, Zhorov B, Dong K. Identification of a cluster of residues in transmembrane segment 6 of domain III of the cockroach sodium channel essential for the action of pyrethroid insecticides. *Biochem J*. 2009; 419:377–385. [PubMed: 19154185]

41. Liu Z, Tan J, Valles SM, Dong K. Synergistic interaction between two cockroach sodium channel mutations and a tobacco budworm sodium channel mutation in reducing channel sensitivity to a pyrethroid insecticide. *Insect Biochem Molec Biol.* 2002; 32:397–404. [PubMed: 11886774]
42. Lee S, Soderlund DM. The V410M mutation associated with pyrethroid resistance in *Heliothis virescens* reduces the pyrethroid sensitivity of house fly sodium channels expressed in *Xenopus* oocytes. *Insect Biochem Mol Biol.* 2001; 31:19–29. [PubMed: 11102831]
43. Vais H, Williamson MS, Devonshire AL, Usherwood PNR. The molecular interactions of pyrethroid insecticides with insect and mammalian sodium channels. *Pest Manag Sci.* 2001; 57:877–888. [PubMed: 11695180]
44. Tan J, Liu Z, Wang R, Huang ZY, Chen AC, Gurevitz M, Dong K. Identification of amino acid residues in the insect sodium channel critical for pyrethroid binding. *Molec Pharmacol.* 2005; 67:513–522. [PubMed: 15525757]
45. Smith TJ, Lee SH, Ingles PJ, Knipple DC, Soderlund DM. The L1014F point mutation in the house fly *Vssc1* sodium channel confers knockdown resistance to pyrethroids. *Insect Biochem Molec Biol.* 1997; 27:807–812. [PubMed: 9474777]
46. Dong K. A single amino acid change in the para sodium channel protein is associated with knockdown-resistance (kdr) to pyrethroid insecticides in German cockroach. *Insect Biochem Molec Biol.* 1997; 27:93–100. [PubMed: 9066120]
47. Burton MJ, Mellor IR, Duce IR, Davies TGE, Field LM, Williamson MS. Differential resistance of insect sodium channels with kdr mutations to deltamethrin, permethrin and DDT. *Insect Biochem Mol Biol.* 2011; 41:723–732. [PubMed: 21640822]
48. Usherwood PNR, Vais H, Khambay BPS, Davies TGE, Williamson MS. Sensitivity of the *Drosophila para* sodium channel to DDT is not lowered by the *super-kdr* mutation M918T on the IIS4–S5 linker that profoundly reduces sensitivity to permethrin and deltamethrin. *FEBS Lett.* 2005; 579:6317–6325. [PubMed: 16263118]
49. Hu Z, Du Y, Nomura Y, Dong K. A sodium channel mutation identified in *Aedes aegypti* selectively reduces cockroach sodium channel sensitivity to type I, but not type II pyrethroids. *Insect Biochem Molec Biol.* 2011; 41:9–13. [PubMed: 20869441]
50. Lee SH, Smith TJ, Knipple DC, Soderlund DM. Mutations in the house fly *Vssc1* sodium channel gene associated with *super-kdr* resistance abolish the pyrethroid sensitivity of *Vssc1*/tipE sodium channels expressed in *Xenopus* oocytes. *Insect Biochem Molec Biol.* 1999; 29:185–194. [PubMed: 10196741]
51. Yoon KS, Symington SB, Lee SH, Soderlund DM, Clark JM. Three mutations identified in the voltage-sensitive sodium channel α -subunit gene of permethrin-resistant human head lice reduce the permethrin sensitivity of house fly *Vssc1* sodium channels expressed in *Xenopus* oocytes. *Insect Biochemistry and Molecular Biology.* 2008; 38:296–306. [PubMed: 18252244]
52. Vais H, Atkinson S, Pluteanu F, Goodson SJ, Devonshire AL, Williamson MS, Usherwood PNR. Mutations of the *para* sodium channel of *Drosophila melanogaster* identify putative binding sites for pyrethroids. *Mol Pharmacol.* 2003; 64:914–922. [PubMed: 14500748]
53. Davies T, O'Reilly A, Field L, Wallace B, Williamson M. Knockdown resistance to DDT and pyrethroids: from target-site mutations to molecular modeling. *Pest Manag Sci.* 2008; 64:1126–1130. [PubMed: 18561309]
54. Hopkins BW, Piertrantonio PV. The *Helicoverpa zea* (Boddie) (Lepidoptera: Noctuidae) voltage-gated sodium channel and mutations associated with pyrethroid resistance in field-collected adult males. *Insect Biochem Molec Biol.* 2010; 40:385–393. [PubMed: 20362057]
55. Yoon KS, Kwon DH, Strucharx JP, Hollingsworth CS, Lee SH, Clark JM. Biochemical and molecular analysis of deltamethrin resistance in the common bed bug (Hemiptera: Cimicidae). *J Med Entomol.* 2008; 45:1092–1101. [PubMed: 19058634]
56. Park Y, Taylor MFJ, Feyereisen R. A Valine421 to methionine mutation in IS6 of the *hscp* voltage-gated sodium channel associated with pyrethroid resistance in *Heliothis virescens* F. *Biochem Biophys Res Comm.* 1997; 239:688–691. [PubMed: 9367829]
57. Lee SH, Gao JR, Yoon KS, Mumcuoglu KY, Taplin D, Edman JD, Takano-Lee M, Clark JM. Sodium channel mutations associated with knockdown resistance in the human head louse, *Pediculus capitis* (De Geer). *Pestic Biochem Physiol.* 2003; 75:79–91.

58. Drali R, Benkouiten S, Badiaga S, Bitam I, Rolain JM, Brouqui P. Detection of a knockdown resistance mutation associated with permethrin resistance in the body louse *Pediculus humanus corporis* by use of melting curve analysis genotyping. *J Clin Microbiol.* 2012; 50:2229–2233. [PubMed: 22573588]
59. Sonoda S, Igaki C, Tsumuki H. Alternatively spliced sodium channel transcripts expressed in field strains of the diamondback moth. *Insect Biochem Molec Biol.* 2008; 38:883–890. [PubMed: 18692135]
60. Carletto J, Martin T, Vanlerberghe-Masutti F, Brevault T. Insecticide resistance traits differ among and within host races in *Aphis gossypii*. *Pest Manag Sci.* 2009; 66:301–307. [PubMed: 19908228]
61. Karatolos N, Gorman K, Williamson MS, Denholm I. Mutations in the sodium channel associated with pyrethroid resistance in the greenhouse whitefly, *Trialeurodes vaporariorum*. *Pest Manag Sci.* 2012; 68:834–838. [PubMed: 22228578]
62. Marshall KL, Moran C, Chen Y, Herron GA. Detection of *kdr* pyrethroid resistance in the cotton aphid, *Aphis gossypii* (Hemiptera: Aphididae), using a PCR-RFLP assay. *J Pestic Sci.* 2012; 37:169–172.
63. Guerrero FD, Jamroz RC, Kammlah D, Kunz SE. Toxicological and molecular characterization of pyrethroid-resistant horn flies, *Haematobia irritans*: Identification of *kdr* and *super-kdr* point mutations. *Insect Biochem Molec Biol.* 1997; 27:745–755. [PubMed: 9443375]
64. Eleftherianos I, Foster SP, Williamson MS, Denholm I. Characterization of the M918T sodium channel gene mutation associated with strong resistance to pyrethroid insecticides in the peach-potato aphid, *Myzus persicae* (Sulzer). *Bulletin of Entomological Research.* 2008; 98:183–191. [PubMed: 18076786]
65. Toda S, Morishita M. Identification of three point mutations on the sodium channel gene in pyrethroid-resistant *Thrips tabaci* (Thysanoptera: Thripidae). *J Econ Entomol.* 2009; 102:2296–2300. [PubMed: 20069860]
66. Haddi K, Berger M, Bielza P, Cifuentes D, Field LM, Gorman K, Rapisarda C, Williamson MS, Bass C. Identification of mutations associated with pyrethroid resistance in the voltage-gated sodium channel of the tomato leaf miner (*Tuta absoluta*). *Insect Biochem Molec Biol.* 2012; 42:506–513. [PubMed: 22504519]
67. Morin S, Williamson MS, Goodson SJ, Brown JK, Tabashnik BE, Dennehy TJ. Mutations in the *Bemisia tabaci para* sodium channel gene associated with resistance to a pyrethroid plus organophosphate mixture. *Insect Biochem Molec Biol.* 2002; 32:1781–1791. [PubMed: 12429130]
68. Morgan JAT, Corley SW, Jackson LA, Lew-Tabor AE, Moolhuijzen PM, Jonsson NN. Identification of a mutation in the *para*-sodium channel gene of the cattle tick *Rhipicephalus (Boophilus) microplus* associated with resistance to synthetic pyrethroid acaricides. *Intl J Parasitol.* 2009; 39:775–779.
69. Forcioli D, Frey B, Frey JE. High nucleotide diversity in the *para*-like voltage-sensitive sodium channel gene sequence in the western flower thrips (Thysanoptera: Thripidae). *J Econ Entomol.* 2002; 95:838–848. [PubMed: 12216829]
70. Bao WX, Sonoda S. Resistance to cypermethrin in melon thrips, *Thrips palmi*, (Thysanoptera: Thripidae), is conferred by reduced sensitivity of the sodium channel and CYP450-mediated detoxification. *Appl Entomol Zool.* 2012
71. Rinkevich FD, Su C, Lazo T, Hawthorne D, Tingey W, Naimov S, Scott JG. Multiple evolutionary origins of knockdown resistance (*kdr*) in pyrethroid-resistant Colorado potato beetle, *Leptinotarsa decemlineata*. *Pestic Biochem Physiol.* 2012; 104:192–200.
72. Araujo RA, Williamson MS, Bass C, Field LM, Duce IR. Pyrethroid resistance in *Sitophilus zeamais* is associated with a mutation (T929I) in the voltage-gated sodium channel. *Insect Molec Biol.* 2011; 20:437–445. [PubMed: 21496128]
73. Lee SH, Yoon KS, Williamson MS, Goodson SJ, Takano-Lee M, Edman JD, Devonshire AL, Clark JM. Molecular analysis of *kdr*-like resistance in permethrin-resistant strains of head lice. *Pediculus capitis Pestic Biochem Physiol.* 2000; 66:130–143.
74. Schuler TH, Martinez-Torres D, Thompson AJ, Denholm I, Devonshire AL, Duce IR, Williamson MS. Toxicological, electrophysiological, and molecular characterization of knockdown resistance

- to pyrethroid insecticides in the diamondback moth, *Plutella xylostella* (L.). Pestic Biochem Physiol. 1998; 59:169–182.
75. Roditakis E, Tsagkarakou A, Vontas J. Identification of mutations in the para sodium channel of *Bemisia tabaci* from Crete, associated with resistance to pyrethroids. Pestic Biochem Physiol. 2006; 85:161–166.
76. Bass C, Schroeder I, Turberg A, Field LM, Williamson MS. Identification of mutations associated with pyrethroid resistance in the *para-type* sodium channel of the cat flea, *Ctenocephalides felis*. Insect Biochem Molec Biol. 2004; 34:1305–1313. [PubMed: 15544944]
77. Brengues C, Hawkes NJ, Chandre F, McCarroll L, Duchon S, Guillet P, Manguin S, Morgan JC, Hemingway J. Pyrethroid and DDT cross-resistance in *Aedes aegypti* is correlated with novel mutations in the voltage-gated sodium channel gene. Medical and Veterinary Entomology. 2003; 17:87–94. [PubMed: 12680930]
78. Saavedra-Rodriguez K, Urdaneta-Marquez L, Rajatileka S, Moulton M, Flores AE, Fernandez-Salas I, Bisset J, Rodriguez M, McCall PJ, Donnelly MJ, Ranson H, Hemingway J, Black WCI. A mutation in the voltage-gated sodium channel gene associated with pyrethroid resistance in Latin American *Aedes aegypti*. Insect Molecular Biology. 2007; 16:785–798. [PubMed: 18093007]
79. Kim H, Baek JH, Lee WJ, Lee SH. Frequency detection of pyrethroid resistance allele in *Anopheles sinensis* populations by real-time PCR amplification of specific allele (rtPASA). Pestic Biochem Physiol. 2007; 87:54–61.
80. Wang ZM, Li CX, Xing D, Yu YH, Liu N, Xue RD, Dong YD, Zhao TY. Detection and widespread distribution of sodium channel alleles characteristic of insecticide resistance in *Culex pipiens* complex mosquitoes in China. Med Vet Entomol. 2012; 26:228–232. [PubMed: 22070231]
81. Martinez-Torres D, Chandre F, Williamson M, Darriet F, Berge J, Devonshire A, Guillet P, Pasteur N, Pauron D. Molecular characterization of pyrethroid knockdown resistance (*kdr*) in the major malaria vector *Anopheles gambiae* s.s. Insect Mol Biol. 1998; 7:179–184. [PubMed: 9535162]
82. Enayati AA, Vatandoost H, Ladonni H, Townson H, Hemingway J. Molecular evidence for a *kdr*-like pyrethroid resistance mechanism in the malaria vector mosquito *Anopheles stephensi*. Med Vet Entomol. 2003; 17
83. Karunaratne SHPP, Hawkes NJ, Perera MDB, Ranson H, Hemingway J. Mutated sodium channel genes and elevated monooxygenases are found in pyrethroid resistant populations of Sri Lankan malaria vectors. Pestic Biochem Physiol. 2007; 88:108–113.
84. Martinez-Torres D, Chevillon C, Brun-Barale A, Berge JB, Pasteur N, Pauron D. Voltage-dependant Na⁺ channels in pyrethroid resistant *Culex pipiens* mosquitoes. Pestic Sci. 1999; 55:1012–1020.
85. Chen L, Zhong D, Zhang D, Shi L, Zhou G, Gong M, Zhou H, Sun Y, Ma L, He J, Hong S, Zhou D, Xiong C, Chen C, Zou P, Zhu C, Yan G. Molecular ecology of pyrethroid knockdown resistance in *Culex pipiens pallens* mosquitoes. PLoS One. 2010; 5:e11681. [PubMed: 20657783]
86. Xu Q, Liu H, Zhang L, Liu N. Resistance in the mosquito, *Culex quinquefasciatus*, and possible mechanisms for resistance. Pest Manag Sci. 2005; 61:1096–1102. [PubMed: 16032654]
87. Brun-Barale A, Bouvier JC, Pauron D, Berge JB, Sauphanor B. Involvement of a sodium channel mutation in pyrethroid resistance in *Cydia pomonella* L., and development of a diagnostic test. Pest Manag Sci. 2005; 61:549–554. [PubMed: 15657957]
88. Rothwell JT, Morgan JAT, James FJ, Brown GW, Guerrero FD, Jorgensen WK. Mechanism of resistance to synthetic pyrethroids in buffalo flies in south-east Queensland. Aust Vet J. 2011; 89:70–72. [PubMed: 21323651]
89. Lee SH, Dunn JB, Clark JM, Soderlund DM. Molecular analysis of *kdr*-like resistance in a permethrin-resistant strain of Colorado potato beetle. Pestic Biochem Physiol. 1999; 63:63–75.
90. Nauen R, Zimmer CT, Andrews M, Slater R, Bass C, Ekbom B, Gustafsson G, Hansen LM, Kristensen M, Zebitz CPW, Williamson MS. Target-site resistance to pyrethroids in European populations of pollen beetle, *Meligethes aeneus* F. Pestic Biochem Physiol. 2012; 103:173–180.
91. Martinez-Torres D, Foster SP, Field LM, Devonshire AL, Williamson MS. A sodium channel point mutation is associated with resistance to DDT and pyrethroid insecticides in the peach-potato

- aphid, *Myzus persicae* (Sulzer) (Hemiptera: Aphididae). Insect Molec Biol. 1999; 8:339–346. [PubMed: 10469251]
92. Fabro J, Sterkel M, Capriotti N, Mougabure-Cueto G, Germano M, Rivera-Pomar R, Ons S. Identification of a point mutation associated with pyrethroid resistance in the *para*-type sodium channel of *Triatoma infestans*, a vector of Chagas' disease. Inf Genet Evol. 2012; 12:487–491.
93. Park Y, Taylor MFJ. A novel mutation L1029H in sodium channel gene *hscp* associated with pyrethroid resistance for *Heliothis virescens* (Lepidoptera: Noctuididae). Insect Biochem Molec Biol. 1997; 27:9–13. [PubMed: 9061924]
94. Liu N, Pridgeon JW. Metabolic detoxication and the *kdr* mutation in pyrethroid resistant house flies, *Musca domestica* (L.). Pestic Biochem Physiol. 2002; 73:157–163.
95. Rinkevich FD, Zhang L, Hamm RL, Brady SG, Lazzaro BP, Scott JG. Frequencies of the pyrethroid resistance alleles of *Vssc1* and *CYP6D1* in house flies from the eastern United States. Insect Molec Biol. 2006; 15:157–167. [PubMed: 16640726]
96. Olafson PU, Pitzer JB, Kaufman PE. Identification of a mutation associated with permethrin resistance in the *para*-type of sodium channel of the stable fly (Diptera: Muscidae). J Econ Entomol. 2011; 104:250–257. [PubMed: 21404865]
97. Stump AD, Atielli FK, Vulule JM, Besansky NJ. Dynamics of the pyrethroid knockdown resistance allele in western Kenyan populations of *Anopheles gambiae* in response to insecticide-treated bed net trials. Am J Trop Med Hyg. 2004; 70:591–596. [PubMed: 15210997]
98. Singh OP, Dykes CL, Das MK, Pradhan S, Bhatt RM, Agrawal OP, Adak T. Presence of two alternative *kdr*-like mutations, L1014F and L1014S, and a novel mutation V1010L, in the voltage-gated Na⁺ channel of *Anopheles culicifacies* from Orissa, India. Malaria J. 2010; 9:146.
99. Ranson H, Jensen B, Vulule JM, Wang X, Hemingway J, Collins FH. Identification of a point mutation in the voltage-gated sodium channel gene of Kenyan *Anopheles gambiae* associated with resistance to DDT and pyrethroids. Insect Mol Biol. 2000; 9:491–497. [PubMed: 11029667]
100. Verhaeghen K, van Bortel W, Trung HD, Sochantha T, Keokenchanh K, Coosemans M. Knockdown resistance in *Anopheles vagus*, *An. sinensis*, *An. paraliae* and *An. peditaeniatus* populations of the Mekong region. Parasit Vect. 2010; 3:59.
101. Luleyap HU, Alptekin D, Kasap H, Kasap M. Detection of knockdown resistance mutations in *Anopheles sacharovi* (Diptera: Culicidae) and genetic distance with *Anopheles gambiae* (Diptera: Culicidae) using cDNA sequencing of the voltage-gated sodium channel gene. J Med Entomol. 2002; 39:870–874. [PubMed: 12495185]
102. Tan WL, Li CX, Wang ZM, Liu MD, Dong YD, Feng XY, Wu ZM, Guo XX, Xing D, Zhang YM, Wang ZC, Zhao TY. First detection of multiple knockdown resistance (*kdr*)-like mutations in voltage-gated sodium channel using three new genotyping methods in *Anopheles sinensis* from Guangxi Province, China. J Med Entomol. 2012; 49:1012–1020. [PubMed: 23025181]
103. Pridgeon JW, Appel AG, Moar WJ, Liu N. Variability of resistance mechanisms in pyrethroid resistant German cockroaches (Dictyoptera: Blattellidae). Pestic Biochem Physiol. 2002; 73:149–156.
104. Endersby NM, Viduka K, Baxter SW, Saw J, Heckel DG, McKechnie SW. Widespread pyrethroid resistance in Australian diamondback moth, *Plutella xylostella* (L.), is related to multiple mutations in the para sodium channel gene. Bulletin of Entomological Research. 2011; 101:393–405. [PubMed: 21342603]
105. Kawada H, Higa Y, Komagata O, Kasai S, Tomita T, Yen NT, Loan LL, Sánchez RAP, Takagi M. Widespread distribution of a newly found point mutation in voltage-gated sodium channel in pyrethroid-resistant *Aedes aegypti* populations in Vietnam. PLoS Neglected Tropical Diseases. 2009; 3:e0000527.
106. Kasai S, Ng LC, Lam-Phua SG, Tang CS, Itokawa K, Komogata O, Kobayashi M, Tomita T. First detection of a putative knockdown resistance gene in major mosquito vector, *Aedes albopictus*. Jpn J Infect Dis. 2011; 64:217–221. [PubMed: 21617306]
107. He H, Chen AC, Davey RB, Ivie GW, George JE. Identification of a point mutation in the para-type sodium channel gene. Biochem Biophys Res Comm. 1999; 261:558–561. [PubMed: 10441465]

108. Feng YN, Zhao S, Sun W, Li M, Lu WC, He L. The sodium channel gene in *Tetranychus cinnabarinus* (Boisduval): identification and expression analysis of a mutation associated with pyrethroid resistance. *Pest Manag Sci.* 2011; 67:904–912. [PubMed: 21370394]
109. Tsagkarakou A, Van Leeuwen T, Khajehali J, Ilias A, Grispou M, Williamson MS, Tirry L, Vontas J. Identification of pyrethroid resistance associated mutations in the *para* sodium channel of the two-spotted spider mite *Tetranychus urticae* (Acari: Tetranychidae). *Insect Molec Biol.* 2009; 18:583–593. [PubMed: 19754737]
110. Head D, McCaffery A, Callaghan A. Novel mutations in the para-homologous sodium channel gene associated with phenotypic expression of nerve insensitivity resistance to pyrethroids in Heliothine lepidoptera. *Insect Molec Biol.* 1998; 7:191–196. [PubMed: 9535164]
111. Pittendrigh B, Reenan R, ffrench-Constant RH, Ganetzky B. Point mutations in the *Drosophila* sodium channel gene *para* associated with resistance to DDT and pyrethroid insecticides. *Mol Gen Genet.* 1997; 256:602–610. [PubMed: 9435785]
112. Liu Z, Valles SM, Dong K. Novel point mutations in the German cockroach para sodium channel gene are associated with knockdown resistance (kdr) to pyrethroid insecticides. *Insect Biochem Molec Biol.* 2000; 30:991–997. [PubMed: 10899465]
113. Sonoda S. Molecular analysis of pyrethroid resistance conferred by target insensitivity and increased metabolic detoxification in *Plutella xylostella*. *Pest Manag Sci.* 2010; 66:572–575. [PubMed: 20146259]
114. Jonsson NN, Cutulle C, Corley SW, Seddon JM. Identification of a mutation in the *para*-sodium channel gene of the cattle tick *Rhipicephalus microplus* associated with resistance to flumethrin but not to cypermethrin. *Intl J Parasitol.* 2010; 40:1659–1664.
115. Fallang A, Denholm I, Horsberg TE, Williamson MS. Novel point mutation in the sodium channel gene of pyrethroid-resistant sea lice *Lepeophtheirus salmonis* (Crustacea: Copepoda). *Dis Aquat Org.* 2005; 65:129–136. [PubMed: 16060266]
116. Cassanelli S, Cerchiari B, Giannini S, Bizzaro D, Mazzoni E, Manicardi GC. Use of the RFLP-PCR diagnostic test for characterizing MACE and *kdr* insecticide resistance in the peach potato aphid *Myzus persicae*. *Pest Manag Sci.* 2005; 61:91–96. [PubMed: 15593078]
117. Srisawat R, Komalamisra N, Eshita Y, Zheng M, Ono K, Itoh TQ, Matsumoto A, Petmitr S, Rongsriyam Y. Point mutations in domain II of the voltage-gated sodium channel gene in deltamethrin-resistant *Aedes aegypti* (Diptera: Culicidae). *Appl Entomol Zool.* 2010; 45:275–282.
118. Jones CM, Liyanapathirana M, Agossa FR, Weetman D, Ranson H, Donnelly MJ, Wilding CS. Footprints of positive selection associated with a mutation (N1575Y) in the voltage-gated sodium channel of *Anopheles gambiae*. *Proc Natl Acad Sci.* 2012; 109:6614–6619. [PubMed: 22493253]
119. Chang C, Shen WK, Wang TT, Lin YH, Hsu EL, Dai SM. A novel amino acid substitution in a voltage-gated sodium channel is associated with knockdown resistance to permethrin in *Aedes aegypti*. *Insect Biochem Molec Biol.* 2009; 39:272–278. [PubMed: 19171193]
120. Kwon DH, Clark JM, Lee SH. Cloning of a sodium channel gene and identification of mutations putatively associated with fenprothrin resistance in *Tetranychus urticae*. *Pestic Biochem Physiol.* 2010; 97:93–100.
121. Wang R, Liu Z, Dong K, Elzen PJ, Pettis J, Huang Z. Association of novel mutations in a sodium channel gene with fluvinate resistance in the mite, *Varroa destructor*. *J Apic Res.* 2002; 41:17–25.
122. Liu Z, Tan J, Huang Z, Dong K. Effect of a fluvalinate-resistance-associated sodium channel mutation from varroa mites on cockroach sodium channel sensitivity to fulvalinate, a pyrethroid insecticide. *Insect Biochem Molec Biol.* 2006; 36:885–889. [PubMed: 17046602]
123. Lee D, Park Y, Brown TM, Adams ME. Altered properties of neuronal sodium channels associated with genetic resistance to pyrethroids. *Molecular Pharmacology.* 1999; 55:584–593. [PubMed: 10051544]
124. Du Y, Khambay B, Dong K. An important role of a pyrethroid-sensing residue F1519 in the action of the N-alkylamide insecticide BTG 502 on the cockroach sodium channel. *Insect Biochem Molecular Bio.* 2011; 41:446–450.

Highlights

- Update the inventory of mutations associated with pyrethroid resistance
- Discuss both their diversity and convergence among diverse arthropod pest species.
- Summarize progress on the elucidation of the pyrethroid receptor(s) on insect sodium

Fig. 1.

Position of pyrethroid resistance-associated sodium channel mutations that are detected in more than one species. The mutations with solid circles have been confirmed to reduce sodium channel sensitivity to pyrethroids in *Xenopus* oocytes. The mutations with open circles have not been examined in *Xenopus* oocytes yet. The information on these mutations is presented in Table 1 and Table 3. The sodium channel protein contains four homologous repeats (I–IV), each having six transmembrane segments (S1–S6). Mutations positions are designated based on house fly sodium channel numbering (Genbank accession number: X96668).

Fig. 2.

Position of pyrethroid resistance-associated sodium channel mutations that are detected only in one species. The mutations with solid circles have been confirmed to reduce sodium channel sensitivity to pyrethroids in *Xenopus* oocytes. The mutations with open circles have not been examined in *Xenopus* oocytes yet. The information on these mutations is presented in Table 2 and Table 3. The sodium channel protein contains four homologous repeats (I–IV), each having six transmembrane segments (S1–S6). Mutations positions are designated based on house fly sodium channel numbering (Genbank accession number: X96668).

Table 1

Resistance-associated sodium channel mutations that are found in more than one species.

Mutation	Species	Common Name	Original Numbering	Reference
V410A	<i>Helicoverpa zea</i>	Corn earworm	V421A	[54]
V410G	<i>Helicoverpa zea</i>	Corn earworm	V421G	[54]
V410L	<i>Cimex luctularis</i>	Common bed bug	V419L	[55]
V410M	<i>Helicoverpa zea</i>	Corn earworm	V421M	[54]
	<i>Heliothis virescens</i>	Tobacco budworm	V421M	[56]
M827I + T929I + L932F	<i>Pediculus humanus capitis</i>	Human head louse	M815I + T917I + L920F	[57]
	<i>Pediculus humanus corporis</i>	Human body louse	M815I + T917I + L920F	[58]
M918I + L1014F	<i>Plutella xylostella</i>	Diamondback moth		[59]
M918L	<i>Aphis gossypii</i>	Melon and cotton aphid		[60]
M918L + L925I	<i>Trialeurodes vaporariorum</i>	Greenhouse whitefly		[61]
M918T	<i>Aphis gossypii</i>	Melon and cotton aphid		[62]
	<i>Tetranychus evansi</i>	Tomato red spider mite		[14]
M918T + L1014F	<i>Haematobia irritans irritans</i>	Horn fly		[63]
	<i>Liriomyza huidobrensis</i>	South American leaf miner		[12]
	<i>Musca domestica</i>	House fly		[8]
	<i>Myzus persicae</i>	Green peach aphid		[64]
	<i>Thrips tabaci</i>	Onion thrips		[65]
	<i>Tuta absoluta</i>	Tomato leaf miner		[66]
M918V	<i>Bemisia tabaci</i>	Sweet potato whitefly		[67]
L925I	<i>Bemisia tabaci</i>	Sweet potato whitefly		[67]
	<i>Cimex luctularis</i>	Common bedbug		[55]
	<i>Trialeurodes vaporariorum</i>	Greenhouse whitefly		[61]
	<i>Rhipicephalus microplus</i>	Southern cattle tick		[68]
T929C	<i>Frankliniella occidentalis</i>	Western flower thrips		[69]
T929C + L1014F	<i>Frankliniella occidentalis</i>	Western flower thrips		[69]
T929I	<i>Thrips tabaci</i>	Onion thrips		[65]
	<i>Thrips palmi</i>	Melon thrips		[70]
	<i>Trialeurodes vaporariorum</i>	Greenhouse whitefly		[61]
	<i>Leptinotarsa decemlineata</i>	Colorado potato beetle		[71]
	<i>Sitophilus zeamais</i>	Maize weevil		[72]
T929I + L932F	<i>Pediculus humanus capitis</i>	Human head louse		[73]
T929I + L1014F	<i>Frankliniella occidentalis</i>	Western flower thrips		[69]
	<i>Leptinotarsa decemlineata</i>	Colorado potato beetle		[71]
	<i>Plutella xylostella</i>	Diamondback moth		[74]
	<i>Tuta absoluta</i>	Tomato leaf miner		[66]
T929N + L1014F	<i>Leptinotarsa decemlineata</i>	Colorado potato beetle		[71]
T929V	<i>Bemisia tabaci</i>	Sweet potato whitefly		[75]
	<i>Ctenocephalides felis</i>	Cat flea		[76]
	<i>Frankliniella occidentalis</i>	Western flower thrips		[69]

Mutation	Species	Common Name	Original Numbering	Reference
T929V + L1014F	<i>Ctenocephalides felis</i>	Cat flea		[76]
I1011M	<i>Aedes aegypti</i>	Yellow fever mosquito	I104M	[77]
I1011V	<i>Aedes aegypti</i>	Yellow fever mosquito		[78]
L1014C	<i>Anopheles sinensis</i>	Malaria mosquito		[79]
	<i>Culex pipiens pipiens</i>	Southern house mosquito		[80]
L1014F	<i>Anopheles gambiae</i>	African malaria mosquito		[81]
	<i>Anopheles stephensi</i>	Malaria mosquito	L31F	[82]
	<i>Anopheles subpictus</i>	Malaria mosquito		[83]
	<i>Aphis gossypii</i>	Melon and cotton aphid		[62]
	<i>Blattella germanica</i>	German cockroach	L993F	[9; 46]
	<i>Ctenocephalides felis</i>	Cat flea		[76]
	<i>Culex pipiens pipiens</i>	House mosquito		[84]
	<i>Culex pipiens pallens</i>	Mosquito		[85]
	<i>Culex pipiens quinquefasciatus</i>	Southern house mosquito		[86]
	<i>Cydia pomonella</i>	Codling moth		[87]
	<i>Frankliniella occidentalis</i>	Western flower thrips		[69]
	<i>Haematobia irritans irritans</i>	Horn fly		[63]
	<i>Haematobia irritans exigua</i>	Buffalo fly		[88]
	<i>Leptinotarsa decemlineata</i>	Colorado potato beetle		[89]
	<i>Liriomyza huidobrensis</i>	South American leaf miner		[12]
	<i>Liriomyza sativae</i>	Vegetable leaf miner		[12]
	<i>Meligethes aeneus</i>	Pollen beetle		[90]
	<i>Musca domestica</i>	House fly		[7; 8; 9]
	<i>Myzus persicae</i>	Green peach aphid		[91]
	<i>Triatoma infestans</i>	Kissing bug		[92]
L1014H	<i>Helicoverpa zea</i>	Corn earworm	L1029H	[54]
	<i>Heliothis virescens</i>	Tobacco budworm	L1029H	[93]
	<i>Liriomyza trifolii</i>	American serpentine leaf miner		[12]
	<i>Musca domestica</i>	House fly		[94; 95]
	<i>Stomoxys calcitrans</i>	Stable fly		[96]
L1014S	<i>Anopheles arabiensis</i>	Malaria mosquito		[97]
	<i>Anopheles culicifacies</i>	Malaria mosquito		[98]
	<i>Anopheles gambiae</i>	African malaria mosquito		[99]
	<i>Anopheles parilae</i>	Malaria mosquito		[100]
	<i>Anopheles pedtaeniatus</i>	Malaria mosquito		[100]
	<i>Anopheles sacharovi</i>	Malaria mosquito		[101]
	<i>Anopheles sinensis</i>	Malaria mosquito		[100]
	<i>Anopheles vagus</i>	Malaria mosquito		[100]
	<i>Culex pipiens pallens</i>	Mosquito		[85]
	<i>Culex pipiens pipiens</i>	House mosquito		[84]
L1014W	<i>Anopheles sinensis</i>	Malaria mosquito		[102]
V1016G	<i>Aedes aegypti</i>	Yellow fever mosquito	V109G	[77]

Mutation	Species	Common Name	Original Numbering	Reference
V1016I	<i>Aedes aegypti</i>	Yellow fever mosquito		[78]
F1020S	<i>Blattella germanica</i>	German cockroach	F999S	[103]
	<i>Plutella xylostella</i>	Diamondback moth		[104]
F1534C	<i>Aedes aegypti</i>	Yellow fever mosquito	F1269C	[105]
	<i>Aedes albopictus</i>	Asian tiger mosquito		[106]
F1538I	<i>Rhipicephalus microplus</i>	Southern cattle tick	F1550I	[107]
	<i>Tetranychus cinnabarinus</i>	Carmine spider mite		[108]
	<i>Tetranychus urticae</i>	Two-spotted spider mite		[109]
D1549V + E1553G	<i>Helicoverpa armigera</i>	Cotton bollworm	D1561V + E1565G	[110]
	<i>Heliothis virescens</i>	Tobacco budworm	D1561V + E1565G	[110]

Mutation positions are designated based on house fly sodium channel numbering (Genbank accession number: X96668).

The “Original Numbering” column refers to the numbering of the mutation in the original paper, if different than house fly numbering.

Reference indicates the first report in the literature.

Table 2

Resistance-associated sodium channel mutations that are found only in one species.

Mutation	Species	Common Name	Original Numbering	Reference
I254N	<i>Drosophila melanogaster</i>	Common fruit fly	I286N	[111]
E435K + C785R + L1014F ³	<i>Blattella germanica</i>	German cockroach	E434K + C764R + L993F	[112]
M827I ³	<i>Pediculus humanus capitis</i>	Human head louse	M815I	[13]
M827I + T929I ³	<i>Pediculus humanus capitis</i>	Human head louse	M815I + T917I	[13]
M827I + L932F ³	<i>Pediculus humanus capitis</i>	Human head louse	M815I + L920F	[13]
T929I + L1014F + A1060T ^I + P1879S	<i>Plutella xylostella</i>	Diamondback moth	A1101T	[113]
G933V ^{3, 4}	<i>Rhipicephalus microplus</i>	Southern cattle tick	G72V	[114]
I936V ³	<i>Helicoverpa zea</i>	Corn earworm	I951V	[54]
Q945R	<i>Lepeophtheirus salmonis</i>	Sea flea		[115]
F979S + L1014F	<i>Myzus persicae</i>	Green peach aphid		[116]
S989P + V1016G	<i>Aedes aegypti</i>	Yellow fever mosquito		[117]
V1010L + L1014S	<i>Anopheles culicifacies</i>	Malaria mosquito		[98]
N1013S	<i>Anopheles sinensis</i>	Malaria mosquito		[102]
L1014F + A1060T ^I + P1879S	<i>Plutella xylostella</i>	Diamondback moth	A1101T	[113]
L1014F + N1575Y	<i>Anopheles gambiae</i>	African malaria mosquito		[118]
V1016G + D1763Y	<i>Aedes aegypti</i>	Yellow fever mosquito	D1794Y	[119]
L1024V	<i>Tetranychus urticae</i>	Two-spotted spider mite	L1022V	[120]
A1060T ^I + P1879S	<i>Plutella xylostella</i>	Diamondback moth	A1101T	[59]
A1215D ^I	<i>Tetranychus urticae</i>	Two-spotted spider mite		[109]
A1410V	<i>Drosophila melanogaster</i>	Common fruit fly	A1549V	[111]
A1494V	<i>Drosophila melanogaster</i>	Common fruit fly	A1648V	[111]
M1524I	<i>Drosophila melanogaster</i>	Fruit fly	Not numbered	[111]
F1528L + L1596P ⁵ + I1752V + M1823I ²	<i>Varroa destructor</i>	Varroa mite	F758L, L826P, I982V, M1055I	[121]

^IThis residue aligns poorly to MdNa_V1. Numbering relative to original numbering is retained.

²M in VdNa_V1 is V in MdNa_V1.

³Reduced sensitivity to pyrethroids has been confirmed in *Xenopus* oocytes (Table 3).

⁴C993V is resistant to deltamethrin [39].

⁵P1596L increases sensitivity to fluvalinate [122].

Mutation positions are designated based on house fly sodium channel numbering (Genbank accession number: X96668).

The “Original Numbering” column refers to the numbering of the mutation in the original paper, if different than house fly numbering.

Reference indicates the first report in the literature.

Table 3

Resistance-associated sodium channel mutations that have been confirmed to reduce sodium channel sensitivity to pyrethroids in *Xenopus* oocytes.

Mutation	Location	Reference
V410M	IS6	[27; 41; 42; 123]
V410M + E435K + C785R	IS6 + LI-II ¹ + LI-II	[41]
E485K + C785R + L1014F	LI-II + LI-II + IIS6	[25]
E485K + L1014F	LI-II + IIS6	[25]
C785R + L1014F	LI-II + IIS6	[25]
M827I	Linker IIS1-S2	[51]
M827I + T929I	Linker IIS1-S2 + IIS5	[51]
M827I + T929I + L932F	Linker IIS1-S2 + IIS5 + IIS5	[51]
M827I + L932F	Linker IIS1-S2 + IIS5	[51]
M918T	Linker IIS4-S5	[43; 48; 50]
M918T + L1014F	Linker IIS4-S5 + IIS6	[26; 48; 50]
L925I	Linker IIS4-S5	[39]
T929I	IIS5	[39; 43; 51]
T929I + L1014F	IIS5 + IIS6	[43]
T929I + L932F	IIS5 + IIS5	[51]
L932F	IIS5	[39; 51]
C933A ²	IIS5	[39]
I936V	IIS5	[39]
L1014F	IIS6	[25; 26; 44; 45; 47; 48]
L1014H	IIS6	[27; 47]
L1014S	IIS6	[47]
F1534C	IIIS6	[49]
F1538I	IIIS6	[124]
L1596P ³	LIII-IV	[122]

¹ the linker connecting repeats I and II of the sodium channel.

² G933A is associated with resistance in *Rhipicephalus microplus* [114].

³ P1596L increases sensitivity to fluvalinate [122].