Published in final edited form as:

Org Lett. 2013 May 17; 15(10): 2410–2413. doi:10.1021/ol400872h.

Synthesis of a Highly Functionalized Core of Verrillin

Alec Saitman and Emmanuel A. Theodorakis

Department of Chemistry and Biochemistry, University of California, San Diego, 9500 Gilman Drive MC: 0358, La Jolla, California 92093-0358, United States

Emmanuel A. Theodorakis: etheodor@ucsd.edu

Abstract

An efficient stereoselective synthesis of furanoverrillin (5), a highly functionalized core of verrillin (1), is reported. The synthetic strategy is based on constructing bicyclic lactone 17 prior to the 10-membered ring macrocyclization. The effect of the C₄ methyl group on the furan reactivity is also discussed.

Gorgonian octocorals and soft corals supply a diverse array of diterpenes and C₄-norditerpenes that biosynthetically derive from a 14-membered cembrane skeleton.¹ Among them, verrillin (1), bielschowskyisin (2), ineleganolide (3) and plumarellide (4) are highlighted by a combination of intricate structural density and potent bioactivity (Figure 1).^{1a,2} For example, bielschowskyisin (2),³ a multifarious hexacyclic cembranoid, was found to exhibit significant anti-malarial activity against *Plasmodium falciparum* (IC₅₀ ca. 10 µg/ml). In addition, potent cytotoxicity data regarding 2 have been reported against non-small cell lung cancer (EKVX, GI₅₀ ca. 0.01 µM) and renal cancer cells (CAKI-1, GI₅₀ ca. 0.50 µM).³ Along these lines, ineleganolide (3) displays strong cytotoxicity against P-388 cancer cells (ED₅₀ 3.82 µg/ml).⁴ On the other hand, plumarellide (4), exhibits moderate hemolytic activity in mice with 50% hemolysis of mouse blood erythrocytes at 140 µM⁵ while the bioactivities of verrillin (1)⁶ have not yet been studied.

A commonality shared by these polycyclic cembranoids is an oxidative ring contraction of the 14-membered cembrane ring across the C₇ and C₁₁ carbons thereby producing the verrillane core (6) (Figure 2).^{3,7} Subsequent ring contractions across the C₆ and C₁₂ or the C₆ and C₁₄ centers would produce the bielschowskyane (9)^{1c,3,7} and the plumarane core (7),^{1c} respectively. On the other hand, oxidative demethylation followed by nucleophilic ring contraction at the C₄ and C₁₃ centers could assemble the inelegane core (8).^{1c}

Although several strategies towards the synthesis of bielschowskyisin⁸ (2) and plumarellide⁹ (4) have been reported, none have generated a true cembrane scaffold with relevant carbon connections and oxidation patterns. Inspired by the challenge, we directed our efforts^{7,10}

Correspondence to: Emmanuel A. Theodorakis, etheodor@ucsd.edu.

Supporting Information Available Detailed experimental procedures, spectral characterization and copies of ¹H and ¹³C NMR data. This material is available free of charge via the Internet at <http://pubs.acs.org>.

towards the synthesis of motif **5** that contains an appropriately functionalized verrillane core (Figure 2). Referred to herein as furanoverrillin, compound **5** represents a branching node toward the synthesis of verrillin (**1**) and its more intricate brethren.

Scheme 1 highlights the key disconnections of furanoverrillin (**5**) and identifies the essential synthetic fragments. We envisioned that construction of the fused bicyclic lactone, present in our target, could be furnished via an Eschenmoser-Claisen rearrangement.¹¹ This disconnection reveals allylic alcohol **13** and commercially available amide acetal **12** as the coupling partners. Claisen alkylation of this adduct with aldehyde **11** could then create the desired C₁₂-C₁₃ connection as well as provide the important C₁₃ oxidation present in this family of compounds.⁷ Pd(0)-coupling could then be used to join furan derivative **10** by creating the C₆-C₇ bond, while macrocyclization under Nozaki-Hiyama-Kishi (NHK) conditions¹² could afford the final C₁-C₂ linkage.

The synthesis of furanoverrillin (**5**) commenced with the formation of lactone **17** (Scheme 2). Furfuryl alcohol **14** was rearranged to a cyclopentenone derivative under microwave irradiation¹³ that, after TBS protection and regioselective iodination,¹⁴ gave rise to iodo enone **15** (3 steps, 40% overall yield). Nucleophilic addition onto the carbonyl with methyl magnesium bromide yielded tertiary alcohol **13** as a single diastereomer in good yield (52%). To our satisfaction, the key Eschenmoser-Claisen rearrangement¹⁵ of **13** with **12** proceeded smoothly and in a diastereoselective manner, to afford amide **16** (70% yield). Treatment of the crude amide **16** with excess acid led to rapid cleavage of the TBS group and cyclization of the resulting alcohol at the amide carbonyl center to produce lactone **17** in 91% yield (Scheme 2).

Deprotonation of **17** with freshly prepared LiHMDS followed by quick addition of the previously described aldehyde **11**⁷ supplied alcohol **18** as a 1:1.7 mixture of C₁₃ diastereomers.⁷ These isomers, **18a** and **18b**, were separated by column chromatography and used separately in subsequent manipulations.¹⁶ TBS protection of the newly installed alcohol functionality was markedly slow and low yielding. Protection of the secondary alcohols **18a** and **18b** with MOMCl¹⁷ was also slow at room temperature. Microwave irradiation conditions at 80 °C for 10 minutes, however, successfully yielded the desired MOM protected alcohols. Selective TBS deprotection of this crude material using PPTS in ethanol afforded primary alcohols **19a** and **19b** (ca. 71% yield over 2 steps).¹⁸

The synthesis of methyl furfuryl stannane **10** was accomplished as shown in Scheme 4. Commercially available furan derivative **26** was reduced to the corresponding primary alcohol¹⁹ that, after oxidation with manganese dioxide²⁰ under microwave irradiation produced aldehyde **27** (2 steps, 85% yield). Stannylation of a transiently protected aldehyde **27** with trimethyltin chloride afforded stannane **10** in 65% yield.²¹ In a similar fashion, aldehyde **28** was converted to stannane **29**.^{7,21}

To our chagrin, coupling of stannane **10** with primary alcohols **19a** and **19b** under the previously established Stille conditions (Pd(PPh₃)₄/CuI)^{7,10,21} was not successful. However, switching of reagents to catalytic Pd₂(dba)₃ and triphenylarsine²² cleanly afforded compounds **20a** and **20b** in 65% average yield (Scheme 3). Appel bromination conditions²³ were employed to convert alcohols **20a** and **20b** to their corresponding bromides **22a** and **22b** (ca. 84% yield).

The final macrocyclization event that would create the 10-membered macrocycle²⁴ also proved to be more problematic than previously described^{7,10,21} leading to a complex mixture of products. Changing the solvent from THF to DMF²⁵ greatly improved the reaction outcome to afford furanoverrillin (**5**), containing the desired verrillin

stereochemistry at C₁₃, together with its C₁₃ α-isomer **24a** in 30% combined yield. The diastereoselectivity of the macrocyclization was 4/1: anti/syn with respect to the orientation of the C₂ alcohol and the C₁ isopropene.^{24a}

Compounds **5** and **24a** were subjected to standard C₂ deoxygenation conditions using triethylsilane and trifluoroacetic acid²⁶ to produce deoxygenated products **30a** and **30b** in lower than expected yields (ca. 40% yield) (Scheme 5).^{7,10,21} Acid-induced MOM deprotection under microwave conditions²⁷ afforded the corresponding alcohols **32a** and **32b** in an average yield of 72%.

During this study, we observed modest yields in converting **22a** and **22b** to **24a** and **5**, respectively (Scheme 3). This may be due to the inherent tendency of these compounds to undergo carbocationic rearrangements at the C₂ center, as previously reported for certain C₂ hydroxy-furanocembrenolides and pseudotanolides.²⁸ An alternative explanation is to consider that, due to its high electronic density, the furan ring becomes sensitive to oxidative decomposition. To test this hypothesis we repeated the aforementioned sequence using the non-methylated furfuraldehyde **28** instead of its methylated analogue **27**. To this end, we coupled the previously synthesized furfural stannane **29**^{7,10} with alcohols **19a** and **19b** to afford **21a** and **21b** in comparable yields (ca. 65% yield). Appel conditions^{7,10,21,23} also smoothly afforded bromides **23a** and **23b** again, in nearly identical yields (ca. 84% yield). When optimized NHK¹² conditions were then applied using DMF as a solvent²⁵, a clean conversion to the desired nor-furanoverrillin intermediates **25a** and **25b** was observed in good yields (ca. 60% yield). Deoxygenation at the C₂ center using triethylsilane and trifluoroacetic acid²⁶ also cleanly produced deoxygenated intermediates **31a** and **31b** in noticeably increased yields. Finally, acidic removal of the MOM group²⁷ created alcohols **33a** and **33b** in similar yields (ca. 72% yield). Along with a striking improvement in reaction yields of the nor-furanoverrillin derivatives was the ability to then obtain X-ray structures for compounds **25b** and **33b** which confirmed the proposed stereochemical assignments²⁹.

In conclusion, we present herein a stereoselective synthesis of furanoverrillin (**5**), a branching node toward the synthesis of verrillin (**1**) and related polycyclic cembrenoids. Our approach relies on introduction of the C₇-C₁₁ cyclopentene ring prior to the macrocyclization of the fully functionalized verrillin core. We observed that the presence of a C₄ methyl group at the furan ring, increases its susceptibility toward oxidative decomposition. In turn, this attests to the role of the C₄ methyl group in oxidative rearrangements that account for the plethora of polycyclic cembranes isolated thus far. The described strategy produces, for the first time, the entire carbon framework of verrillin and allows for further functionalization toward verrillin (**1**) and structurally related natural products.³⁰

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We gratefully acknowledge the National Institutes of Health (NIH) for financial support of this work through Grant Number R01 GM081484. We thank the National Science Foundation for instrumentation grants CHE9709183 and CHE0741968. We also thank Dr. Anthony Mrse (UCSD NMR Facility), Dr. Yongxuan Su (UCSD MS Facility) and Dr. Arnold L. Rheingold and Dr. Curtis E. Moore (UCSD X-Ray Facility). We also thank Danaya Pratchayanan and Parawin Adisayathepkul (UCSD, Summer Undergraduate Internship) for their efforts in synthesizing early-stage intermediates.

References

1. (a) Roethle PA, Trauner D. *Nat Prod Rep.* 2008; 25:298–317. [PubMed: 18389139] (b) Li Y, Pattenden G. *Nat Prod Rep.* 2011; 28:429–440. [PubMed: 21212904] (c) Li Y, Pattenden G. *Nat Prod Rep.* 2011; 28:1269–1310. [PubMed: 21637894]
2. Lakshmi V, Kumar R. *Nat Prod Res.* 2009; 23:801–850. [PubMed: 19488923]
3. Marrero J, Rodriguez AD, Baran P, Raptis RG, Sanchez JA, Ortega-Barria E, Capson TL. *Org Lett.* 2004; 6:1661–1664. [PubMed: 15128261]
4. Duh CY, Wang SK, Chia MC, Chiang MY. *Tetrahedron Lett.* 1999; 40:6033–6035.
5. Stonik VA, Kapustina II, Kalinovsky AI, Dmitrenok PS, Grebnev BB. *Tetrahedron Lett.* 2002; 43:315–317.
6. Rodriguez AD, Shi YP. *J Org Chem.* 2000; 65:5839–5842. [PubMed: 10970333]
7. Saitman A, Sullivan SDE, Theodorakis EA. *Tetrahedron Lett.* 2013; 54:1612–1615. [PubMed: 23626379]
8. (a) Doroh B, Sulikowski GA. *Org Lett.* 2006; 8:903–906. [PubMed: 16494470] (b) Miao R, Gramani SG, Lear MJ. *Tetrahedron Lett.* 2009; 50:1731–1733. (c) Nicolaou KC, Adsool VA, Hale CRH. *Angew Chem Int Ed.* 2011; 50:5149–5152. (d) Farcet J, Himmelbauer M, Mulzer J. *Org Lett.* 2012; 14:2195–2197. [PubMed: 22506798] (e) Jana A, Mondal S, Hossain MF, Ghosh S. *Tetrahedron Lett.* 2012; 53:6830–6833.
9. (a) Li Y, Pattenden G. *Tetrahedron Lett.* 2011; 52:2088–2092. (b) Palframan M, Pattenden G. *Tetrahedron Lett.* 2013; 54:324–328.
10. Saitman A, Rulliere P, Sullivan SDE, Theodorakis EA. *Org Lett.* 2011; 13:5854–5857. [PubMed: 21973225]
11. (a) Wick AE, Felix D, Steen K, Eschenmoser A. *Helv Chim Acta.* 1964; 47:2425–2429. (b) Castro AMM. *Chem Rev.* 2004; 104:2939–3002. [PubMed: 15186185]
12. (a) Okude Y, Hirano S, Hiyama T, Nozaki H. *J Am Chem Soc.* 1979; 99:3179–3181. (b) Takai K, Kimura K, Kuroda T, Hiyama T, Nozaki T. *Tetrahedron Lett.* 1983; 24:5281–5284. (c) Fürster A. *Chem Rev.* 1999; 99:991–1045. [PubMed: 11848998]
13. Kruger G, Harde C, Bohlmann F. *Tetrahedron Lett.* 1985; 49:6027–6030.
14. Otero MP, Santin EP, Rodriguez-Barrios F, Vaz B, De Lera AR. *Biorg Med Chem Lett.* 2009; 19:1883–1886.
15. Mulzer J, Giester G, Gilbert M. *Helv Chim Acta.* 2005; 88:1560–1579.
16. Lautens M, Paquin JF. *Org Lett.* 2003; 5:3391–3394. [PubMed: 12967282] Bourcet E, Fache F, Piva O. *Tetrahedron.* 2010; 66:1319–1326. (c) The stereochemistry of the C₁₃ hydroxyl group was unambiguously determined after macrocyclization via single crystal X-ray diffraction analysis of compounds 25a and 33b.
17. Maegawa T, Otake K, Hirosawa K, Goto A, Fujioka H. *Org Lett.* 2012; 14:4798–4801. [PubMed: 22931431]
18. Crouch RD. *Tetrahedron.* 2004; 60:5833–5871.
19. Aldington RM, Baldwin JE, Mayweg AVW, Pritchard GJ. *Org Lett.* 2002; 4:3009–3011. [PubMed: 12182611]
20. Hashami ASK, Haffner T, Yang W, Pankajakshan S, Schafer S, Schultes L, Rominger F, Frey W. *Chem Eur J.* 2012; 18:10480–10486. [PubMed: 22807391]
21. Roethle P, Trauner D. *Org Lett.* 2006; 8:345–347. [PubMed: 16408911]
22. (a) Farina V, Krishnan B. *J Am Chem Soc.* 1991; 113:9585–9595. (b) Farina V, Kapadia S, Krishnan B, Wang C, Liebeskind LS. *J Org Chem.* 2000; 65:3192–3199. [PubMed: 10814214] (c) Roth GP, Farina V. *Tetrahedron Lett.* 1995; 36:2191–2194.
23. Appel R. *Angew Chem Int Ed.* 1975; 14:801–811.
24. (a) Huang QH, Rawal VH. *Org Lett.* 2006; 8:543–545. [PubMed: 16435880] (b) Paquette LA, Astles PC. *J Org Chem.* 1993; 58:165–169. (c) Rayner CM, Astles PC, Paquette LA. *J Am Chem Soc.* 1992; 114:3926–3936.
25. Maguire RJ, Mulzer J, Bats JW. *Tetrahedron Lett.* 1996; 37:5487–5490.
26. Kursanov DN, Parnes ZN, Loim NM. *Synthesis.* 1974; 9:633–651.

27. Yadav JS, Premalatha K, Harshavardhan SJ, Reddy BVS. *Tetrahedron Lett.* 2008; 49:6765–6767.
28. (a) Hyde EG, Thornhill SM, Boyer AJ, Abramson SN. *J Med Chem.* 1995; 38:4704–4709. [PubMed: 7473598] (b) Rodriguez AD, Shi JG, Shi YP. *J Org Chem.* 2000; 65:3192–3199. [PubMed: 10814214]
29. CCDC 929062 (25a) and 929063 (33b) contain the supplementary crystallographic data. This information can be obtained free of charge from the Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/products/csd/request/. In the figures some hydrogen atoms have been removed for clarity
30. (a) Roethle PA, Hernandez PT, Trauner D. *Org Lett.* 2006; 8:5901–5904. [PubMed: 17134301] (b) Tang B, Bray CD, Pattenden G. *Tetrahedron Lett.* 2008; 47:6401–6404. (c) Kimbrough TJ, Roethle PA, Mayer P, Trauner D. *Angew Chem Int Ed.* 2010; 49:1–4. (d) Marshall JA, Griot CA, Chobanian HR, Myers WH. *Org Lett.* 2010; 12:4328–4331. [PubMed: 20795632] (e) Li Y, Pattenden G. *Tetrahedron.* 2011; 67:10045–10052.

1: verrillin **2: bielschowskysin** **3: ineleganolide** **4: plumarellide**

Figure 1.
Structures of selected polycyclic cembrenoids

Figure 2.
Carbocyclic scaffolds related to verrillane core **6**

Scheme 1.
Retrosynthetic analysis of furanoverrillin (5)

Scheme 2.
Synthesis of fragment 17

Scheme 3.
Synthesis of furanoverrillin **5** and derivatives

Scheme 4.
Synthesis of furfuryl stannanes **10** and **29**

Scheme 5.
 Synthesis of **32** and **33**