Measurement of Bone Mineral Density in Children With Cancer

To the Editor:

The article by Aldhafiri et al¹ raises several clinically relevant issues. As addressed in the principle objective given in the study, measurement of bone mineral density in children by dual-energy x-ray absorptiometry (DXA) requires adjustment for height and weight, therefore providing validation of our observations reported in 2009.² Secondly, it is important to establish local norms with DXA and not rely on the ranges provided by the manufacturers of densitometers.³ Thirdly, body mass index is an imperfect measure of nutritional status in children with cancer,⁴ in whom arm anthropometry appears to provide a better estimate.⁵ Finally, DXA offers accurate determinations of body composition: lean body mass (fat-free mass), fat mass, and whole-body bone mineral content, summing up to body weight.³ It would be of interest to know whether, in children examined by Aldhafiri and colleagues, there was a relationship between lean body mass and bone mineral density corrected for height and weight.

Ronald D. Barr, MB, ChB, MD

Department of Pediatrics, Division of Hematology-Oncology, McMaster University, Hamilton, ON

REFERENCES

1. Aldhafiri F, Al-Nasser A, Al-Sugair A, et al. Importance of adjusting dual-energy X-ray output for body size: an example from survivors of childhood acute lymphoblastic leukemia. *J Pediatr Hematol Oncol*. 2013;35:e27.
2. Webber CE, Sala A, Barr RD. Accounting for body size deviations when

reporting bone mineral density variables in children. *Osteoporos Int*. 2009;20:113–121.

3. Sala A, Webber CE, Morrison J, et al. Whole body bone mineral content, lean body mass, and fat mass measured by dual-energy X-ray absorptiometry in a population of normal Canadian children and adolescents. *Can Assoc Radiol J*. 2007;58:46–52.
4. Sala A, Pencharz P, Barr RD. Children, cancer, and nutrition—a dynamic triangle in review. *Cancer*. 2004;100:677–687.
5. Sala A, Rossi E, Antillon F, et al. Nutritional status at diagnosis is related to clinical outcomes in children and adolescents with cancer: a perspective from Central America. *Eur J Cancer*. 2012;48:243–252.

In Response:

We welcome the letter by Prof Barr.¹ Indeed, we consider our study on a sample of adolescent survivors of ALL² to provide empirical support for the prediction made in a study by Webber et al³: interpreting dual energy x-ray (DXA) bone health data requires caution. In particular, failure to adjust DXA bone mineral density (BMD) data for body size risks erroneous judgments about bone health, which could be clinically important,³ particularly in children and adolescents with chronic disease.^{1,2} Although DXA is a valuable component of the assessment of bone health,⁴ and knowledge of the need to adjust BMD values appropriately for body size is increasing, this point is not as widely known as it should be at present.² Barr¹ also highlights the potentially valuable role for DXA in the assessment of body composition. We agree that DXA has great potential as an accurate and precise measure of fat and lean mass in children and adolescents. However, DXA is not a “gold standard” for body composition measurement,⁵ and we

would add that caution is also necessary here. Different combinations of DXA hardware and software are likely to have different accuracy, and the errors inherent in DXA estimates of body composition can be very large.⁶ In summary, DXA, when used appropriately, can be extremely valuable in the assessment of both bone health and body composition, and the technique can inform both research and clinical management.

John J. Reilly, PhD*

Sheila K. Shepherd, PhD†

Fahad K. Aldhafiri, MSc‡

Faisal Ahmed, MD, RCPCH‡

*School of Psychological Sciences and Health, University of Strathclyde

†Yorkhill Hospitals

‡College of Medical, Veterinary, and Life Sciences, University of Glasgow, Yorkhill Hospitals, Glasgow, Scotland

REFERENCES

1. Barr RD. Response to Aldhafiri et al. Importance of adjusting dual energy x-ray output for body size: an example from survivors of childhood acute lymphoblastic leukemia' JPHO 35.5 (in press).
2. Aldhafiri F, Al-Nasser A, Al-Sugair A, et al. Importance of adjusting dual energy x-ray output for body size: an example from survivors of childhood acute lymphoblastic leukaemia. *J Pediatr Hematol Oncol*. 2013;35:e27–e29.
3. Webber CE, Sala A, Barr RD. Accounting for body size deviations when reporting bone mineral density variables in children. *Osteoporos Int*. 2009;20:113–121.
4. Wong SC, Khanna S, Rashid R, et al. Auditing bone densitometry and fractures in children with chronic disease. *Arch Dis Child*. 2008;93:705–707.
5. Wells JCK, Fewtrell MS. Measuring body composition. *Arch Dis Child*. 2006;91:612–617.
6. Reilly JJ, Gerasimidis K, Papararcleous N, et al. Validation of dual energy x-ray absorptiometry and foot-foot impedance against deuterium dilution measures of fatness in children. *Int J Pediatr Obes*. 2010;5:111–115.