Published in final edited form as:

Tetrahedron Lett. 2013 July 31; 54(31): 4083–4085. doi:10.1016/j.tetlet.2013.05.101.

Chemoselective hydroamination of vinyl arenes catalyzed by an NHC-amidate-alkoxide Pd(II) complex and *p*-TsOH

Richard Giles, Justin O'Neill, Joo Ho Lee, Michael K. Chiu, and Kyung Woon Jung

Loker Hydrocarbon Research Institute and Department of Chemistry, University of Southern California, Los Angeles, California 90089, USA

Abstract

The hydroamination of various substituted vinyl arenes with benzenesulfonamide was explored using an NHC-amidate-alkoxide palladium catalyst in conjunction with *p*-TsOH. Utilizing halide-substituted and electron-rich vinyl arenes, this methodology selectively furnished the cross-coupled hydroamination products in moderate to excellent yields in a Markovnikov fashion while greatly reducing undesired acid-catalyzed homocoupling of the vinyl arenes. Electron-rich vinyl arenes typically required milder conditions than electron-poor ones. While most effective for para-substituted substrates, the catalyst system also furnished the desired products from ortho- and meta-substituted vinyl arenes with high chemoselectivities.

Keywords

Hydroamination; Vinyl arenes; Palladium catalyst; Brønsted acid; Sulfonamide

Introduction

Hydroamination is an expeditious method to form carbon-nitrogen bonds from olefins. While this process is typically a challenging reaction due to its high barrier of activation, various methods have been successfully developed.¹ In particular, late transition metal catalysts have been extensively employed in hydroamination reactions due to their high reactivity, resistance towards degradation by water, and high functional group tolerance. Homogeneous intermolecular hydroamination of vinyl arenes with weakly basic nitrogen sources such as sulfonamides has been catalyzed by the salts of metals including platinum, copper, bismuth, hafnium, iron, gold, silver, and zirconium.^{2a-f} However, most of these methods have shown limitations, especially with electron-rich vinyl arenes. These substrates, including methyl or methoxy-substituted styrenes, often furnished hydroamination products in low-to-intermediate yields along with large amounts of side products. In addition, some of these procedures were only effective under inert atmospheric conditions.

Other methods utilized to catalyze the homogeneous intermolecular hydroamination of vinyl arenes with sulfonamides include the use of N-bromosuccinimide,³ molecular iodine,⁴ N-

© 2013 Elsevier Ltd. All rights reserved.

Correspondence to: Kyung Woon Jung.

Corresponding author. Tel.: 1-213-740-8768; fax: 1-213-740-6270; kwjung@usc.edu..

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

fluorobenzenesulfonimide,⁵ and ionic liquids.⁶ Heterogeneous methods using gallium⁷ and Amberlyst-15⁸ have also been reported. Notably, the few homogeneous Brønsted acid-catalyzed examples that were reported were limited to simple unsubstituted styrenes.^{9a-b} There have been several reasons for the dearth of these reactions, including the tendency of the nitrogen sources to buffer the effect of the acids as well as the acid-catalyzed formation of undesired homocoupling products from vinyl arene substrates.^{2d,8}

In order to avert these shortcomings, we actively pursued the development of new conditions by employing dual catalysts such as our NHC-amidate-alkoxide palladium complex **1** (figure 1)¹⁰ and *p*-toluenesulfonic acid. When we used acids as the sole catalysts, 4-methylstyrene **2** underwent homocoupling to furnish dimer **5** predominantly (scheme 1). However, our hybrid method facilitated the Markovnikov-type hydroamination efficiently and improved the yields of the hydroamination products (e.g., **4** in scheme 1) while minimizing undesired homocoupling side products (e.g., **5** in scheme 1).

Results and Discussion

As shown in table 1, we embarked on a comparative study on hydroamination and homocoupling. Using DMF as an internal NMR standard, we recorded the conversion yields of **4** based on the sulfonamide as the limiting reagent and the mole ratios of **4** and **5** as the measure of chemoselectivity. Among several acids examined including TfOH and TFA, *p*-TsOH turned out to be the most effective for hydroamination but still gave the dimer **5** as the major product (entry 1). We then investigated the use of palladium complexes as co-catalysts. When known palladium complexes were used (entries 2 and 3), both the yield of hydroamination product and the chemoselectivity remained similar to those of the acid-only conditions (entry 1). In sharp contrast, the addition of catalyst **1** led to a great improvement in hydroamination conversion and considerable reversal of chemoselectivity. Comparable results were observed with MeSO₃H (entry 5). Unlike known catalysts, our catalyst **1** inhibited undesired Brønsted acid-catalyzed homocoupling of the olefin while promoting the desired hydroamination, implying that the structure and electronic properties of the ligand, rather than the simple presence of a palladium salt, could be the driving force behind the unique reactivity and selectivity.

With these trends noted, we endeavored to determine the influence of several other factors including solvent and substrate ratios (table 2). Three distinct outcomes based on solvent effects were observed; the reaction was very active but not selective (1,2-dichloroethane, entry 1), no reaction occurred (dioxane, entry 2), and the reaction was efficient and highly selective towards the hydroamination product (toluene, entry 3). When we decreased the amount of the olefin to limit formation of dimer **5**, we observed much higher selectivities as anticipated. However, the yields of **4** diminished significantly, revealing the importance of using an excess of the olefin for optimal results (entries 4 – 5).

Subsequently, we investigated the effect of lowering catalyst **1** and acid loading at different temperatures (table 3). When we used 5 mol% of *p*-TsOH, the chemoselectivities improved dramatically but the reactions were sluggish and provided much lower yields of the hydroamination product (entries 1 and 2). With 10 mol% of the acid and 2 – 5 mol% of the Pd catalyst, we observed reasonably high yield and selectivity in favor of the hydroamination product (entry 3). Lower temperature improved the selectivity despite slightly lower yield (entry 4) while higher temperatures gave comparable yields with slightly lower selectivity (entry 5). Thus depending on the nature of the substrate, the temperature can be varied to improve either selectivity or yield as desired.

Under the optimized catalytic conditions, the substrate scope of this methodology was examined by screening various different vinyl arenes (scheme 2 and table 4).¹¹ Electron-rich vinyl arenes, including 4-methylstyrene (entry 1), 2,4-dimethylstyrene (entry 2), and 4-methoxystyrene (entry 3), furnished the hydroamination products in good yields and selectivities at relatively low temperatures (40 – 60 °C). 2,4-Dimethylstyrene in particular reacted in essentially quantitative yield under these conditions. 4-Methoxystyrene, in comparison, was much more sensitive to polymerization than any other substrates and required very mild conditions, including increased catalyst loading and a lower temperature, to give the hydroamination product in a decent yield. When the substitution pattern was changed from 4-methoxy to 3-methoxy (entry 4), the reactivity decreased dramatically and higher temperatures were required to afford comparable yields. 3-Methoxystyrene behaved similarly to unsubstituted and electron-neutral substrates encompassing styrene (entry 5) and 2-vinylnaphthalene (entry 6), which reacted smoothly at higher temperatures than electron-rich styrenes. Higher temperatures or increased amounts of *p*-TsOH resulted in significantly lower chemoselectivities, yielding increased amounts of homocoupled product **8**.

4-F-, Cl-, and Br-substituted vinyl arenes were subjected to the developed catalytic conditions and each of these substrates provided high yields and selectivities under conditions similar to those used for electron-neutral examples (entries 7-9).

As expected based on the aforementioned examples, the reactivities dropped significantly when the substitution pattern was changed from *para* to *ortho*. These substrates required higher temperatures in chlorobenzene solvent with higher loading of *p*-TsOH (entries 10-12). 2-Fluoro- and 2-bromostyrenes provided decent yields and selectivities at 130 °C. In contrast, 2-chlorostyrene underwent significant polymerization as well as hydroamination and dimerization (entry 11). 3-Chlorostyrene was a poor substrate (entry 13) while additional styrenes with electron withdrawing groups were also resistant to hydroamination conditions. These results suggest that the vinyl arenes play an electrophilic role in the hydroamination reaction and the electron-withdrawing substituents on the styrenes would slow their reactivities. In pursuit of wider applications, we sought to conduct hydroamination on disubstituted styrenes but failed to acquire satisfactory results. For instance, *cis*-4-methylstyrene barely generated the desired product because of almost complete isomerization to the more stable *trans*-olefin (entry 14) which was largely resistant to these conditions.

Conclusion

The combination of the tridentate NHC-amidate-alkoxide palladium complex **1** and *p*-TsOH efficiently catalyzed the Markovnikov-type hydroamination of electronically varying vinyl arenes with benzenesulfonamide. Our hybrid method improved the yields of the hydroamination products while minimizing the undesired homocoupling side products, which existing methods commonly produced in the presence of strong acids. In addition, while existing methods were generally limited to unsubstituted or *para*-substituted substrates, our method was highly efficient for vinyl arenes with *ortho*-, *meta*-, and *para*-substitutions. Electron-rich and electron-neutral vinyl arenes were well suited for the intended selective hydroamination. We hope that our ongoing studies could elucidate the mechanism of this transformation in due course.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We acknowledge generous financial support from the Hydrocarbon Research Foundation and the National Institute of Health (S10 RR025432).

References and notes

1. Muller TE, Hultzsck KC, Yus M, Foubelo F, Tada M. Chem. Rev. 2008; 108:3795–3892. [PubMed: 18729420]
2. a Qian H, Widenhoefer RA. Org. Lett. 2005; 7:2635–2638. [PubMed: 15957909] b Taylor JG, Whittall N, Hii KK. Org. Lett. 2006; 8:3561–3564. [PubMed: 16869660] c Qin H, Yamagiwa N, Matsunaga S, Shibasaki M. Chem. Asian. J. 2007; 2:150–154. [PubMed: 17441148] d Michaux J, Terrasson V, Marque S, Wehbe J, Prim D, Campagne J-M. Eur. J. Org. Chem. 2007:2601–2603. e Zhang X, Corma A. Dalton. Trans. 2008; 7:397–403. [PubMed: 18411849] f Yang L, Xu L-W, Zhou W, Gao Y-H, Sun W, Xia C-G. Synlett. 2009; 7:1167–1171. g Dal Zotto C, Michaux J, Zarate-Ruiz A, Gayon E, Virieux D, Campagne J-M, Terrasson V, Pieters G, Gaucher A, Prim D. J. Organomet. Chem. 2011; 696:296–304. h Giner X, Nájera C, Kovács G, Lledós A, Ujaque G. Adv. Synth. Catal. 2011; 353:3451–3466.
3. Talluri SK, Sudalai A. Org. Lett. 2005; 7:855–857. [PubMed: 15727458]
4. Yadav JS, Reddy BVS, Rao TS, Krishna BBM. Tetrahedron. Lett. 2009; 50:5351–5353.
5. Xu T, Qiu S, Liu G. J. Organomet. Chem. 2011; 696:46–49.
6. Yang L, Xu L-W, Xia C-G. Synthesis. 2009; 12:1969–1974.
7. Jaspers D, Kubiak R, Doye S. Synlett. 2010; 8:1268–1272.
8. Qureshi ZS, Deshmukh KM, Tambade PJ, Dhake KP, Bhanage BM. Eur. J. Org. Chem. 2010:6233–6238.
9. a Rosenfeld DC, Shekhar S, Takemiya A, Utsunomiya M, Hartwig JF. Org. Lett. 2006; 8:4179–4182. [PubMed: 16956181] b Li Z, Zhang J, Brouwer C, Yang C-G, Reich NW, He C. Org. Lett. 2006; 8:4175–4178. [PubMed: 16956180]
10. Lee JH, Yoo KS, Park CP, Olsen J, Sakaguchi S, Prakash GKS, Mathew T, Jung KW. Adv. Synth. Catal. 2009; 351:563–568. [PubMed: 20221298]
11. General conditions: to a mixture of **3** (0.25 mmol), **1** (10 mol%), and *p*-TsOH (5.00×10^{-2} mmol) in a 2 dram vial were added toluene (3 mL) and **2** (0.75 mmol). After stirring for 18 h at 100 °C, the reaction mixture was diluted with CH₂Cl₂ (3 mL), neutralized with Et₃N (5.00×10^{-2} mmol), and filtered through Celite. The resulting solution was concentrated *in vacuo* then purified using flash column chromatography with silica gel (4:1 hexanes/EtOAc).

Figure 1.
NHC-amidate-alkoxide Pd(II) catalyst **1**.

Scheme 1.
Hydroamination of 4-methylstyrene.

Scheme 2.
Hydroamination of vinyl arenes.

Table 1Effect of acids and palladium complexes on hydroamination of 4-methylstyrene.^a

entry	Pd catalyst ^b	Acid	4 (yield)	4 : 5 (mol ratio)
1	-	<i>p</i> -TsOH	71%	1 : 1.6
2	Pd(PPh ₃) ₂ Cl ₂	<i>p</i> -TsOH	81%	1 : 1.6
3	Pd(OAc) ₂	<i>p</i> -TsOH	71%	1 : 1.7
4	1	<i>p</i> -TsOH	95%	7.9 : 1
5	1	MeSO ₃ H	91%	5.1 : 1

^a All reactions were run using 0.25 mmol of benzenesulfonamide, 0.75 mmol of 4-methylstyrene, and 20 mol% of acids in toluene at 60 °C.^b 10 mol% of Pd catalysts were used.

Table 2Effect of solvents and substrate ratios.^a

entry	Ratio (2 : 3)	Solvent	4 (yield)	4 : 5 (mol ratio)
1	3 : 1	DCE	81%	1.2 : 1
2	3 : 1	dioxane	0%	0 : 0
3	3 : 1	toluene	81%	20 : 1
4	1 : 1	toluene	67%	67 : 1
5	1 : 3	toluene	61%	61 : 1

^a All reactions were run using 10 mol% of the catalyst and acid each at 60 °C.

Table 3

Effect of catalyst and acid at different temperatures.^a

entry	1 (mol%)	<i>p</i> -TsOH (mol%)	temp (°C)	4 (yield)	4 : 5 (mol ratio)
1	5	5	60	55%	55 : 1
2	10	5	60	30%	30 : 1
3	5	10	60	93%	13 : 1
4	5	10	40	80%	40 : 1
5	5	10	100	91%	7 : 1

^a All reactions were run using 0.25 mmol of benzenesulfonamide and 0.75 mmol of 4-methylstyrene in toluene.

Table 4

Hydroamination of various vinyl arenes with benzenesulfonamide.

entry	vinyl arene	<i>p</i> -TsOH (mol%)	solvent	temp (°C)	products	yield 7 (%)	ratio 7 : 8
1	4-methylstyrene (6a)	10	toluene	60	7a, 8a	93	13 : 1
2	2,4-dimethylstyrene (6b)	10	toluene	60	7b, 8b	99	9 : 1
3	4-methoxystyrene ^a (6c)	5	toluene	40	7c, 8c	70	7.8 : 1
4	3-methoxystyrene (6d)	15	toluene	100	7d, 8d	73	12 : 1
5	styrene (6e)	15	toluene	100	7e, 8e	84	9.3 : 1
6	2-vinylnaphthalene (6f)	10	toluene	100	7f, 8f	57	14 : 1
7	4-fluorostyrene (6g)	15	toluene	100	7g, 8g	95	9.5 : 1
8	4-chlorostyrene (6h)	15	toluene	100	7h, 8h	83	28 : 1
9	4-bromostyrene (6i)	15	toluene	100	7i, 8i	87	15 : 1
10	2-fluorostyrene (6j)	20	chlorobenzene	130	7j, 8j	70	23 : 1
11	2-bromostyrene (6k)	20	chlorobenzene	130	7k, 8k	63	16 : 1
12	2-chlorostyrene ^b (6l)	20	chlorobenzene	130	7l, 8l	31	4.4 : 1
13	3-chlorostyrene (6m)	20	chlorobenzene	130	7m, 8m	41	8.2 : 1
14	cis- -methylstyrene (6n)	20	chlorobenzene	130	7n, 8n	13	13 : 1

^a 10% catalyst loading was required.^b Significant polymerization of starting material was observed.