Cardamonin Inhibits Osteoclastogenesis Induced by Tumor Cells Through Interruption of the Signaling Pathway Activated by Receptor Activator of NF- κ B Ligand

Vivek R. Yadav, Sahdeo Prasad, Simone Reuter, Bokyoung Sung, and Bharat B. Aggarwal
Cytokine Research Laboratory, Department of Experimental Therapeutics, The University of Texas MD Anderson Cancer Center, Houston, Texas

Abstract

Bone loss/resorption or osteoporosis is a disease that is accelerated with aging and age-associated chronic diseases such as cancer. Bone loss has been associated with human multiple myeloma, breast cancer, and prostate cancer and is usually treated with a bisphosphonate. Because of the numerous side effects of the currently available drugs, the search continues for safe and effective therapies for bone loss. Recently, receptor activator of NF- κ B ligand (RANKL), a member of the TNF superfamily, has emerged as a major mediator of bone loss via activation of osteoclastogenesis. We have identified cardamonin, a chalcone first isolated from grass cardamom (*Alpinia katsumadai* Hayata), that can affect osteoclastogenesis through modulation of RANKL. We found that treatment of monocytes with cardamonin suppressed RANKL-induced NF- κ B activation and this suppression correlated with inhibition of I κ B kinase and of phosphorylation and degradation of I κ B, an inhibitor of NF- κ B. Cardamonin suppressed the differentiation of monocytes to osteoclasts in a dose-dependent and time-dependent manner. We also found that an NF- κ B-specific inhibitory peptide blocked RANKL-induced osteoclastogenesis, indicating a direct link with NF- κ B. Finally, osteoclastogenesis induced by human breast cancer cells or human multiple myeloma cells was completely suppressed by cardamonin. Collectively, our results indicate that cardamonin suppresses osteoclastogenesis induced by RANKL and tumor cells by suppressing activation of the NF- κ B pathway.

Keywords

Osteoclastogenesis; RANKL; NF- κ B; Cancer; Cardamonin

1. Introduction

Bone undergoes constant turnover and is kept in balance (homeostasis) by osteoblasts (creating bone) and osteoclasts (destroying bone). The osteoclast is a unique bone-resorbing cell derived from cells of monocyte-macrophage lineage. Osteoclastogenesis comprises many stages, including commitment, differentiation, multinucleation, and activation of

© 2011 Elsevier Ireland Ltd. All rights reserved.

*To whom correspondence should be addressed: Bharat B. Aggarwal, The University of Texas M. D. Anderson Cancer Center, 1515 Holcombe Blvd., Box 143, Houston, TX 77030. Phone: 713-794-1817; Fax: 713-745-6339; aggarwal@mdanderson.org.

Conflict of interest: The authors have no conflicts of interest related to this manuscript.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

immature osteoclasts. Within the bone microenvironment, T and B lymphocytes, bone marrow stromal cells, macrophages, and osteoblasts all can produce cytokines that have an impact on osteoclastogenesis. A variety of both systemic hormones and cytokines regulate osteoclast differentiation and function [1].

Receptor activator of nuclear factor- κ B ligand (RANKL), a member of the tumor necrosis factor (TNF) superfamily, has emerged as a major osteoclastogenic cytokine. Binding of RANKL to its receptor, RANK, on macrophages prompts them to assume the osteoclast phenotype [2; 3]. RANKL is expressed on the surface of osteoblastic/stromal cells and by various types of cancer cells, and is directly involved in the differentiation of monocyte macrophages into osteoclasts [4]. When RANKL binds to RANK, it undergoes trimerization and then binds to an adaptor molecule, TNF receptor-associated factor 6 (TRAF6), which then sequentially activates I κ B kinase (IKK), nuclear factor- κ B (NF- κ B), and nuclear factor of activated T cells, cytoplasm 1 (NFATc1), thus leading to osteoclastogenesis [5; 6].

Several types of cancer, both solid and hematopoietic, are deeply linked with the skeleton and cause an increase in osteoclast formation, either systemically, as in humoral hypercalcemia of malignancy, or locally, as in bone metastasis. Bone metastasis makes bone more fragile and leads to pathologic fractures and spinal compression. This osteolysis is associated with severe bone pain, which may be intractable. Bone metastasis represents a common cause of morbidity in patients with many types of cancer, occurring in as many as 70% of patients with advanced breast or prostate cancer and in about 15% to 30% of patients with lung, colon, kidney, thyroid, or stomach carcinoma [7; 8]. It is more common in patients with advanced multiple myeloma or breast, prostate, or lung cancer, as these tumors have a remarkable propensity to metastasize to the bone [9; 10].

RANKL has been shown to play a major role in bone metastasis [9], and thus is an important therapeutic target. Agents that suppress RANKL signaling have potential for inhibition of osteoclastogenesis and bone metastasis. As reported here, we investigated whether cardamonin, a chalcone isolated from grass cardamom (*A. katsumadai* Hayata) and black cardamom (*Amomum subulatum*), can modulate RANKL-induced signaling and osteoclastogenesis. We found that cardamonin can suppress RANKL-induced NF- κ B activation through inhibition of IKK and inhibits osteoclastogenesis induced by RANKL and by breast cancer and multiple myeloma cells.

2. Materials and Methods

2.1. Reagents

Cardamonin was isolated and supplied by Dr. Akira Murakami (Kyoto University, Kyoto, Japan) was prepared as 20 mM solution in dimethylsulfoxide and then further diluted in cell culture medium [11]. Dulbecco modified essential medium (DMEM)/F12, RPMI 1640, DMEM, fetal bovine serum, 0.4% trypan blue vital stain, and antibiotic-antimycotic mixture were obtained from Invitrogen (Carlsbad, CA). RANKL protein was kindly provided by Dr. Bryant Darnay of The University of Texas MD Anderson Cancer Center (Houston, TX). Antibodies against IKK α , IKK β and I κ B were purchased from Imgenex (San Diego, CA), while cell-permeable NF- κ B essential modulator (NEMO; also called IKK γ)-binding domain peptide (NBP) were kind gifts from Imgenex. Antibody against phospho-I κ B (Ser32/36) was purchased from Cell Signaling Technology (Danvers, MA). Goat anti-rabbit and goat anti-mouse horseradish peroxidase conjugates were purchased from Bio-Rad (Hercules, CA). α -actin antibody and leukocyte acid phosphatase kit (387-A) for tartrate-resistant acid phosphatase (TRAP) staining were purchased from Sigma-Aldrich (St. Louis, MO). Protein A/G-agarose beads were obtained from ThermoScientific (Rockford, IL). [γ - ^{32} P]ATP was purchased from MP Biomedicals (Solon, OH).

2.2. Cell lines

RAW 264.7 (mouse macrophage) cells were kindly provided by Dr. Bryant Darnay. For these studies, we used a single clone (#28) that has been selected after limited dilution. RAW 264.7 cells were cultured in DMEM/F12 supplemented with 10% fetal bovine serum and antibiotics. This cell line is a well-established osteoclastogenic cell system that has been shown to express RANK and to differentiate into functional TRAP-positive osteoclasts when cultured with soluble RANKL [12]. Moreover, RANKL has been shown to activate NF- κ B in RAW 264.7 cells [13]. MDA-MB-231 (human breast adenocarcinoma) and U266 (human multiple myeloma) cells were obtained from the American Type Culture Collection (Manassas, VA, USA). MDA-MB-231 cells were cultured in DMEM and U266 cells in RPMI 1640 with 10% fetal bovine serum.

2.3. Osteoclast differentiation assay

RAW 264.7 cells were cultured in 24-well plates at a density of 1×10^4 per well and allowed to adhere overnight. The medium was then replaced, and the cells were treated with 5 nM RANKL for 5 days. All cells were subjected to TRAP staining using the leukocyte acid phosphatase kit. For co-culture experiments with cancer cells, RAW 264.7 cells were seeded at 5×10^3 per well and allowed to adhere overnight. The following day, U266 or MDA-MB-231 cells, at 1×10^3 per well, were added to the RAW 264.7 cells, treated with cardamonin, and co-cultured for 5 days before being subjected to TRAP staining.

2.4. Electrophoretic mobility shift assay (EMSA) for NF- κ B

Nuclear extracts were prepared as described previously [14]. Briefly, nuclear extracts from RANKL-treated cells were incubated with ^{32} P-end-labeled 45-mer double-stranded NF- κ B oligonucleotide (15 μ g protein with 16 fmol DNA) from the HIV long terminal repeat, 5'-TTGTTACAAGGGACTTTCCGCTGGGGACTTTCCAGGGGGAGGCGTGG-3' (boldface indicates NF- κ B-binding sites), for 30 min at 37°C, and the DNA-protein complex formed was separated from free oligonucleotide on 6.6% native polyacrylamide gels. The dried gels were visualized with a Storm820, and radioactive bands were quantified using a densitometer and Image Quant software (GE Healthcare, Piscataway, NJ).

2.5. Western blot analysis

To determine the levels of protein expression in the cytoplasm and nucleus, we prepared extracts [14] and fractionated them by 10% sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE). After electrophoresis, the proteins were electrotransferred to nitrocellulose membranes, blotted with each antibody, and detected with a chemiluminescence reagent (GE Healthcare).

2.6. IKK assay

To determine the effect of cardamonin on RANKL-induced IKK activation, IKK assay was performed by a method described previously [15]. Briefly, the IKK complex from whole-cell extracts (600 μ g protein) was precipitated with antibody against IKK, followed by treatment with protein A/G-agarose beads. After 2 h of incubation, the beads were washed with lysis buffer and assayed in a kinase assay mixture containing 50 mM HEPES (pH 7.4), 20 mM MgCl₂, 2 mM dithiothreitol, 20 μ Ci [γ - ^{32} P]ATP, 10 μ M unlabeled ATP, and 2 μ g of substrate glutathione S-transferase (GST)-I κ B (amino acids 1–54). After incubation at 30°C for 30 min, the reaction was terminated by boiling with SDS sample buffer for 5 min. Finally, the protein was resolved on 10% SDS-PAGE, the gel was dried, and the radioactive bands were visualized with a PhosphorImager. To determine the total amounts of IKK α and IKK β in each sample, the whole-cell protein was resolved on 10% SDS-PAGE,

electrotransferred to a nitrocellulose membrane, and blotted with anti-IKK or anti-IKK antibody.

2.7. Trypan blue exclusion assay

Cells were harvested from the 24-well plates by treatment with 0.2% trypsin-EDTA, centrifuged, and suspended in one ml culture medium. Cell suspension was mixed with equal volume of 0.4% isotonic trypan blue solution. Total cell number and fraction of nonviable, dye-accumulating cells were counted after 2 min in Fuchs-Rosenthal hemocytometer under light microscope.

3. Results

The aim of the study presented here was to investigate the effect of cardamonin (Fig. 1A) on RANKL signaling and on osteoclastogenesis. Whether cardamonin could modulate osteoclastogenesis induced by tumor cells was another focus of these studies. We used the RAW 264.7 cell (murine macrophage) system because it is a well-established model for osteoclastogenesis and contains no osteoblast/bone marrow stromal cells or cytokine-like macrophage colony-stimulating factor, thus allowing us to focus on RANKL signaling in preosteoclast cells [16].

3.1. Cardamonin inhibits RANKL-induced osteoclastogenesis

The effect of cardamonin on RANKL-induced osteoclastogenesis was examined. RAW 264.7 cells were incubated with different concentrations of cardamonin in the presence of RANKL and allowed to differentiate into osteoclasts. As shown in Figure 1B, RANKL induced formation of osteoclasts in the absence of cardamonin at day 5. In contrast, differentiation into osteoclasts was significantly decreased in the presence of cardamonin. Moreover, the formation of osteoclasts decreased with increasing concentration of cardamonin (Fig. 1B and 1C). As little as 50 nM cardamonin had a significant effect on RANKL-induced osteoclast formation. Under these conditions, the viability of cells was not significantly affected (Fig. 1D).

RAW 264.7 cells were incubated with cardamonin and RANKL for 3, 4, or 5 days and allowed to differentiate into osteoclasts. Morphological observations clearly demonstrated that RAW 264.7 cells differentiated into osteoclasts after RANKL addition, and that cardamonin inhibited this differentiation (Fig. 2A). The extent of suppression was measured by counting the number of TRAP-positive osteoclasts per well (Fig. 2B). We found that RANKL induced osteoclast differentiation in a time-dependent manner, with a maximum of TRAP-positive osteoclasts per well at day 5 (Fig. 2A and 2B). On the other hand, cardamonin decreased the number of TRAP-positive osteoclasts in a dose-dependent manner, with a strong inhibition at 500 nM at all days examined (Fig. 2B).

3.2. Cardamonin acts at an early step in the pathway leading to RANKL-induced osteoclastogenesis

It normally takes as long as 5 days for RAW 264.7 cells to differentiate into osteoclasts in response to RANKL. To elucidate at what point in this pathway cardamonin acts, RAW 264.7 cells were treated initially with RANKL, cardamonin was added 1, 2, 3, and 4 days after RANKL treatment, and then its effect on osteoclast formation was determined (Fig. 3A). As determined by observation (Fig. 3A, right panel) and by counting of TRAP-positive osteoclasts per well (Fig. 3B), cardamonin markedly inhibited osteoclast formation when the cells were exposed to the compound for 1 or 2 days after RANKL stimulation. By days 3 and 4 after RANKL addition, however, osteoclast formation was no longer completely

prevented by cardamonin (Fig. 3A and 3B), indicating that cardamonin probably acts at an early step in the osteoclast differentiation pathway.

3.3 Cardamonin inhibits osteoclastogenesis induced by tumor cells

Osteoclastogenesis is commonly linked with breast cancer [17] and multiple myeloma [18] through activation of NF- κ B [19]. Whether cardamonin also inhibits tumor cell-induced osteoclastogenesis of RAW 264.7 cells was investigated. As shown in Figures 4A and 4B, incubating RAW 264.7 cells with MDA-MB-231 breast cancer cells or U266 multiple myeloma cells induced osteoclast differentiation, and cardamonin suppressed this differentiation. The results presented in Figures 4A and 4B indicate that certain cytokines secreted by tumor cells are responsible for osteoclast differentiation. These results indicate that osteoclastogenesis induced by tumor cells are significantly suppressed by the presence of cardamonin.

3.4. Cardamonin abrogates RANKL-induced NF- κ B activation

To determine the concentration of cardamonin required to suppress RANKL-induced NF- κ B activation, cells were pretreated with various concentrations of cardamonin and then exposed to RANKL. The chalcone almost completely suppressed NF- κ B activation at 20 μ M (Fig. 5A). Treatment of cells with 20 μ M of cardamonin for 12 h had no effect on cell viability as determined by the trypan blue exclusion method.

To investigate whether cardamonin modulates RANKL-induced NF- κ B activation in RAW 264.7 cells, cells were either pretreated with cardamonin for 12 h or left untreated and then exposed to RANKL for indicated times, nuclear extracts were prepared, and NF- κ B activation was assayed by EMSA. As shown in Figure 5B, RANKL activated NF- κ B in a time-dependent manner; however, cardamonin completely abrogated RANKL-induced NF- κ B activation.

3.5. Cardamonin inhibits RANKL-induced I κ B α phosphorylation and degradation

Because the translocation of NF- κ B to the nucleus requires proteolytic degradation of I κ B, we next sought to determine whether cardamonin-induced NF- κ B inhibition was due to inhibition of I κ B degradation. Therefore, we examined NF- κ B activation in the nucleus by EMSA (Fig. 5B) and I κ B degradation in the cytoplasm by western blot (Fig. 6A), after RANKL stimulation for various times. As shown in Figure 5B, RANKL activated NF- κ B. Results, western blot analysis showed that RANKL induced I κ B degradation in control cells after 5 min and returned to normal level within 60 min (Fig. 6A). On the other hand, cells pretreated with cardamonin showed no degradation of I κ B.

Given that I κ B phosphorylation is necessary for I κ B degradation, we next investigated the effect of cardamonin on I κ B phosphorylation by using the proteasome inhibitor *N*-acetyl-leu-leu-norleucinal (ALLN), which prevents RANKL-induced I κ B degradation (Fig. 6B). Western blot analysis showed that co-treatment with RANKL plus ALLN induced phosphorylation of I κ B at serines 32 and 36, and that cardamonin pretreatment inhibited this activation in RAW 264.7 cells. Only RANKL alone degraded I κ B protein. These results clearly indicate that cardamonin inhibits both RANKL-induced NF- κ B activation as well as I κ B degradation and phosphorylation.

3.6. Cardamonin inhibits RANKL-induced IKK activation

Because cardamonin inhibits the phosphorylation and degradation of I κ B, we next checked whether cardamonin alters the activity or level of IKK, which leads to I κ B phosphorylation. Immunocomplex kinase assay on cells treated with RANKL showed a sharp increase in IKK activity as indicated by the phosphorylation of GST-I κ B within 2

min. In contrast, cells pretreated with cardamonin could not phosphorylate GST-I β after treatment with RANKL (Fig. 6C). To check whether the apparent loss of IKK activity was due to the loss of IKK protein expression, the levels of the IKK subunits IKK α and IKK β were tested by western blot analysis. Results in Figure 6C clearly show that cardamonin treatment did not alter the expression of IKK α and IKK β proteins.

3.7. Inhibition of osteoclastogenesis by cardamonin is NF- κ B specific

To ascertain the specificity of NF- κ B in osteoclastogenesis, we used a specific inhibitor of the regulatory subunit of the IKK complex, IKK γ , also known as the NF- κ B essential modulator (NEMO). While the serine kinases IKK α and IKK β target serines 32 and 36 of the I β protein, NEMO regulates the IKK complex activity through its binding to the carboxyl-terminal region of the IKK α and IKK β subunits, called the NEMO-binding domain. In this regard, a cell-permeable peptide that blocks the NEMO-binding domain would inhibit the association of NEMO with the IKK complex and consequently suppress NF- κ B activation and, most likely, osteoclastogenesis.

To determine the effect of the NEMO-binding domain peptide (NBP) on RANKL-induced osteoclastogenesis, we pretreated RAW 264.7 cells with 100 μ M of NBP for 2 h and then treated the cells with RANKL for 5 days. Our results show that the peptide, which targets the NEMO-binding domain and thus inhibits the IKK complex activity, strongly inhibited osteoclast differentiation (Fig. 7A and 7B). Furthermore, when we treated nuclear extracts from RAW 264.7 cells with 100 μ M NBP for 2 h and then with RANKL for 30 min, RANKL-induced NF- κ B activation was completely inhibited (Fig. 7C). These results confirm that NF- κ B is responsible for osteoclast differentiation of RAW 264.7 cells, and that inhibition of NF- κ B by either cardamonin or NBP prevents osteoclastogenesis.

4. Discussion

Osteoclasts, which are responsible for bone resorption, are rare; each 1 mm³ of bone contains only two or three osteoclasts. Despite these small numbers, loss of function of osteoclasts or decrease in their number leads to osteosclerosis/osteopetrosis. On the other hand, an increase in their number or function induces bone osteoporosis, indicating that osteoclasts play a pivotal role in bone loss [20; 21]. Systemic hormones and cytokines provide the molecular cues that control osteoclastogenesis and thus maintain homeostasis [1]. Major problems associated with aging, such as arthritis, cancer, and other chronic inflammatory illnesses, disturb this balance, however, and safe, efficacious, and affordable compounds that can inhibit bone loss are needed. Cardamonin is one such compound that has been shown to suppress inflammatory pathways. The goal of this study was to investigate the effect of cardamonin, a bioactive chalcone, on RANKL-induced NF- κ B activation and on osteoclastogenesis induced by both RANKL and tumor cells.

A number of inflammatory cytokines produced during different types of inflammation have a synergic role in osteoclastogenesis, include interleukins 1 and 6, TNF- α , and oncostatin M. These have been reported to stimulate osteoclastic differentiation and bone resorption [22; 23; 24]. Chemokines are chemotactic signals for monocytes that can facilitate the fusion of monocytes into multinucleated osteoclasts [25]. Our results indicate that RANKL activates NF- κ B in osteoclast precursor cells through activation of IKK and subsequent I β phosphorylation and degradation. The RANKL-induced NF- κ B activation signaling pathway differs from the TNF-induced pathway. For instance, NIK, which may function as an activator of IKK α , is necessary in RANKL-induced NF- κ B activation [13]; however, it is dispensable for TNF-induced NF- κ B activation [26]. Thus NIK-deficient osteoclast precursors have been reported to not respond to RANKL in an *in vitro* differentiation system

devoid of osteoblasts [27]. We found that cardamonin inhibited RANKL-induced IKK activation, leading to suppression of NF- κ B activation.

Other studies have shown that IKK α , but not IKK β , is a potent regulator of inflammation-induced bone loss and is required for osteoclastogenesis and inflammatory arthritis [28]. The study reported here is the first showing that cardamonin can suppress RANKL-induced IKK activation and consequently NF- κ B activation. How cardamonin inhibits RANKL-induced IKK activation is not clear.

By recruiting the adapter proteins TRAF 2, 3, 5, and 6 and NIK, RANK activates NF- κ B and the JNK, p38 MAPK, and p44/p42 MAPK signaling pathways [5; 12]. In addition to its role in inflammatory diseases, the NF- κ B signaling pathway has been demonstrated to be a major mediator of bone loss [29]. It is already proven that NF- κ B p50 and p52 expression are essential for the differentiation of RANK-expressing-osteoclast precursors into TRAP-positive osteoclasts in response to RANKL and other osteoclastogenic cytokines [30].

RANKL has been shown to play a major role in cancer-associated osteoclast differentiation. Furthermore, a series of electrolytes and degradative enzymes have been implicated in osteoclastogenesis, bone resorption, and calcium homeostasis [4; 31]. Mice deficient in the *rankl* gene have been shown to display severe osteopetrosis, stunted growth, defective tooth eruption, and osteoblasts that cannot support osteoclastogenesis [31]. Thus agents that can inhibit RANKL signaling have a great potential for inhibiting osteoclastogenesis.

Our results indicate that cardamonin effectively inhibits RANKL-induced osteoclastogenesis. A kinetic study indicated that cardamonin acts at an early step in the osteoclast differentiation process. To further confirm that inhibition of the NF- κ B signaling pathway is responsible for arrest of the osteoclastogenesis process in our system, we used a cell-permeable peptide that targets the NEMO-binding domain of the IKK α and IKK β kinases and so prevents NF- κ B activation. This NEMO-binding domain peptide has been shown to inhibit osteoclastogenesis *in vivo*, and also delayed the onset, lowered the incidence, and decreased the severity of rheumatoid arthritis [28]. Moreover, previous studies demonstrated that pharmacological or genetic inactivation of IKK α and/or IKK β is sufficient for inhibition of osteoclastogenesis and prevention of inflammation and osteolytic bone loss [32; 33]. Our results show that the NF- κ B inhibitor NEMO-binding domain peptide completely blocked RANKL-induced osteoclastogenesis in the same manner as cardamonin. Interestingly, the inhibitory effect of 100 μ M of this peptide was as potent as 500 nM cardamonin, which suggests that cardamonin is 200 times more potent, at least *in vitro*. These findings indicate that cardamonin's inhibitory effect on osteoclastogenesis is probably specific to NF- κ B inhibition.

A major health problem today that affects over 350,000 patients in the United States annually is malignant tumors of skeleton as the primary site as well as metastatic bone lesions. Among them, 70% to 95% of multiple myeloma patients and up to 75% of patients with advanced breast cancer or prostate cancer develop bone metastasis. A major complication in metastatic breast cancer and multiple myeloma is osteoclast-mediated bone destruction [7; 34]. Breast cancers commonly cause osteolytic metastasis that depends on osteoclast-mediated bone resorption [35], but the mechanism responsible for this has not yet been clarified. We showed in this study that osteoclastogenesis induced by breast cancer cells inhibited by cardamonin. Bone resorption is also associated with multiple myeloma [35], and we found that multiple myeloma cell-induced osteoclastogenesis was also suppressed by cardamonin. As IKK activation is known to accelerate the proliferation and metastasis of cancer cells [36; 37], its inhibitors, such as cardamonin, might have potential in the treatment of cancers that metastasize to the bone. These tumors have been shown to

express RANKL [18; 38] and exhibit constitutive NF- κ B activation [39; 40]. Thus, it can be concluded that these tumors activate osteoclastogenesis through RANKL expression.

The bisphosphonates are the only drugs now available for treatment of bone metastasis or cancer-related bone diseases. These drugs are highly toxic, however, and adverse effects such as renal impairment or osteonecrosis of the jaw have been reported [41]. An antibody to RANKL, denosumab (Prolia), has been approved recently for treatment of osteoporosis [42]. Cardamonin is derived from the seed of grass cardamom (*A. katsumadai* Hayata) and the fruit of black cardamom (*A. subulatum*), and should have minimum toxicity, as it is used routinely for traditional medicine [11; 43]. Thus cardamonin could be used in the treatment of secondary bone lesions associated with cancer and also nonmalignant diseases such as postmenopausal osteoporosis, Paget disease, and rheumatoid arthritis. Further studies are needed, however, to confirm whether cardamonin can suppress osteoclastogenesis *in vivo*.

Acknowledgments

We thank Kathryn Hale for editorial review of this manuscript and Dr. Bryant Darnay for providing RAW 264.7 cells and RANKL protein. Dr. Aggarwal is the Ransom Horne, Jr., Professor of Cancer Research. This work was supported by supported by a core grant (CA-16672), and a program project grant from National Institutes of Health (NIH CA-124787-01A2), and a grant from the Center for Targeted Therapy of MD Anderson Cancer Center.

References

1. Yavropoulou MP, Yovos JG. Osteoclastogenesis--current knowledge and future perspectives. *J Musculoskelet Neuronal Interact.* 2008; 8:204–216. [PubMed: 18799853]
2. Anderson DM, Maraskovsky E, Billingsley WL, Dougall WC, Tometsko ME, Roux ER, Teepe MC, DuBose RF, Cosman D, Galibert L. A homologue of the TNF receptor and its ligand enhance T-cell growth and dendritic-cell function. *Nature.* 1997; 390:175–179. [PubMed: 9367155]
3. Jin W, Chang M, Paul EM, Babu G, Lee AJ, Reiley W, Wright A, Zhang M, You J, Sun SC. Deubiquitinating enzyme CYLD negatively regulates RANK signaling and osteoclastogenesis in mice. *J Clin Invest.* 2008; 118:1858–1866. [PubMed: 18382763]
4. Lacey DL, Timms E, Tan HL, Kelley MJ, Dunstan CR, Burgess T, Elliott R, Colombero A, Elliott G, Scully S, Hsu H, Sullivan J, Hawkins N, Davy E, Capparelli C, Eli A, Qian YX, Kaufman S, Sarosi I, Shalhoub V, Senaldi G, Guo J, Delaney J, Boyle WJ. Osteoprotegerin ligand is a cytokine that regulates osteoclast differentiation and activation. *Cell.* 1998; 93:165–176. [PubMed: 9568710]
5. Darnay BG, Haridas V, Ni J, Moore PA, Aggarwal BB. Characterization of the intracellular domain of receptor activator of NF- κ B (RANK). Interaction with tumor necrosis factor receptor-associated factors and activation of NF- κ B and c-Jun N-terminal kinase. *J Biol Chem.* 1998; 273:20551–20555. [PubMed: 9685412]
6. Takayanagi H, Kim S, Koga T, Nishina H, Isshiki M, Yoshida H, Saiura A, Isobe M, Yokochi T, Inoue J, Wagner EF, Mak TW, Kodama T, Taniguchi T. Induction and activation of the transcription factor NFATc1 (NFAT2) integrate RANKL signaling in terminal differentiation of osteoclasts. *Dev Cell.* 2002; 3:889–901. [PubMed: 12479813]
7. Coleman RE. Metastatic bone disease: clinical features, pathophysiology and treatment strategies. *Cancer Treat Rev.* 2001; 27:165–176. [PubMed: 11417967]
8. Coleman RE. Clinical features of metastatic bone disease and risk of skeletal morbidity. *Clin Cancer Res.* 2006; 12:6243s–6249s. [PubMed: 17062708]
9. Onishi T, Hayashi N, Theriault RL, Hortobagyi GN, Ueno NT. Future directions of bone-targeted therapy for metastatic breast cancer. *Nat Rev Clin Oncol.* 2010; 7:641–651. [PubMed: 20808302]
10. Sturge J, Caley MP, Waxman J. Bone metastasis in prostate cancer: emerging therapeutic strategies. *Nat Rev Clin Oncol.* 2011; 8:357–368. [PubMed: 21556025]
11. Takahashi A, Yamamoto N, Murakami A. Cardamonin suppresses nitric oxide production via blocking the IFN- γ /STAT pathway in endotoxin-challenged peritoneal macrophages of ICR mice. *Life Sci.* 2011; 89:337–342. [PubMed: 21784080]

12. Hsu H, Lacey DL, Dunstan CR, Solovyev I, Colombero A, Timms E, Tan HL, Elliott G, Kelley MJ, Sarosi I, Wang L, Xia XZ, Elliott R, Chiu L, Black T, Scully S, Capparelli C, Morony S, Shimamoto G, Bass MB, Boyle WJ. Tumor necrosis factor receptor family member RANK mediates osteoclast differentiation and activation induced by osteoprotegerin ligand. *Proc Natl Acad Sci U S A*. 1999; 96:3540–3545. [PubMed: 10097072]
13. Wei S, Teitelbaum SL, Wang MW, Ross FP. Receptor activator of nuclear factor-kappa b ligand activates nuclear factor-kappa b in osteoclast precursors. *Endocrinology*. 2001; 142:1290–1295. [PubMed: 11181547]
14. Sung B, Pandey MK, Aggarwal BB. Fisetin, an inhibitor of cyclin-dependent kinase 6, down-regulates nuclear factor-kappaB-regulated cell proliferation, antiapoptotic and metastatic gene products through the suppression of TAK-1 and receptor-interacting protein-regulated IkappaBalpha kinase activation. *Mol Pharmacol*. 2007; 71:1703–1714. [PubMed: 17387141]
15. Reuter S, Prasad S, Phromnoi K, Kannappan R, Yadav VR, Aggarwal BB. Embelin suppresses osteoclastogenesis induced by receptor activator of NF-kappaB ligand and tumor cells in vitro through inhibition of the NF-kappaB cell signaling pathway. *Mol Cancer Res*. 2010; 8:1425–1436. [PubMed: 20826545]
16. Kobayashi Y, Mizoguchi T, Take I, Kurihara S, Udagawa N, Takahashi N. Prostaglandin E2 enhances osteoclastic differentiation of precursor cells through protein kinase A-dependent phosphorylation of TAK1. *J Biol Chem*. 2005; 280:11395–11403. [PubMed: 15647289]
17. Chikatsu N, Takeuchi Y, Tamura Y, Fukumoto S, Yano K, Tsuda E, Ogata E, Fujita T. Interactions between cancer and bone marrow cells induce osteoclast differentiation factor expression and osteoclast-like cell formation in vitro. *Biochem Biophys Res Commun*. 2000; 267:632–637. [PubMed: 10631114]
18. Lai FP, Cole-Sinclair M, Cheng WJ, Quinn JM, Gillespie MT, Sentry JW, Schneider HG. Myeloma cells can directly contribute to the pool of RANKL in bone bypassing the classic stromal and osteoblast pathway of osteoclast stimulation. *Br J Haematol*. 2004; 126:192–201. [PubMed: 15238139]
19. Park BK, Zhang H, Zeng Q, Dai J, Keller ET, Giordano T, Gu K, Shah V, Pei L, Zarbo RJ, McCauley L, Shi S, Chen S, Wang CY. NF-kappaB in breast cancer cells promotes osteolytic bone metastasis by inducing osteoclastogenesis via GM-CSF. *Nat Med*. 2007; 13:62–69. [PubMed: 17159986]
20. Marie PJ. Human endosteal osteoblastic cells: relationship with bone formation. *Calcif Tissue Int*. 1995; 56(1):S13–16. [PubMed: 7719974]
21. Miyamoto T, Suda T. Differentiation and function of osteoclasts. *Keio J Med*. 2003; 52:1–7. [PubMed: 12713016]
22. Hanazawa S, Amano S, Nakada K, Ohmori Y, Miyoshi T, Hirose K, Kitano S. Biological characterization of interleukin-1-like cytokine produced by cultured bone cells from newborn mouse calvaria. *Calcif Tissue Int*. 1987; 41:31–37. [PubMed: 3113699]
23. Ishimi Y, Miyaura C, Jin CH, Akatsu T, Abe E, Nakamura Y, Yamaguchi A, Yoshiki S, Matsuda T, Hirano T, et al. IL-6 is produced by osteoblasts and induces bone resorption. *J Immunol*. 1990; 145:3297–3303. [PubMed: 2121824]
24. Zou W, Bar-Shavit Z. Dual modulation of osteoclast differentiation by lipopolysaccharide. *J Bone Miner Res*. 2002; 17:1211–1218. [PubMed: 12096834]
25. Kim MS, Magno CL, Day CJ, Morrison NA. Induction of chemokines and chemokine receptors CCR2b and CCR4 in authentic human osteoclasts differentiated with RANKL and osteoclast like cells differentiated by MCP-1 and RANTES. *J Cell Biochem*. 2006; 97:512–518. [PubMed: 16211583]
26. Uhlik M, Good L, Xiao G, Harhaj EW, Zandi E, Karin M, Sun SC. NF-kappaB-inducing kinase and IkappaB kinase participate in human T-cell leukemia virus I Tax-mediated NF-kappaB activation. *J Biol Chem*. 1998; 273:21132–21136. [PubMed: 9694868]
27. Novack DV, Yin L, Hagen-Stapleton A, Schreiber RD, Goeddel DV, Ross FP, Teitelbaum SL. The IkappaB function of NF-kappaB2 p100 controls stimulated osteoclastogenesis. *J Exp Med*. 2003; 198:771–781. [PubMed: 12939342]

28. Dai S, Hirayama T, Abbas S, Abu-Amer Y. The IkappaB kinase (IKK) inhibitor, NEMO-binding domain peptide, blocks osteoclastogenesis and bone erosion in inflammatory arthritis. *J Biol Chem.* 2004; 279:37219–37222. [PubMed: 15252035]
29. Xu J, Wu HF, Ang ES, Yip K, Woloszyn M, Zheng MH, Tan RX. NF-kappaB modulators in osteolytic bone diseases. *Cytokine Growth Factor Rev.* 2009; 20:7–17. [PubMed: 19046922]
30. Xing L, Bushnell TP, Carlson L, Tai Z, Tondravi M, Siebenlist U, Young F, Boyce BF. NF-kappaB p50 and p52 expression is not required for RANK-expressing osteoclast progenitor formation but is essential for RANK- and cytokine-mediated osteoclastogenesis. *J Bone Miner Res.* 2002; 17:1200–1210. [PubMed: 12096833]
31. Kong YY, Yoshida H, Sarosi I, Tan HL, Timms E, Capparelli C, Morony S, Oliveira-dos-Santos AJ, Van G, Itie A, Khoo W, Wakeham A, Dunstan CR, Lacey DL, Mak TW, Boyle WJ, Penninger JM. OPGL is a key regulator of osteoclastogenesis, lymphocyte development and lymph-node organogenesis. *Nature.* 1999; 397:315–323. [PubMed: 9950424]
32. Chaisson ML, Branstetter DG, Derry JM, Armstrong AP, Tometsko ME, Takeda K, Akira S, Dougall WC. Osteoclast differentiation is impaired in the absence of inhibitor of kappa B kinase alpha. *J Biol Chem.* 2004; 279:54841–54848. [PubMed: 15485831]
33. Ruocco MG, Maeda S, Park JM, Lawrence T, Hsu LC, Cao Y, Schett G, Wagner EF, Karin M. I{kappa}B kinase (IKK) {beta}, but not IKK {alpha}, is a critical mediator of osteoclast survival and is required for inflammation-induced bone loss. *J Exp Med.* 2005; 201:1677–1687. [PubMed: 15897281]
34. Carlin BI, Andriole GL. The natural history, skeletal complications, and management of bone metastases in patients with prostate carcinoma. *Cancer.* 2000; 88:2989–2994. [PubMed: 10898342]
35. Mundy GR. Metastasis to bone: causes, consequences and therapeutic opportunities. *Nat Rev Cancer.* 2002; 2:584–593. [PubMed: 12154351]
36. Jourdan M, Moreaux J, Vos JD, Hose D, Mahtouk K, Abouladze M, Robert N, Baudard M, Reme T, Romanelli A, Goldschmidt H, Rossi JF, Dreano M, Klein B. Targeting NF-kappaB pathway with an IKK2 inhibitor induces inhibition of multiple myeloma cell growth. *Br J Haematol.* 2007; 138:160–168. [PubMed: 17542984]
37. Kishida Y, Yoshikawa H, Myoui A. Parthenolide, a natural inhibitor of Nuclear Factor-kappaB, inhibits lung colonization of murine osteosarcoma cells. *Clin Cancer Res.* 2007; 13:59–67. [PubMed: 17200339]
38. Martin TJ, Gillespie MT. Receptor activator of nuclear factor kappa B ligand (RANKL): another link between breast and bone. *Trends Endocrinol Metab.* 2001; 12:2–4. [PubMed: 11137033]
39. Bharti AC, Takada Y, Aggarwal BB. Curcumin (diferuloylmethane) inhibits receptor activator of NF-kappa B ligand-induced NF-kappa B activation in osteoclast precursors and suppresses osteoclastogenesis. *J Immunol.* 2004; 172:5940–5947. [PubMed: 15128775]
40. Biswas DK, Shi Q, Baily S, Strickland I, Ghosh S, Pardee AB, Iglehart JD. NF-kappa B activation in human breast cancer specimens and its role in cell proliferation and apoptosis. *Proc Natl Acad Sci U S A.* 2004; 101:10137–10142. [PubMed: 15220474]
41. Kyle RA, Yee GC, Somerfield MR, Flynn PJ, Halabi S, Jagannath S, Orlowski RZ, Roodman DG, Twilte P, Anderson K. American Society of Clinical Oncology 2007 clinical practice guideline update on the role of bisphosphonates in multiple myeloma. *J Clin Oncol.* 2007; 25:2464–2472. [PubMed: 17515569]
42. Kearns AE, Khosla S, Kostenuik PJ. Receptor activator of nuclear factor kappaB ligand and osteoprotegerin regulation of bone remodeling in health and disease. *Endocr Rev.* 2008; 29:155–192. [PubMed: 18057140]
43. Bheemasankara Rao C, Namosiva Rao T, Suryaprakasam S. Cardamonin and alpinetin from the seeds of *Amomum subulatum*. *Planta Med.* 1976; 29:391–392. [PubMed: 959382]

Figure 1. Cardamonin inhibits RANKL-induced osteoclastogenesis

(A) The structure of cardamonin. (B) RAW 264.7 cells (1×10^4 /mL) were incubated with cardamonin (500 nM) or RANKL (5 nM) alone, or with RANKL plus cardamonin (0, 50, 100, 250, 500 nM), for 5 days, and then stained for TRAP expression. TRAP-positive cells were photographed. Original magnification, $\times 100$. (C) Quantification of multinucleated osteoclasts (i.e., those containing three nuclei) after treatment with medium, RANKL (5 nM) alone, or RANKL plus cardamonin (0, 50, 100, 250, 500 nM) for 5 days. Columns, mean of 3 measurements; bars, standard deviation; *, $P < 0.01$; **, $P < 0.001$ vs. RANKL alone. (D) RAW 264.7 cells (1×10^4 cells) were incubated with indicated concentration of cardamonin for 12 hours, and then treated with 5 nmol/L RANKL for 5 days. To determine cell viability, cells were trypsinized, and then subjected to trypan blue exclusion assay.

Figure 2. Cardamonin inhibits RANKL-induced osteoclastogenesis

(A) RAW 264.7 cells ($1 \times 10^4/\text{mL}$) were incubated with cardamonin (500 nM) or RANKL (5 nM) alone, or with RANKL plus cardamonin (0, 50, 100, 250, 500 nM), for 3, 4, or 5 days, and then stained for TRAP expression. TRAP-positive cells were photographed. Original magnification, $\times 100$. (B) Quantification of multinucleated osteoclasts (i.e., those containing three nuclei) after treatment with medium (C, Control), or RANKL plus cardamonin (0, 50, 100, 250, 500 nM) for 3, 4, or 5 days. Columns, mean of 3 measurements; bars, standard deviation; *, $P < 0.01$; **, $P < 0.001$ vs. RANKL alone.

Figure 3. Cardamonin inhibits RANKL-induced osteoclastogenesis 24 h after stimulation
(A) RAW 264.7 cells (1×10^4 /mL) were incubated with RANKL (5 nM), cardamonin (500 nM), or both for the indicated times and stained for TRAP expression. **(B)** Multinucleated osteoclasts (i.e., those containing three nuclei) were counted. “M” stands for cells treated with medium alone (control). *Columns*, mean of 3 measurements; *bars*, standard deviation; *, $P < 0.01$; **, $P < 0.001$ vs. medium alone.

Figure 4. Cardamonin suppresses osteoclastogenesis induced by tumor cells

(A) RAW 264.7 cells (1×10^4 /mL) were incubated in the presence of MDA-MB-231 cells (1×10^3 /mL) and exposed to cardamonin (500 nM) for 5 days, and finally stained for TRAP expression. Multinucleated osteoclasts were counted (right panel). **(B)** RAW264.7 cells (1×10^4 /mL) were incubated in the presence of U266 cells (1×10^3 /mL) and exposed to cardamonin (500 nM) for 5 days, and finally stained for TRAP expression. Multinucleated osteoclasts were counted (right panel). *Columns*, mean of 3 measurements; *bars*, standard deviation; *, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$.

Figure 5. RANKL induces NF- B activation and cardamonin inhibits it in a dose- and time-dependent manner

(A) RAW 264.7 cells ($1.5 \times 10^6/\text{mL}$) were incubated with different concentrations of cardamonin for 12 h, treated with 10 nM RANKL for 30 min, and examined for NF- B activation by EMSA. Fold value is based on the value for medium (control), arbitrarily set at 1. (B) RAW 264.7 cells ($1.5 \times 10^6/\text{mL}$) were incubated with 20 μM of cardamonin for 12 h, treated with 10 nM RANKL for the indicated times (min), and reexamined for NF- B activation by EMSA. Fold value is based on the value for medium (control), arbitrarily set at 1. CV, cell viability.

Figure 6. Cardamomin suppresses RANKL-induced I κ B degradation and phosphorylation through inhibition of IKK activity

(A) RAW 264.7 cells ($1.5 \times 10^6/\text{mL}$) were incubated with $20 \mu\text{M}$ of cardamomin for 12 h and then treated with 10 nM RANKL for the indicated times (min). Cytoplasmic extracts were examined for I κ B degradation by western blot using an anti-I κ B antibody. Anti- β -actin antibody was used as a loading control. (B) RAW 264.7 cells ($1.5 \times 10^6/\text{mL}$) were pretreated with cardamomin ($20 \mu\text{M}$) for 12 h, then incubated with ALLN ($50 \mu\text{g}/\text{mL}$) for 30 min, and then treated with RANKL (10 nM) for 15 min. Cytoplasmic extracts were prepared and analyzed by western blot using an anti-phospho-I κ B antibody. The same membrane was re probed with I κ B and β -actin antibody. (C) RAW 264.7 cells ($1.5 \times 10^6/\text{mL}$) were pretreated with cardamomin ($20 \mu\text{M}$) for 12 h and then incubated with RANKL (10 nM) for the indicated times (min). Whole-cell extracts were immunoprecipitated using an antibody against IKK and analyzed by an immune complex kinase assay using recombinant GST-I κ B as described in Materials and Methods. To examine the effect of cardamomin on the level of IKK proteins, whole-cell extracts were analyzed by western blot using anti-IKK and anti-IKK antibodies.

Figure 7. A peptide that targets the NEMO-binding domain inhibits RANKL-induced osteoclastogenesis

(A) RAW 264.7 cells (1×10^4 /ml) were pretreated with 100 μ M of the NEMO-binding domain peptide (NBP) for 2 h, medium was changed, and then RANKL (5 nM) was added for 5 days. (B) Multinucleated osteoclasts (i.e., those containing three nuclei) were counted. Columns, mean of 3 measurements; bars, standard deviation. (C) RAW 264.7 cells (1.5×10^6 /mL) were incubated with 100 μ M of NBP for 2 h, and then incubated with 10 nM RANKL for 30 min and examined for NF- κ B activation by EMSA. Fold value is based on the value for medium (control), arbitrarily set at 1. CV, cell viability.