Published in final edited form as:

Mol Microbiol. 2013 September ; 89(5): . doi:10.1111/mmi.12326.

VeA and MvIA repression of the cryptic orsellinic acid gene cluster in *Aspergillus nidulans* involves histone 3 acetylation

Jin Woo Bok¹, Alexandra A. Soukup², Elizabeth Chadwick¹, Yi-Ming Chiang^{3,4}, Clay C. C. Wang^{4,5}, and Nancy P. Keller^{1,6,*}

¹Department of Medical Microbiology and Immunology, University of Wisconsin-Madison, WI

²Department of Genetics, University of Wisconsin-Madison, WI

³Graduate Institute of Pharmaceutical Science, Chia Nan University of Pharmacy and Science, Tainan, Taiwan, ROC 71710

⁴Department of Pharmacology and Pharmaceutical Sciences, University of Southern California, 1985 Zonal Avenue, Los Angeles, CA, USA 90033

⁵Department of Chemistry, University of Southern California, 1985 Zonal Avenue, Los Angeles, CA, USA 90033

⁶Department of Bacteriology, University of Wisconsin-Madison, WI

Summary

A perplexing aspect of fungal secondary metabolite gene clusters is that most clusters remain 'silent' under common laboratory growth conditions where activation is obtained through gene manipulation or encounters with environmental signals. Few proteins have been found involved in repression of silent clusters. Through multicopy suppressor mutagenesis, we have identified a novel cluster suppressor in *Aspergillus nidulans*, MvIA (modulator of *veA* loss). Genetic assessment of MvIA mutants revealed the role of both itself and VeA (but not the VeA partner LaeA) in the suppression of the cryptic *ors* gene cluster producing orsellinic acid and its F9775 derivatives. Loss of *veA* upregulates F9775A and F9775B production and this increase is reduced 4~5 fold when an overexpression *mvIA* (*OE::mvIA*) allele is introduced into the *veA* background. Previous studies have implicated a positive role for GcnE (H3K9 acetyltransferase of the SAGA/ADA complex) in *ors* cluster expression and here we find expression of *gcnE* is upregulated in *veA* and suppressed by *OE::mvIA* in the *veA* background. H3K9 acetylation levels of *ors* cluster genes correlated with *gcnE* expression and F9775 production in *veA* and *OE::mvIA veA* strains. Finally, deletion of *gcnE* in the *veA* background abolishes *ors* cluster activation and F9775 production. Together, this work supports a role for VeA and MvIA in modifying SAGA/ADA complex activity.

Keywords

LaeA; Velvet Complex; F9775A; Urc4; suppressor; CclA; SAGA/ADA

Introduction

Secondary metabolism is a process by which plants, fungi, and bacteria naturally produce chemicals that are used in niche specialization (Demain & Fang, 2000). As opposed to primary metabolites, secondary metabolites are not necessary for laboratory growth

*Corresponding author: Phone: (608) 262-9795; fax: (608) 262-8418, npkeller@wisc.edu.

conditions but appear to afford defensive/protective properties to the producing organism (Yin *et al.*, 2012, Rohlfs & Churchill, 2011). Secondary metabolites are of great interest to humankind because they can be both beneficial (e.g. pharmaceuticals) and harmful (e.g. mycotoxins) to human and animal health. This interest has led to many studies aimed at understanding the genetic regulation of these bioactive metabolites.

Secondary metabolism in filamentous fungi is globally regulated by a conserved heterotrimeric protein complex called the Velvet Complex (Bayram *et al.*, 2008, Wiemann *et al.*, 2010, Wu *et al.*, 2012, Hoff *et al.*, 2010, López-Berges *et al.*, 2012, Yang *et al.*, 2012, Amaike & Keller, 2009). This complex is composed of the LaeA, VeA, and VelB proteins. Deletion of these genes decreases secondary metabolite production but the phenotypes of the deletants are not equivalent. Loss of *laeA* and *veA* – the two genes most studied – have a major impact on secondary metabolism in *A. nidulans* (Bayram *et al.*, 2008). *laeA* encodes a putative nuclear methyltransferase which positively regulates metabolic gene clusters and controls expression of metabolites such as sterigmatocystin, aflatoxin, penicillin, lovastatin, terraquinone, and mycelial pigments, with deletion strains reduced and overexpression strains increased in metabolite production (Bok & Keller, 2004). Loss of *veA*, also called the “*velvet*” gene and encoding a light regulated protein (Calvo *et al.*, 2002), results in decrease of many of the same metabolites but increase in mycelial pigmentation. In contrast to *laeA* overexpression, *veA* overexpression leads to decreased penicillin production (Spröte & Brakhage, 2007) and oppositely regulates certain penicillin cluster genes (Kato *et al.*, 2003).

These latter studies suggest that VeA can negatively regulate some secondary metabolite clusters through an unknown mechanism. Recent insight into positive regulation of secondary metabolite clusters by VeA and LaeA have arisen from genetic screens of deletants of either protein. Chemical mutagenesis of a *veA* strain resulted in the identification of RtfA, a RNA-pol II transcription elongation factor-like protein, which when deleted in a *veA* background resulted in the remediation of sterigmatocystin in a medium dependent manner (Ramamoorthy *et al.*, 2012). Loss of *rtfA* did not, however, rescue sterigmatocystin in a *laeA* strain. In a multicopy genetic suppressor screen of a *laeA* strain, overexpression of two genes was found to partially complement sterigmatocystin production. One gene encoded the bZIP transcription factor RsmA (restorer of secondary metabolism A) that specifically bound to the promoter of *afIR*, the sterigmatocystin specific transcription factor (Yin *et al.*, 2012). Overexpression of *rsmA* also restored sterigmatocystin production in a *veA* background (Shaaban *et al.*, 2010). The other gene encoded EsaA, a histone 4 acetyltransferase involved in early activation of multiple secondary metabolite gene clusters (Soukup *et al.*, 2012a). This latter study supported earlier research correlating chromatin modifications with secondary metabolite cluster activation or repression (Roze *et al.*, 2007, Reyes-Dominguez *et al.*, 2010, Strauss & Reyes-Dominguez, 2011, Bok *et al.*, 2009).

Here, using the same multicopy suppressor mutagenesis strategy which identified RsmA and EsaA in the *laeA* background, we looked for genes impacting secondary metabolism in a *veA* genetic background with the thought that some of the genes would be identical to those found in the *laeA* screen. Although this was not the case, our screen revealed a novel protein termed MvIA (modulator of *veA* loss A). Analysis of both *mvIA* deletion and overexpression alleles in several genetic backgrounds allowed us to characterize both VeA and MvIA as negative regulators of the cryptic orsellinic acid (*ors*) secondary metabolite gene cluster. This regulation was associated with histone 3 modifications, particularly increased H3K9 acetylation by the histone 3 acetyltransferase SAGA/ADA complex member GcnE. Furthermore, *gcnE* deletion eliminated *ors* expression and product formation in the *veA* background. This is the first genetic screen, to our knowledge, to identify a

novel negative regulator of a cryptic gene cluster as well as provide some insight into a mechanism for VeA negative regulation of a secondary metabolite cluster.

Results

Identification of *mvIA* as a moderator of *veA* loss

Transformation of RJW113.4 (*veA*, *pyrG89*) with the AMAI library generated 30,000 transformants, three of which partially remediated the *veA* phenotype as detected by pigment loss in mycelium as observed from the bottom side of petri plates (Fig. S1). One of the plasmids rescued from one of the three transformants contained a single full-length ORF of the gene AN8797, termed *mvIA* (modulator of VeA loss A), with a functional promoter and termination cassette, which maps to chromosome 3. *mvIA* encodes a 480 aa protein with considerable identity (43%) to Urc4, a protein recently characterized as important in uracil degradation in the yeast *Saccharomyces (Lachnaceae) kluyveri* (Andersen *et al.*, 2008, Björnberg *et al.*, 2010). Further BLAST searching identified another *A. nidulans* gene, AN12027, encoding a 429 aa protein with 44% identity to both MvIA and Urc4. All sequenced *Aspergilli* contain homologs of these two similar proteins.

To characterize *mvIA*, we overexpressed and deleted *mvIA* in wild type and overexpressed *mvIA* in a *veA* background (Fig. S2A). With the thought that AN12027 could possibly be playing a similar role as *mvIA*, we also deleted and overexpressed this gene, hence called *ssmA* (similar sequence to *MvIA*), in wild type and also overexpressed *ssmA* in a *veA* background (Fig. S2B). Northern data confirmed high expression of *mvIA* and *ssmA* by their respective overexpression alleles (Fig. 1). Similar to the original AMAI mutant, overexpression of *mvIA* in the *veA* background decreased pigmentation but overexpression of *ssmA* had no effect on pigmentation (Fig. 2) leading us to conclude the two proteins did not fulfill the same roles in the fungus. Localization of MvIA was determined by microscopy of a *mvIA::eGFP* strain. The protein was evenly dispersed throughout the cytoplasm (Fig. S3)

Both *mvIA* and *veA* negatively regulate the cryptic *ors* gene cluster

Loss of either *veA* or *laeA* results in decreased levels of multiple secondary metabolites including the well-known metabolites sterigmatocystin and penicillin (Bok & Keller, 2004, Kato *et al.*, 2003). Neither of these metabolites was greatly impacted by either *mvIA* overexpression or *mvIA* loss in wild type, *veA* or *laeA* backgrounds as determined by TLC and bioassay (Fig. S4 and data not shown). As *veA* and *laeA* phenotypes themselves are not equivalent, with *veA* but not *laeA* exhibiting a dark pigmentation, there was no change of pigmentation in *OE::mvIA laeA* as compared to *laeA* alone (data not shown).

Postulating that natural products were responsible for the pigmentation of the *veA* strain, we next examined extracts from WT, *veA*, *OE::mvIA veA*, *OE::mvIA*, and *mvIA* by LC/MS for impact on other secondary metabolites (Fig. 3). This data clearly showed a dramatic increase in the production of the polyketide orsellinic acid and its derivatives, the cathepsin K inhibitors F9775A and F9775B, in the *veA* strain. This increase was dampened in the *OE::mvIA veA* strain. The increase in orsellinic acid and F9775 production in *veA* was accompanied by an increase in gene expression of the *ors* secondary metabolite gene cluster (*orsA-E*) responsible for the production of orsellinic acid and F9775 (Fig. 4). Paralleling the chemistry, *ors* cluster gene expression was delayed in the *OE::mvIA veA* strain. The dark pigmentation of the *veA* strain thus appeared to be due to in part to high production of *ors* cluster derived products. This was confirmed by deleting *orsA* encoding the polyketide synthase required for production of orsellinic acid and F9775 (Sanchez *et al.*, 2010) in the *veA* background. The double *veA orsA* strain yielded a much lighter mycelial

pigmentation phenotype similar in appearance to the *OE::mvIA veA* strain (Fig. S5A). Also *veA* loss did not increase expression of other genes associated with fungal pigmentation, further supporting the contribution of *ors* products to the dark coloration of *veA* (Fig. S6).

MvIA and VeA connection with histone 3 modification and GcnE activity

The *ors* gene cluster encoding for F9775 was one of the first cryptic gene clusters to be characterized in *A. nidulans* (Bok *et al.*, 2009, Schroeckh *et al.*, 2009). In these previous studies, a role for chromatin remodeling enzymes had been implicated in *ors* expression. Loss of CclA, a member of the Compass Complex required for histone 3 lysine 4 (H3K4) methylation, had been found to result in derepression of several secondary metabolite clusters including the *ors* cluster, the latter leading to increased F9775 production (Bok *et al.*, 2009). This study also established crosstalk with H3K4 and H3K9 modifications as methylation of both lysine residues was decreased in the derepressed clusters. *Streptomyces rapamycinicus* induction of the *ors* cluster was found to be dependent on the histone 3 acetyltransferase (HAT) complex SAGA/ADA in *A. nidulans* (Nützmann *et al.*, 2011). We thus thought it possible that either MvIA or VeA might be involved in crosstalk with these complexes and/or chromatin remodeling of the *ors* cluster.

We assessed gene expression of two members of the SAGA/ADA complex, *gcnE* and *adaB* (Reyes-Dominguez *et al.*, 2008) that were found to be important for *S. rapamycinicus* activation of the *ors* cluster (Nützmann *et al.*, 2011). Both genes, *gcnE* more so than *adaB*, were modestly upregulated in the *veA* background but only *gcnE* subsequently dampened in the *OE::mvIA veA* strain (Fig. 5A). ChIP analysis of these strains reflected differential *gcnE* expression and activity as demonstrated by altered acetylation of the target residue H3K9 on two genes in the *ors* cluster (Fig. 5B). Analysis of the *orsD* promoter showed that the *veA* mutant displayed significantly increased levels of the activating H3K9ac mark. This increase was abolished in the *OE::mvIA veA* strain. Similarly, acetylation of the *orsA* promoter in the *OE::mvIA veA* is depressed relative to *veA*. This is in contrast to acetylation levels of the control *tubA* promoter, where there is no significant difference in acetylation among strains.

This data suggested that GcnE activity was important for *ors* cluster expression and F7995 production in the *veA* background. To more concretely assess this possibility, we next examined two *veA gcnE* double mutants. Both *ors* expression and F7995 production were lost in these strains (Fig. 6). Notably, the dark coloration of mycelium of the *veA* strain was remediated in the double mutants (Fig. S5B).

Since both H3K9 and K4 levels are impacted in the *cclA* background, we also asked if *mvIA* overexpression could limit F9775 production in the *cclA* background. As expected, and shown previously (Bok *et al.*, 2009), *cclA* loss resulted in an increase in F9775 production compared to wild type (Fig. 3D). Introduction of the overexpression *mvIA* allele into the *cclA* background (*OE::mvIA cclA*) resulted in a reduction of F9775 production as compared to *cclA* alone. The fold reduction was similar to that of *OE::mvIA* in the *veA* background.

veA removal does not allow for increased F9775 production by two cluster activators

We were then interested to determine if *veA* loss – e.g. removal of a negative regulator of the *ors* cluster – would allow for further activation of the *ors* cluster by positive secondary metabolite regulators. We chose two regulators which act through different mechanisms. RsmA is a bZIP transcription factor that increases sterigmatocystin and asperthecin production through promoter binding and activation (Yin *et al.*, 2012) whereas EsaA, a histone 4 acetyltransferase, has a more general mechanism in activating multiple secondary

metabolite clusters through increased H4K12 acetylation (Soukup *et al.*, 2012a). Unexpectedly, examination of *OE::rsmA veA* and *OE::esaA veA* strains compared to their respective controls showed that both EsaA and RsmA overexpression resulted in an approximately 2 fold decrease in F9775 production in the *veA* background compared to *veA* alone (Fig. 3D).

***mvIA* and *ssmA* are not required for uracil catabolism**

As noted earlier, both MvIA and SsmA shared significant identity to Urc4, a protein required for uracil utilization in *S. kluyveri*. Following yeast procedures (Andersen *et al.*, 2008), the single and double *mvIA* and *ssmA* deletion mutants were examined for their ability to grow on uracil as a sole nitrogen source. Loss of *mvIA* was associated with decreased sporulation but this phenotype was independent of uracil supplementation (Fig. S7).

Discussion

The nature of fungal secondary metabolite gene cluster regulation has been the topic of many recent investigations (Yin & Keller, 2011, Strauss & Reyes-Dominguez, 2011). Central to our understanding of cluster regulation has been the characterization of the conserved Velvet Complex whose members, particularly *laeA* and *veA*, are required for activation of dozens of secondary metabolite gene clusters (Bayram *et al.*, 2008, Wiemann *et al.*, 2010, Wu *et al.*, 2012, Hoff *et al.*, 2010, López-Berges *et al.*, 2012, Yang *et al.*, 2012, Amaike & Keller, 2009). While generally considered to be positive regulators of such clusters, a few studies have implicated these proteins in negative regulation of certain secondary metabolites, for example a possible negative impact of VeA on penicillin production in *A. nidulans* (Kato *et al.*, 2003, Spröte & Brakhage, 2007) and Vel1 (the *Fusarium fujikuroi* VeA ortholog) on bikaverin production (Wiemann *et al.*, 2010). Here we describe our finding of a novel gene, *mvIA*, that together with *veA* negatively regulates the cryptic *ors* gene cluster.

The dark pigmentation of *veA* allowed for identification of *mvIA* in the multicopy suppressor mutagenesis screen (Fig. 2 and Fig. S1). We found that increased *ors* metabolite production contributed to this dark coloration. Both *OE::mvIA* and deletion of the polyketide synthase OrsA in the *veA* background yielded a decrease in pigmentation (Figs. 2 and S5). The remaining pigmentation may be attributable to other mycelial pigments not regulated by VeA such as Met13 associated pigments (Fig. S6 and Frandsen *et al.*, 2010).

MvIA and VeA act at the transcriptional level to negatively regulate *ors* gene cluster expression. This was demonstrated by *ors* gene overexpression in *veA* and a delay in the overexpression when *OE::mvIA* was introduced into the *veA* background (Fig. 4). This regulation was reflected in product synthesis (Fig. 3). As examined here, this repressive impact of both proteins on the *ors* clusters appears to be at the level of chromatin modification, notably of histone 3 (Fig. 5). H3K9 methylation is a common heterochromatin mark (Strauss and Reyes-Dominguez, 2011) and decreased H3K9-2/3 methylation was also important in the induction of the *ors* cluster by *S. rapamycinicus* (Nützmann *et al.*, 2011). Specifically acetylation of this residue – the opposing euchromatin mark to methylation – by the SAGA/ADA complex was critical for activation of the *ors* cluster. This complex is not fully delineated in *A. nidulans* but at least two members, GcnE and AdaB, have been identified (Reyes-Dominguez *et al.*, 2008). A requirement for GcnE for an increase in *ors* cluster activation and subsequent F9775 production was supported by loss of both transcript and metabolites in the double *gcnE veA* strain.

A regulatory link between *veA* and the SAGA/ADA complex may occur in part at the transcriptional level. *veA* loss was associated with a modest increased transcript level of both *gcnE* and *adaB*; however only *gcnE* appeared dampened by *mvIA* overexpression in the *veA* background. Transcript levels of *ors* genes are increased in the *veA* strain, but absent in the *gcnE veA* double mutant, further suggesting that regulation of SAGA complex activity may be contributing to differential *ors* induction. To confirm that these alterations in transcript level reflected post translational activity of the complex, we used H3K9ac as a marker for SAGA/ADA complex activity. H3K9ac was strongly enriched at the promoters of *orsA* and *orsD* in the *veA* strain, but decreased to wild type levels in the double mutant, mirroring what was seen at the transcript and metabolite levels. These data confirm that acetylation of the *ors* cluster by the SAGA complex is necessary for cluster activation.

Interestingly, increased acetylation of *orsA* was also seen in both the *OE::mvIA* and *veA* strains, although transcript levels were unaltered in this strain. Therefore, H3K9 acetylation does not appear to be sufficient to activate transcription of this gene on its own. This confirms previous works of study on the orsellinic acid and sterigmatocystin clusters, where changes in histone modifications have been described that either failed to activate gene transcription (Nuetzmann et al., 2011; Soukup et al., 2012), or dramatic changes in transcript levels did not correspond with altered modification of histone residues (Reyes Dominguez et al., 2010). Notably, the majority of these studies found modification of additional residues that correlated strongly with transcription, affirming the complexity of regulation, particularly regarding histone crosstalk.

The *ors* cluster was also shown to be activated by loss of CclA, a member of the H3K4 methylation Compass Complex (Bok et al., 2009). Here the *cclA* mutant not only resulted in decreased H3K4-2/3 methylation – typically thought of as an activating mark – but also decreased H3K9-2/3 methylation. We also propose that decreased F9775 in the *OEmvIA cclA* strain as compared to *cclA* alone reflects possible crosstalk in H3K4 and H3K9 modification activity.

Removal of the negative *ors* cluster regulator VeA did not allow for increased F9775 synthesis by two positive-acting secondary metabolite cluster regulators. Like GcnE and CclA, EsaA also is involved in chromatin modification, specifically acetylation of histone 4 residues (Soukup et al., 2012a). A previous study showed that whereas *esaA* overexpression increased cluster gene transcription and product formation of the sterigmatocystin, terraquinone and penicillin gene clusters, there was no impact on the *ors* cluster nor its products. Our prediction that *OE::esaA* would increase F9775 production when *veA* was deleted was not borne out, at least not in the conditions utilized here. This may suggest that histone 4 acetylation does not impact *ors* cluster expression. Nor did overexpression of RsmA, known to increase production of sterigmatocystin and asperthecin, increase F9775 production in the *veA* background. This was not unexpected as RsmA has been found to increase sterigmatocystin, and hypothetically asperthecin, through activation of the sterigmatocystin specific transcription factor AfIR (Yin et al., 2012). However, the decrease in F9775 production for both double mutants was not predicted. One explanation for this reduction in F9775 could involve decreased availability of metabolite precursors to *ors* cluster products as several of the RsmA and EsaA regulated metabolites draw on the acyl CoA pool. Indeed, studies have shown that inactivation of one secondary metabolite can result in increased production of other secondary metabolites in both fungi and plants (Bok et al., 2009, Post et al., 2012, Wiemann et al., 2012).

Beyond its apparent repressive affect on GcnE activity and H3K9 acetylation levels in the *veA* background, the mechanism of MvIA remains enigmatic. Despite its fairly high identity to Urc4, a protein involved in uracil metabolism in *L. kluyveri* (Andersen et al.,

2008, Björnberg *et al.*, 2010), our study did not indicate a role for MvIA, or SsMA, in pyrimidine metabolism in *A. nidulans*. MvIA is conserved in fungi and it will be interesting to see if there may be a role for this protein in VeA interactions in other fungi.

Experimental procedures

Strains and Growth Conditions

Strains used or created in this study are listed in Table S1. All strains unless otherwise noted were grown on glucose minimal media, GMM (Shimizu & Keller, 2001), with additional supplements for auxotrophic strains (pyrodoxin, riboflavin, uridine, uracil, or para aminobenzoic acid as needed). The genetic background of the primary strains used in this study are as follows: RDIT9.32 (WT) and RJW112.2 (*-veA*) are isogenic prototrophs used for expression and physiology studies. RJW113.4 (*-veA*) is an auxotroph (*pyrG89*) used for sexual crossing and mutagenesis. RJW112.2 and RJW113.4 were obtained from sexual cross between RDIT47.32 (*-veA*, *trpC80I*) and RTMH207.13 (*pyrG89*, *veA+*). RDIT47.32 is a sexual cross progeny of DVARI X RTMK30.23. RDIT30.23 is from FGSC237 X RDIT1.1. RDIT1.1 is from pW1 X FGSC773. All strains are maintained as glycerol stocks at -80°C .

Transformation with AMAI genomic DNA library

The recipient strain, RJW113.4, was transformed with the pRG3-AMA1-based *A. nidulans* wild type genomic DNA plasmid library with *pyr4* of *N. crassa* as the selective marker (Shaaban *et al.*, 2010). Transformation techniques followed previously described procedures (Shaaban *et al.*, 2010). Thousands of transformants were screened for pigment reduction on the underside of the fungal colonies. Total DNA from individual transformants was used to amplify insert by polymerase chain reaction (PCR) with primers flanking the DNA insert in the AMAI plasmid (AMAFWD and AMAREV, Table S2). The PCR amplicons were sequenced, and the assembled sequences were used for BLASTn search of the *A. nidulans* genome (www.broad.harvard.edu/annotation/genome/aspergillus_group/). Plasmid pJW100 was rescued from one of the transformants and re-transformed into strain RJW113.4 to confirm reduction of pigmentation. The sequence of the insert in pJW100 was confirmed one more time by BLAST searching of the *Aspergillus* comparative database.

Construction of *mvIA* deletion, overexpression and double mutants

The *mvIA* deletion strain (TJW98) was created in RJMP1.49 by replacing *AN8797* (*mvIA*) open reading frame with *A. parasiticus pyrG* using modified double joint PCR (Yu *et al.*, 2004) consisting of the following: 1 kb DNA fragment upstream of the *mvIA* start codon (primers 8797F5 and 8797PR5), a 2 kb DNA fragment of *A. parasiticus pyrG* (primers parapyrGF and parapyrGR) via pJW24 (Calvo *et al.*, 2004), and a 1 kb DNA fragment downstream of the *mvIA* stop codon (primers 8797PF3 and 8797R3). First round PCR protocol started with an initial activation of the Pfu Ultra fusion II polymerase (Agilent) at 95°C for 3 min with hot start. For the next 36 PCR cycles, the temperature was held at 95°C for 30 s denaturing, then ramped up to 58°C for 30 s annealing, and to 68°C for 2 min extension. Finally, a post extension step was performed at 68°C for 10 min. Second round PCR utilized gel-purified DNA templates from first round PCR with Pfu Ultra fusion II polymerase at 95°C for 3 min with hot start. For the next 11 PCR cycles, the temperature was held at 95°C for 40 s denaturing, then ramped to 60°C for 3 min annealing, and to 68°C for 4 min extension. Finally, a post extension step was performed at 68°C for 10 min. Third round PCR started with Expand long template PCR system (Roche) at 94°C for 3 min. For the next 36 PCR cycles, the temperature was held at 94°C for 20 s denaturing, then ramped to 59°C for 30 s annealing, and to 68°C for 4 min extension. Finally, a post extension step was performed at 68°C for 10 min. 30 μl of G50 purified third round PCR product was used for fungal transformation. *mvIA* deletants were confirmed by PCR and

Southern blot and one correct transformant, TJW98, used for sexual crosses. TJW98 was sexually crossed with RTMH207.13 to obtain the prototroph REC1C. The recombinants were confirmed with PCR.

A *mvIA* overexpression (OE) cassette, pJW111.58, was created by inserting a single joint PCR product (1.4 kb DNA fragment of *A. nidulans gpdA* promoter (primers 8797GPDF and 8797PR5), a 2.4 kb DNA fragment containing the *mvIA* open reading frame (ORF) and termination cassette (primers 8797OEF and 8797OER) into the *HindIII* and *SpeI* sites of pJW53 which resulted in the fusion of the *mvIA* gene with the *gpdA* promoter. pJW111.58 was used to transform RJMP1.31 to create an overexpression *mvIA* allele in the *pyrA* locus. The resulting strain was called TJW99. We confirmed TJW99 by PCR and Southern blotting and then sexually crossed TJW99 with RJW113.4 to create a prototrophic strain (REC2C). Subsequent northern analysis was done using radiolabeled probes for the corresponding transcript (1 kb DNA fragment, primers 8797IF and 8797IR).

The *veAOE::mvIA* double mutant strain was created by crossing TJW99 with RJW113.4 to obtain RJW221.5 and confirmed with selective media and PCR. All of primers used in this study are listed in Table S2. *laeAOE::mvIA* and *cclAOE::mvIA* double mutants were created by crossing TJW99 with RJW135.1 and RJW122.2 to obtain RJW223.2 and RJW254.12 respectively. Strain RAAS44.15 was obtained through mating of TAAS16.13 (Soukup *et al.*, 2012a) with DVARI (Kim *et al.*, 2002).

Construction of *ssmA* deletion, overexpression, and double mutants

AN12027(ssmA) was deleted by replacing it with *A. fumigatus riboB* in *A. nidulans* strain TNO2A7 using modified double joint PCR. Construction of *ssmA* deletion, overexpression and double mutants consisted of the following: 1 kb DNA fragment upstream of the *ssmA* start codon (primers 12027F5 and 12027R5), a 2.3 kb DNA fragment of *A. fumigatus riboB* (primers 12027rb5 and 12027rb3), and a 1 kb DNA fragment downstream of the *ssmA* stop codon (primers 12027F3 and 12027R3). The deletion mutants were confirmed by PCR and Southern blot and one deletion, TJW124.13, was sexually crossed with RDIT2.1 to get a prototrophic strain (RJW215.25) and a *metG* auxotrophic strain (RJW215.9). RJW215.9 was sexually crossed with TJW98 to obtain a *mvIA ssmA* double deletion mutant (RJW229.2). A *ssmA* overexpression mutant was created with *A. parasiticus pyrG* with double joint PCR product consisting of the following: 1 kb DNA fragment upstream of the *ssmA* start codon (primers 12027F5 and 12027oe5 R), a 3.6 kb DNA fragment of *A. parasiticus pyrG* (primers parapyrGF and parapyrGR) with *A. nidulans gpdA* promoter via pJMP9 (Soukup *et al.*, 2012b) and a 1 kb DNA fragment downstream of the *ssmA* start codon (primers 12027oe3 F and 120273 R). The transformants were screened by PCR and Southern blot to obtain TJW125.1. This mutant was sexually crossed with RDIT2.1 and RJW113.4 to get a overexpression prototroph RJW216.15 and a *OE::ssmA veA* double mutant (RJW226.4), respectively. Subsequent northern analysis was done using radiolabeled probes for the corresponding transcript (1 kb DNA fragment, primers 12027IF and 12027IR). All primers used in this study are listed in Table S2.

Construction of *veA* and *gcnE* double deletion mutants

The double deletion strain (TJW142) was created in RJW113.4 by replacing AN3621 (*gcnE*) open reading frame with *A. parasiticus pyrG* using modified double joint PCR consisting of the following: 1 kb DNA fragment upstream of the *gcnE* start codon (primers gcnE5F and gcnE5R), a 2 kb DNA fragment of *A. parasiticus pyrG* (primers parapyrGF and parapyrGR) via pJW24 (Calvo *et al.*, 2004), and a 1 kb DNA fragment downstream of the *gcnE* stop codon (primers gcnE3F and gcnE3R). 30 µl of G50 purified third round PCR product was

used for fungal transformation. Double deletants were confirmed by PCR and two correct transformants, TJW142.1 and 142.2 used for subsequent experiments.

Construction of green fluorescent tagged *mvIA*

A N-terminus GFP tagged *mvIA* cassette was created through two steps. Firstly, a DNA fragment of the *mvIA* open reading frame (1.7kb, primers 8797gfp_{gpdF} and 8797gfp_{gpdR}) was inserted in between the *gpdA* promoter and *gfp* of pJMP50 (*A. fumigatus pyroA* with *A. nidulans gpdA* promoter and a fused *gfp*) to make pJW147.17 using Quick change method (Bok & Keller, 2012). For creation of pJMP50, the *ptrA* gene was removed from pSK505 by restriction digestion with *SphI* and *PstI*. Subsequently, the *A. fumigatus pyroA* gene was amplified using the oligonucleotides JP Af pyroA *SphI* For (5' ACATCAGcATGCTGGATTACTAAGGTAATG) and JP Af pyroA *PstI* Rev (5' AGTA_cGCAGCATCCACATGATC) and cloned into pCR TOPO Blunt II (Invitrogen) to construct pJMP61. A 1.8 kb *SphI-PstI* fragment corresponding to the *A. fumigatus pyroA* gene was cut out of pJMP61 and ligated into the *SphI-PstI* sites of pSK505 to construct pJMP50. The *gpdA* promoter of pJW147.17 was then replaced with a *mvIA* promoter (1kb, primers 8797gfp_{natF} and 8797gfp_{natR}) by Quick change method to create the plasmid pJW149.7. This plasmid was confirmed by sequence analysis to contain a *mvIA* promoter, N-terminus *gfp* and *mvIA* ORF, and then used to transform TJW98 to create the strain TJW106. Dark cultured germlings were viewed at 40X magnification using a fluorescence microscope (Zeiss AxioImager.A1) and AxioVision image capturing software. Germlings were grown for over night at 30 °C in liquid GMM over a cover slip. The cover slip was turned over onto a slide and viewed immediately. Hoechst dye was used to identify nuclei for the N-terminus GFP tagged *MvIA* strain TJW106.

Nucleic acid analysis

DNA extraction, restriction enzyme digestion, gel electrophoresis, blotting, hybridization, and probe preparation were performed by standard methods (Sambrook & Russell, 2001). RNA extractions were made from mycelia of cultures where 10⁵ spores/ml were grown in 50 ml liquid GMM at 37 °C with shaking at 250 rpm for 48 hr. For pigment and pigment associated gene (*yA* Aramayo and Timberlake, 1990; *cfwA* Marquez-Fernandez *et al.*, 2007; *met13* Frandsen RJ *et al.* 2010) expression studies, 1 × 10⁶ spores/ml of strains was inoculated into 50 ml of liquid GMM and grown for 24 h at 37 °C and 250 rpm shaking. Mycelia were harvested by filtration through Miracloth (Calbiochem) and transferred onto solid GMM for additional 6 hr and 20 hr. Total RNA was extracted using Isol-RNA Lysis Reagent (5 Prime) according to the manufacturer's instructions with approximately 30 µg of total RNA for RNA blot analysis. RNA blots were hybridized with *mvIA*, *ssmA*, *veA*, *orsA*, *orsB*, *orsC*, *orsD*, *orsE*, *gcnE*, *adaB*, *yA*, *cfwA*, *met13* and *gpdA* DNA fragments which were generated by PCR using gene-specific primers as shown in Table S2. Expressed gene intensity was quantified using ImageJ software.

Metabolite Assays

Thin layer chromatography (TLC) was used to assess sterigmatocystin (ST) production. ST was extracted from GMM solid medium culture, which was point-inoculated with 10³ spores/plate and grown for 5 days at 37 °C. One cm cores were punched from the center of point inoculated plates and homogenized with 3 ml ddH₂O. Three ml of chloroform were added and samples were centrifuged for 10 min. The organic layer was removed and put into a 3 ml glass vial and left to sit and dry in a fume hood overnight. Dried extracts were resuspended with 100 µl chloroform and 5 or 10 µl were loaded onto a non-UV coated TLC plate. ST was spotted as a standard. The plates were run in chloroform: acetone (8:2) solvent and stained with 15 % aluminum chloride in 95 % ethanol. TLC plates were viewed under 254 nm UV light.

Penicillin bioassays were carried out with fungal culture filtrate (10 ml) harvested from 10^5 spores/ml liquid-shaking culture after 120 hours at 37 °C, 225 rpm. Freeze-dried culture filtrate was resuspended in 1 ml ddH₂O and 200 µl of resuspended sample was added into 1 cm diameter wells in a TSA + *Micrococcus luteus* bacteria plate (15 g Bacto Tryptone, 5 g Bacto Soytone, 5 g NaCl, 10 g agar per Liter of media, with 500 µl *M. luteus* (OD = 1) added to 40 ml TSA per plate). Plates were left at 4 °C for 3 hours then incubated at 37 °C overnight. Where indicated, 12 U of penicillinase was added to 200 µl of extracts and incubated for 3 hours at 37 °C before adding to wells as a control.

Fermentation and LC/MS analysis

A. nidulans wild type and mutant strains were cultivated at 37 °C, 225 rpm in 50 ml of liquid GMM at 10^5 spores/ml. After 48 hours or 5 days, culture broth was harvested by filtration and 40 ml lyophilized. The lyophilized sample was suspended in H₂O (20 ml) and partitioned with ethyl acetate (20 ml). In order to extract acidic phenolic metabolites, the water layer was acidified to pH 2 and partitioned with ethyl acetate (20 ml) a second time. The combined ethyl acetate layers were evaporated in vacuo, re-dissolved in 0.5 ml of DMSO/methanol (1:4) and injected with 10 µl for HPLC-DAD-MS analysis. LC/MS was carried out in negative mode using a ThermoFinnigan LCQ Advantage ion trap mass spectrometer with an RP C18 column (Alltech Prevail C18 3 µm 2.1 × 100mm) at a flow rate of 125 µl/min as described (Bok *et al.*, 2009).

For F9775A and B quantitation, linear curves were generated by using UV-Vis 430 nm as detection. The extracts were prepared so that the concentration of F9775A and B lay within the standard curves.

Chromatin immunoprecipitation and real time qPCR analysis

Fifty milliliter cultures of liquid GMM were inoculated with 1×10^6 spores per ml and incubated at 250 rpm and 37 °C for 24 hours under light. Triplicate cultures were performed for each strain. Chromatin immunoprecipitation was carried out as described previously (Boedi *et al.*, 2012). Antibodies used for ChIP were: rabbit polyclonal to histone H3 acetyl K9, Abcam, ab10812 and rabbit polyclonal to C-terminus histone H3 antibody, Abcam, ab1791. Two µg of antibody was used per reaction of 200 mg total protein. Amplification and detection of precipitated DNA in real-time qPCR was performed with iQTM SYBR® Green Supermix (Bio-Rad, Cat#170-8880) following the manufacturer's instructions using primers listed in Table S2. Relative amounts of DNA were calculated by dividing the immunoprecipitated DNA by the input DNA. Each PCR reaction was replicated. To normalize the amount of DNA precipitated with histone H3-acetyl K9 the quantities from precipitation with this antibody was divided by the previously calculated ratio of the anti-C-terminus histone H3 precipitation to input DNA. All values are shown relative to wild type. To ensure that levels of precipitated DNA were above background, Rabbit IgG Abcam, ab27478 was used as a nonspecific control. In all cases, precipitated DNA levels were at least 100 fold above background (data not shown).

Pyrimidine utilization assay

Uracil or uridine (2 g/L) was supplied as a nitrogen source in the solid medium containing Difco yeast nitrogen base without amino acid or Difco yeast nitrogen base without amino acid and ammonium sulfate 2.5 g/L, glucose 10 g/L, *A. nidulans* trace element 1 ml/L (1000 fold stock) and 1.5 % agar as modified ingredient published by Andersen *et al* (2008). One µl of 10^6 conidia/ml was inoculated on the center of media and strains cultured 10 days in a 37 °C incubator.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work was supported in part by NIH grant PO1GM084077 from the National Institute of General Medical Sciences to N.P.K. and C.C.W. and in part by the Cargill-Benevenga Fund and Food Research Institute to N.P.K. and E.C. and by NIH T32 GM07133 and NIH NRSA AI55397 to A.A.S.

References

- Amaike S, Keller NP. Distinct roles for VeA and LaeA in development and pathogenesis of *Aspergillus flavus*. *Eukaryot Cell*. 2009; 8:1051–1060. [PubMed: 19411623]
- Andersen G, Björnberg O, Polakova S, Pynyaha Y, Rasmussen A, Møller K, Hofer A, Moritz T, Sandrini MP, Merico AM, Compagno C, Akerlund HE, Gojkovi Z, Piskur J. A second pathway to degrade pyrimidine nucleic acid precursors in eukaryotes. *J Mol Biol*. 2008; 380:656–666. [PubMed: 18550080]
- Aramayo R, Timberlake WE. Sequence and molecular structure of the *Aspergillus nidulans* *yA* (laccase *I*) gene. *Nucleic Acids Res*. 1990; 18(11):3415. [PubMed: 2192364]
- Bayram O, Krappmann S, Ni M, Bok JW, Helmstaedt K, Valerius O, Braus-Stromeier S, Kwon NJ, Keller NP, Yu JH, Braus GH. VelB/VeA/LaeA complex coordinates light signal with fungal development and secondary metabolism. *Science*. 2008; 320:1504–1506. [PubMed: 18556559]
- Björnberg O, Vodnala M, Domkin V, Hofer A, Rasmussen A, Andersen G, Piskur J. Ribosylurea accumulates in yeast *urc4* mutants. *Nucleosides Nucleotides Nucleic Acids*. 2010; 29:433–437. [PubMed: 20544532]
- Boedi S, Reyes-Dominguez Y, Strauss J. Chromatin immunoprecipitation analysis in filamentous fungi. *Methods Mol Biol*. 2012; 944:221–236. [PubMed: 23065620]
- Bok JW, Chiang YM, Szewczyk E, Reyes-Dominguez Y, Davidson AD, Sanchez JF, Lo HC, Watanabe K, Strauss J, Oakley BR, Wang CC, Keller NP. Chromatin-level regulation of biosynthetic gene clusters. *Nat Chem Biol*. 2009; 5:462–464. [PubMed: 19448638]
- Bok JW, Keller NP. LaeA, a regulator of secondary metabolism in *Aspergillus* spp. *Eukaryot Cell*. 2004; 3:527–535. [PubMed: 15075281]
- Bok JW, Keller NP. Fast and easy method for construction of plasmid vectors using modified quick-change mutagenesis. *Methods Mol Biol*. 2012; 944:163–174. [PubMed: 23065615]
- Calvo AM, Bok J, Brooks W, Keller NP. veA is required for toxin and sclerotial production in *Aspergillus parasiticus*. *Appl Environ Microbiol*. 2004; 70:4733–4739. [PubMed: 15294809]
- Calvo AM, Wilson RA, Bok JW, Keller NP. Relationship between secondary metabolism and fungal development. *Microbiol Mol Biol Rev*. 2002; 66:447–459. table of contents. [PubMed: 12208999]
- Demain AL, Fang A. The natural functions of secondary metabolites. *Adv Biochem Eng Biotechnol*. 2000; 69:1–39. [PubMed: 11036689]
- Frandsen RJ, Albertsen KS, Stougaard P, Sørensen JL, Nielsen KF, Olsson S, Giese H. Methylene-tetrahydrofolate reductase activity is involved in the plasma membrane redox system required for pigment biosynthesis in filamentous fungi. *Eukaryot Cell*. 2010; 9:1225–35. [PubMed: 20543064]
- Hoff B, Kamerewerd J, Sigl C, Mitterbauer R, Zadra I, Kürnsteiner H, Kück U. Two components of a velvet-like complex control hyphal morphogenesis, conidiophore development, and penicillin biosynthesis in *Penicillium chrysogenum*. *Eukaryot Cell*. 2010; 9:1236–1250. [PubMed: 20543063]
- Kato N, Brooks W, Calvo AM. The expression of sterigmatocystin and penicillin genes in *Aspergillus nidulans* is controlled by *veA*, a gene required for sexual development. *Eukaryot Cell*. 2003; 2:1178–1186. [PubMed: 14665453]
- Kim H, Han K, Kim K, Han D, Jahng K, Chae K. The *veA* gene activates sexual development in *Aspergillus nidulans*. *Fungal Genet Biol*. 2002; 37:72–80. [PubMed: 12223191]

- López-Berges MS, Hera C, Sulyok M, Schäfer K, Capilla J, Guarro J, Di Pietro A. The velvet complex governs mycotoxin production and virulence of *Fusarium oxysporum* on plant and mammalian hosts. *Mol Microbiol.* 2012; 87:49–65. [PubMed: 23106229]
- Marquez-Fernandez O, et al. Phosphopantetheinyl transferase CfwA/NpgA is required for *Aspergillus nidulans* secondary metabolism and asexual development. *Eukaryot Cell.* 2007; 6:710–20. [PubMed: 17277172]
- Nützmam HW, Reyes-Dominguez Y, Scherlach K, Schroeckh V, Horn F, Gacek A, Schumann J, Hertweck C, Strauss J, Brakhage AA. Bacteria-induced natural product formation in the fungus *Aspergillus nidulans* requires Saga/Ada-mediated histone acetylation. *Proc Natl Acad Sci U S A.* 2011; 108:14282–14287. [PubMed: 21825172]
- Post J, van Deenen N, Fricke J, Kowalski N, Wurbs D, Schaller H, Eisenreich W, Huber C, Twyman RM, Prüfer D, Gronover CS. Laticifer-specific cis-prenyltransferase silencing affects the rubber, triterpene, and inulin content of *Taraxacum brevicorniculatum*. *Plant Physiol.* 2012; 158:1406–1417. [PubMed: 22238421]
- Ramamoorthy V, Shantappa S, Dhingra S, Calvo AM. *veA*-dependent RNA-pol II transcription elongation factor-like protein, RtfA, is associated with secondary metabolism and morphological development in *Aspergillus nidulans*. *Mol Microbiol.* 2012; 85:795–814. [PubMed: 22783880]
- Reyes-Dominguez Y, Bok JW, Berger H, Shwab EK, Basheer A, Gallmetzer A, Scazzocchio C, Keller N, Strauss J. Heterochromatic marks are associated with the repression of secondary metabolism clusters in *Aspergillus nidulans*. *Mol Microbiol.* 2010; 76:1376–1386. [PubMed: 20132440]
- Reyes-Dominguez Y, Narendja F, Berger H, Gallmetzer A, Fernandez-Martin R, Garcia I, Scazzocchio C, Strauss J. Nucleosome positioning and histone H3 acetylation are independent processes in the *Aspergillus nidulans* *prnD-prnB* bidirectional promoter. *Eukaryot Cell.* 2008; 7:656–663. [PubMed: 18296621]
- Rohlf M, Churchill AC. Fungal secondary metabolites as modulators of interactions with insects and other arthropods. *Fungal Genet Biol.* 2011; 48:23–34. [PubMed: 20807586]
- Roze LV, Arthur AE, Hong SY, Chanda A, Linz JE. The initiation and pattern of spread of histone H4 acetylation parallel the order of transcriptional activation of genes in the aflatoxin cluster. *Mol Microbiol.* 2007; 66:713–726. [PubMed: 17919289]
- Sambrook, J.; Russell, DW. Molecular cloning: a laboratory manual. Cold Spring Harbor Laboratory Press; Cold Spring Harbor, N.Y.: 2001.
- Sanchez JF, Chiang YM, Szewczyk E, Davidson AD, Ahuja M, Elizabeth Oakley C, Woo Bok J, Keller N, Oakley BR, Wang CC. Molecular genetic analysis of the orsellinic acid/F9775 gene cluster of *Aspergillus nidulans*. *Mol Biosyst.* 2010; 6:587–593. [PubMed: 20174687]
- Schroeckh V, Scherlach K, Nützmam HW, Shelest E, Schmidt-Heck W, Schuemann J, Martin K, Hertweck C, Brakhage AA. Intimate bacterial-fungal interaction triggers biosynthesis of archetypal polyketides in *Aspergillus nidulans*. *Proc Natl Acad Sci U S A.* 2009; 106:14558–14563. [PubMed: 19666480]
- Shaaban MI, Bok JW, Lauer C, Keller NP. Suppressor mutagenesis identifies a velvet complex remediator of *Aspergillus nidulans* secondary metabolism. *Eukaryot Cell.* 2010; 9:1816–1824. [PubMed: 20935144]
- Shimizu K, Keller NP. Genetic involvement of a cAMP-dependent protein kinase in a G protein signaling pathway regulating morphological and chemical transitions in *Aspergillus nidulans*. *Genetics.* 2001; 157:591–600. [PubMed: 11156981]
- Soukup AA, Chiang YM, Bok JW, Reyes-Dominguez Y, Oakley BR, Wang CC, Strauss J, Keller NP. Overexpression of the *Aspergillus nidulans* histone 4 acetyltransferase EsaA increases activation of secondary metabolite production. *Mol Microbiol.* 2012a; 86:314–30. [PubMed: 22882998]
- Soukup AA, Farnoodian M, Berthier E, Keller NP. NosA, a transcription factor important in *Aspergillus fumigatus* stress and developmental response, rescues the germination defect of a *laeA* deletion. *Fungal Genet Biol.* 2012b; 49:857–65. [PubMed: 23022264]
- Spröte P, Brakhage AA. The light-dependent regulator velvet A of *Aspergillus nidulans* acts as a repressor of the penicillin biosynthesis. *Arch Microbiol.* 2007; 188:69–79. [PubMed: 17375284]
- Strauss J, Reyes-Dominguez Y. Regulation of secondary metabolism by chromatin structure and epigenetic codes. *Fungal Genet Biol.* 2011; 48:62–69. [PubMed: 20659575]

- Wiemann P, Albermann S, Niehaus EM, Studt L, von Bargen KW, Brock NL, Humpf HU, Dickschat JS, Tudzynski B. The Sfp-type 4 -phosphopantetheinyl transferase Ppt1 of *Fusarium fujikuroi* controls development, secondary metabolism and pathogenicity. *PLoS One*. 2012; 7:e37519. [PubMed: 22662164]
- Wiemann P, Brown DW, Kleigrewe K, Bok JW, Keller NP, Humpf HU, Tudzynski B. FfVel1 and FfLae1, components of a velvet-like complex in *Fusarium fujikuroi*, affect differentiation, secondary metabolism and virulence. *Mol Microbiol*. 2010 [Epub ahead of print].
- Wu D, Oide S, Zhang N, Choi MY, Turgeon BG. ChLae1 and ChVel1 regulate T-toxin production, virulence, oxidative stress response, and development of the maize pathogen *Cochliobolus heterostrophus*. *PLoS Pathog*. 2012; 8:e1002542. [PubMed: 22383877]
- Yang Q, Chen Y, Ma Z. Involvement of BcVeA and BcVelB in regulating conidiation, pigmentation and virulence in *Botrytis cinerea*. *Fungal Genet Biol*. 2013; 50:63–71. [PubMed: 23147398]
- Yin W, Keller NP. Transcriptional regulatory elements in fungal secondary metabolism. *J Microbiol*. 2011; 49:329–339. [PubMed: 21717315]
- Yin WB, Amaike S, Wohlbach DJ, Gasch AP, Chiang YM, Wang CC, Bok JW, Rohlf M, Keller NP. An *Aspergillus nidulans* bZIP response pathway hardwired for defensive secondary metabolism operates through *afIR*. *Mol Microbiol*. 2012; 83:1024–1034. [PubMed: 22283524]
- Yu JH, Hamari Z, Han KH, Seo JA, Reyes-Domínguez Y, Scazzocchio C. Double-joint PCR: a PCR-based molecular tool for gene manipulations in filamentous fungi. *Fungal Genet Biol*. 2004; 41:973–981. [PubMed: 15465386]

Figure 1.

veA, *mvIA*, and *ssmA* gene expression in *A. nidulans* wild type (WT), *veA*, *OE::mvIA*, *OE::mvIA veA*, *mvIA*, *OE::ssmA*, *OE::ssmA veA*, and *ssmA* strains (RDIT9.32, RJW112.2, RJW220.17, RJW221.5, REC1C, RJW216.15, RJW226.4, and RJW215.25, respectively) grown in liquid shaking GMM for 48 hours at 37° C. RNA was extracted from independent biological duplicated samples. Ethidium bromide-stained rRNA and *gpdA* are indicated for loading.

Figure 2.

Mycelial pigmentation patterns of *A. nidulans* WT, ΔveA , *OE::mvIA*, *OE::mvIA* ΔveA , $\Delta mvIA$, *OE::ssmA*, *OE::ssmA* ΔveA , $\Delta ssmA$, $\Delta mvIA$ $\Delta ssmA$, and *OE::mvIA* *OE::ssmA* strains (RDIT9.32, RJW112.2, RJW220.17, RJW221.5, REC1C, RJW216.15, RJW226.4, RJW215.25, RJW229.2, and RJW228.3, respectively) strains grown on solid GMM at light and dark for 5 days at 37 °C.

Figure 3.

Overexpression of *mvIA* suppresses *ors* gene cluster activated in *veA* strain. (A) Total scan HPLC profile of secondary metabolites extracted from the culture medium of *A. nidulans* WT, *veA*, *OE::mvIA*, *OE::mvIA veA*, and *mvIA* (RDIT9.32, RJW112.2, RJW220.17, RJW221.5, and REC1C, respectively) strains. (B) Metabolites identified from *veA* strain. (C) The yield of F9775A and B from *veA* and *OE::mvIA veA* strains in 48 hours GMM liquid culture. (D) Relative yield of F9775A and B from extracts obtained from 48 hours liquid culture media of the listed strains as compared to *veA*.

Figure 4.

Gene expression analysis of *mvIA*, *veA*, and *ors* cluster genes of *A. nidulans* WT, *veA*, *OE::mvIA*, *OE::mvIA veA*, and *mvIA* (RDIT9.32, RJW112.2, RJW220.17, RJW221.5, and REC1C, respectively) strains. 36, 60 and 48 represent hours of growth at 37°C, 250rpm after inoculation to GMM liquid media. Ethidium bromide-stained rRNA and *gpdA* are indicated for loading. The *ors* gene cluster is depicted at the bottom of the figure.

Figure 5.

Transcriptional regulation of *gcnE*. *veA* deletion strain is decreased by overexpressed *mvIA*. (A) *adaB* and *gcnE* expression of *A. nidulans* WT, *veA*, OE::mvIA and OE::mvIA *veA* strains (RDIT9.32, RJW112.2, RJW220.17, and RJW221.5, respectively) strains grown on liquid GMM for 48 hours at 37° C. RNA was extracted from independent biological triplicated samples. Ethidium bromide-stained rRNA and *gpdA* are indicated for loading. (B) ChIP analysis of H3K9 acetylation levels at the *orsA*, *orsD*, and *tubA* promoters for 48 hour cultures. Acetylation data has been normalized to total H3 in all cases. Letters represent significant differences at $p < 0.05$. Numbers below each lane are *gpdA*-normalized relative expression levels of the mutant strains in relation to the *gpdA*-normalized wild-type control, which has been arbitrarily set to 1.

Figure 6.

Deletion of *gcnE* suppresses *ors* gene cluster expression and subsequent metabolite production in *veA*. (A) *orsA* and *orsE* expression in *A. nidulans veA* and *veA gcnE* (RJW112.2, TJW142.1, and TJW142.2, respectively) strains grown in liquid GMM for 48 hours at 37° C. RNA was extracted from independent biological triplicated samples. Ethidium bromide-stained *rRNA* and *gpdA* are indicated for loading. (B) Total scan HPLC profile of secondary metabolites extracted from the culture medium of *A. nidulans veA* and *veA gcnE* strains. Compound 1, 2 and 3 are the same structures shown as Fig. 3.