Published in final edited form as:

Psychol Med. 2013 February ; 43(2): 293–302. doi:10.1017/S0033291712000918.

Heterogeneity of Amygdala Response in Major Depressive Disorder: The Impact of Lifetime Sub-Threshold Mania

Jay C. Fournier^a, Matthew T. Keener^a, Benjamin C. Mullin^b, Danella M. Hafeman^a, Edmund J. LaBarbara^a, Richelle S. Stiffler^a, Jorge Almeida^a, Dina M. Kronhaus^c, Ellen Frank^a, and Mary L. Phillips^a

^aDepartment of Psychiatry, University of Pittsburgh School of Medicine

^bDepartment of Psychiatry and Behavioral Sciences, Children's Hospital Colorado

^cUniversity of Cambridge, St. Catharine's College and Computer Laboratory

Abstract

Background—Patients with major depressive disorder (MDD) present with highly heterogeneous symptom profiles. We aimed to examine whether individual differences in amygdala activity to emotionally-salient stimuli were related to heterogeneity in lifetime levels of depressive and sub-threshold manic symptoms among adults with MDD.

Methods—We compared age- and gender-matched adults with MDD (N=26) with healthy controls (HC, N=28). While undergoing fMRI, participants performed an implicit emotional faces task: they labeled a color flash superimposed upon initially neutral faces that dynamically morphed into one of four emotions (angry, fearful, sad, happy). Region of interest analyses examined group differences in amygdala activity. For conditions in which adults with MDD displayed abnormal amygdala activity versus HC, within-group analyses examined amygdala activity as a function of scores on a continuous measure of lifetime depression-related and mania-related pathology.

Results—Adults with MDD showed significantly greater right-sided amygdala activity to angry and happy conditions than HC ($p < 0.05$, corrected). Multiple regression analyses revealed that greater right amygdala activity to the happy condition in adults with MDD was associated with higher levels of sub-threshold manic symptoms experienced across the lifespan ($p = 0.002$).

Conclusions—Among depressed adults with MDD, lifetime features of sub-threshold mania were associated with abnormally elevated amygdala activity to emerging happy faces. These findings are a first step toward identifying biomarkers that reflect individual differences in neural mechanisms in MDD, and challenge conventional mood disorder diagnostic boundaries by suggesting that some adults with MDD are characterized by pathophysiologic processes that overlap with bipolar disorder.

Introduction

Major depressive disorder (MDD) is a highly heterogeneous disorder, with large inter-individual differences in symptom, illness course, and treatment response profiles. The research agenda for the National Institute of Mental Health (NIMH) emphasizes a translation of basic and clinical neuroscience research findings into a new classification system for all psychiatric disorders based upon biomarkers that reflect pathophysiologic and

Reprint request address: Jay C. Fournier, University of Pittsburgh School of Medicine, 3811 O'Hara St., Pittsburgh, PA 15213.

All other authors report no biomedical financial interests or potential conflicts of interest.

etiological processes (Charney *et al.*, 2002, Hasler *et al.*, 2006, Hasler *et al.*, 2004, Phillips and Frank, 2006). Despite years of research, however, the search for reliable, consistently present biomarkers of MDD has proven elusive.

Perhaps the best-established neuroimaging marker of MDD is abnormally elevated amygdala activity to negative emotional information (Fitzgerald *et al.*, 2008, Kessler *et al.*, 2011, Peluso *et al.*, 2009), including fear (Sheline *et al.*, 2001) and sadness (Fu *et al.*, 2004, Suslow *et al.*, 2010, Victor *et al.*, 2010). Despite broad agreement that abnormalities in amygdala function are associated with MDD, these findings are not always obtained (e.g., Fitzgerald *et al.*, 2008). For example, some researchers report an absence of abnormally elevated amygdala activity to negative emotional stimuli in depressed individuals with MDD (Almeida *et al.*, 2010, Surguladze *et al.*, 2005), while others describe heterogeneity in amygdala activity in depressed groups (Abler *et al.*, 2007, Canli *et al.*, 2005, Dannlowski *et al.*, 2007, Lee *et al.*, 2007, Siegle *et al.*, 2006, Siegle *et al.*, 2007). Abnormal amygdala activity to positive emotional stimuli is less consistently observed, with some researchers (Sheline *et al.*, 2001) reporting abnormally elevated amygdala activity to happy faces in depressed individuals with MDD, others (Surguladze *et al.*, 2005) failing to find this association, and still others (Suslow *et al.*, 2010, Victor *et al.*, 2010) finding the opposite pattern.

The inconsistent findings regarding abnormal amygdala activity in MDD may reflect the heterogeneity of affective psychopathology among individuals diagnosed with the disorder. Standard measures of *current* symptom severity, however, may not be sufficiently sensitive to explain the large inter-individual variation in neural activity among individuals with MDD. Individual differences in neural circuitry function may, instead, be more strongly associated with measures that assess the full continuum of *lifetime* manifestations of affective psychopathology.

Recognizing the need to develop a clinical, lifetime measure capturing the full range of symptoms and associated features across the spectrum of mood disorders, Cassano and colleagues developed the Mood Spectrum Self-Report Instrument (MOODS-SR; Dell'Osso *et al.*, 2002). The instrument, and the model of psychopathology from which it originated, posits that subtle, subthreshold signs and symptoms experienced over one's lifetime may represent a clinically-meaningful underlying diathesis that is shared with individuals meeting diagnostic criteria for the relevant disorder. This approach assumes that individuals with a particular diagnosis, e.g., MDD, may nevertheless have an underlying pathology that is on a continuum with other, traditionally separate illnesses, e.g., bipolar disorder, panic disorder, OCD, etc. The MOODS-SR measures dimensions of affective psychopathology, including manic and depressive symptoms, that characterize the lifetime presence of affective dysregulations comprising both fully syndromal and sub-threshold mood disturbance (Dell'Osso *et al.*, 2002).

Using a structured interview version of the MOODS instrument, Cassano and colleagues (Cassano *et al.*, 2004) observed substantial heterogeneity among adults with MDD regarding the presence of lifetime mania-related psychopathology, despite careful clinical screening to ensure that no individual diagnosed with MDD met diagnostic criteria for any bipolar disorder. Depressed individuals with higher levels of lifetime mania-related psychopathology had earlier onsets of depression, were more likely to experience suicidal ideation, and experienced increased paranoia (Cassano *et al.*, 2004). These findings suggest that differences in underlying pathology may exist among individuals with MDD and that some depressed individuals may have an underlying pathology that is closer to that of bipolar disorder. Given this, it is possible that those individuals with MDD with higher levels of lifetime mania-related pathology may show patterns of amygdala activity reported

more consistently in individuals with bipolar disorder than in individuals with MDD: for example, abnormally elevated amygdala activity to positive emotional (happy) faces (Blumberg *et al.*, 2005, Lawrence *et al.*, 2004, Pavuluri *et al.*, 2007).

The present study's primary goal was to examine individual differences in abnormal amygdala functioning among depressed adults with MDD as a function of a dimensional measure of lifetime affective psychopathology. To do this, we first ascertained under which conditions amygdala functioning was abnormal for the depressed adults with MDD by comparing amygdala activity in depressed and healthy control adults. All participants performed an implicit emotion processing task in which they labeled a color flash that was superimposed upon negative (fear, sad, anger) and positive (happy) emotional, dynamically changing emotional face stimuli. Given the findings reviewed above, we expected to observe that adults with MDD would show functional abnormalities in the amygdala, specifically, significantly elevated amygdala activity relative to healthy control adults to both positive and negative emotional faces. We also expected to observe substantial heterogeneity among adults with MDD in the level of abnormal amygdala activation. We hypothesize that for those emotional conditions during which adults with MDD displayed abnormal amygdala activity: 1) individual differences in the magnitude of amygdala activity to the negative emotional stimuli will be associated with levels of lifetime depression-related psychopathology; and 2) individual differences in the magnitude of amygdala activity to positive emotional stimuli will be associated with levels of lifetime mania-related psychopathology.

Methods

Participants

We recruited sixty-three right-handed, native English-speaking individuals: 32 currently depressed adults diagnosed with MDD, and 31 healthy control participants (HC) with no personal or family history of psychiatric illness. Adults with MDD were carefully screened to ensure that they did not meet diagnostic criteria for bipolar disorder. Psychiatric diagnoses were made using the Structured Clinical Interview for Psychiatric Disorders (SCID-P; First *et al.*, 1995). Exclusion criteria were: history of head injury (from medical records and participant report), systemic medical illness, cognitive impairment (score < 24 Mini-Mental State Examination; Folstein *et al.*, 1975), premorbid IQ estimate < 85 (National Adult Reading Test; Blair and Spreen, 1989), Axis-II borderline personality disorder, and standard MRI exclusion criteria (e.g. presence of metallic objects in the body). Adults with MDD were also excluded if they met criteria for an alcohol/substance use disorder within 2 months before the scan. For HC, additional exclusion criteria included current/previous alcohol or substance abuse/dependence (determined by SCID-I, saliva and urine screen), and any personal or family history of Axis I disorder. The two groups were age- and gender-matched. Six depressed patients were excluded from the analyses (three for movement > 2mm between two successive scans; two for < 75% color labeling accuracy during the scan; one for scoring 2.5 standard deviations above the mean level of depression severity), three HC were excluded because of motion > 2mm between two successive scans. The final sample included 26 adults with MDD and 28 HC (Table 1). The study protocol was approved by the University of Pittsburgh Institutional Review Board. After complete description of the study to participants, written informed consent was obtained.

Measures

Current depression severity was assessed using the 17-item Hamilton Rating Scale for Depression (HRSD; Hamilton, 1960). Current manic symptoms were assessed using the Young Mania Rating Scale (YMRS; Young *et al.*, 1978). Clinical and demographic

information was collected through self-report questionnaires and clinical interview using the SCID-P. 65% of adults with MDD had a history of anxiety disorder, and 35% had a history of substance abuse. Lifetime depression-related and mania-related psychopathology were assessed using the MOODS-SR–Lifetime Version (Dell'Osso *et al.*, 2002). This 161-item questionnaire assesses the presence of symptoms and features for periods of at least 3–5 days during one's lifetime. The measure is organized into three domains for mania-related psychopathology and three domains for depression-psychopathology (mood, energy, and cognition), and a seventh domain assessing disturbances in rhythmicity. In the present study, because the three domain scores for depression and for mania were highly correlated with their respective total scores (all r s > 0.90), the total scores for depression and mania were used.

Paradigm

Participants completed a 12.5-minute emotional dynamic faces task during fMRI. Stimuli comprised faces from the NimStim set (Tottenham *et al.*, 2009) that were morphed in 5% increments, from neutral (0% emotion) to 100% emotion for 4 emotions: happy, sad, angry and fear (Figure 1). Morphed faces were collated into 1s movies progressing from 0% to 100% emotional display. In control trials, movies comprised a simple shape (dark oval) superimposed on a light-grey oval, with similar structural characteristics to the face stimuli, which subsequently morphed into a larger shape, approximating the movement of the morphed faces. There were three blocks for each of the four emotional conditions, with twelve stimuli per block, and six control blocks with six stimuli per block. Emotional and control blocks were presented in a pseudorandomized order so that no two blocks of any condition were presented sequentially. Participants were asked to use one of three fingers to press a button indicating the color of a semi-transparent foreground color flash (orange, blue, or yellow) that appeared during the mid 200ms–650ms of the 1s presentation of the dynamically-changing face. The emotional faces were task-irrelevant and processed by the participants implicitly.

Data acquisition

Neuroimaging data were collected using a 3.0 Tesla Siemens Trio MRI scanner at the Magnetic Resonance Research Center in the University of Pittsburgh Medical Center. Structural 3D axial MPRAGE images were acquired in the same session (TR/TE=2200/3.29ms; Flip angle 9°, FOV: 256×192mm²; Slice thickness:1mm; Matrix: 256×256;192 continuous slices). BOLD images were then acquired with a gradient echo EPI sequence during approximately thirteen minutes (378 successive brain volumes) covering 39 axial slices (3.2 mm thick; TR/TE=2000/28 ms/ms; FOV=205×205 mm²; matrix=64×64; Flip angle 90°).

Functional Neuroimaging Data Analyses

Data were preprocessed and analyzed with statistical parametric mapping software (SPM8; <http://www.fil.ion.ucl.ac.uk/spm>). During preprocessing, data were corrected for differences in acquisition time between slices, coregistered, realigned, resampled to 2×2×2mm³ voxels, spatially normalized into standard stereotactic space (Montreal Neurologic Institute, MNI), and spatially smoothed using a 6mm FWHM Gaussian kernel. A first-level fixed-effect model was constructed in which each of the four emotion conditions (anger, fear, sad, and happy) were entered as separate conditions in a block design and contrasted with the shape condition, which served as the baseline in the design matrix. Movement parameters from the preprocessing procedure were entered as covariates of no interest to control for subject movement. Trials were modeled with the Canonical Hemodynamic Response Function. The four emotion contrasts (i.e., anger-minus-shape, fear-minus-shape, sad-minus-shape, and

happy-minus-shape) were entered into second-level analyses with the relevant t-contrast images.

Between Groups Region of Interest Analyses

Region of interest (ROI) analyses were conducted to examine the effect of group in left and right a-priori amygdala ROIs, as defined in the Wake Forest Toolbox PickAtlas Talairach Daemon template (Maldjian *et al.*, 2003). (No filter was used in the construction of these ROIs). To control for multiple statistical testing we maintained a cluster-level false positive detection rate at $p < 0.05$ by using a voxel threshold of $p < 0.05$ with a cluster (k) extent empirically determined by Monte Carlo simulations implemented in AlphaSim of 26 voxels, computed separately for left and right amygdalae. This well-validated technique accounts for spatial correlations between blood oxygen level dependent (BOLD) signal changes in neighboring voxels (Ward, 2002). We utilized the between-groups analysis to identify those conditions in which the MDD and healthy control adults differed (that is, in which the adults with MDD demonstrated abnormal amygdala function) so as to guide the within-group analyses described below.

Within Group Region of Interest Analyses

To examine the extent to which abnormal amygdala activity was associated with dimensions of mood-related psychopathology, we extracted amygdala BOLD response for each emotion condition associated with abnormal activity in adults with MDD relative to HC. Rather than extracting only those voxels for which significant MDD-HC differences were observed, we extracted all voxels from the amygdala ROI and averaged them. Thus, mean BOLD response in the entire anatomically-defined amygdala masks were used as the dependent variables. Within adults with MDD, we examined associations between neural activity and lifetime mania-related and lifetime depression-related psychopathology. Mean amygdala response was modeled as a function of mania-related and depression-related MOODs spectrum scores, controlling for scores on the YMRS, HRSD, gender, age, illness duration, history of prior substance abuse/dependence, history of anxiety disorder, and current medications (3 categories: no psychiatric medication (N=8), antidepressant medication only (N=11), and antidepressant medication and benzodiazepines, mood stabilizer, and/or antipsychotic medication (N=7). Owing to the large number of variables for which we were controlling, we adopted a strategy by which we tested the effect of mania-related and depression-related psychopathology in the full model (all 10 variables), and in a final model that was reduced using backwards stepwise regression (using a significance criterion for variable retention of $p = .10$). Regression analyses were performed using SAS9.2 (SAS Institute Inc, Cary, NC).

Exploratory Whole-Brain Analyses

We conducted exploratory whole-brain analyses to identify those regions in which adults with MDD displayed greater activity than HC to the four emotion-minus-shape contrasts. To control for multiple voxelwise tests, we set a voxelwise threshold of $p < .001$ and a minimum cluster (k) extent of 20 voxels.

Results

Task performance

Mean color labeling accuracy and mean reaction time were calculated for each participant across all conditions. Overall, task accuracy was high (HC = 96%; MDD = 94%), and there were no significant differences between groups in accuracy or reaction times (between

groups variances were unequal for both comparisons, thus the Satterthwaite method was used: $t(39.7)=1.08$, $p=0.29$ for accuracy; $t(43.0)=-0.78$, $p=0.44$ for reaction times).

Activity

Between Groups Region of Interest Analysis—A significant main effect of group was observed in the right amygdala, by which activity was higher for adults with MDD than for HC ($t(208)=2.40$; $p=0.009$; $k=26$ voxels).¹ Subsequent contrasts revealed that this effect resulted from significantly greater activity for adults with MDD during the anger ($t(208)=2.35$; $p=0.01$; $k=31$ voxels) and happy ($t(208)=3.19$; $p=0.001$; $k=83$ voxels) conditions relative to HC (Figure 2; Supplemental Table 1). To test for a possible effect of medications on these findings, we compared right amygdala activity to the anger and happy conditions in those adults with MDD taking psychiatric medication ($N=18$) and those not taking medication ($N=8$). Neither between-group contrast met the pre-specified criteria for significance (both $t_s < 1.9$, $k_s < 1$ voxels).

Within Group Region of Interest Analysis—To investigate the extent to which abnormally elevated right amygdala activity to the anger and happy conditions in adults with MDD was associated with individual differences in levels of lifetime affective psychopathology, mean amygdala BOLD signal during each of these conditions was extracted from the anatomically-defined right amygdala mask. Separate regression models were estimated for each emotion condition. To correct for the two parallel tests (one regression model each for right amygdala to happy and anger conditions), a corrected p -value of 0.025 was used. In each model, mean right amygdala activity was modeled as a function of depression-related and mania-related MOODS spectrum scores, age, gender, illness duration, HRSD scores, YMRS scores, history of substance abuse/dependence, history of anxiety disorder, and psychiatric medication. To the happy condition only, there was a significant relationship between lifetime mania-related psychopathology and right amygdala activity in adults with MDD in both the full model that controlled for all of the covariates ($F(1, 14)=8.77$, $p=0.01$; Supplementary Table 2), and in the final, reduced model (4 variables) for which a backwards-stepwise regression framework was applied ($F(1, 21)=12.26$, $p=0.002$; Figure 3; Table 2). In the final reduced model, there was also a significant effect of gender: women with MDD displayed greater amygdala activity to the happy condition than men with MDD ($F(1,21)=7.57$, $p=0.01$). (See Supplemental Materials for additional regression model details).

We conducted two additional analyses to assess the robustness and specificity of findings regarding the association between elevated right amygdala activity during the happy condition and lifetime subthreshold mania. First, we divided adults with MDD into two groups of 15 patients (Here, we increased the sample of adults with MDD from 26 to 30 by relaxing the conservative exclusion criteria described above and including the participant with the high HRSD score and additional participants whose movement during the scan was < 6 mm). Previous research (Fagiolini *et al.*, 2007; Rucci personal communication 09/02/2011) identified a cut-off score of 22 on this measure as reflecting clinically significant subthreshold mania. This value was used to divide adults with MDD into Low-

¹The group-by-emotion interaction was not hypothesized to be significant, and it did not reach the AlphaSim corrected threshold for significance. Rather, we hypothesized that adults with MDD would show abnormal amygdala activity for the negative and for the positive emotional conditions, which the main effect of group and the absence of a group-by-emotion interaction effect corroborated. We further hypothesized that different sets of individual difference variables would be associated with the abnormalities for positive and negatively valenced stimuli, which the within-group analyses partially supported. The presence or absence of a group-by-condition interaction was irrelevant to this second hypothesis. One set of patient characteristics may be associated with abnormalities in one emotional condition whereas a different set could be associated with abnormalities in another emotion condition, regardless of whether the magnitude of the abnormalities (defined as MDD-HC differences) between the conditions was the same (no interaction effect) or different (significant interaction effect).

and High-Mania subgroups. The mean mania score for the Low-Mania subgroup ($N=15$, $M=10.1$, $SD=5.7$, $Range=1-18$) was two standard deviations below the cutoff. The mean score for the High-Mania subgroup ($N=15$, $M=32.0$, $SD=7.5$, $Range=23-48$) was one standard deviation above the cutoff. Right amygdala activity to the happy condition was higher in the High-Mania versus the Low-Mania subgroup ($t(112)=2.48$, $p=0.007$, $k=43$ voxels).

Second, to examine whether the results reported above were specific to the happy condition, we calculated the degree to which activity in the right-amygdala was correlated among the four emotional conditions for adults with MDD and for HC. For HC, right-amygdala activity to the happy condition was uncorrelated with activity to any other emotion (all $r_s < .25$, $p_s > .20$). For adults with MDD, right-amygdala activity was uncorrelated with activity during the anger and sad conditions (all $r_s < .10$, $p_s > .63$), and correlated at the level of a non-significant trend with activity to the fear condition ($r=.37$, $p=.06$).

Regarding individual differences in amygdala activity in response to the anger condition, neither the effect of lifetime depression-related psychopathology ($| \beta | = .05$, $t(14) = .20$, $p = .84$), mania-related ($| \beta | = .04$, $t(14) = .13$, $p = .90$) psychopathology, nor any of the other variables entered into the regression model (all $F_s \leq 1.60$, $p_s > 0.22$; Supplemental Table 3) were significantly associated with right amygdala activity to the anger condition.

Exploratory Whole Brain Analysis—There was a significant positive effect of group in bilateral occipitotemporal, frontal, and parietal regions, resulting from adults with MDD showing significantly greater activity than HC in these regions to anger and happy – but not fear or sad – conditions (Supplemental Table 4).

Discussion

The goal of the current study was to examine the extent to which continuous measures of lifetime depression-related and mania-related psychopathology could explain individual differences among adults with MDD in abnormal amygdala activity to negative and positive emotional stimuli. We observed abnormally elevated amygdala activity to angry and happy faces in adults with MDD relative to HC. Furthermore, the level of endorsed lifetime, sub-threshold mania-related psychopathology was positively associated with greater amygdala activity to happy faces in depressed adults.

Few previous studies of emotion processing in MDD examined amygdala activity to angry faces. Many studies (Almeida *et al.*, 2010, Sheline *et al.*, 2001) examined responses to other negative emotional faces (e.g., fearful or sad), and those that did include angry faces typically did not parse the results as a function of the individual emotional condition (Hariri *et al.*, 2002) or did not include a control group (Dannowski *et al.*, 2007). Other studies reported functional abnormalities to angry faces in orbitofrontal cortical regions in individuals with MDD (Lee *et al.*, 2008) and in participants with a prior history of both MDD and suicide attempt relative to individuals with a history of MDD alone (Jollant *et al.*, 2008). To our knowledge, the present study represents the first report of abnormally elevated amygdala activity to angry faces in individuals with MDD.

It is not clear why individuals with MDD in the current study did not demonstrate abnormal activity to the fear or sad conditions. These findings are consistent with at least one prior study that likewise observed no statistical difference between depressed and healthy individuals to these stimuli (Almeida *et al.*, 2010); however, they are inconsistent with other published findings in which such abnormalities have been reported (e.g., Fu *et al.*, 2004, Sheline *et al.*, 2001, Suslow *et al.*, 2010, Victor *et al.*, 2010). Although we cannot say with

certainly why we did not observe abnormalities in the MDD group during these two conditions, we concur with Mayberg (2003) that inconsistent reports regarding abnormalities among depressed patients require further study. Furthermore, we believe that the presence of inconsistent findings like these underscores the need for efforts to explicitly examine individual differences among individuals with depression that might help to resolve discrepancies regarding abnormal neural function. Indeed, a recent report by Grant and colleagues (2011) found that the association between depression and amygdala activity to sad facial displays was driven largely by the prior experience of childhood maltreatment. Those depressed individuals in the study who did not also have a history of childhood maltreatment did not differ from healthy controls in response to sad facial expressions. We were not able to examine this association in the current study. Future work should examine other possible patient characteristics that could help both to resolve discrepant findings in the literature and to explain the heterogeneity of abnormal amygdala function among depressed patients in response to fearful and sad facial displays.

Our finding of abnormally elevated amygdala activity in adults with MDD to the happy condition is consistent with one prior report (Sheline *et al.*, 2001). In the present study, this functional abnormality was strongly and positively associated with the level of lifetime subthreshold mania/hypomania symptoms. Furthermore, when we relaxed our strict inclusion criteria, thus rendering the sample more representative of the patients who presented to the study, and divided the sample into those with substantial subthreshold lifetime mania-related pathology and those with minimal mania-related psychopathology, the same pattern emerged: those with high lifetime mania displayed greater amygdala activity to happy faces than those with low lifetime mania. Previous studies of adults and youth with bipolar disorder similarly reported abnormally elevated amygdala activity to happy faces (Blumberg *et al.*, 2005, Lawrence *et al.*, 2004, Pavuluri *et al.*, 2007). Thus, increased amygdala activity to happy faces may represent a potentially important biomarker, reflecting individual differences in pathophysiology among individuals with MDD. The present findings support the Research Domain Criteria initiative of the NIMH (Insel *et al.*, 2010) and other commentators who have argued for the need to move beyond a categorical classification system for psychiatric disorders, and instead to identify continuous dimensions of psychopathology that are more closely linked to underlying neural mechanisms that cut across the defined categories (Charney *et al.*, 2002, Hasler *et al.*, 2006, Hasler *et al.*, 2004, Phillips and Frank, 2006).

It is unclear why lifetime depression-related psychopathology was not significantly related to abnormal amygdala function in adults with MDD. One possibility is that, by definition, all depressed adults met criteria for MDD. This design feature no doubt restricted the range of scores on this measure, which could be expected to have reduced the power of our statistical tests of this relationship. Furthermore, it is possible that the presence of current depressive symptoms may have interfered with our ability to detect a relationship between lifetime depressive features and amygdala function, although it should be noted that lifetime depressive features were unrelated to amygdala activity whether or not current symptoms were controlled in the statistical model. Future work on this topic should recruit participants from across the normal and pathological range who present with different levels of lifetime depression-related pathology.

Limitations

Features of the current study may limit the generalizability of some of the findings. No remitted depressed participants took part. Thus, we were unable to determine whether abnormalities in amygdala function represented trait or state effects. Despite this, within-group analyses that used dimensional measures of lifetime psychopathology suggested that elevated amygdala activity to the happy condition was associated with lifetime, trait-like

sub-threshold mania-related psychopathology. Many depressed adults were taking psychotropic medications. We were not able to control for medication load (or any other clinically relevant variable) in between-group analyses, as all clinical variables were co-linear with group membership. We did, however, examine the effect of medications among the depressed individuals. There were no significant relationships between medications and amygdala activity. The lack of association between psychiatric medication and neuroimaging markers associated with mood disorders is common (see Phillips *et al.*, 2008 for a more detailed discussion of the issues involving neuroimaging research with actively medicated participants).

Because this was the first study to examine associations between continuous, lifetime measures of affective psychopathology and neural activity to emotionally-salient stimuli, our analyses intentionally focused on the amygdala. Exploratory whole-brain analyses did, however, reveal that adults with MDD showed greater activity than HC in occipitotemporal, frontal, and parietal regions supporting visual attentional processing to the anger and happy conditions. These results parallel the right amygdala findings. An additional limitation to our analysis of the amygdala is that we were not able to control for other variables that have been shown to affect amygdala activity, such as genetic polymorphisms of the serotonin transporter gene (Hariri *et al.*, 2002) and abuse experienced in childhood (Dannlowski *et al.*, 2012). Finally, there were relatively few male participants. Despite this, we found that women with MDD demonstrated greater right amygdala activity than men with MDD to the happy condition. Future studies with larger samples should examine gender differences in relationships between amygdala activity and lifetime measures of sub-threshold affective psychopathology in MDD.

Conclusion

The present findings suggest that neural mechanisms underlying mood disorders may be more nuanced than the current categorical diagnostic system captures. Our finding of greater amygdala activity to positive emotional stimuli in adults with MDD may represent one biomarker that reflects individual differences in the extent to which underlying affective pathophysiology overlaps with that of bipolar disorder. These findings challenge conventional mood disorder diagnostic boundaries, and this knowledge may help to inform more personalized approaches to the treatment of MDD based upon a better understanding of individual differences in underlying pathophysiologic processes.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Dr. Ellen Frank has served as a consultant to Servier and Vanda Pharmaceuticals, has received grant/research support from The Fine Foundation, The Pittsburgh Foundation, and Forest Research Institute, and has received royalties from Guilford Press and the American Psychological Association.

All work was carried out within the Department of Psychiatry, University of Pittsburgh. Neuroimaging data were collected at the Brain Imaging Research Center, University of Pittsburgh, and Carnegie Mellon University. We thank Dr. K.J. Jung, S. Kurdilla, and D. Vizslay for their help acquiring neuroimaging data. We also thank Satish Iyengar for providing statistical advice and guidance.

This research was supported by grant MH076971 and T32 MH018269 from the National Institute of Mental Health, Bethesda, MD.

References

- Abler B, Erk S, Herwig U, Walter H. Anticipation of aversive stimuli activates extended amygdala in unipolar depression. *Journal of Psychiatric Research*. 2007; 41:511–522. [PubMed: 17010993]
- Almeida JRC, Versace A, Hassel S, Kupfer DJ, Phillips ML. Elevated amygdala activity to sad facial expressions: A state marker of bipolar but not unipolar depression. *Biological Psychiatry*. 2010; 67:414–421. [PubMed: 19931855]
- Blair J, Spreen O. Predicting premorbid IQ: A revision of the National Adult Reading Test. *The Clinical Neuropsychologist*. 1989; 3:129–136.
- Blumberg HP, Donegan NH, Sanislow CA, Collins S, Lacadie C, Skudlarski P, Gueorguieva R, Fulbright RK, McGlashan TH, Gore JC, Krystal JH. Preliminary evidence for medication effects on functional abnormalities in the amygdala and anterior cingulate in bipolar disorder. *Psychopharmacology*. 2005; 183:308–313. [PubMed: 16249909]
- Canli T, Cooney RE, Goldin P, Shah M, Sivers H, Thomason ME, Whitfield-Gabrieli S, Gabrieli JD, Gotlib IH. Amygdala reactivity to emotional faces predicts improvement in major depression. *NeuroReport*. 2005; 16:1267–1270. [PubMed: 16056122]
- Cassano GB, Rucci P, Frank E, Fagioli A, Dell'Osso L, Shear MK, Kupfer DJ. The mood spectrum in unipolar and bipolar disorder: Arguments for a unitary approach. *American Journal of Psychiatry*. 2004; 161:1264–1269. [PubMed: 15229060]
- Charney, D.; Barlow, D.; Botteron, K.; Cohen, J.; Goldman, D.; Gur, R.; Lin, K.; Lopez, J.; Meador-Woodruff, J.; Moldin, S.; Nestler, E.; Waton, S.; Zalcman, S. Neuroscience research agenda to guide development of a pathophysiologically based classification system. In: Kupfer, D.; First, M.; Regier, D., editors. *A Research Agenda for DSM-V*. Washington, DC: American Psychiatric Association; 2002. p. 31-83.
- Dannlowski U, Ohrmann P, Bauer J, Kugel H, Arolt V, Heindel W, Kersting A, Baune B, Suslow T. Amygdala reactivity to masked negative faces is associated with automatic judgmental bias in major depression: a 3 T fMRI study. *Journal of Psychiatry and Neuroscience*. 2007; 32:423–429. [PubMed: 18043766]
- Dannlowski U, Stuhrmann A, Beutelmann V, Zwanzger P, Lenzen T, Grotegerd D, Domschke K, Hohoff C, Ohrmann P, Bauer J, Lindner C, Postert C, Konrad C, Arolt V, Heindel W, Suslow T, Kugel H. Limbic Scars: Long-Term Consequences of Childhood Maltreatment Revealed by Functional and Structural Magnetic Resonance Imaging. *Biological Psychiatry*. 2012; 71:286–293. [PubMed: 22112927]
- Dell'Osso L, Armani A, Rucci P, Frank E, Fagioli A, Corretti G, Shear MK, Grochocinski VJ, Maser JD, Endicott J, Cassano GB. Measuring mood spectrum: Comparison of interview (SCI-MOODS) and self-report (MOODS-SR) instruments. *Comprehensive Psychiatry*. 2002; 43:69–73. [PubMed: 11788923]
- Fagioli A, Frank E, Rucci P, Cassano G, Turkin S, Kupfer D. Mood and anxiety spectrum as a means to identify clinically relevant subtypes of bipolar I disorder. *Bipolar Disorders*. 2007; 9:462–467. [PubMed: 17680916]
- First, M.; Spitzer, R.; Gibbon, M.; Williams, J.; Benjamin, L. *Structured Clinical Interview for DSM-IV Axis I Disorders (SCID, version 2.0)*. New York: Biometric Research Department, New York State Psychiatric Institute; 1995.
- Fitzgerald PB, Laird AR, Maller J, Daskalakis ZJ. A meta-analytic study of changes in brain activation in depression. *Human Brain Mapping*. 2008; 29:683–695. [PubMed: 17598168]
- Folstein MF, Folstein SE, McHugh PR. "Mini-mental state". A practical method for grading the cognitive state of patients for the clinician. *Journal of Psychiatric Research*. 1975; 12:189–198. [PubMed: 1202204]
- Fu CHY, Williams SCR, Cleare AJ, Brammer MJ, Walsh ND, Kim J, Andrew CM, Pich EM, Williams PM, Reed LJ, Mitterschiffthaler MT, Suckling J, Bullmore ET. Attenuation of the neural response to sad faces in major depression by antidepressant treatment: A prospective, event-related functional magnetic resonance imaging study. *Archives of General Psychiatry*. 2004; 61:877–889. [PubMed: 15351766]

- Grant MM, Cannistraci C, Hollon SD, Gore J, Shelton R. Childhood trauma history differentiates amygdala response to sad faces within MDD. *Journal of Psychiatric Research*. 2011; 45:886–895. [PubMed: 21276593]
- Hamilton MA. A rating scale for depression. *Journal of Neurology, Neurosurgery, and Psychiatry*. 1960; 23:56–62.
- Hariri AR, Mattay VS, Tessitore A, Kolachana B, Fera F, Goldman D, Egan MF, Weinberger DR. Serotonin transporter genetic variation and the response of the human amygdala. *Science*. 2002; 297:400–403. [PubMed: 12130784]
- Hasler G, Drevets WC, Gould TD, Gottesman II, Manji HK. Toward Constructing an Endophenotype Strategy for Bipolar Disorders. *Biological Psychiatry*. 2006; 60:93–105. [PubMed: 16406007]
- Hasler G, Drevets WC, Manji HK, Charney DS. Discovering endophenotypes for major depression. *Neuropsychopharmacology*. 2004; 29:1765–1781. [PubMed: 15213704]
- Insel T, Cuthbert B, Garvey M, Heinssen R, Pine DS, Quinn K, Sanislow C, Wang P. Research domain criteria (RDoC): toward a new classification framework for research on mental disorders. *American Journal of Psychiatry*. 2010; 167:748–751. [PubMed: 20595427]
- Jollant F, Lawrence NS, Giampietro V, Brammer MJ, Fullana MA, Drapier D, Courtet P, Phillips ML. Orbitofrontal cortex response to angry faces in men with histories of suicide attempts. *American Journal of Psychiatry*. 2008; 165:740–748. [PubMed: 18346998]
- Kessler H, Taubner S, Buchheim A, Munte TF, Stasch M, Kachele H, Roth G, Heinecke A, Erhard P, Cierpka M, Wiswede D. Individualized and clinically derived stimuli activate limbic structures in depression: an fMRI study. *PLoS One*. 2011; 6:e15712. [PubMed: 21283580]
- Lawrence NS, Williams AM, Surguladze S, Giampietro V, Brammer MJ, Andrew C, Frangou S, Ecker C, Phillips ML. Subcortical and Ventral Prefrontal Cortical Neural Responses to Facial Expressions Distinguish Patients with Bipolar Disorder and Major Depression. *Biological Psychiatry*. 2004; 55:578–587. [PubMed: 15013826]
- Lee B-T, Seok J-H, Lee B-C, Cho SW, Yoon B-J, Lee K-U, Chae J-H, Choi I-G, Ham B-J. Neural correlates of affective processing in response to sad and angry facial stimuli in patients with major depressive disorder. *Progress in Neuro-Psychopharmacology & Biological Psychiatry*. 2008; 32:778–785. [PubMed: 18207298]
- Lee BT, Seong Whi C, Hyung Soo K, Lee BC, Choi IG, Lyoo IK, Ham BJ. The neural substrates of affective processing toward positive and negative affective pictures in patients with major depressive disorder. *Progress in Neuro-Psychopharmacology & Biological Psychiatry*. 2007; 31:1487–1492. [PubMed: 17688985]
- Maldjian J, Laurienti P, Kraft R, Burdette J. An automated method for neuroanatomic and cytoarchitectonic atlas-based interrogation of fMRI datasets. *Neuroimage*. 2003; 19:1233–1239. [PubMed: 12880848]
- Mayberg HS. Modulating dysfunctional limbic-cortical circuits in depression: towards development of brain-based algorithms for diagnosis and optimised treatment. *British Medical Bulletin*. 2003; 65:193–207. [PubMed: 12697626]
- Pavuluri MN, O'Connor MM, Harral E, Sweeney JA. Affective neural circuitry during facial emotion processing in pediatric bipolar disorder. *Biological Psychiatry*. 2007; 62:158–167. [PubMed: 17097071]
- Peluso MA, Glahn DC, Matsuo K, Monkul ES, Najt P, Zamarripa F, Li J, Lancaster JL, Fox PT, Gao JH, Soares JC. Amygdala hyperactivation in untreated depressed individuals. *Psychiatry Research: Neuroimaging*. 2009; 173:158–161.
- Phillips ML, Frank E. Redefining Bipolar Disorder: Toward DSM-V. *American Journal of Psychiatry*. 2006; 163:1135–1136. [PubMed: 16816214]
- Phillips ML, Travis MJ, Fagiolini A, Kupfer DJ. Medication Effects in Neuroimaging Studies of Bipolar Disorder. *American Journal of Psychiatry*. 2008; 165:313–320. [PubMed: 18245175]
- Sheline YI, Barch DM, Donnelly JM, Ollinger JM, Snyder AZ, Mintun MA. Increased amygdala response to masked emotional faces in depressed subjects resolves with antidepressant treatment: An fMRI study. *Biological Psychiatry*. 2001; 50:651–658. [PubMed: 11704071]

- Siegle GJ, Carter CS, Thase ME. Use of fMRI to predict recovery from unipolar depression with cognitive behavior therapy. *American Journal of Psychiatry*. 2006; 163:735–738. [PubMed: 16585452]
- Siegle GJ, Thompson W, Carter CS, Steinhauer SR, Thase ME. Increased amygdala and decreased dorsolateral prefrontal BOLD responses in unipolar depression: related and independent features. *Biological Psychiatry*. 2007; 61:198–209. [PubMed: 17027931]
- Surguladze S, Brammer MJ, Keedwell P, Giampietro V, Young AW, Travis MJ, Williams SCR, Phillips ML. A Differential Pattern of Neural Response Toward Sad Versus Happy Facial Expressions in Major Depressive Disorder. *Biological Psychiatry*. 2005; 57:201–209. [PubMed: 15691520]
- Suslow T, Konrad C, Kugel H, Rumstadt D, Zwitterlood P, Schoning S, Ohrmann P, Bauer J, Pyka M, Kersting A, Arolt V, Heindel W, Dannlowski U. Automatic mood-congruent amygdala responses to masked facial expressions in major depression. *Biological Psychiatry*. 2010; 67:155–160. [PubMed: 19748075]
- Tottenham N, Tanaka J, Leon A, McCarry T, Nurse M, Hare T, Marcus D, Westerlund A, Casey B, Nelson C. The NimStim set of facial expressions: judgments from untrained research participants. *Psychiatry Research*. 2009; 168:242–249. [PubMed: 19564050]
- Victor T, Furey M, Fromm S, Öhman A, Drevets W. Relationship between amygdala responses to masked faces and mood state and treatment in major depressive disorder. *Archives of General Psychiatry*. 2010; 67:1128–1138. [PubMed: 21041614]
- Ward, B. AlphaSim. National Institute of Mental Health; 2002.
- Young RC, Biggs JT, Ziegler VE, Meyer DA. A rating scale for mania: Reliability, validity and sensitivity. *British Journal of Psychiatry*. 1978; 133:429–435. [PubMed: 728692]

Figure 1. A single happy trial of the dynamic emotional faces task.^a

^a Over a 1-second duration, the participants viewed a movie of a face that changed in 5% increments from neutral (0% emotion) to a happy, sad, angry, or fearful (100% emotion) face. Participants were asked to identify the color flash presented in the middle of the dynamic change.

A Response to Angry Faces (MDD > HC)

B: Response to Happy Faces (MDD > HC)

Figure 2. Between group differences (adults with MDD > healthy controls) in right amygdala activity during processing of angry and happy emotional conditions.^a

^a The bar graphs represent the mean neural activity in the clusters within the amygdala ROI showing greater activity for adults with MDD relative to healthy controls.

Figure 3. Activity in the right amygdala during the happy emotional condition among adults with MDD.^a

^a The line graph represents mean activity in the right amygdala (anatomically defined) as a function of adjusted lifetime sub-threshold mania-related psychopathology. These values have been adjusted by first regressing out the following covariates identified for inclusion in the final regression model: Young Mania Rating Scale scores, age, and history of anxiety disorder. The graph thus represents the semi-partial correlation (i.e., the unique contribution) of mania-related psychopathology to activity in the right amygdala accounting for all of the covariates. The squared correlation coefficient from this relationship, $R^2 = .29$, is equivalent to the change in R^2 obtained by adding mania-related psychopathology to a regression model that has already accounted for the variance that could be explained by the covariates. Without mania-related psychopathology, R^2 for the covariates = .21. The change in R^2 obtained by adding mania-related psychology = .29, which is significant, $p = .002$, resulting in a model $R^2 = .50$.

Table 1

Demographic, Behavioral, and Clinical Variables

Variable	MDD (N=26)	HC (N=28)	t	z	P
Demographic					
% Female	69%	57%	-	0.84	0.36
Age	30.6 (7.8)	32.6 (6.4)	1.06	-	0.29
Behavioral					
Total % Correct	94% (6%)	96% (3%)	1.08 ^a	-	0.29
Total Reaction Time	956.2 (153.0)	928.54 (101.8)	-0.78 ^a	-	0.44
Clinical ^b					
Lifetime Mania-related Psychopathology	19.5 (12.5)	7.8 (10.0)			
Lifetime Depression-related Psychopathology	42.1 (10.1)	4.0 (6.9)			
Hamilton Depression Rating Scale	21.4 (3.9)	1.5 (2.2)			
Young Mania Rating Scale	3.7 (2.0)	0.5 (1.2)			
Psychiatric Medication:					
None	31%	100%			
Antidepressants Only	42%	-			
Antidepressants + Augmentation	27%	-			
Duration of Illness (years)	12.2 (7.4)	-			
History of Substance Abuse/Dependence	35%	0%			
History of Anxiety Disorder	65%	0%			

^a Satterthwaite method was used to correct for unequal variances between groups^b As expected, the two groups differed on all clinical variables at $p < .001$

Table 2

Reduced Regression Model of Right Amygdala Activity in Response to the Happy Condition.

Variable	t(21)		p
Lifetime Mania-related Psychopathology	0.57	3.50	0.002
Young Mania Rating Scale	−0.29	−1.85	0.08
Gender	0.43	2.75	0.01
History of Anxiety Disorder	0.31	1.96	0.06