Modulation by EEG features of BOLD responses to interictal epileptiform discharges

Pierre LeVan*, Louise Tyvaert, and Jean Gotman

Montreal Neurological Institute, McGill University, Montreal, Canada

Abstract

Introduction—EEG-fMRI of interictal epileptiform discharges (IEDs) usually assumes a fixed hemodynamic response function (HRF). This study investigates HRF variability with respect to IED amplitude fluctuations using independent component analysis (ICA), with the goal of improving the specificity of EEG-fMRI analyses.

Methods—We selected EEG-fMRI data from 10 focal epilepsy patients with a good quality EEG. IED amplitudes were calculated in an average reference montage. The fMRI data were decomposed by ICA and a deconvolution method identified IED-related components by detecting time courses with a significant HRF time-locked to the IEDs (F -test, $p < 0.05$). Individual HRF amplitudes were then calculated for each IED. Components with a significant HRF/IED amplitude correlation (Spearman test, $p < 0.05$) were compared to the presumed epileptogenic focus and to results of a general linear model (GLM) analysis.

Results—In 7 patients, at least one IED-related component was concordant with the focus, but many IED-related components were at distant locations. When considering only components with a significant HRF/IED amplitude correlation, distant components could be discarded, significantly increasing the relative proportion of activated voxels in the focus ($p = 0.02$). In the 3 patients without concordant IED-related components, no HRF/IED amplitude correlations were detected inside the brain. Integrating IED-related amplitudes in the GLM significantly improved fMRI signal modeling in the epileptogenic focus in 4 patients ($p < 0.05$).

Conclusion—Activations in the epileptogenic focus appear to show significant correlations between HRF and IED amplitudes, unlike distant responses. These correlations could be integrated in the analysis to increase the specificity of EEG-fMRI studies in epilepsy.

Keywords

Independent component analysis; EEG-fMRI; Epilepsy; Spikes

Introduction

Simultaneously acquired electroencephalogram (EEG) and functional magnetic resonance imaging (fMRI) constitutes a non-invasive modality that may be used to localize cerebral

*Corresponding author. EEG Department, Room 786, Montreal Neurological Institute, McGill University, 3801 University Street, Montreal (Qc), Canada, H3A 2B4. Fax: +1 514 398 8106. pierre.levan@mail.mcgill.ca (P. LeVan).

regions of activation in epileptic patients, which could then be used for presurgical evaluation purposes (Zijlmans et al., 2007). The method consists of detecting significant blood oxygenation level dependent (BOLD) signal changes following interictal discharges seen on the EEG recorded inside the scanner. So far, several studies have successfully detected plausible regions of activations in many patients (Al Asmi et al., 2003; Federico et al., 2005; Salek-Haddadi et al., 2006; Aghakhani et al., 2006). However, they also report a large proportion of cases without any significant response, or with responses distant from the presumed generator of epileptic activity. In general, these studies have used the general linear model (GLM) framework (Worsley and Friston, 1995; Worsley et al., 2002). This assumes that the hemodynamic response function (HRF) to interictal discharges follows a fixed canonical shape, such as the response averaged from brief auditory stimuli (Glover, 1999).

However, the HRF to sensory events may not be an appropriate representation of the HRF to pathological events such as epileptic spikes (Benar et al., 2002). Moreover, it has been shown that the HRF may vary between subjects, brain regions, and across time (Aguirre et al., 1998; Handwerker et al., 2004; Menz et al., 2006). More flexible HRF models may thus be more appropriate to detect responses with a different delay or shape than the canonical HRF (Bagshaw et al., 2004; Lu et al., 2006).

While model-driven frameworks require the prior specification of the time courses of interest, data-driven methods offer an alternative means of investigating relevant fluctuations in the BOLD signal. Independent component analysis (ICA) has been applied to fMRI data to extract patterns of activation related to specific tasks (McKeown et al., 1998). This exploratory method can extract consistent regions of activations and associated time courses based on a criterion of spatial independence, with minimal prior assumptions on the data. Careful examination of the potentially large number of extracted components may reveal regions of activation related to the epileptic discharges (Rodionov et al., 2007; LeVan and Gotman, 2009). ICA does not impose any constraints on the HRF; rather, an analysis of the component time courses may yield information about the BOLD response that had not been initially hypothesized. In the context of EEG-fMRI acquisitions of epilepsy patients, ICA could also be used to decompose the EEG data prior to the fMRI analysis (Mirsattari et al., 2006; Jann et al., 2008; Marques et al., 2009), but the current study will focus on the use of ICA directly on the fMRI data.

Advancements in artifact correction techniques now result in a high quality EEG in the course of simultaneous EEG-fMRI studies (Allen et al., 2000; Benar et al., 2003; Srivastava et al., 2005). This not only allows the identification of interictal epileptiform discharges (IEDs), but also their characterization. Features of IEDs such as amplitude may modulate the BOLD response, yielding variability in the HRF that could contribute to a loss of sensitivity in model-driven analyses. The current study thus proposes to use ICA to investigate the variability of the HRF in relation to IED amplitudes measured on the EEG.

Methods

Data acquisition

Subjects were selected from a database of focal epilepsy patients who underwent continuous EEG-fMRI recordings in a 3T Siemens Trio scanner with a CP head coil between March 2007 and May 2008. After a standard T1-weighted anatomical scan (TR=23 ms, TE=7.4 ms, flip angle of 30°, 1 mm isotropic voxel size, 256×256 matrix, 176 sagittal slices), BOLD-EPI data were acquired using the following scanning parameters: TR=1.75 s, TE=30 ms, flip angle of 90°, 5 mm isotropic voxel size, 64×64 matrix, 25 transverse slices. The session lasted up to 90 min, with short pauses every 6 min to inquire about the patient's comfort and willingness to continue with the acquisition. We discarded any 6-min run in which the patient moved by more than 1 mm, as determined by subsequent motion correction. During the fMRI scanning, simultaneous EEG, low-pass filtered at 1 kHz and sampled at 5 kHz, was recorded using 25 MR-compatible Ag/AgCl electrodes and a BrainAmp amplifier (Brain Products, Gilching, Germany). The electrodes were positioned according to the 10–20 system (19 standard scalp locations), with additional electrodes at F9-T9-P9 and F10-T10-P10, and referenced to FCz. Two additional electrodes were placed on the back to record the electrocardiogram. The electrodes were adjusted until all impedances were below 5kΩ; this was monitored throughout the scanning session. After the acquisition, scanner gradient artifact was removed by an averaged subtraction method (Allen et al., 2000) implemented in BrainVision Analyzer software (Brain Products, Gilching, Germany). Ballistocardiographic artifact was eliminated by an ICA method (Benar et al., 2003; Srivastava et al., 2005).

A neurologist reviewed the EEGs of consecutive patients, without knowledge of the fMRI analysis results, until 10 patients were obtained who satisfied the following inclusion criteria: EEG of good quality without residual gradient or ballistocardiographic artifacts and at least 20 IEDs of similar morphology as those recorded in previous EEG investigations outside the scanner. It should also be noted that patients had been initially recruited for the EEG-fMRI acquisition on the basis of having frequent IEDs. The neurologist carefully marked, in an average reference montage, the start, peak, and end of each IED, on the channel in which they were the most prominent (see Fig. 1). In case the peak marker was not exactly at the IED peak, a computer program was used to automatically move the marker to the nearest local maximum within 25 ms. IED amplitude was then defined as the average of the start-to-peak and peak-to-end amplitude. In the case of spike-and-waves, only the spike was marked. If different IED types were present (for example, left and right temporal spikes occurring independently), only the most frequent IED type was considered in the following analysis. The description of the patients and their IEDs is shown in Table 1. No major differences were noted in any patient between the morphology and frequency of the IEDs recorded inside and outside the scanner.

Data analysis

ICA decomposition—The fMRI data were motion-corrected and smoothed with a Gaussian kernel of full-width at half-maximum of 6 mm. The resulting preprocessed data then underwent an ICA decomposition using the MELODIC toolbox (Beckmann and Smith,

2004) implemented in FSL (Smith et al., 2004). This involved correcting for slice timing differences using sinc interpolation, high-pass filtering to remove scanner drift, masking to remove voxels outside the brain (Smith, 2002), and calculating the independent components by an iterative fixed-point method maximizing the non-Gaussianity of the sources, which has been shown to be equivalent to maximizing statistical independence (Hyvarinen and Oja, 2000). The number of extracted sources was determined by performing 20 repetitions of the ICA decomposition with random initialization, as true sources are likely to appear consistently in multiple ICA decompositions (Himberg et al., 2004; LeVan and Gotman, 2009). The reproducible components were then identified using the RAICAR algorithm (Yang et al., 2008), which matches components across ICA realizations based on their pairwise spatial cross-correlation coefficient. The goal was then to identify components related to the IEDs and quantify the amplitude of the BOLD response to each individual IED.

Identification of components related to IEDs—While ICA is purely data-driven, some modeling assumptions are necessary to detect components representing BOLD changes related to IEDs. To minimize the constraints on the HRF, the BOLD responses were modeled as a convolution of the events with a windowed Fourier basis (Josephs and Henson, 1999; Jacobs et al., 2009). The Fourier basis consisted of a linear combination of 5 sine and cosine functions multiplied by a Tukey window spanning an interval from 10 s before to 20 s after the marked events. The sine–cosine pairs had periods equal to submultiples of the 30-s window, and their number was limited to 5 so that only relatively low-frequency components of the HRF would be fitted, as higher-frequency sinusoids would be more likely to fit noise. The 30-s interval was chosen to accommodate the wide variety of HRF peak delays that have been observed in previous studies, including HRFs preceding the EEG discharges (Bagshaw et al., 2004; Hawco et al., 2007; Moeller et al., 2008; Jacobs et al., 2009). The HRF was assumed to have the same shape for each IED, and the amplitude of each basis function was fitted using a least-squares criterion.

This approach resulted in a deconvolved HRF for each component time course. In components related to IEDs, it was hypothesized that this HRF would account for a large amount of signal variance. This was statistically tested using an F -test ($p < 0.05$, corrected for the number of components), with the motion parameters used as confounds. Noise was modeled as an autoregressive (AR) process estimated from the residuals using the Yule-Walker equations. The AR order was determined by performing the estimation using gradually increasing AR orders, until the residuals no longer contained significant autocorrelations (Ljung-Box test, $p < 0.01$). The AR parameters were used to compute an adjusted F -statistic with appropriate effective degrees of freedom (Krugger et al., 2002). Statistically significant F -values were then indicative of components significantly related to the IEDs.

Unlike the GLM analysis, in which the analyzed signals correspond to individual voxels, ICA components are associated with a spatial map encompassing the entire brain. It is thus necessary to threshold these maps to identify in each component the voxels that are implicated. This was accomplished using the methods implemented in the MELODIC toolbox (Beckmann and Smith, 2004), in which the spatial maps are modeled by a mixture

of one Gaussian and two Gamma distributions, representing background noise, positive, and negative voxel intensities, respectively. A voxel was considered as activated if it had a 99.9% or greater probability of not belonging to background noise. The results were then displayed in a single summary map for each patient indicating, at each activated voxel, the variance accounted by all components that were significantly related to the IEDs. Moreover, two different color scales were used to distinguish positive (activation) from negative (deactivation) response. The sign of the response was determined from the direction of the average signal change of the deconvolved HRFs.

Determination of the response amplitude in individual IEDs—Up to now, the components related to IEDs were identified with the assumption that the HRF, although of arbitrary shape, was identical for each discharge. This assumption was then relaxed by fixing the HRF shape, but allowing a variable amplitude for each individual response (Fig. 2). The amplitudes were determined by least-squares fitting of the following model:

$$Y(t) = \sum \alpha_i h(t - t_i) + \sum \beta_j x_j(t) + \varepsilon,$$

where $Y(t)$ is the component time course, $h(t)$ is the deconvolved HRF shape calculated previously, the t_i s are the times of occurrence of IEDs, the x_j s represent confounds, the α_i s and β_j s are the coefficients to be fitted, and ε is zero-mean Gaussian noise. Confounds included temporal trends and motion parameters.

The HRF amplitudes α_i s were then plotted against the amplitudes of the corresponding IEDs on the EEG. The relationship between HRF and IED amplitudes was assessed using Spearman's correlation coefficient, with the significance level set at $p < 0.05$. New summary maps were then created for each patient using only the components that showed a significant correlation.

It should be noted that the fitted HRF amplitudes are unlikely to be accurate representations of individual response amplitudes for closely spaced IEDs, as the HRFs would then overlap in a non-linear fashion (Friston et al., 1998a). Nevertheless, although a canonical HRF commonly used in fMRI studies (Glover, 1999) only resolves to baseline after approximately 25 s, overlapping HRFs have been found to superpose approximately linearly for intervals as short as 2 s (Friston et al., 1999). The correlations between HRF and IED amplitudes were thus only calculated for IEDs separated from any other IED by at least 2 s.

GLM processing—For comparison purposes, the fMRI data were also analyzed using the GLM implemented in the FMRISTAT software package (Worsley et al., 2002). The BOLD response to IEDs was modeled using 4 HRFs peaking at 3, 5, 7, and 9 s (Bagshaw et al., 2004). Polynomial baseline drifts and motion parameters were treated as confounds. The resulting statistical t -maps were thresholded at $|t| > 3.1$, with an additional spatial extent threshold of 5 voxels to obtain a corrected family-wise error p -value of 0.05 (Friston et al., 1994; Worsley et al., 2002).

To assess whether integrating information about the IEDs could improve the GLM results, a second analysis was performed with a model accounting for HRF fluctuations related to IED

amplitudes. This model actually used two regressors of interest. The first regressor used HRFs with fixed amplitudes, and was thus identical to the regressor of interest used in the previous analysis. The second regressor used HRFs whose amplitudes were proportional to the IED amplitudes, and then orthogonalized to the first regressor (Fig. 3). This model structure ensured that the two analyses would be nested. In this way, statistically significant t -values associated with the second regressor of this second analysis would indicate voxels where the BOLD response showed a significant additional effect that could not be detected in the first analysis.

Results

ICA decomposition

The ICA decomposition identified a number of reproducible components ranging from 172 to 373 (see Table 2). After identifying IED-related components using the HRF deconvolution method, brain areas corresponding to these components were visually classified as concordant or discordant with the presumed epileptogenic focus of each patient. This classification was performed on the basis of the available electro-clinical data (shown in Table 1). Four patients (2, 3, 5, and 6) had focal MRI findings (either dysplastic lesions or mesial temporal abnormalities, see Table 1) consistent with the IED topography and clinical semiology. For these patients, components were considered concordant if their spatial topography included lesional or peri-lesional areas. For the remaining 6 patients, one (patient 7) had discordant findings (left hippocampal atrophy, but right-sided spikes), one (patient 4) had generalized cerebral atrophy, and the remaining four had negative MRI findings. For these 6 patients, the localization was only defined at the lobar scale based on the spatial distribution of the IEDs. Components were considered concordant if their spatial topography was completely or partially within the cerebral lobe containing the presumed generator of IEDs.

One patient (patient 8) did not have any IED-related components. In the remaining 9 patients, between 1 and 72 IED-related components were identified. In 7 of these 9 patients, there was at least one IED-related concordant component, while the other 2 patients (patients 9 and 10) only had discordant components (see Table 2).

HRF and IED amplitude correlation in patients with concordant components

Among the 7 patients with concordant components, the identified concordant regions accounted for an average of 33.8% of the total volume of IED-related components (Table 2). This indicates that a large proportion of the detected responses were distant from the presumed epileptogenic focus. In particular, patients with a large number of IED-related components had corresponding widespread patterns of BOLD response. An example of such a patient is shown in Fig. 4. In this example, the identified brain regions include the presumed left central epileptogenic focus, but lack specificity. However, when considering components for which HRF amplitudes were significantly correlated to IED amplitudes, most of the distant responses could be discarded. Further examples of components with significant HRF and IED amplitude correlation are shown in Fig. 5. Other patients will be presented in the following sections.

In 6 of the 7 patients with concordant components, at least one of the IED-related components showed a significant correlation between HRF and IED amplitudes (see Table 2). When considering only components with such correlation, concordant components now accounted for an average of 64.8% of the volume of the response. This represented a significant increase from before (paired t -test, $p=0.02$). Although there were still some significant responses outside the presumed epileptogenic focus, the proportion of discordant regions was much smaller than in the original analysis. In fact, there was only one patient (patient 6) for whom the proportion of concordant areas did not increase when considering only components with a significant correlation between HRF and IED amplitudes (see Fig. 6). In this patient, the epileptogenic focus was presumed to be located in a left parietal cortical dysplasia, but significant correlations were only found more anteriorly in central regions, bilaterally (positive correlation on the right, and negative on the left). In the remaining 5 patients, concordant components showed a significant positive correlation between HRF and IED amplitudes.

For the 2 patients (patients 4 and 6) that did not show significant correlations between HRF and IED amplitudes in concordant regions, the smallest p -value of the correlation was 0.2, which is far from the threshold of significance. The absence of significant correlations thus does not appear to be due to a thresholding issue.

HRF and IED amplitude correlation in patients without concordant components

As mentioned previously, 1 patient (patient 8) did not have any IED-related components, and 2 patients (patients 9 and 10) only had discordant components. Patient 9 had a normal MRI and left temporal spikes, but only had a single IED-related component in the caudate nuclei. That component did not show a significant HRF and IED amplitude correlation (see Table 2). Patient 10 had a normal MRI and right temporal spikes, but no IED-related components were identified in the right temporal lobe. Only one component showed a significant IED and HRF amplitude correlation. However, that component was likely artifactual, as it involved voxels outside the brain, in the sphenoidal sinuses, that had not been discarded by the automatic brain masking procedure. This region is known to be affected by susceptibility artifacts that may violate fMRI modeling assumptions, yielding spurious activations (Wu et al., 1997; Parrish et al., 2000). Therefore, in all 3 patients who did not have concordant components, the method did not misleadingly identify clusters in discordant brain regions.

To determine whether the absence of concordant components was due to a statistical thresholding issue, it was also attempted to lower the significance threshold from $p < 0.05$ (corrected) to $p < 0.001$ (uncorrected). This resulted in the identification of additional components for all 3 patients. All components were in discordant areas for patients 8 and 9, while a concordant component in the right temporal lobe was found for patient 10 ($p=0.0005$, uncorrected). However, that component did not show a significant correlation between HRF and IED amplitudes ($\rho=0.25$, $p=0.2$).

Comparison with GLM

Analysis with the GLM framework resulted in concordant activations in 7 patients: 6 of 7 patients who had such activations with the ICA method, and 1 of 3 patients who did not have

concordant ICA results (see Table 3). In 2 of the patients with concordant activations, adjacent concordant deactivations were present as well. Similar to the ICA analysis, the maps included clusters of significant BOLD response distant from the presumed epileptogenic focus in all 10 patients. In 4 patients, the maximum positive peak of the statistical map was part of a concordant activation cluster, and in one other patient, the maximum negative peak of the statistical map was part of a concordant deactivation cluster. In addition, a special case is patient 2, for whom the maximum positive peak was part of a discordant activation cluster in the ventricles. This activation was clearly artifactual; when ignoring this cluster, the highest remaining positive peak was now in a concordant right temporal activation cluster.

The analysis was then repeated with an additional regressor modeling HRF amplitude fluctuations as proportional to IED amplitudes. Statistical maps showing areas where this regressor modeled a significant amount of additional signal variance revealed regions in the presumed epileptogenic focus in 4 patients (patients 1, 2, 3, and 10), including 1 (patient 10) in whom the concordant regions had not been identified in the original analysis (see Table 3). These maps also identified fewer discordant regions than originally. Examples are shown in Figs. 7 and 8. However, there were also 4 patients (patients 4, 5, 7, and 8) for whom the regressor modeling IED amplitude fluctuations was not significant in the epileptogenic focus, even though this area was detected by the original GLM analysis.

Discussion

Data-driven vs model-driven methods

The relationship between the BOLD response to IEDs and the amplitude of these IEDs on EEG was studied using ICA. The detection of BOLD responses using model-driven methods such as the GLM is only optimal if the modeling assumptions are correct. GLM analyses of event-related fMRI data commonly employ a canonical HRF shape, sometimes with multiple HRF delays (Bagshaw et al., 2004) or a temporal derivative regressor (Friston et al., 1998b) to account for temporal shifts. More flexible basis sets may also be employed to model arbitrary HRF shapes (Josephs and Henson, 1999; Lu et al., 2006). However, these models do not account for inter-trial variability. Model-driven analyses also require an accurate characterization of the noise confounding the signal of interest. Some sources of structured noise such as patient movement (Friston et al., 1996) or physiological artifacts (Glover et al., 2000) can now be integrated into the analysis. It has also been demonstrated that the state of vigilance, as measured by EEG rhythms, significantly affects the BOLD signal (Tyvaert et al., 2008a). However, it may become impractical to explicitly model all sources of BOLD signal variability.

In contrast, data-driven methods such as ICA do not require strong constraints on the BOLD signal. The various sources composing the fMRI data are automatically separated, so that sources of structured noise do not need to be explicitly specified (Thomas et al., 2002). However, modeling assumptions are still required to identify components of interest, namely the IED-related fluctuations. Our method used flexible constraints to identify HRFs of arbitrary shapes and inter-trial amplitude fluctuations. This resulted in a non-linear model, so the shape and amplitudes were fitted separately in successive linear least-squares iterations

(Hinrichs et al., 2000). It would be possible to perform additional iterations, but it was found that this did not result in significant changes in the parameter estimates.

HRF amplitude fluctuations

The method was able to detect significant correlations between the HRF and IED amplitudes in the presumed epileptogenic focus. While such a relationship was expected, given the coupling of the BOLD signal with neuronal activity, this demonstrates that artifact correction on the EEG recorded inside the scanner is sufficient to perform meaningful measurements of IED features. A previous study could only detect positive, but not statistically significant correlations between IED and HRF amplitudes (Benar et al., 2002). That study defined the HRF amplitude as the energy of the detrended BOLD signal in the time period following IEDs. This calculation required the HRFs to be non-overlapping, so only IEDs separated by at least 30 s were considered. After applying this strict constraint, only a few IEDs remained, resulting in low statistical power. In contrast, the current method used a deconvolution approach that could separate overlapping HRFs. This assumes that linear superposition applies, which was shown to be the case for events separated by at least 2 s (Friston et al., 1999).

While the current study investigated the correlation between IED and HRF amplitudes within the brain regions represented by the ICA components, there are several factors that could confound this relationship. IED amplitudes depend not only on the number of active neurons, but also on their level of synchrony. However, a change in synchrony alone would not necessarily be associated with increased metabolic demand and thus a BOLD response. As for an increase in the number of active neurons, this may manifest itself as an increase in active cortical area (Tao et al., 2005). However, neither the GLM, which examines individual voxels, nor ICA, which extracts fixed spatial patterns, are ideally suited for the investigation of fluctuating areas of activation.

Nevertheless, there are indications that IED amplitude fluctuations are also associated with local BOLD signal changes within the scale of one MRI voxel or one ICA component, and can thus be detected by the proposed method. In a study with simultaneous MEG and SEEG recordings, some patients showed a significant correlation between MEG spikes and intracranial spikes (Santiuste et al., 2008); the potentials recorded by the SEEG presumably represent the activity in a very small volume (<1 cm) around each depth electrode contact. Also, studies with simultaneous scalp EEG and ECoG recordings have shown that while large scalp IEDs are associated with large consistent areas of active cortex, small-amplitude scalp IEDs may result from small intracranial sources with variable localizations (Tao et al., 2005; Ray et al., 2007). On average, a given MRI voxel would thus show a smaller BOLD response to small IEDs.

The identification of a correlation between IED and HRF amplitudes has some important implications for the analysis of EEG-fMRI data of epileptic patients. The current study has demonstrated that the modeling of the BOLD response could be improved by integrating IED amplitude information in the GLM. Brain areas have been identified in which a regressor modeling HRF amplitude fluctuations accounted for a statistically significant amount of variance of the BOLD signal, additional to the variance already explained by a

regressor modeling constant HRF amplitudes. Statistical maps based on the regressor modeling IED-related HRF amplitude fluctuations may provide a more specific identification of the epileptogenic focus. Alternatively, an F -statistic map testing the effect of the combination of the 2 regressors should show better sensitivity in detecting IED-related BOLD changes in the focus. The current study only looked at amplitude correlations, but future investigations could further characterize the relationship between HRFs and IEDs. For example, the BOLD signal may follow a complex, possibly non-linear, function of IED amplitudes or other IED features, which could then be integrated into the GLM analysis (Buchel et al., 1998).

It may also not be necessary to go back to a model-driven analysis, as the ICA method could already detect the IED-related HRF fluctuations. There were some slight differences between the GLM and ICA maps; they may be due to the distinct thresholding methods, but there is also the effect of the pre-specification of the HRF shape and amplitude in the GLM. In contrast, the ICA method allows for arbitrary HRFs. It is especially well-suited as an exploratory analysis to characterize the BOLD response from the time course of IED-related components (Rodionov et al., 2007; LeVan and Gotman, 2009). It is only after characterizing the relationship between IEDs and HRFs that it is then possible to integrate it into the model. A possible methodology for the analysis of EEG-fMRI data could thus consist of a standard GLM analysis to provide a first estimate of the activation, after which an ICA could be used to further characterize the relationship between IEDs and HRFs. Improvements in modeling artifacts and other confounding factors may, for example, allow the identification of non-linearities that are unknown a priori and thus cannot be modeled in the GLM. New hypotheses based on the ICA results may then be integrated in the GLM framework to refine the initial activations.

The results from this study have also shown that HRF fluctuations specifically related to IED amplitudes occur mainly in the epileptogenic focus. In general, BOLD signal changes to IEDs often involve the focus, but may also be present in distant regions (Salek-Haddadi et al., 2006; Benar et al., 2006). Some brain regions, such as the “default-mode” network (Raichle et al., 2001), may show significant BOLD signal changes, even though they are only indirectly related to IEDs and are not part of the epileptogenic network (Gotman et al., 2005; Laufs et al., 2007). Deactivations were identified in some of those regions (precuneus, posterior cingulate, bilateral inferior parietal lobes, and medial prefrontal cortex) in 4 patients in the current study (see Table 2).

Many studies have simplified the evaluation of widespread BOLD response clusters by considering mainly the location of the voxel with maximum t -statistic value in the GLM analysis (Salek-Haddadi et al., 2006; Tyvaert et al., 2008b). However, such interpretation should be treated with caution, as hemodynamic coupling shows spatial variability, implying that differences in BOLD signal levels between two regions do not necessarily correspond to different levels of neuronal activity (Ances et al., 2008). Nevertheless, in the current study, the maximum positive peak of the GLM map could identify a concordant activation cluster in 5 patients (when ignoring clearly artifactual clusters outside the brain), and the maximum negative peak could identify a concordant deactivation cluster in 1 patient. However, in the remaining 4 patients, the location of the peaks indicated misleading discordant regions. In

contrast, ICA components showing a significant correlation between HRF and IED amplitudes could also be used to identify concordant clusters and reject distant responses, but with better specificity. There was only one patient for whom the ICA method only identified discordant clusters inside the brain with a significant HRF/IED amplitude correlation. In the remaining 9 patients, either the clusters with a significant HRF/IED amplitude correlation included concordant areas with fewer distant regions (5 cases), or no clusters inside the brain showed a significant HRF/IED correlation (4 cases). The method could thus detect regions that are specifically related to IEDs while discarding distant regions that are only indirectly related to the epileptic network. This resulted in a reduced proportion of discordant areas, but some distant clusters still remained.

Further studies would be necessary to investigate how clusters apparently distant from the epileptogenic focus could show consistent BOLD changes following IEDs, and whose HRF amplitudes are related to IED amplitudes. Some of the distant clusters may result from resting-state networks such as the default-mode areas, which are only indirectly related to the IEDs. Moreover, artifacts may cause spurious correlations, particularly if they involve signal transients that violate the stationarity assumptions of the statistical model. This problem may be alleviated by the use of ICA classification methods that have been developed to detect and discard artifactual components (De Martino et al., 2007; Perlberg et al., 2007). Further investigations may also be necessary to determine whether significant negative correlations between HRF and IED amplitudes, such as those found for patients 5 and 6, represent a true physiological phenomenon, for example related to inhibition, or whether these correlations are artifactual. In any case, while the proposed method appears to improve the specificity of EEG-fMRI analyses, some discordant results were still present. Interpretation of the results should thus still be done in combination with the other available electro-clinical modalities.

Conclusion

ICA can be used as an exploratory method in EEG-fMRI data from epilepsy patients to investigate the relationship between EEG features of IEDs and the BOLD response. In particular, BOLD activations in the epileptogenic focus appear to show significant correlations between HRF and IED amplitudes, unlike distant responses. An analysis based on these correlations could increase the specificity of EEG-fMRI studies in epilepsy, and might also be integrated into the model-driven GLM framework.

Acknowledgments

This project was supported by the Canadian Institutes of Health Research grant MOP-38079 and the Natural Sciences and Engineering Research Council of Canada scholarship CGSD.

References

- Aghakhani Y, Kobayashi E, Bagshaw A, Hawco C, Benar C, Dubeau F, Gotman J. Cortical and thalamic fMRI responses in partial epilepsy with focal and bilateral synchronous spikes. *Clin Neurophysiol.* 2006; 117:177–191. [PubMed: 16314143]
- Aguirre G, Zarahn E, D'Esposito M. The variability of human, BOLD hemodynamic responses. *NeuroImage.* 1998; 8:360–369. [PubMed: 9811554]

- Al Asmi A, Benar C, Gross D, Khani Y, Andermann F, Pike B, Dubeau F, Gotman J. fMRI activation in continuous and spike-triggered EEG-fMRI studies of epileptic spikes. *Epilepsia*. 2003; 44:1328–1339. [PubMed: 14510827]
- Allen P, Josephs O, Turner R. A method for removing imaging artifact from continuous EEG recorded during functional MRI. *NeuroImage*. 2000; 12:230–239. [PubMed: 10913328]
- Ances B, Leontiev O, Perthen J, Liang C, Lansing A, Buxton R. Regional differences in the coupling of cerebral blood flow and oxygen metabolism changes in response to activation: implications for BOLD-fMRI. *NeuroImage*. 2008; 39:1510–1521. [PubMed: 18164629]
- Bagshaw A, Aghakhani Y, Benar C, Kobayashi E, Hawco C, Dubeau F, Pike G, Gotman J. EEG-fMRI of focal epileptic spikes: analysis with multiple haemodynamic functions and comparison with gadolinium-enhanced MR angiograms. *Hum Brain Mapp*. 2004; 22:179–192. [PubMed: 15195285]
- Beckmann C, Smith S. Probabilistic independent component analysis for functional magnetic resonance imaging. *IEEE Trans Med Imaging*. 2004; 23:137–152. [PubMed: 14964560]
- Benar C, Aghakhani Y, Wang Y, Izenberg A, Al Asmi A, Dubeau F, Gotman J. Quality of EEG in simultaneous EEG-fMRI for epilepsy. *Clin Neurophysiol*. 2003; 114:569–580. [PubMed: 12705438]
- Benar C, Gross D, Wang Y, Petre V, Pike B, Dubeau F, Gotman J. The BOLD response to interictal epileptiform discharges. *NeuroImage*. 2002; 17:1182–1192. [PubMed: 12414258]
- Benar C, Grova C, Kobayashi E, Bagshaw A, Aghakhani Y, Dubeau F, Gotman J. EEG-fMRI of epileptic spikes: concordance with EEG source localization and intracranial EEG. *NeuroImage*. 2006; 30:1161–1170. [PubMed: 16413798]
- Buchel C, Holmes A, Rees G, Friston K. Characterizing stimulus-response functions using nonlinear regressors in parametric fMRI experiments. *NeuroImage*. 1998; 8:140–148. [PubMed: 9740757]
- De Martino F, Gentile F, Esposito F, Balsi M, Di Salle F, Goebel R, Formisano E. Classification of fMRI independent components using IC-fingerprints and support vector machine classifiers. *NeuroImage*. 2007; 34:177–194. [PubMed: 17070708]
- Federico P, Archer J, Abbott D, Jackson G. Cortical/Subcortical BOLD changes associated with epileptic discharges: an EEG-fMRI study at 3 T. *Neurology*. 2005; 64:1125–1130. [PubMed: 15824333]
- Friston K, Fletcher P, Josephs O, Holmes A, Rugg M, Turner R. Event-related fMRI: characterizing differential responses. *NeuroImage*. 1998b; 7:30–40. [PubMed: 9500830]
- Friston K, Josephs O, Rees G, Turner R. Nonlinear event-related responses in fMRI. *Magn Reson Med*. 1998a; 39:41–52. [PubMed: 9438436]
- Friston K, Williams S, Howard R, Frackowiak R, Turner R. Movement-related effects in fMRI time-series. *Magn Reson Med*. 1996; 35:346–355. [PubMed: 8699946]
- Friston K, Worsley K, Frackowiak R, Mazziotta J, Evans A. Assessing the significance of focal activations using their spatial extent. *Hum Brain Mapp*. 1994; 1:210–220. [PubMed: 24578041]
- Friston K, Zarahn E, Josephs O, Henson R, Dale A. Stochastic designs in event-related fMRI. *NeuroImage*. 1999; 10:607–619. [PubMed: 10547338]
- Glover G. Deconvolution of impulse response in event-related BOLD fMRI. *NeuroImage*. 1999; 9:416–429. [PubMed: 10191170]
- Glover G, Li T, Ress D. Image-based method for retrospective correction of physiological motion effects in fMRI: RETROICOR. *Magn Reson Med*. 2000; 44:162–167. [PubMed: 10893535]
- Gotman J, Grova C, Bagshaw A, Kobayashi E, Aghakhani Y, Dubeau F. Generalized epileptic discharges show thalamocortical activation and suspension of the default state of the brain. *Proc Natl Acad Sci U S A*. 2005; 102:15236–15240. [PubMed: 16217042]
- Handwerker D, Ollinger J, D'Esposito M. Variation of BOLD hemodynamic responses across subjects and brain regions and their effects on statistical analyses. *NeuroImage*. 2004; 21:1639–1651. [PubMed: 15050587]
- Hawco C, Bagshaw A, Lu Y, Dubeau F, Gotman J. BOLD changes occur prior to epileptic spikes seen on scalp EEG. *NeuroImage*. 2007; 35:1450–1458. [PubMed: 17399999]
- Himberg J, Hyvarinen A, Esposito F. Validating the independent components of neuroimaging time series via clustering and visualization. *NeuroImage*. 2004; 22:1214–1222. [PubMed: 15219593]

- Hinrichs H, Scholz M, Tempelmann C, Woldorff M, Dale A, Heinze H. Deconvolution of event-related fMRI responses in fast-rate experimental designs: tracking amplitude variations. *J Cogn Neurosci*. 2000; 12 (Suppl 2):76–89. [PubMed: 11506649]
- Hyvarinen A, Oja E. Independent component analysis: algorithms and applications. *Neural Netw*. 2000; 13:411–430. [PubMed: 10946390]
- Jacobs J, LeVan P, Moeller F, Boor R, Stephani U, Gotman J, Siniatchkin M. Hemodynamic changes preceding the interictal EEG spike in patients with focal epilepsy investigated using simultaneous EEG-fMRI. *NeuroImage*. 2009; 45:1220–1231. [PubMed: 19349236]
- Jann K, Wiest R, Hauf M, Meyer K, Boesch C, Mathis J, Schroth G, Dierks T, Koenig T. BOLD correlates of continuously fluctuating epileptic activity isolated by independent component analysis. *NeuroImage*. 2008; 42:635–648. [PubMed: 18585061]
- Josephs O, Henson R. Event-related functional magnetic resonance imaging: modelling, inference and optimization. *Philos Trans R Soc Lond B Biol Sci*. 1999; 354:1215–1228. [PubMed: 10466147]
- Kruggel F, Pelegrini-Issac M, Benali H. Estimating the effective degrees of freedom in univariate multiple regression analysis. *Med Image Anal*. 2002; 6:63–75. [PubMed: 11836135]
- Laufs H, Hamandi K, Salek-Haddadi A, Kleinschmidt A, Duncan J, Lemieux L. Temporal lobe interictal epileptic discharges affect cerebral activity in “default mode” brain regions. *Hum Brain Mapp*. 2007; 28:1023–1032. [PubMed: 17133385]
- LeVan P, Gotman J. Independent component analysis as a model-free approach for the detection of BOLD changes related to epileptic spikes: a simulation study. *Hum Brain Mapp*. 2009; 30:2021–2031. [PubMed: 18726909]
- Lu Y, Bagshaw A, Grova C, Kobayashi E, Dubeau F, Gotman J. Using voxel-specific hemodynamic response function in EEG-fMRI data analysis. *NeuroImage*. 2006; 32:238–247. [PubMed: 16774839]
- Marques J, Rebola J, Figueiredo P, Pinto A, Sales F, Castelo-Branco M. ICA decomposition of EEG signal for fMRI processing in epilepsy. *Hum Brain Mapp*. 2009; 30:2986–2996. [PubMed: 19172633]
- McKeown M, Makeig S, Brown G, Jung T, Kindermann S, Bell A, Sejnowski T. Analysis of fMRI data by blind separation into independent spatial components. *Hum Brain Mapp*. 1998; 6:160–188. [PubMed: 9673671]
- Menz M, Neumann J, Muller K, Zysset S. Variability of the BOLD response over time: an examination of within-session differences. *NeuroImage*. 2006; 32:1185–1194. [PubMed: 16857390]
- Mirsattari S, Wang Z, Ives J, Bihari F, Leung L, Barth A, Menon R. Linear aspects of transformation from interictal epileptic discharges to BOLD fMRI signals in an animal model of occipital epilepsy. *NeuroImage*. 2006; 30:1133–1148. [PubMed: 16414283]
- Moeller F, Siebner H, Wolff S, Muhle H, Boor R, Granert O, Jansen O, Stephani U, Siniatchkin M. Changes in activity of striato-thalamo-cortical network precede generalized spike wave discharges. *NeuroImage*. 2008; 39:1839–1849. [PubMed: 18082429]
- Parrish T, Gitelman D, LaBar K, Mesulam M. Impact of signal-to-noise on functional MRI. *Magn Reson Med*. 2000; 44:925–932. [PubMed: 11108630]
- Perlberg V, Bellec P, Anton J, Pelegrini-Issac M, Doyon J, Benali H. CORSICA: correction of structured noise in fMRI by automatic identification of ICA components. *Magn Reson Imaging*. 2007; 25:35–46. [PubMed: 17222713]
- Raichle M, MacLeod A, Snyder A, Powers W, Gusnard D, Shulman G. A default mode of brain function. *Proc Natl Acad Sci USA*. 2001; 98:676–682. [PubMed: 11209064]
- Ray A, Tao J, Hawes-Ebersole S, Ebersole J. Localizing value of scalp EEG spikes: a simultaneous scalp and intracranial study. *Clin Neurophysiol*. 2007; 118:69–79. [PubMed: 17126071]
- Rodionov R, De Martino F, Laufs H, Carmichael D, Formisano E, Walker M, Duncan J, Lemieux L. Independent component analysis of interictal fMRI in focal epilepsy: comparison with general linear model-based EEG-correlated fMRI. *NeuroImage*. 2007; 38:488–500. [PubMed: 17889566]
- Salek-Haddadi A, Diehl B, Hamandi K, Merschhemke M, Liston A, Friston K, Duncan J, Fish D, Lemieux L. Hemodynamic correlates of epileptiform discharges: an EEG-fMRI study of 63 patients with focal epilepsy. *Brain Res*. 2006; 1088:148–166. [PubMed: 16678803]

- Santiuste M, Nowak R, Russi A, Tarancon T, Oliver B, Ayats E, Scheler G, Graetz G. Simultaneous magnetoencephalography and intracranial EEG registration: technical and clinical aspects. *J Clin Neurophysiol*. 2008; 25:331–339. [PubMed: 18997623]
- Smith S. Fast robust automated brain extraction. *Hum Brain Mapp*. 2002; 17:143–155. [PubMed: 12391568]
- Smith S, Jenkinson M, Woolrich M, Beckmann C, Behrens T, Johansen-Berg H, Bannister P, De Luca M, Drobnjak I, Flitney D, Niazy R, Saunders J, Vickers J, Zhang Y, De Stefano N, Brady J, Matthews P. Advances in functional and structural MR image analysis and implementation as FSL. *NeuroImage*. 2004; 23 (Suppl 1):S208–S219. [PubMed: 15501092]
- Srivastava G, Crottaz-Herbette S, Lau K, Glover G, Menon V. ICA-based procedures for removing ballistocardiogram artifacts from EEG data acquired in the MRI scanner. *NeuroImage*. 2005; 24:50–60. [PubMed: 15588596]
- Tao J, Ray A, Hawes-Ebersole S, Ebersole J. Intracranial EEG substrates of scalp EEG interictal spikes. *Epilepsia*. 2005; 46:669–676. [PubMed: 15857432]
- Thomas C, Harshman R, Menon R. Noise reduction in BOLD-based fMRI using component analysis. *NeuroImage*. 2002; 17:1521–1537. [PubMed: 12414291]
- Tyvaert L, Hawco C, Kobayashi E, LeVan P, Dubeau F, Gotman J. Different structures involved during ictal and interictal epileptic activity in malformations of cortical development: an EEG-fMRI study. *Brain*. 2008b; 131:2042–2060. [PubMed: 18669486]
- Tyvaert L, LeVan P, Grova C, Dubeau F, Gotman J. Effects of fluctuating physiological rhythms during prolonged EEG-fMRI studies. *Clin Neurophysiol*. 2008a; 119:2762–2774. [PubMed: 18977169]
- Worsley K, Friston K. Analysis of fMRI time-series revisited—again. *NeuroImage*. 1995; 2:173–181. [PubMed: 9343600]
- Worsley K, Liao C, Aston J, Petre V, Duncan G, Morales F, Evans A. A general statistical analysis for fMRI data. *NeuroImage*. 2002; 15:1–15. [PubMed: 11771969]
- Wu D, Lewin J, Duerk J. Inadequacy of motion correction algorithms in functional MRI: role of susceptibility-induced artifacts. *J Magn Reson Imaging*. 1997; 7:365–370. [PubMed: 9090592]
- Yang Z, Laconte S, Weng X, Hu X. Ranking and averaging independent component analysis by reproducibility (RAICAR). *Hum Brain Mapp*. 2008; 29:711–725. [PubMed: 17598162]
- Zijlmans M, Huiskamp G, Hersevoort M, Seppenwoolde J, van Huffelen A, Leijten F. EEG-fMRI in the preoperative work-up for epilepsy surgery. *Brain*. 2007; 130:2343–2353. [PubMed: 17586868]

Fig. 1.

Marking of IED amplitudes. On the artifact-corrected EEG, the start, peak, and end of IEDs are marked on the channel where they are most prominent. In this case, right temporal spikes are marked on channel F10.

Fig. 2.

Determination of amplitudes of HRFs for individual IEDs. (A) An example component time course (blue trace) shows visible BOLD changes following the IED timings (indicated by vertical dotted lines). The red trace shows the HRFs modeled by a Fourier basis, assuming identical HRF shapes and amplitudes for each IED. The fitted shape is also shown on the right. (B) Determination of HRF amplitudes, by fixing the HRF shape calculated in panel A and allowing arbitrary amplitude fluctuations. Note how larger responses for the 1st and 3rd IEDs and a smaller response for the 4th IED in the component time course (blue trace) are now fitted by corresponding variable amplitude HRFs (red trace).

Fig. 3.

Regressors used in the GLM analysis. (A) IED timings and amplitudes. (B) Standard regressor based on IED timings, convolved with a canonical HRF of constant amplitude peaking 5 s after the IED. This is the regressor used in the original GLM analysis. (C) Regressor in which the HRF amplitudes fluctuate according to IED amplitudes. (D) Regressor from panel C, orthogonalized from the original regressor from panel B. It is equivalent to the difference between the 2 regressors. The new GLM analysis performed statistical testing on the effect of this new regressor, with the original regressor from panel B treated as a confounding factor.

Fig. 4.

ICA results for patient 5. Spike-and-waves on channels Cz–Pz–C3–P3 presumably originate from a subtle dysplastic lesion visible on T2 FLAIR MRI. IED-related components involve the area around the lesion, but also widespread distant regions. However, almost all components distant from the lesion do not show a significant correlation between HRF and IED amplitudes.

Fig. 5.

Examples of concordant IED-related components with a significant correlation between HRF and IED amplitudes. For patient 2, with a presumed right temporal epileptogenic focus, a right temporal component has a deconvolved HRF with a canonical shape with an early peak at around 2.5 s. The correlation coefficient between HRF and IED amplitudes was 0.52 ($p=0.002$). For patient 7, also with a right temporal focus, a concordant right temporal component has a fitted HRF peaking at around 4 s. The correlation coefficient between HRF and IED amplitudes was 0.51 ($p=0.01$).

Fig. 6.

ICA results for patient 6. Spikes on channels P3–Pz–O1 presumably originate from a left parietal focal cortical dysplasia. There is an ICA activation in the lesion (red arrow), along with more distant frontal and contralateral clusters. In this patient, the lesional component does not show a significant correlation between HRF and IED amplitudes.

Fig. 7.

ICA and GLM results for patient 3. Spike-and-waves on channels F3–C3 presumably originate from a left frontal cortical dysplasia. IED-related ICA components involve the lesion and widespread distant areas. However, only left frontal and parietal components show a significant correlation between HRF and IED amplitudes. The GLM map is similar to the ICA map, with widespread clusters of significant BOLD responses. However, areas related to IED amplitude fluctuations include regions around the lesion, partially overlapping the original activation, with fewer remote clusters.

Fig. 8.

ICA and GLM results for patient 1. Right temporal spikes on channels F8–T4–F10–T10 are not associated with any MRI findings. The ICA analysis reveals a concordant right temporal activation. Part of this activation also shows a significant correlation between HRF and IED amplitudes. The GLM analysis also reveals a right temporal activation, along with small scattered distant clusters. Similar to the ICA analysis, a right temporal cluster is shown to be related to IED fluctuations. This cluster partially overlaps the original activation, and the map has fewer distant clusters than in the original analysis.

Table 1
Description of the patients, including habitual seizure semiology, MRI findings, and interictal EEG.

Patient	Age/Gender	Seizure semiology	MRI	Interictal EEG	No. IEDs during scanning
1	15/M	Flashes in L visual field, feeling of déjà vu, confusion	Normal	RTP sharp waves, F8–T4–F10–T10	47
2	40/F	Aura with coldness, bad smell, descending sensation, staring, vocalization, eye blinking, oral automatisms; also one GTC	R amygdala abnormality	RT sharp waves, F8–T4–F10–T10	39
3	36/F	R arm stiffness, then R leg, then L hemibody and GTC	LFC cortical dysplasia with macrogyria	LFC spike-and-waves, F3–C3	63
4	52/F	Aura: L rising epigastric sensation followed by auditory manifestation; anxiety, right facial twitch, lip smacking and then absence; dysphasia and confusion	Generalized atrophy	LT spike-and-waves, T3–T9	101
5	34/M	Two types: 1) abrupt falls (stiffness), 2) feeling of tightness, leg movements and stiffness, R arm tonic posturing	Dysplastic lesion CP vertex	LC spike-and-waves, Cz–Pz–C3–P3	33
6	33/M	R arm sensation, then arm movement (elevation)	LCP cortical dysplasia	LP spikes, P3–Pz–O1	114
7	48/F	Aura: anxiety, feeling of déjà vu. Staring, restlessness, confusion	LH atrophy	RT sharp waves, F8–T4–F10–T10	27
8	13/F	Clonus in shoulder spreading to arms, raising of left arm and head turn to the right; body stiffness	Normal	BiCPO spike-and-waves predominantly left, C3–P3–T5–O1	21
9	34/F	Epigastric aura, vision of bright lights, LOC, scream and fear; then stiffness with lip smacking.	Normal	LT spikes, F7–T3–F9–T9	48
10	50/M	Feeling of motion, dizziness, loss of feeling on L hemiface then upper and lower limbs, staring, automatisms of R fingers	Normal	BiIT independent spikes, most are RT, F8–F10–T10	27

Abbreviations: LOC: loss of consciousness, GTC: generalized tonic-clonic seizures, R: right, L: left, BiI: bilateral, F: frontal, C: central, T: temporal, P: parietal, O: occipital.

Table 2

Results of the ICA decomposition and correlation between HRF and IED amplitudes. The presumed location of the epileptogenic focus (column 2) is determined from the available electro-clinical data. For IED-related components (columns 4 to 7) and for IED-related components with significant HRF/IED amplitude correlation (columns 8 to 11), the table shows the number and locations of the components, as well as the percentage of volume of significant response that is accounted for by concordant components. Concordant locations are indicated in bold type.

Patient	Focus	No. of ICs	No. of IED-related ICs	Percent volume of concordant ICs	ICA map	Deactivation	No. of ICs with HRF/IED amplitude correlation	HRF/IED amplitude correlation	
								Positive	Negative
1	RT	217	3	62.6	RT	RCer	1	RT	None
2	RT	185	28	17.4	BHT , RE, RP, Bil thal., brainstem, BilCer	BHT , BilP, posterior cing., precuneus, RF	5	BHT , thal.	None
3	LF	313	60	36.7	BilCP mostly L , BilTO, Bil caudate, BilCer	BilC , BilP, Bil insula	7	LC , LP	None
4	LT	294	11	41.9	LT , BilO	LT , BilO, posterior cing.	0	None	None
5	LC	223	72	12.7	BilCP , BilTPO	BilFT, Bil thal., ventricles, brainstem, BilCer	2	LCP	RP
6	LP	334	7	11.8	LCP , RT, BilF	RC	2	RC	LC
7	RT	252	5	53.2	RT , L insula, BilC	LF	1	RT	None
8	LP	205	0	–	None	None	0	None	None
9	LT	373	1	0	Bil caudate	None	0	None	None
10	RT	172	8	0	BilO, BilCer	Bil putamen, L posterior cing., sphenoidal sinus	1	Sphenoidal sinus	None

Abbreviations: ICs: independent components, R: right, L: left, Bil: bilateral, F: frontal, C: central, T: temporal, P: parietal, O: occipital, Cer: cerebellum, cing: cingulate, thal: thalamus.

Table 3

Results of the GLM analysis. The presumed location of the epileptogenic focus is repeated from Table 2 to facilitate interpretation. Columns 3 and 4 show GLM clusters of significant BOLD response when assuming a constant HRF amplitude. The locations of the maximum activation and deactivation are shown in columns 5 and 6. Columns 7 and 8 show GLM clusters revealed by an additional regressor modeling HRF amplitudes as proportional to IED amplitudes. Concordant locations are indicated in bold type.

Patient	Focus	GLM results		Regressor modeling IED amplitude fluctuations			
		Activation	Deactivation	Maximum activation	Maximum deactivation	Activation	Deactivation
1	RT	RT , BilP	LT, posterior cingulate, RCer	RT	Posterior cingulate	RT , LC	RO, LCer
2	RT	RT , BilF, BilO, brainstem, ventricles	BilP, BilO	L lateral ventricle	LP	BiIT , Bil insula, medial F	RO
3	LF	LF CP, BiIT, LO, BilFC, BilCer	BiIT, BilO, BilP, Bil insula, Bil thalamus	LF	RP	LCP , RT, LO, BilCer	BilO
4	LT	LT	BiIT , BilO	LT	LO	None	LF, RCer
5	LC	BiICP , BiIT, BiO, BilF	BilF, BiICer	LC	RC	RF, brainstem	BilF, LT
6	LP	RO, RT	RF, RP	RO	RP	RF, LO	BilF, LT
7	RT	RT , Bil insula, BiICP, RCer	BilF, BiIT, BilP, posterior cingulate	R insula	LP	None	LCer
8	LP	LTP , LF, Bil insula	LPO , RF, precuneus	LF	LPO	LT	None
9	LT	BilC, Bil insula, Bil putamen, BilCer	BilF	L insula	RF	None	LCer
10	RT	BilO, precuneus	BiIT, BilC, Bil thalamus, RCer	Precuneus	R thalamus	RT , BilO, LP, R putamen	BiIT, BilF

Abbreviations: R: right, L: left, Bil: bilateral, F: frontal, C: central, T: temporal, P: parietal, O: occipital, Cer: cerebellum.