Published in final edited form as:

Org Lett. 2013 August 2; 15(15): 3966–3969. doi:10.1021/ol401747u.

Nitroethylation of Vinyl Triflates and Bromides

Rosaura Padilla-Salinas, Ryan R. Walvoord, Sergei Tcyrulnikov, and Marisa C. Kozlowski

Penn Merck High Throughput Experimentation Laboratory, Department of Chemistry, University of Pennsylvania, Philadelphia, Pennsylvania 19104-6323, United States

Marisa C. Kozlowski: marisa@sas.upenn.edu

Abstract

A two-carbon homologation of vinyl triflates and bromides for the synthesis of homoallylic nitro products is described. This palladium-catalyzed double coupling of nitromethane exploits the anion stabilizing and leaving group properties of nitromethane, generating the homo allyl nitro products via a tandem cross-coupling/ -allylation sequence. The resultant process provides a mild and convenient entry of nitroethylated products, which are versatile precursors to α,β -unsaturated carbonyls, homoallylic amines, and nitrile oxides.

Recently, we reported a highly versatile and robust cross-coupling reaction of aryl halides with nitromethane.¹ This work provided simple access to a range of useful aryl nitromethane building blocks and exploited nitromethane as a formyl equivalent to directly afford aryl aldehydes or oximes (Scheme 1a). While exploring the use of vinyl bromides to further expand the scope, an unexpected, two-carbon homologated reaction product was encountered. Herein, we communicate the results from this investigation wherein palladium catalyzed coupling of vinyl halides and enol triflates with nitromethane affords nitroethylated products (Scheme 1b).

Nitromethane has been shown to react as a nucleophile for the Pd-catalyzed substitution reactions with allyl acetates, phenyl ethers, esters, and carbonate derivatives.² In addition, allyl nitro compounds have been observed to undergo ionization to form a Pd-allyl complex.^{3,4} However, accessing homoallylic nitroalkanes directly from vinyl triflates or halides has not been previously reported.^{2c,2f,5} Such a tandem process provides access to an array of nitroethylated compounds, which are versatile precursors to α,β -unsaturated carbonyls, homoallylic amines, and other useful functionality (Scheme 2).

Our studies began with the discovery that the palladium-catalyzed nitromethylation of β -bromostyrene gave rise to an unexpected product (Scheme 3). Instead of obtaining the allylnitro species, the nitroethylated product was isolated in moderate yield (54%).

The formation of the nitroethylated product is postulated to arise from a tandem cross-coupling/ -allylation mechanism (Scheme 4). Notably, the intermediate allylnitro compound

Correspondence to: Marisa C. Kozlowski, marisa@sas.upenn.edu.

Supporting Information Available. Experimental procedures and spectral data. This material is available free of charge via the Internet at <http://pubs.acs.org>.

was never observed in any of the studies conducted, indicating that it undergoes facile oxidative ionization under the reaction conditions. Utilizing nitromethane as a traceless coupling partner to homologate the vinyl moiety by one carbon and to incorporate a leaving group to allow formation of π -allyl intermediate **V** (Scheme 4) and a second homologation is a unique feature of this process.

Notably, the two palladium-catalyzed C-C bond forming events proceed via different mechanisms (oxidative addition/transmetallation/reductive elimination vs oxidative ionization/outer sphere nucleophilic attack), and there is no expectation that a single palladium-ligand species would effectively catalyze both portions.⁶ For example, with the exception of CataCXium POMetB, bidentate and bis-*tert*-butyl-ligands performed poorly in the nitromethylation of aryl halides.¹ Indeed, typical π -allylation conditions employing trialkylphosphine ligands do not catalyze this initial cross coupling step.^{2a}

Further study of the transformation from (Scheme 3) revealed that the nitroethylated product decomposed under the reaction conditions over time as observed by GC analysis (see Supporting Information). Additionally, subjecting the nitroethylated product to the reaction conditions for 6 hours led to only a 68% mass recovery. The low stability of the product coupled with the potential multiple pathways available from the Pd π -allyl intermediate (Scheme 5) and the need to simultaneously optimize two very different transformations (cross-coupling vs π -allylation, see above) prompted us to utilize parallel microscale (10 μ mol vinyl halide, 100 μ L reaction volume) experimentation to screen conditions. In addition to being able to rapidly assess the impact of different combinations of base, solvent, palladium source, and ligand on product formation, LCMS analysis of the reaction profiles provides insight into which sets of conditions give rise to undesired reaction pathways.

Due to the low stability of π -bromostyrene and its nitroethylated products, the screening focused on the more stable **3a** as a test substrate (Table 1). For a general process, vinyltrifluoromethanesulfonates are also preferred because they are more easily obtained (via enolization of the ketone) than the corresponding vinyl bromides. A variety of bis-*tert*-butyl and biscyclohexyl ligands were selected for screening known to promote the first step of the process (nitromethylation of the aryl halide).¹ Based on the efficiency of PPh₃ in the C-alkylation of allylic acetates with (phenylsulfonyl)nitromethane,^{2a} a combination PPh₃ with Xantphos, BINAP, CataCXium POMetB, JohnPhos, BrettPhos, or XPhos were also examined. With these ligands, 6 different bases (2.4 equiv) were screened.

Select results from this screen are listed in Table 1. In general, K₃PO₄ and Cs₂CO₃ were the most efficient bases (Table 1, entries 5–8). Bis-*tert*-butyl ligands (Table 1, entries 1 and 8) and mixed ligand system (Table 1, entry 11) performed poorly (<3.0 product:internal standard). Biscyclohexyl (DavePhos and SPhos) phosphine ligands and CataCXium POMetB were superior (Table 1, entries 5–7), even though they are not highly effective in typical π -allylations. The top three results were scaled up (0.360 mmol scale) using Cs₂CO₃ and afforded the desired product in 51–72% isolated yield (Table 1). Complete conversion was observed with DavePhos and SPhos while incomplete conversion was observed with CataCXium POMetB. The SPhos ligand proved most effective, offering enhanced reactivity and cleaner reaction profiles in comparison to DavePhos and CataCXium POMetB.

The use of chloride ion additives was explored since it is well documented that chloride ions are critical for coupling enol triflates.^{7,8} The allylPdOTfL_n intermediate formed after oxidative addition of the enol triflate is proposed to undergo a counterion exchange.⁹ Indeed, performing the reaction in the presence of excess chloride ions (10 equiv KCl) led to improved isolated yields and rates providing good results for both vinyl triflates and bromides (Scheme 6).

The optimized conditions did not prove to be broadly applicable to other alkene, diene, - and -styrene substrates affording the desired product in <50% (see Supporting Information). Incomplete conversion of starting material and byproducts were observed in most cases. Thus, lead ligands from the initial screen were assessed with a variety of alkene substrates to draw out reactivity trends (Table 2). Indeed, the yields could be improved to >50% yield for each substrate class.¹⁰

Dienes, disubstituted alkenes, and di- and tri-substituted -bromostyrene derivatives fall into class **A** where the XPhos ligand performed best (Table 2). -Bromo styrenes and -tetralone substrates comprise class **B**, for which the CataCXium POMeTB and DavePhos ligands are superior. SPhos is most effective for -tetralone substrates (class **C**). As expected, the addition of potassium chloride additive is most effective for enol triflates derived from -tetralones and did not improve the outcome for vinyl bromides.

In an effort to understand the discrete reaction steps, the allyl nitro species for the corresponding -tetralone was synthesized and subjected to the reaction conditions. The desired nitroethylated product was isolated in 14% yield (Scheme 7) suggesting that slow formation of the allyl nitro intermediate is necessary. Oxidative ionization of the allyl nitro ostensibly competes with other pathways such as dimerization, the products from which could be identified in LCMS of the HTE screen. Once the -allyl species does form, however, nucleophilic attack is highly regioselective for the primary position. The branched product was not observed here (Scheme 7) or with any of the other substrates studied.

To probe whether a similar reaction pathway is feasible with benzylic¹¹ vs allylic nitro intermediates, 2-bromo naphthalene was subjected to the reaction conditions (Scheme 8). Small amounts of the anticipated nitroethylated product were observed indicating the potential of such substrates.

In conclusion, a tandem cross-coupling/ -allylation of alkenyl bromides and triflates with nitromethane has been developed. Examining the reactivity of a variety of alkene substrates with selected conditions obtained from a test substrate proved to be an effective strategy for rapid optimization of different substrate classes. This method allows for facile nitroethylation of both enol triflates, readily available from the corresponding ketones, and bromides to afford versatile homoallylic nitro building blocks.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We are grateful to National Institutes of Health (GM-087605) and the National Science Foundation (CHE-105544) for financial support of this research. R.P.S gratefully acknowledges a Gates Millennium Scholar Fellowship. The NSF provided for funding of the High Throughput Laboratory (GOALI CHE-0848460). Partial instrumentation support was provided by the NIH for MS (1S10RR023444) and NMR (1S10RR022442) and by the NSF for an X-Ray Diffractometer (CHE 0840438).

References

1. Walvoord RR, Berritt S, Kozlowski MC. *Org Lett*. 2012; 14:4086–4089. [PubMed: 22839593]
2. a) Wade PA, Hinney HR, Amin NV, Vail PD, Morrow SD, Hardinger SA, Saft MS. *J Org Chem*. 1981; 46:765–770. b) Wade PA, Morrow SD, Hardinger SA. *J Org Chem*. 1982; 47:365–367. c) Deardoff DR, Savin KA, Justman CJ, Karanjawala ZE, Sheppeck JE II, Hager DC, Aydin N. *J Org Chem*. 1996; 61:3616–3622. [PubMed: 11667207] d) Rieck H, Helmchen G. *Angew Chem, Int Ed*. 1996; 34:2687–2689. e) Giambastiani G, Giovanni P. *J Org Chem*. 1998; 63:9608–9609. f) Trost

- BM, Surivet J. *Angew Chem, Int Ed.* 2000; 39:3122–3124.g) Uozumi Y, Suzuka T. *J Org Chem.* 2006; 71:8644–8646. [PubMed: 17064049]
3. a) Tamura R, Hegedus LS. *J Am Chem Soc.* 1982; 104:3727–3729.b) Ono N, Hamamoto I, Kaji A. *J Chem Soc, Chem Commun.* 1982:821–822.
4. Tamura R, Kamimura A, Ono N. *Synthesis.* 1991; 6:423–434.
5. Select examples Aleksandrowicz P, Piotrowska H. Sas Wojciech *Tetrahedron.* 1982; 38:1321–1327.Maki K, Kanai M, Shibaski M. *Tetrahedron.* 2007; 63:4250–4257.Grenning AJ, Tunge JA. *Org Lett.* 2010; 12:740–742. [PubMed: 20088536] Grenning AJ, Tunge JA. *Angew Chem, Int Ed.* 2011; 50:1688–1691.
6. For reviews of sequential catalyzed processes, see: Fogg DE, dos Santos EN. *Coord Chem Rev.* 2004; 248:2365–2379.Lee J, Na Y, Han H, Chang S. *Chem Soc Rev.* 2004; 33:302–312. [PubMed: 15272370] Wasilke J, Obrey SJ, Baker T, Bazan GC. *Chem Rev.* 2005; 105:1001–1020. [PubMed: 15755083]
7. Solvation of the metal ion to enhance the nucleophilicity of nitromethane with the use of ethereal solvents did not improve the yield.
8. Scott WJ, Crisp GT, Stille JK. *J Am Chem Soc.* 1984; 106:4630–4632.
9. a) Jutand A, Mosleh A. *Organometallics.* 1995; 14:1810–1817.b) Jutand A, Négri S. *Organometallic.* 2003; 22:4229–4237.c) Jutand A. *Appl Organometal Chem.* 2004; 18:574–582.
10. 3 ÅMS suppressed aldehyde formation from water. See ref 1.
11. a) Kawatsura M, Hartwig JF. *J Am Chem Soc.* 2000; 122:9546–9547.b) Nettekoven U, Hartwig JF. *J Am Chem Soc.* 2002; 124:1166–1167. [PubMed: 11841274] c) Johns AM, Utsunomiya M, Incarvito CD, Hartwig JF. *J Am Chem Soc.* 2006; 128:1828–1839. [PubMed: 16464081] d) Kuwano R, Kondo Y, Matsuyama Y. *J Am Chem Soc.* 2003; 125:12104–12105. [PubMed: 14518991] e) Trost BM. *J Am Chem Soc.* 2010; 132:15534–15536. [PubMed: 20961054] f) Trost BM, Czabaniuk LC. *J Am Chem Soc.* 2012; 134:5778–5781. [PubMed: 22420708]

a) Prior Work: One Carbon Homologation **b) This Work: Two Carbon Homologation**

Scheme 1.
Nitromethane Homologation.

Scheme 2.
Utilization of Nitroethylated Products

Scheme 3.

Scheme 4.
Proposed Mechanism

Scheme 5.
Potential Reaction Pathways from the -Allyl Nitro Intermediate.

Scheme 6.
Use of KCl in the Nitroethylation

Scheme 7.

One Carbon Homologation from the Putative Allylnitro Intermediate.

Scheme 8.
Nitroethylation of an Aryl Halide.

Product:internal standard ratios from initial screening.

Org Lett. Author manuscript; available in PMC 2014 August 02.

Table 2

Palladium-Catalyzed Nitroethylation.

		$ \begin{array}{c} \text{R}^1 \\ \\ \text{R}^2-\text{C}=\text{C}-\text{X} \\ \\ \text{R}^3 \end{array} \xrightarrow[\substack{\text{Cs}_2\text{CO}_3 \text{ (2.4 equiv)} \\ 0.1 \text{ M MeNO}_2 \\ 3 \text{ \AA MS, } 65^\circ\text{C, 4-8 h}}]{\substack{\text{Pd}_2(\text{dba})_3 \text{ (5 mol \%)} \\ \text{ligand (12 mol \%)}}} \begin{array}{c} \text{R}^1 \\ \\ \text{R}^2-\text{C}=\text{C}-\text{CH}_2\text{CH}_2\text{NO}_2 \\ \\ \text{R}^3 \end{array} $			
class	substrate	Yield ^a			
		SPhos	XPhos	CataCXium PONetB	DavePhos
A		26%	47%		
	5
	42% ^b	76%	46%	59%
	6
	69%	62%	27%
B	1
	64%	47%	
	7
	47%	43%	60%	33%
	8				

$ \begin{array}{c} \text{R}^1 \\ \\ \text{R}^2 - \text{C} = \text{C} - \text{X} \\ \\ \text{R}^3 \end{array} \xrightarrow[\substack{\text{Cs}_2\text{CO}_3 \text{ (2.4 equiv)} \\ 0.1 \text{ M MeNO}_2 \\ 3 \text{ \AA MS, 65 }^\circ\text{C, 4-8 h}}]{\substack{\text{Pd}_2(\text{dba})_3 \text{ (5 mol \%)} \\ \text{ligand (12 mol \%)}}} \begin{array}{c} \text{R}^1 \\ \\ \text{R}^2 - \text{C} = \text{C} - \text{CH}_2\text{CH}_2\text{NO}_2 \\ \\ \text{R}^3 \end{array} $						
$ \begin{array}{c} \text{X} = \text{OTf, Br} \end{array} $						
class	substrate	Yield ^a				
		SPhos	XPhos	CataCXium	POMeB	DavePhos
C		16%	66%	74%		68%
	83%
	80% ^c
	93% ^c
	67%

<div>$\begin{array}{c} \text{R}^1 \\ \\ \text{R}^2 - \text{C} = \text{C} - \text{X} \\ \\ \text{R}^3 \end{array} \xrightarrow[\begin{array}{c} \text{Pd}_2(\text{dba})_3 \text{ (5 mol \%)} \\ \text{ligand (12 mol \%)} \\ \text{Cs}_2\text{CO}_3 \text{ (2.4 equiv)} \\ 0.1 \text{ M MeNO}_2 \\ 3 \text{ \AA MS, } 65^\circ\text{C, 4-8 h} \end{array}]{}$</div>		<div>$\begin{array}{c} \text{R}^1 \\ \\ \text{R}^2 - \text{C} = \text{C} - \text{CH}_2\text{CH}_2\text{NO}_2 \\ \\ \text{R}^3 \end{array}$</div>			
$\text{X} = \text{OTf, Br}$					
class	substrate	Yield ^a			
		SPhos	XPhos	CataCXium POMetB	DavePhos
	55% ^b	49%		
	11				

^a Isolated yields.

^b Conversion determined by ¹H NMR.

^c 10 equiv KCl added.