Published in final edited form as:

Bioconjug Chem. 2013 July 17; 24(7): 1186–1190. doi:10.1021/bc400062d.

Tb³⁺-tRNA for LRET studies of Protein Synthesis

Dulce Alonso^{1,†}, Wei Liu¹, Gabriel Rosenblum^{1,2}, Tomoyasu Mani³, Yale E. Goldman², and Barry S. Cooperman^{1,*}

¹Department of Chemistry, University of Pennsylvania, Philadelphia, PA, 19104-6323

²Pennsylvania Muscle Institute, Perlman School of Medicine, University of Pennsylvania, Philadelphia, PA 19104-6083

³Department of Biochemistry and Biophysics, Perlman School of Medicine, University of Pennsylvania, Philadelphia, PA 19104-6059

Abstract

When suitably labeled bulk tRNAs are transfected into cells they give rise to FRET (fluorescence resonance energy transfer) signals via binding to ribosomes that provide a measure of total protein synthesis. Application of this approach to monitoring rates of specific protein synthesis requires achieving very high signal-to-noise ratio. Such high ratios may be attainable using LRET (luminescence resonance energy transfer) in place of FRET. Lanthanide complexes containing an antenna chromophore are excellent LRET donors. Here we describe the synthesis of a Phe-tRNA^{Phe} labeled with a Tb³⁺ complex, denoted Tb³⁺-Phe-tRNA^{Phe} that, notwithstanding the bulkiness of the Tb³⁺ complex, is active in protein synthesis.

INTRODUCTION

We and our collaborators recently have demonstrated that when fluorescently-labeled tRNAs (fl-tRNAs), both FRET (fluorescence resonance energy transfer) donors and acceptors, are transfected into cells, they give rise to FRET signals.¹ Such signals are expected to arise when donor-labeled and acceptor-labeled tRNAs come into close proximity (< 6 nm), as they do on the ribosome during the course of polypeptide elongation.^{2,3} In accord with this expectation, the intensity of the FRET signal was found to correlate with the number of ribosomes engaged in protein synthesis, providing a real-time, live-cell assay for measuring rates of protein synthesis. In this prior work, fl-tRNAs were synthesized by condensing fluorescent weak-base amines, including hydrazides, with the tetrahydrouridine (THU) residues in bulk tRNA that are formed by specific reduction of dihydrouridine (DHU) residues within the D-loops of tRNA, as described.^{4–6} The FRET signal generated using bulk fl-tRNAs provided a measure of the rate of total protein synthesis within transfected cells.¹

Current efforts are focused on applying this approach to measure the intracellular rates of synthesis of specific proteins, which will be facilitated by higher signal-to-background ratios. Background interference from unbound donor and acceptor fl-tRNAs, autofluorescent compounds and light scattering, while minimized by excitation wavelength selection and

*To whom correspondence should be addressed.

†present address: Department of Molecular, Cellular & Developmental Biology, Yale University, New Haven, CT 06520

SUPPORTING INFORMATION AVAILABLE

A figure comparing luminescence observed for VI and unmodified tRNA^{Phe} on incubation with Tb³⁺. This material is available free of charge via the Internet at <http://pubs.acs.org>.

emission filters, limits the sensitivity and precision of the FRET approach. Substituting gated-detection of LRET (luminescence resonance energy transfer) for FRET offers a plausible solution to the background problem. LRET uses long-lifetime lanthanide (Ln^{3+}) donors (e. g., Tb^{3+} or Eu^{3+}) to temporally separate the delayed ($>100\ \mu\text{s}$) emission of acceptors from the prompt (ns) emission of fluorescent species that are unrelated to resonance energy transfer, with reported increases in signal-to-background ratios of 500-fold.⁷

Ln^{3+} complexes suitable for LRET studies are composed of an “antenna” chromophore, necessary to overcome the inherently low absorbance of Ln^{3+} , attached to a multidentate chelating site for the Ln^{3+} .^{7,8} Here we report the synthesis of a hydrazide derivative of such a Ln^{3+} -complex, its attachment to tRNA^{Phe} , and the activity of this bioconjugate, charged with Phe, both in forming a ternary complex with EF-Tu-GTP that is required for Phe- tRNA^{Phe} uptake by the ribosome, and in supporting synthesis of the full-length protein emerald green fluorescent protein (EmGFP).

EXPERIMENTAL PROCEDURES

Chemicals and Instrumentation

The following reagents were purchased from Aldrich: 1,3-phenylenediamine, diethyl-1,3-acetonedicarboxylate, zinc chloride, *tert*-butyl carbazate, *N*-(3-dimethylaminopropyl)-*N'*-ethylcarbodiimide hydrochloride (EDAC), diethylenetriaminepentaacetic acid (DTPA), terbium (III) chloride, europium (III) chloride. *E. Coli* tRNA^{Phe} was obtained from Chemical Block (Moscow). ^3H -Phenylalanine (Phe) was obtained from Perkin Elmer.

Assaying the translational activity of the Tb^{3+} -tRNA using cell-free protein synthesis (CFPS)

The CFPS kit (5prime) was utilized to synthesize EmGFP according to the manufacturer's manual, with some modifications.⁹ Briefly, the cell lysate was supplemented with protease inhibitors and an inhibitor of tRNA^{Phe} -aminoacyl-synthetase (PheRS) and then dialyzed in order to deplete all amino acids while preventing Phe regeneration and tRNA^{Phe} aminoacylation. Following dialysis, the other CFPS components were added according to the manufacturer's manual, excluding the amino acid Phe. The total ribosome concentration was estimated as $0.3\ \mu\text{M}$ based on A_{260} . The ability of this CFPS- Phe mixture to produce EmGFP was dependent on the addition of exogenous Phe, supplied in the form of either Phe- tRNA^{Phe} or Tb^{3+} -Phe- tRNA^{Phe} ($2\ \mu\text{M}$). The reaction was initiated by addition of 130 ng plasmid DNA per 10 μl reaction volume and fluorescence was recorded using a plate reader (EnVision - Perkin-Elmer) at $30\ ^\circ\text{C}$ using excitation and emission wavelengths of 486 and 535 nm, respectively.

Spectra

^1H NMR spectra were recorded in a Bruker DMX 300 spectrometer in deuterated solvents. ^1H chemical shifts were referenced to internal solvent resonances and reported relative to SiMe_4 . Low-resolution electrospray (ESI) mass spectra were obtained at the Waters SQD equipped with an Acquity UPLC. UV-visible (UV-vis) absorption spectra were recorded using a Hewlett Packard 8452A Diode Array Spectrophotometer. Fluorescence emission spectra were recorded using a Fluorolog-3 spectrofluorometer (Horiba Jobin Yvon). Time-resolved luminescence measurements were performed on a FS900 spectrofluorometer (Edinburgh Instruments, UK), equipped with R2658P photomultiplier tube (Hamamatsu) and a xenon flash lamp as the excitation source. Kinetic experiments were performed on a KinTek SF-300X stopped-flow spectrofluorometer.

Synthesis (Fig. 1)

Compounds **I** and **III** were prepared as described.^{10,11}

Compound II. To a suspension of **I** (70 mg, 0.32 mmol) in DCM (2 mL) and DMF (0.8 mL) was added *tert*-butyl carbazate (85 mg, 0.64 mmol) and *N*-(3-dimethylaminopropyl)-*N*-ethylcarbodiimide hydrochloride (EDAC) (80 mg, 0.41 mmol). The reaction was maintained for 20 h at room temperature. Solvent was removed by evaporation in vacuo. The residue was dissolved in ethyl acetate (15 mL) and extracted with a solution of saturated aqueous NaHCO₃ (3 × 10 mL). The organic layer was dried over anhydrous Na₂SO₄ and evaporated. The product was purified by column chromatography on silica gel using a chloroform/ethanol mixture (8:2) as eluent. Yield 62%. ESIMS (*m/z*) 333 [M+H]⁺. ¹H NMR (300 MHz, DMSO-*d*₆): 1.39 (s, 9H, Boc), 3.52 (s, 2H, -CH₂CO-), 5.74 (broad, 2H, 7 amino), 6.07 (s, 1H, 3H), 6.37 (d, 1H, 8H, *J* = 2.1), 6.43 (dd, 1H, 6H, *J*₁ = 8.7 Hz, *J*₂ = 2.1 Hz), 7.41 (d, 1H, 5H, *J* = 8.7), 9.85 (1H, broad, NH), 11.20 (1H, broad, amide). ¹³C NMR (300 MHz, DMSO-*d*₆): 28.99, 44.06, 80.77, 97.40, 110.68, 111.01, 116.84, 125.92, 140.41, 146.01, 151.35, 156.50, 162.62, 170.35.

Compound III. Pyridine (1.2 mL) and acetonitrile (1.2 mL) warmed to 50 °C were mixed with a solution of DTPA (490 mg, 1.37 mmol) in acetic anhydride (0.8 mL, 8.60 mmol). After 24 h, solvent was removed by evaporation in vacuo and the residue was washed with acetic anhydride and diethyl ether. The solid was dried under vacuum. Yield 92%. ESIMS (*m/z*) 356 [M-H]⁻. ¹H NMR (300 MHz, DMSO-*d*₆): 3.71 (s, 8H, terminal -NCH₂CO₂), 3.30 (s, 2H, central -NCH₂CO₂), 2.74 (t, 4H, *J* = 5.7 Hz, -NCH₂CH₂N-), 2.59 (t, 4H, *J* = 5.7 Hz, -NCH₂CH₂N-).

Compound IV. To a suspension of DTPA dianhydride **III** (54 mg, 0.15 mmol) in DMF (1 mL) was added triethylamine (0.08 mL, 0.6 mmol) under nitrogen. After 10 min a solution of **II** (15 mg, 0.04 mmol) in DMF (0.5 mL) was added and the reaction was maintained at room temperature for 3 h. A few drops of water were added to quench the reaction. The product was purified by preparative thin layer chromatography, using acetonitrile/water (7:3) as eluent. Yield 30%. ESIMS (*m/z*) 706 [M-H]⁻. ¹H NMR (300 MHz, D₂O): 1.39 (s, 9H, Boc), 2.60-3.90 (m, 20H), 6.47 (s, 1H, 3H), 7.16 (m, 1H, 8H), 7.67-7.45 (m, 2H, 5H-6H).

Compound V. A mixture of DCM:TFA (1:1) (100 μL) was added to **IV** (2 mg, 0.003 mmol) at 0 °C and the reaction was maintained at room temperature for 2 h. **V** was used without further purification in the preparation of **VI**. The solvent was evaporated in vacuo. Quantitative yield. ESIMS (*m/z*) 608 [M+H]⁺. 606 [M-H]⁻; $\epsilon_{\text{max}} = 14800 \text{ M}^{-1} \text{ cm}^{-1}$ at 340 nm.¹²

Compound VI. Derivatization of Phe-tRNA^{Phe} with the hydrazide **V** was based on a previously described protocol^{5,6} involving nucleophilic substitution at THU residues. Selective reduction of the two DHU residues at positions 16 and 20 in *E. coli* tRNA^{Phe} to THU residues was carried out by incubating tRNA^{Phe} (2.5 mg/mL), NaBH₄ (10 mg/mL, added from 100 mg/mL in 10 mM KOH) in 40 mM Tris-HCl (pH=7.5), at 0 °C for 60 min, in a total volume of 390 μL. The reaction was quenched by lowering the pH to 4–5 by gradually adding 6 M acetic acid. Unreacted NaBH₄ was removed via three ethanol precipitations of tRNA^{Phe} (reduced). Aminoacylation of tRNA^{Phe}(reduced) (0.5 mg/mL) was performed in a buffer containing 75 μM ³H-phenylalanine (370 cpm/pmol), 100 mM Tris-HCl (pH 8.0), 4 mM ATP (pH 7.8), 20 mM MgCl₂, 1 mM EDTA, 7 mM 2-mercaptoethanol and 100 μM S100 crude synthetase in a total volume of 1.7 mL. The reaction was incubated at 37 °C during 30 min. Following phenol-chloroform extraction and ethanol precipitation, Phe-tRNA^{Phe}(reduced) was loaded onto a FPLC monoQ column.

Chromatography with a gradient of 0.5–0.9 M NaCl in 50 mM NaAc (pH 5.0), yielded a preparation of Phe-tRNA^{Phe}(reduced) in which the tRNA was 30% charged with Phe. This preparation (0.7 mg) was resuspended in 11 μ L of 0.2 M NaCOOH (pH 3.7) and 10 μ L of DMSO. 2 μ L of a DMSO solution of **V** (290 mM) was added and incubation was carried out for 150 min at 37 °C. During this period, additional 1 μ L portions of **V** in DMSO solution were added at 45 and 90 min. Vacuum drying and removal of excess **V** by three rounds of ethanol precipitation gave an overall stoichiometry of labeling of 0.4 **V**/tRNA. Unlabeled Phe-tRNA^{Phe} (reduced) was removed from **V**-labeled Phe-tRNA^{Phe} by RP-HPLC using a C18/15 μ m column (Waters) with a gradient of 6–21% ethanol in buffer containing 20 mM NH₄Ac (pH 5.0), 10 mM MgAc₂ and 400 mM NaCl, with absorption monitored at 260 and 340 nm. In this gradient, the labeled tRNA elutes after the unlabeled tRNA. This procedure yielded preparations, designated as **VI**, containing 1.3 **V**/Phe-tRNA^{Phe}; i. e., mixtures of Phe-tRNA^{Phe}s containing 1 or 2 **V**/tRNA^{Phe}.

Synthesis of Tb³⁺-Phe-tRNA^{Phe}. **VI** was dissolved in water and 1.5 eq of a 300 μ M TbCl₃ solution, based on the stoichiometry of **V** incorporated into **VI**, was added, with vigorous stirring for 30 min at 0 °C. Excess TbCl₃ was removed by 4 rounds of ethanol precipitation. An analogous procedure was used to prepare Eu³⁺-Phe-tRNA^{Phe}. Suitable control experiments demonstrated that incubation with TbCl₃ of tRNA^{Phe} not labeled with **V** afforded no luminescence at 550 nm on excitation at 345 nm (see Fig. S1).

Cell-free expression of EmGFP using the CFPS^{-Phe} kit

The assay was performed essentially as previously described.⁹ *E. coli* Phe-tRNA^{Phe}s (2 μ M, measured by the amount of charged tRNA) either unlabeled or labeled with **V** and Tb³⁺, were added to a CFPS^{-Phe} kit and EmGFP synthesis was monitored by fluorescence intensity at 535 nm.

Ternary Complex (TC) formation

TCs were formed by combining elongation factor Tu (EF-Tu) and Phe-tRNA^{Phe} in buffer A (50 mM Tris-HCl [pH7.5], 70 mM NH₄Cl, 30 mM KCl, 7 mM MgCl₂ and 1 mM DTT, 1 mM GTP, 1.5 mM phosphoenolpyruvate and 0.5 mg/L of pyruvate kinase) as previously described¹³ except that the ratio of EF-Tu/tRNA employed was 4:1 unless otherwise specified. Time-dependent TC formation was performed at 25 °C by mixing 0.1 μ M of Tb³⁺-Phe-tRNA^{Phe} with 0.4 μ M of EF-Tu^{Cy5} with excitation of 345 nm. Static fluorescence was recorded with excitation of 345 nm.

RESULTS

Synthesis of the probe, **V**

Compound **I** was obtained by condensing 1,3-phenylenediamine and diethyl-1,3 acetonedicarboxylate, followed by saponification of the ethyl ester according to previously reported procedure.¹⁰ We introduced the Boc-protected hydrazide function by treatment of **I** with *tert*-butyl carbazate and *N*-(3-dimethylaminopropyl)-*N*'-ethylcarbodiimide hydrochloride (EDAC), yielding **II**. An excess of DTPA dianhydride **III** was reacted with **II** in Et₃N and DMF at room temperature for 3 hours, yielding **IV**, which was purified by thin layer chromatography. Deprotection of **IV** with TFA:DCM (1:1) afforded the DTPA-based chelate hydrazide (**V**).

Labeling of Phe-tRNA^{Phe}(reduced) with **V**

E. coli tRNA^{Phe} has DHU residues at positions 16 and 20. Synthesis of charged, **VI** labeled tRNA^{Phe} for use in functional studies was carried out using the three-step procedure

previously employed to label DHU positions, involving 1) NaBH₄ reduction of the DHU residues to THU to form tRNA^{Phe}(reduced); 2) charging of tRNA^{Phe}(reduced) with Phe by Phe-RS and 3) labeling of Phe-tRNA^{Phe}(reduced) with a large molar excess of hydrazide.^{5,6} Such labeling proceeds via a substitution at the THU residues to yield *N*-substituted tetrahydrocytidine (THC) residues within the tRNA. Previous work⁶ showed that Cy3 hydrazide labeled Phe-tRNA^{Phe}(reduced) at both positions 16 and 20 to near equal extents (relative labeling 1.0:1.2, respectively) and we assume a similar distribution is obtained for labeling of tRNA^{Phe}(reduced) by **VI**. In the present case, the overall labeling stoichiometry was 0.4 **V**/tRNA^{Phe}. A more enriched preparation containing 1.3 **V**/tRNA^{Phe}, denoted **VI**, was obtained by separation of derivatized tRNA from underivatized tRNA by RP-HPLC. Within **VI**, 24–30% of the tRNA^{Phe} was charged with Phe.

Formation and characterization of the Tb³⁺-VI Complex

Treatment of **VI** with aqueous lanthanide trichlorides afforded Tb³⁺ and Eu³⁺ complexes. It was also possible to label Phe-tRNA^{Phe} with a preformed Ln³⁺-**V** complex. However, resolution of the resulting reaction mixture by RP-HPLC was less successful than the procedure described in Experimental above. We denote the charged form of Tb³⁺-**VI** as Tb³⁺-Phe-tRNA^{Phe}. On excitation of the 7-aminoquinolone moiety at 345 nm, Tb³⁺-**VI** shows the multiple luminescence bands characteristic of Tb³⁺ complexes in the range 485 – 620 nm, corresponding to the ⁵D₄ to ⁷F_n (n = 3 – 6) transitions indicated (Fig. 2A). The following luminescence lifetimes were measured: Tb³⁺-**VI**: 1.06 ms, Eu³⁺-**VI**: 0.6 ms, Tb³⁺-Phe-tRNA: 1.08 ms. All lifetime measurements were well fit by single exponential decays.

Functional assays

Ternary complex (TC) formation—Binding of aminoacyl-tRNA (aa-tRNA) to the ribosome occurs via the ternary complex aa-tRNA·EF-Tu·GTP, where EF-Tu is a translation elongation factor. We have previously shown that an EF-Tu mutant labeled with Cy5 at position 348, denoted EF-Tu^{Cy5}, retains functional activity in protein synthesis.¹⁴ Within the TC, this position is within 34 Å and 24 Å of positions 16 and 20, respectively, of aminoacyl-tRNA [calculated from pdb1OB2], making it suitable for LRET studies with Tb³⁺-**VI**. Formation of the TC between Tb³⁺-Phe-tRNA^{Phe} and EF-Tu^{Cy5}·GTP is accompanied by a decrease in Tb³⁺-Phe-tRNA^{Phe} luminescence as EF-Tu^{Cy5} is increased (Fig. 2A) and an increase in EF-Tu^{Cy5} fluorescence as Tb³⁺-Phe-tRNA^{Phe} is increased (Fig. 2B). The K_d value estimated from the results in Figs. 2A, B is approximately 0.3 μM. Additional evidence for LRET between Tb³⁺ and Cy5 within the TC comes from the difference in luminescence lifetimes of the TC complexes made from Tb³⁺-Phe-tRNA^{Phe} and either EF-Tu^{Cy5}·GTP (0.45 ms, t_L) or unlabeled EF-Tu·GTP (1.08 ms, t_L⁰). An estimated LRET efficiency, φ, equal to 0.58, could be calculated from the measured luminescence lifetime values via eq 1.

$$f = (t_L^{-1} - t_L^{0^{-1}}) / t_L^{-1} \quad (1)$$

The transient changes in Tb³⁺-Phe-tRNA^{Phe} luminescence and EF-Tu^{Cy5} fluorescence (Fig. 2C) on rapid mixing of Tb³⁺-Phe-tRNA^{Phe} and EF-Tu^{Cy5}·GTP allows calculation of a 2nd order rate constant for TC formation of 1.8 × 10⁶ M⁻¹s⁻¹.

EmGFP Synthesis—We recently described a real-time coupled transcription–translation assay that continuously monitors formation of the fluorescent protein Emerald GFP (EmGFP) and can be manipulated so that EmGFP formation is totally dependent on exogenously added Phe-tRNA^{Phe}, through use of a so-called CFPS^{-Phe} kit.⁹ Here we use this assay to demonstrate that Tb³⁺-Phe-tRNA^{Phe} displays activity in supporting EmGFP synthesis that is roughly comparable to that of native Phe-tRNA^{Phe} (Fig. 3). The assay was

carried out under conditions in which the extent of EmGFP synthesis is roughly proportional to added Phe-tRNA^{Phe} concentration. It is possible that the lower amount of EmGFP synthesized with Tb³⁺-Phe-tRNA^{Phe} reflects a functional heterogeneity of this sample. For example, the labeling stoichiometry is 1.3 **VI**/tRNA, and it is possible that the 2:1 labeled tRNA is less active than either of the 1:1 complexes.

DISCUSSION

This work demonstrates that Phe-tRNA^{Phe} derivatized with the rather bulky luminescent Tb³⁺ complex **VI** at THU positions formed by selective NaBH₄ reduction of DHUs at residues 16 and 20 within the D-loop retains high functionality in protein synthesis (Fig. 3). Such retention of functionality is not completely unexpected. Positions 16 and 20 are both quite far from the site of EF-Tu interaction with charged tRNAs, accounting for the readiness with which Tb³⁺-Phe-tRNA^{Phe} reacts with EF-Tu to form a ternary complex (Fig. 2C), and proximal to the variable loop of tRNA, within which insertions of up to 19 nucleotides are well-tolerated by the ribosome during protein translation.

The success of the current work opens up the prospect that Tb³⁺-Phe-tRNA^{Phe}, or molecules like it, can be used to monitor LRET signals in live cells that are generated by juxtaposition on the ribosome of a tRNA labeled with **VI** and a suitable fl-tRNA, labeled, for example, with Cy5, that serves as an LRET acceptor. The antenna in **VI**, an aminooxoquinoline, was chosen based on prior work showing it to have excellent optical and stability properties for LRET experiments.¹⁵ More recently, other antennas have been described that have superior optical properties,^{16,17} but are somewhat larger than 7-aminoquinolones. Further work will determine whether tRNAs containing Ln³⁺ complexes of such antennas will demonstrate functionality comparable to that found with Tb³⁺-Phe-tRNA^{Phe}.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work was funded by NIH grant GM090404 to BSC and by a Human Frontier Science Program fellowship to GR. We thank Profs. Larry Miller (UIC) and Sergei Vinogradov (Penn) for helpful discussions.

REFERENCES

1. Barhoom S, Kaur J, Cooperman BS, Smorodinsky NI, Smilansky Z, Ehrlich M, Elroy-Stein O. Quantitative single cell monitoring of protein synthesis at subcellular resolution using fluorescently labeled tRNA. *Nucl. Acids Res.* 2011; 39:e129. [PubMed: 21795382]
2. Blanchard SC, Kim HD, Gonzalez RL Jr, Puglisi JD, Chu S. tRNA dynamics on the ribosome during translation. *Proc. Natl. Acad. Sci. U. S. A.* 2004; 101:12893–12898. [PubMed: 15317937]
3. Chen C, Stevens B, Kaur J, Cabral D, Liu H, Wang Y, Zhang H, Rosenblum G, Smilansky Z, Goldman Y, Cooperman BS. Single Molecule Fluorescence Measurements of Ribosomal Translocation Dynamics. *Mol. Cell.* 2011; 42:367–377. [PubMed: 21549313]
4. Wintermeyer W, Schleich HG, Zachau HG. Incorporation of amines or hydrazines into tRNA replacing wybutine or dihydrouracil. *Methods Enzymol.* 1979; 59:110–121. [PubMed: 440077]
5. Pan D, Qin H, Cooperman BS. Synthesis and functional activity of tRNAs labeled with fluorescent hydrazides in the D-loop. *RNA.* 2009; 2:346–354. [PubMed: 19118261]
6. Kaur J, Raj M, Cooperman BS. Fluorescent labeling of tRNA dihydrouridine residues: Mechanism and distribution. *RNA.* 2011; 17:1393–1400. [PubMed: 21628433]

7. Rajapakse HA, Gahlaut N, Mohandessia S, Yu D, Turner JR, Miller LW. Time-resolved luminescence resonance energy transfer imaging of protein–protein interactions in living cells. *Proc. Natl. Acad. Sci. U.S.A.* 2010; 107:13582–13587. [PubMed: 20643966]
8. Rajapakse HE, Reddy DR, Mohandessi S, Butlin NG, Miller LW. Luminescent terbium protein labels for time-resolved microscopy and screening. *Angew. Chem. Int. Ed. Engl.* 2009; 48:4990–4992. [PubMed: 19492378]
9. Rosenblum G, Chen C, Kaur J, Cui X, Goldman YE, Cooperman BS. Real-time assay for testing components of protein synthesis. *Nucl. Acids Res.* 2012
10. Reddy DR, Pedró Rosa LE, Miller LW. Luminescent trimethoprim-polyaminocarboxylate lanthanide complex conjugates for selective protein labeling and time-resolved bioassays. *Bioconjugate Chem.* 2011; 22:1402–1409.
11. Ardestani MS, Arabzadeh AJ, Heidari Z, Hosseinzadeh A, Ebrahimi H, Hashemi E, Mosayebnia M, Shafiee-Alavidjeh M, Alavi A, Babai MH, Rahmim A, Ebrahimi SES, Amanlou M. Novel and facile methods for the synthesis of DTPA-mono-amide: a new completely revised strategy in the radiopharmaceutical chemistry. *J. Radioanal Nucl. Chem.* 2010; 283:447–455.
12. Krasnoperov LN, Marras SAE, Kozlov M, Wirpsza L, Mustaev A. Luminescent probes for ultrasensitive detection of nucleic acids. *Bioconjugate Chem.* 2010; 21:319–327.
13. Pan D, Kirillov SV, Cooperman BS. Kinetically competent intermediates in the translocation step of protein synthesis. *Mol. Cell.* 2007; 25:519–529. [PubMed: 17317625]
14. Cooperman, BS.; Goldman, YE.; Chen, C.; Farrell, I.; Kaur, J.; Liu, H.; Liu, W.; Rosenblum, G.; Smilansky, Z.; Stevens, V.; Zhang, H. Mechanism and Dynamics of the Elongation Cycle. In: Rodnina, MV., et al., editors. *Ribosomes: Structure, Function and Dynamics*. Springer Verlag; 2011. p. 339-348.
15. Li M, Selvin PR. Amine-reactive forms of a luminescent diethylenetriaminepentaacetic acid chelate of terbium and europium: attachment to DNA and energy transfer measurements. *Bioconjugate Chem.* 1997; 8:127–132.
16. Moore EG, Samuel AP, Raymond KN. From antenna to assay: lessons learned in lanthanide luminescence. *Acc. Chem. Res.* 2009; 42:542–552. [PubMed: 19323456]
17. Xu J, Comeillie TM, Moore EG, Law GL, Butlin NG, Raymond KN. Octadentate cages of Tb(III) 2-hydroxyisophthalamides: a new standard for luminescent lanthanide labels. *J. Am. Chem. Soc.* 2011; 133:19900–19910. [PubMed: 22010878]

Fig. 1. Synthetic Scheme for VI

Fig. 2. TC formation between EF-Tu^{Cy5}-GTP and Tb³⁺-Phe-tRNA^{Phe}

A. Luminescence spectrum of Tb³⁺-Phe-tRNA^{Phe} (0.2 μ M) with varying amounts of EF-Tu^{Cy5} on excitation at 345 nm. **B.** Fluorescence spectrum of EF-Tu^{Cy5} (0.2 μ M) with varying amounts of Tb³⁺-Phe-tRNA^{Phe} on excitation at 345 nm. **C.** Decrease in Tb³⁺-Phe-tRNA^{Phe} luminescence (550 $\pm$ 10 nm) and increase in EF-Tu^{Cy5} fluorescence (680 $\pm$ 10 nm) on excitation at 345 nm following rapid mixing of Tb³⁺-Phe-tRNA^{Phe} (0.1 μ M, measured as Phe), containing 0.24 Phe/tRNA, and EF-Tu^{Cy5}-GTP (0.4 μ M) (final concentrations). In parts A. and B. excitation frequency doubling in the Fluorolog-3 spectrofluorometer resulted in a very large peak at 690 nm. This artifact was not a problem for the KinTek SF-300X stopped-flow spectrofluorometer employed in part C.

Fig. 3. Phe-tRNA^{Phe}-dependent Cell-Free Protein Synthesis of EmGFP

Experiments were performed using the CFPS^{Phe} mixture, supplemented with 2 μ M of the indicated tRNAs, initiated with 130 ng plasmid DNA. Recordings of EmGFP fluorescence were performed every 45 seconds using the plate reader, set to 30 °C using excitation and emission wavelengths of 486 and 535 nm, respectively.