CORRESPONDENCE

Bleeding Complications and Liver Injuries During Phenprocoumon Treatment: a Multicentre Prospective Observational Study in Internal Medicine Departments

by Dr. med. Sven Schmiedl, Dipl.-Stat. Marietta Rottenkolber, Jacek Szymanski, Prof. Dr. med. Werner Siegmund, PD Dr. rer. nat. Marion Hippus, PD Dr. med. Katrin Farker, Prof. Dr. med. Bernd Drewelow, Prof. Dr. med. Joerg Hasford, Prof. Dr. med. Petra Thürmann in volume 14/2013

Excessive Treatment With Anticoagulants

Over recent years I have observed an extreme increase in anticoagulation therapy among my patients. Consistent with guidelines, almost all patients with a diagnosis of atrial fibrillation are started on therapy with phenprocoumon, warfarin, and also the new oral anticoagulants (NOACs) by cardiologists in private practice as well as in hospitals. A patient's life expectancy, their mental status, and practicability are not taken into consideration when determining the indication for treatment.

In addition to the enormous logistical complexities, too many patients in my practice have experienced serious complications—such as subdural, gastrointestinal, and retroperitoneal hemorrhage—as a result of this treatment, which is given as a merely prophylactic measure. In the same context, the less serious complications—such as hemarthrosis, epistaxis, and hematuria—should also be mentioned.

I thank the authors for their valuable and overdue multicenter observational study (1). I hope that the study will rapidly prompt a rethink and modification of current guidelines. In my long years of experience in clinical practice, the potential risk of an embolic cerebrovascular accident is not weighed up by the serious, occasionally fatal complications of prophylactic anticoagulation therapy. DOI: 10.3238/arztebl.2013.0541a

REFERENCES

1. Schmiedl S, Rottenkolber M, Szymanski J, Siegmund W, Hippus M, Farker K, Drewelow B, Hasford J, Thürmann P: Bleeding complications and liver injuries during phenprocoumon treatment—a multicentre prospective observational study in internal medicine departments. Dtsch Arztebl Int 2013; 110(14): 244–52.

Dr. med. Hubertus Sroka

Primary Care Physician in private practice
dr.med.h.sroka@googlemail.com

Conflict of interest statement

The author declares that no conflict of interest exists.

Education for Patients Taking Anticoagulants

I am a certified provider of an education and treatment program for patients receiving longterm oral anticoagulation therapy and would have appreciated a deeper analysis of the data. Namely: are there any differences

in the rates of complications between the trained and non-trained groups of patients taking phenprocoumon? Participants in the educational training program receive detailed information about and are made aware of the most important complications—which were also mentioned in the excellent original article—such as hemorrhage, interactions with other medications, and liver injuries. If there is a significant difference, it would be worth considering whether by default, all patients in whom long-term oral anticoagulation with phenprocoumon, warfarin, etc. is indicated should undergo such training, in order to prevent the frequency of complications and health related harms at the individual level, on the one hand, and financial burdens on the health-care system on the other hand. The annual therapeutic cost of treatment with phenprocoumon and the NOACs differ by a factor of 10, which means that phenprocoumon costs about €80 and treatment with NOACs €600–1200 per person/year. The education program is an one-off expense at €170 per person, as is the measuring device (€900) and the test strips (56 per year), and in case of long-term treatment this would certainly be more cost effective in times where keeping contributions down takes priority.

A further, important piece of information: for the NOACs (apixaban, rivaroxaban, and dabigatran etexilate) no data from long term studies are available to date, no antagonists exist, and in case of a heart valve replacement they are not suitable as a therapeutic alternative.

DOI: 10.3238/arztebl.2013.0541b

REFERENCES

1. Schmiedl S, Rottenkolber M, Szymanski J, Siegmund W, Hippus M, Farker K, Drewelow B, Hasford J, Thürmann P: Bleeding complications and liver injuries during phenprocoumon treatment—a multicentre prospective observational study in internal medicine departments. Dtsch Arztebl Int 2013; 110(14): 244–52.

Dr. med. Guntram Fischer

MVZ Immenstadt Allgäu GmbH, Immenstadt
g.fischer@t-online.de

Conflict of interest statement

Dr Fischer is the provider of an education and treatment program for patients receiving long-term oral anticoagulation treatment. He has received honoraria for speaking from Roche.

In Reply:

We thank our correspondents and other colleagues for their feedback on their analysis of phenprocoumon related bleeding complications and liver injuries (1).

Dr Sroka reports on patients with atrial fibrillation who developed bleeding complications while taking guideline-conform prophylactic phenprocoumon or new oral anticoagulants (NOACs). At the same time he expresses hope that current guidelines will be rethought and assessed in the near future. As a matter of fact, in

the scenario of guideline-conform therapy, it is primarily older patients who are given a multitude of medications for existing comorbidities, which may trigger interactions and adverse effects (2). The fundamental problem is that guidelines are based on randomized controlled studies that are more likely to include younger, less multimorbid patients who take far fewer additional medications (3). This does not do justice to the age-related higher risks associated with anticoagulant therapy (for example, after a fall), nor the potential interactions, for example, with phenprocoumon. Our analysis of the risk of bleeding complications associated with phenprocoumon treatment and practice conditions also showed a notable increase with older age. Especially the combination with analgesics, which is common in older age, represents a substantial risk factor for gastrointestinal bleeding complications. We have to concede, however, that, although we were able to quantify the risks of bleeding complications in detail, we were not able to calculate the benefit of anticoagulation—that is, the number of prevented thromboembolic complications in the studied population.

Dr Fischer in his reader's letter emphasizes the importance of educational training programs for patients taking anticoagulants. In the context of our study, participation in and, where applicable, the intensity of, training programs were regrettably not documented. However, our results also show that the international normalized ratio (INR) was too high in about two-thirds of patients with bleeding complications. Routine clinical practice will show the extent to which exactly these patients benefit from NOACs (4). The product information for the three currently available NOACs includes dosage reductions and contraindications

taking into consideration renal function—but the question remains about whether such dosage adjustments or stopping/switching medications are even feasible in routine clinical practice in older patients and in case of fluctuating renal function (5).

DOI: 10.3238/arztebl.2013.0541c

REFERENCES

1. Schmiedl S, Rottenkolber M, Szymanski J, Siegmund W, Hippus M, Farker K, Drewelow B, Hasford J, Thürmann P: Bleeding complications and liver injuries during phenprocoumon treatment—a multicentre prospective observational study in internal medicine departments. *Dtsch Arztebl Int* 2013; 110(14): 244–52.
2. Boyd CM, Darer J, Boulton C, Fried LP, Boulton L, Wu AW: Clinical practice guidelines and quality of care for older patients with multiple comorbid diseases. *JAMA* 2005; 294: 716–24.
3. Thürmann PA: Weniger wäre tatsächlich mehr – die Arzneimittelversorgung alter Menschen. *Z Evid Fortbild Qual Gesundheitsw* 2013; 107: 148–52.
4. Schott G, Bräutigam K, Ludwig WD: Orale Antikoagulation bei nicht valvulärem Vorhofflimmern. Empfehlungen zum Einsatz der neuen Antikoagulation Dabigatran (Pradaxa®) und Rivaroxaban (Xarelto®). Arzneimittelkommission der deutschen Ärzteschaft. Version 1.0. September 2012.
5. Helldén A, Odar-Cederlöf I, Nilsson G, Sjövik S, Söderström A, von Euler A, Öhlén G, Bergman U: Renal function estimations and dose recommendations for dabigatran, gabapentin and valaciclovir: a data simulation study focused on the elderly. *BMJ Open* 2013; 3: e002686. doi:10.1136/bmjopen-2013-002686.

Prof. Dr. med. Petra Thürmann

Philipp Klee-Institut für Klinische Pharmakologie, Helios Klinikum Wuppertal
Lehrstuhl für Klinische Pharmakologie, Universität Witten/Herdecke
petra.thuermann@helios-kliniken.de

Conflict of interest statement

Professor Thürmann has received consulting fees from Biotest Pharma AG, Fresenius Kabi, and MYR GmbH and lecture fees from BayerVital, Biotest Pharma AG, Rottapharm Madaus GmbH, and from the German Institute for Quality and Efficiency in Health Care (IQWiG). Third-party funding was provided to her by Biotest Pharma AG, Stada GmbH, and Bayer Schering Pharma AG.