Published in final edited form as:

Curr Cancer Drug Targets. 2012 July ; 12(6): 607–616.

Cyclooxygenase-2 (COX-2) Mediates Arsenite Inhibition of UVB-Induced Cellular Apoptosis in Mouse Epidermal Cl41 Cells

Z. Zuo^{1,2}, W. Ouyang¹, J. Li¹, M. Costa¹, and C. Huang^{1,*}

¹Nelson Institute of Environmental Medicine, New York University School of Medicine, 57 Old Forge Road, Tuxedo, NY 10987, USA

²Key Laboratory of Ministry of Education for Subtropical Wetland Ecosystem Research, School of Life Sciences, Xiamen University, Xiamen, 361005, China

Abstract

Inorganic arsenic is an environmental human carcinogen, and has been shown to act as a co-carcinogen with solar ultraviolet (UV) radiation in mouse skin tumor induction even at low concentrations. However, the precise mechanism of its co-carcinogenic action is largely unknown. Apoptosis plays an essential role as a protective mechanism against neoplastic development in the organism by eliminating genetically damaged cells. Thus, suppression of apoptosis is thought to contribute to carcinogenesis. It is known that cyclooxygenase-2 (COX-2) can promote carcinogenesis by inhibiting cell apoptosis under stress conditions; and our current studies investigated the potential contribution of COX-2 to the inhibitory effect of arsenite in UV-induced cell apoptosis in mouse epidermal Cl41 cells. We found that treatment of cells with low concentration (5 μ M) arsenite attenuated cellular apoptosis upon UVB radiation accompanied with a co-inductive effect on COX-2 expression and nuclear factor- κ B (NF- κ B) transactivation. Our results also showed that the COX-2 induction by arsenite and UVB depended on an NF- κ B pathway because COX-2 co-induction could be attenuated in either p53-deficient or p50-deficient cells. Moreover, UVB-induced cell apoptosis could be dramatically reduced by the introduction of exogenous COX-2 expression, whereas the inhibitory effect of arsenite on UVB-induced cell apoptosis could be impaired in COX-2 knockdown Cl41 cells. Our results indicated that COX-2 mediated the anti-apoptotic effect of arsenite in UVB radiation through an NF- κ B-dependent pathway. Given the importance of apoptosis evasion during carcinogenesis, we anticipated that COX-2 induction might be at least partially responsible for the co-carcinogenic effect of arsenite on UVB-induced skin carcinogenesis.

Keywords

Apoptosis; arsenite; COX-2; NF- κ B; UVB

INTRODUCTION

Inorganic arsenic is a well-known environmental toxin that is distributed throughout the earth's crust and is released into the environment during the process of energy generation upon utilization of coal, oil shale, and geothermal sources [1, 2]. Ingestion of arsenic has been reported to increase the risk of various cancers, including liver, bladder, kidney, lung,

*Address correspondence to this author at the Nelson Institute of Environmental Medicine, New York University School of Medicine, 57 Old Forge Road, Tuxedo, NY 10987, USA; Tel: 845-731-3519; Fax: 845-351-2118; chuanshu.huang@nyumc.org.

CONFLICT OF INTEREST

Declared none.

as well as skin [3]. A dose-response relationship has been established between the prevalence of skin cancers and the arsenic concentration in drinking water [4, 5]. Ultraviolet radiation (UVR) is a main risk factor for skin carcinogenesis [6], and this threat has increased over recent decades because of the depletion of stratospheric ozone, the principal barrier to the biological effect of UVB (290–320nm) on human skin [7, 8]. Instead of being a complete carcinogen, arsenic could enhance the mutagenicity of other carcinogens such as UVR on the mouse skin [9]. However, the molecular mechanisms underlying the co-carcinogenic effect of arsenic with UVR are not well understood [10].

Cyclooxygenase (COX) exists as two distinct isoforms. COX-1 is a constitutive enzyme, whereas COX-2 is the inducible isoform [11]. COX-2 is undetectable in most mammalian cells, but its expression can be induced rapidly (2–6 h) in different mammal cell lines in response to growth factors, tumor promoters and cytokines [12, 13]. Increasing evidence indicates that COX-2 plays an important role in skin carcinogenesis. For example, mice that are deficient in COX-2 experience reduced tumor formation by 75% in comparison to that in wild type littermates in a multistage mouse skin carcinogenesis model [14]. Oral administration of specific COX-2 inhibitors is also effective in the inhibition of UVB-induced skin tumors by up to 90% [15]. COX-2 induction upon arsenite exposure mediates the anti-apoptotic effect in both mouse Cl41 cells and human BEAS-2B cells [10, 16]. It has been reported that apoptosis induction is a key mechanism for the effectiveness of some cancer chemopreventive agents, and failure of apoptosis is now believed to contribute to the development of human cancer [17], whereas COX-2 overexpression increases rat intestinal epithelial cell adhere to the extracellular matrix and be resistant to apoptosis [18]. Thus, we examined our hypothesis that the synergetic induction of COX-2 by arsenite and UVB is responsible for the anti-apoptotic effect of arsenite against UVB radiation-related cell death in mouse epidermal JB6 Cl41 cells.

MATERIALS AND METHODS

Cell Culture and Reagents

Mouse epidermal Cl41 cells and their stable transfectants were cultured in Eagle's Minimal Essential Medium (MEM, Calbiochem San Diego, CA) that was supplemented with 5% fetal bovine serum (FBS), 1% penicillin/streptomycin, and 2 mM L-glutamine (Life Technologies, Inc., Rockville, MD) at 37°C in a humidified atmosphere with 5% CO₂. Mouse embryonic fibroblasts (MEFs) and the MEFs with either p65 deficiency (p65^{-/-}) or p50 deficiency (p50^{-/-}) were cultured in Dulbecco's Modified Eagle's Medium (DMEM) supplemented with 10% FBS, 1% penicillin/streptomycin, and 2 mM L-glutamine at 37°C in a humidified atmosphere with 5% CO₂. The cultures were dissociated with trypsin and transferred to new 75-cm² culture flasks (Fisher) twice a week. FBS was purchased from Nova-Tech, and other cell culture reagents were obtained from Sigma (St. Louis, MO); the luciferase assay substrate was purchased from Promega (Madison, WI); arsenite (As³⁺) was purchased from Aldrich (Milwaukee, WI); and the antibodies against p65, p50 and COX-2 were purchased from Santa Cruz Biotechnology (Santa Cruz, CA), whereas antibody specific for β -Actin was obtained from Sigma-Aldrich (St. Louis, MO, USA) or Sungene Biotech (Tianjin, China).

Stable Transfection

The AP-1-luciferase, NFAT-luciferase, NF- κ B-luciferase, and COX-2-luciferase reporter plasmids have been described in studies [19, 20]. The COX-2 expression vector and the COX-2 siRNA vector as well as IKK γ -KM were used in our previous studies [10]. The plasmid DNA isolation was performed according to Qiagen plasmid MIDI kit protocol. Cell transfection was carried out with Lipofectamine 2000 (Invitrogen, Inc.) according to the

manufacturer's instructions. Briefly, the cells were cultured in a 6-well plate to 75–85% confluence. A mixture of 2 µg plasmid DNA and 6 µL Lipofectamine 2000 reagent was used to transfect each well in the absence of serum. After 4–6 h, the medium was replaced with 5% FBS MEM. Approximately 36–48 h after the beginning of the transfection, the cells were selected with normal culture medium containing 500 µg/mL G418 (Gibco BRL, Rockville, MD). After a 28–45 day selection, the stable transfectants were established and identified using Western blot and/or luciferase assays. All stable transfectants were cultured in G418-free medium for at least two passages before each experiment.

Western Blot

CI41 cells and their transfectants (2×10^5) were seeded and cultured in each well of 6-well plates until 70–80% confluence. The cells were exposed to arsenite and/or UVB for the dosage and time indicated, and then extracted with SDS-sample buffer, as described in our previous studies [10]. The cell extracts were applied for Western blotting using specific antibodies as indicated. The protein band, specifically bound to the primary antibody, was detected using an IgG-AP-linked secondary antibody and an ECF Western blotting system (Amersham Biosciences, Piscataway, NJ). Quantification of COX-2 expression was normalized to α -Actin using the Image J program.

DNA Fragmentation Assay

DNA fragmentation assay was performed as described in our previous study [21]. Briefly, CI41 cells were cultured in monolayers in 100 mm dishes with 5% FBS MEM. The cells were exposed to arsenite and/or UV radiation in 0.1% FBS MEM after cell density reached 75–80%. All the cells were harvested by centrifugation and lysed with a lysis buffer (5 mM Tris-HCl, pH 8.0, 20 mM EDTA, 0.5% Triton X-100) on ice for 45 min. The fragmented DNA in the supernatant after centrifugation at 14,000 rpm for 30 min at 4°C was extracted twice with phenol/chloroform/isoamyl alcohol (25:24:1, v/v) and once with chloroform before precipitation with ethanol and salt. The DNA pellet was washed once with 70% ethanol and resuspended in TE buffer (pH 8.0) with 100 µg/mL RNase at 37°C for 2 h. The DNA fragments were separated by 1.8% agarose gel electrophoresis and visualized under UV light.

Flow Cytometry Assay

Cells (2×10^5) were seeded into each well of 6-well plates and cultured to 75–80% confluence. After exposure, the cells were harvested and fixed with 3 mL of ice-cold 80% ethanol overnight. The fixed cells were then centrifuged (3000 rpm, 3 min), suspended in lysis buffer (100 mM sodium citrate and 0.1% Triton X-100) and incubated for 15 min at room temperature. The cells were incubated with RNase A (10 mg/mL) (Sigma Chemical, St. Louis, MO) for 10 min at room temperature and DNA was stained with propidium iodide (PI, 50 µg/mL) for 1 h at 4°C. The cell proportions in sub-G1 and S phases were determined using flow cytometry with an Epics XL FACS (Beckman Coulter) and EXPO 32 software [22].

Cell Viability Assay

Cell viability was determined using MTT-based cell growth assay. Cells were rinsed with PBS, incubated in DMEM-free medium containing 0.5 mg/mL 3-[4,5-dimethylthiazol-2-yl]-2,5-diphenyltetrazolium bromide (also called thiazolyl blue or MTT) for 4 h. The purple crystals yielded were dissolved in isopropanol/0.1N HCl. The cell viability was calculated by measuring the difference between the optical densities of the resulting purple solution at 570 and 690 nm.

Gene Reporter Assay

A confluent monolayer of AP-1-, NFAT-, NF- κ B-, or COX-2 luciferase reporter stable transfectants were trypsinized, and 8×10^3 viable cells suspended in 100 μ L of medium were added to each well of 96-well plates. After the cell density reached 75–80% confluence, the cells were treated with arsenite and/or UVB for time periods as indicated in the figure legends. The cells were then lysed with 50 μ L lysis buffer, and the luciferase activity was measured using Promega Luciferase assay reagent with a luminometer (Wallac 1420 Victor2 multipliable counter system). The results were expressed as AP-1, NFAT or NF- κ B activation relative to the medium control (relative AP-1, NFAT or NF- κ B activation) or COX-2 induction relative to the medium control (relative COX-2 induction), as described in our earlier studies [23, 24].

Statistical Analysis

The Student's *t* test was used to determine the significance of differences in COX-2 induction and AP-1, NFAT or NF- κ B activation among various groups as indicated. The differences were considered significant at $P < 0.05$.

RESULTS

Exposure of C141 Cells to Low Concentration Arsenite Rendered the Cells Resistant to the Apoptosis Due to UVB Radiation

UVB irradiation induces apoptosis and alters differentiation in primary human keratinocytes, HaCaT cells, and human epithelial corneal cells [25–27]. In an earlier study, we demonstrate that arsenite exerts an anti-apoptotic effect on C141 cells at low concentrations [10]. Since skin tissue is a major target of both UVB and arsenic, and because arsenite at low concentrations shows a co-carcinogenic effect on mouse skin, mouse epidermal C141 cells were employed to test whether low concentrations of arsenite were able to inhibit UVB-induced cell apoptosis. As shown in Fig. (1A), treatment of C141 cells with UVB radiation at 1 KJ/m² led to marked cell death, whereas 5 μ M of arsenite alone did not produce an observable effect on cell death. However, the pretreatment of C141 cells with 5 μ M of arsenite for 30 min showed a dramatic inhibition of UVB-induced cell death under the same experimental conditions (Fig. 1A). These findings were consistent with the results that we obtained from the cell viability assay using MTT (Fig. 1B) and flow cytometry assay with PI staining (Fig. 1C). DNA content analysis using PI staining followed by flow cytometry analysis was used to quantify the cell apoptosis rate upon arsenite and/or UVB radiation. The results showed that UVB radiation led to 73.5% C141 cell apoptosis at 48 h after exposure, while co-treatment of arsenite and UVB radiation only resulted in 32.9% cell death, in comparison to the treatment of cells with arsenite alone, which led to 6.8% cell apoptosis (Fig. 1C). Accordingly, assays for DNA fragmentation and cleavage caspase 3, two classic indicators of apoptosis [21], were further employed to verify the anti-apoptotic effect of arsenite on UVB-treated C141 cells (Figs. 1D and 1E). Kapahi *et al.* reported that arsenite at doses over 10 μ M could inhibit IKK/NF- κ B activation by binding to Cys-179 in the activation loop of the IKK catalytic subunits and promoted cell apoptosis, whereas it did not show observable inhibition on IKK/NF- κ B activation and induction of cell apoptosis at the dose at 6.25 μ M [28]. Those results were consistent with our previous findings that high dose of arsenite treatment led to cell apoptosis *via* p50/JNK pathway [29], while low doses of arsenite exposure resulted in cell transformation [30]. Thus, the differential effects of arsenite on NF- κ B activation might be due to arsenite doses that were applied. Collectively, our results demonstrated that low concentrations of arsenite protected UVB-exposed C141 cells from apoptosis in mouse epidermal C141 cells. This was consistent with the findings demonstrated in human keratinocytes [31].

Arsenite Had a Co-Inductive Effect with UVB Radiation on COX-2 Expression in CI41 Cells

COX-2 is an inducible early responsive gene, and plays a role in the mediation of inflammation and carcinogenesis [11]. Although studies demonstrate that exposure of cells to either UVB or arsenite induces COX-2 expression both *in vivo* and *in vitro* [10, 18, 32–35], no study has been conducted to investigate whether or not arsenite has a co-inductive effect on COX-2 expression in combination with UVB radiation. Thus, we examined the COX-2 induction by arsenite and/or UVB in CI41 cells. Consistent with the previous reports, exposure of CI41 cells with either UVB or arsenite alone resulted in increases in COX-2 expression (Figs 2A and 2B), whereas arsenite at 5 μM had a co-inductive effect with UVB on COX-2 protein expression (Fig. 2C). Moreover, our results indicated that treatment of CI41 with arsenite and 1KJ/m² UVB also resulted in a significant co-induction of COX-2 expression in the COX-2-luciferase reporter assay (Fig. 2D), suggesting that an co-inductive effect of arsenite with UVB on COX-2 expression might occur at transcriptional level.

NF κ B, but not AP-1 nor NFAT, Was Critical for Co-Induction of COX-2 Due to Arsenite and UVB Exposure

The above results, obtained using the COX-2-luciferase reporter, suggested that co-induction of COX-2 by arsenite and UVB radiation might be regulated at the transcriptional level. Studies show that the COX-2 promoter region contains binding sites for multiple transcription factors, including AP-1, NF κ B, and NFAT, and each of these transcription factors is involved in the regulation of COX-2 expression in various experimental systems, depending on cell types and stimulus [18, 36, 37]. To test whether AP-1 was implicated in the regulation of COX-2 co-induction upon exposure to arsenite and UVB radiation, we determined the AP-1-dependent transactivation due to arsenite and/or UVB radiation using CI41 AP-1-luciferase stable transfectants. The results indicated that arsenite at concentrations of 2.5–10 μM showed neither induction of AP-1 activity, nor any co-inductive effect on AP-1 activity with UVB radiation, even though UVB alone induced a high level of AP-1 transactivation (Fig. 3A). Consistent with AP-1 activation, arsenite and/or UVB radiation showed a similar effect on NFAT transactivation in CI41 NFAT-luciferase reporter assay (Fig. 3B). These results suggested that the transcription factor AP-1 and NFAT might not be involved in the co-inductive effect of arsenite on COX-2 expression with UVB radiation. In contrast with the results observed in AP-1 and NFAT activation, either exposure of cells to arsenite at a dosage range between 2.5–10 μM , and UVB radiation, increased NF κ B activity, with similar patterns of their effects on COX-2 induction (Figs. 3C and 2D). Moreover, arsenite showed a significant co-inductive effect with UVB radiation on NF κ B transactivation under the same experimental conditions (Fig. 3C). Those results indicated that arsenite alone showed neither any inductive effect on activation of AP-1 and NFAT, nor additive effect on UVB-induced activation of AP-1 and NFAT, whereas arsenite alone induced NF κ B activation by itself and also showed a synergistic effect on NF κ B activation with UVB radiation. Thus, we anticipate that NFAT and AP-1 might not be responsible for co-induction of COX-2 expression by arsenite with UVB radiation. It should be noted that we can't exclude the potential role of AP-1 and NFAT in mediation of COX-2 expression by UVB radiation alone. Our published studies have shown that IKK- α and IKK- β are coordinately required for induction of AP-1 activation through an NF- κ B-independent pathway upon apoptotic dose (20 μM) of arsenite exposure [38]. To further elucidate role of NF κ B in COX-2 induction by both arsenite and UVB radiation, the stable transfectants of CI41 cells harboring IKK- α -KM, a dominant mutant of IKK- α were used [10]. As shown in Fig. (3D), combination of arsenite exposure and UVB irradiation led to dramatic increase in COX-2 promoter activity in the control cells expressing mock plasmid, while introduction of IKK- α -KM obviously rendered the blockage in the synergetic effect of arsenite and UVB on COX-2 promoter induction. To further verify the subunit of

NF- κ B involved in this process, NF- κ B p65 knockout MEFs (p65^{-/-}) or NF- κ B p50 knockout MEFs (p50^{-/-}) were employed. As shown in Fig. (3E), knockout of either p65 or p50 dramatically inhibited COX-2 induction by UVB and arsenite (Fig. 3E). It should be noted that there was still marginal COX-2 induction in p65^{-/-} MEFs upon exposure of cells with UVB alone or UVB and arsenite (Fig. 3E). This effect may be mediated by activation of other transcription factors, such as AP-1 and/or NFAT, upon UVB radiation alone. Taken together, these data demonstrated that NF- κ B, not AP-1 nor NFAT, was responsible for co-induction of COX-2 protein expression due to arsenite and UVB radiation.

COX-2 Co-Induction Mediated Arsenite Protection of UVB-Exposed Cl41 Cells from Apoptosis

The COX-2 expression is induced in an IKK- α /NF- κ B-dependent manner after treatment with arsenite in Cl41 cells, and plays an important role in antagonizing apoptosis due to arsenite exposure [10]. Thus, we transfected the COX-2 expression construct into Cl41 cells and determined if COX-2 overexpression provided protection from apoptosis to UVB-exposed Cl41 cells. As shown in Fig. (4A), the COX-2 protein was overexpressed in Cl41 cell transfectants in comparison to that in mock vector-transfectants (Fig. 4A). Overexpression of COX-2 as seen in Cl41-COX2 transfectant showed much less susceptibility to UVB radiation for apoptotic induction in comparison to that in mock vector-transfectant (Fig. 4B), suggesting that COX-2 could provide protection from apoptosis in UVB-irradiated Cl41 cells. To determine whether or not COX-2 co-induction by arsenite and UVB could result in the inhibitory effect of arsenite in UVB-induced cell apoptosis, we used a COX-2 shRNA to knockdown endogenous COX-2 expression in Cl41 cells. We found that introduction of COX-2 shRNA knocked down the serum-induced COX-2 protein expression by 70% (Fig. 4C). Deficiency of COX-2 expression resulted in an increase in UVB-induced cell apoptosis in comparison to the control transfectants (20.7% v. 30.5%, Fig. 4D). Moreover, the reduction of COX-2 expression attenuated the arsenite protective effect on UVB-induced cell apoptosis completely (7.1% v. 25.5%, Fig. 4D). These results demonstrated that co-induction of COX-2 expression upon arsenite and UVB radiation mediated arsenite inhibition of UVB-induced cell apoptosis, and might, at least, be one of the major factors responsible for the arsenite co-carcinogenic effect on mouse skin.

DISCUSSION

The co-carcinogenic effect of arsenite with UVR on mouse skin is well established [9], but the molecular basis underlying this is largely unknown. Here, we focused on the co-inductive effect of arsenite with UVB radiation on signaling pathways leading to COX-2 induction in mouse epidermal JB6 Cl41 cells. We found that arsenite had a co-inductive effect with UVB radiation on COX-2 expression, which made a significant contribution to arsenite protection from apoptosis on UVB-exposed cells. We anticipated this arsenite protection might be at least one of the key factors responsible for the skin co-carcinogenic effect of arsenite with UVB radiation.

The carcinogenic effect of arsenite by occupational or environmental exposure is convincing [3, 39]. Levels of arsenic in drinking-water in epidemiological studies ranged from 54 ppb (0.72 μ M) to 12 ppm (160 μ M) [40], and such levels of arsenite exposure are associated with various cancers [40–42]. Although the arsenite doses used for studies in both *in vitro* and *in vivo* vary among different types of cell and animal models, the doses used in the present study (ranging from 2.5–10 μ M) are similar to those used in previous studies in cell culture models [43–45]. The inhibition of DNA repair capacity is one of the potential mechanisms for the co-carcinogenic activity of arsenic [46]. Low concentration arsenic (2.5 μ M) exposure decreases the expression of critical genes involved in nucleotide excision repair of damaged DNA after UV irradiation [47–49]. Ding *et al.*, (2009) demonstrate that arsenite at low

concentrations inhibits oxidative DNA damage repair by interaction with the PARP-1 zinc finger domain, which leads to the inhibition of PARP-1 activity [50]. Apoptosis typically occurs when cellular genetic damage exceeds the repair capacity and suppression of apoptosis, in the face of significant genetic damage, could facilitate accumulation of aberrant cells and may be a critical step in the pathogenesis of malignancy. Dysregulation of apoptosis may lead to various kinds of pathological conditions, such as cancer development [51]. In this study, we demonstrated that exposure of cells to low arsenite concentrations inhibited UVB-induced cellular apoptosis, revealing that arsenite might have initiated anti-apoptotic signaling and/or gene expression, and provided an anti-apoptotic effect on UVB-exposed cells. This anti-apoptotic effect initiated by arsenite might account for the co-carcinogenesis of arsenite with UV radiation demonstrated in the mouse skin carcinogenesis model.

COX-1 is expressed constitutively, whereas COX-2 is inducible in cells upon treatment with growth factors, tumor promoters, and cytokines [52]. Accumulating evidence suggests that cancer cells which express higher levels of COX-2 may obtain a survival advantage that eventually facilitates tumor development and progression [53, 54]. Over-expression of COX-2 in human skin basal cell carcinoma cells increases antiapoptosis, angiogenesis, and tumorigenesis induced by UVB [55], and NFAT-mediated COX-2 induction is crucial for TNF α -induced cell transformation in C141 cells [56]. The promoter region of COX-2 contains a canonical TATA-box and multiple regulatory elements, which can be recognized by transcription factors, such as NF κ B, AP-1, and NFAT [22, 57–59]. Arsenite induces COX-2 expression at the transcriptional level by activating the NFAT pathway in human bronchial epithelial BEAS-2B cells, whereas in C141 cells, arsenite induces COX-2 through the NF κ B-dependent pathway [16, 37]. These published studies from our group suggest that the signaling pathway leading to COX-2 induction depends on the cell types and stimuli. In the present study, we showed that arsenite at low concentration or UVB radiation alone induced COX-2 expression in C141 cells, while co-exposure of cells with UVB and low concentration arsenite resulted in a marked COX-2 co-induction at both protein and transcription levels in C141 cells. We further found that C141 cells transfected with the COX-2 constitutive expression construct were resistant to UVB for apoptotic induction in comparison to that in mock vector-transfected C141 cells. Moreover, the knockdown of endogenous COX-2 expression by its specific siRNA in C141 cells attenuated the protective effect of low concentration arsenite (5 μ M) on cellular apoptosis upon UVB radiation. Collectively, these results demonstrated that low concentrations of arsenite exerted an anti-apoptotic effect by sustaining a relatively high level of COX-2 expression in C141 cells exposed to UVB radiation. It should be noted that many mechanistic studies have shown that a number of molecules and gene products are involved in arsenic-induced cell proliferation and growth inhibition in model cell lines including Bcl-xL, Bcl-2, c-myc, Cyclin D1, HO-1 and NADPH oxidase [60–62]. Although our results indicated an important role of NF κ B/COX-2 pathway in protection of UV-induced cells from apoptosis, we couldn't exclude other genes involved in such protection.

In our current study, we also found that co-treatment of cells with arsenite and UVB substantially induced NF κ B activation in C141 cells, whereas it did not show any notable co-inductive effect on AP-1 and NFAT activation. Our previous studies demonstrate that arsenite induces AP-1 activation both in the mouse C141 cells and in the skin of AP-1-luciferase transgenic mice, and that the AP-1 induction appears to be mediated by the activation of protein kinase family members [22, 57]. UVB induces COX-2 expression by increasing the activation of AP-1 [34, 36], or NFAT [63], and inhibition of UVB-induced NFAT activation in keratinocytes reduces COX-2 protein expression and increases UV-induced apoptosis [63]. In the current study, although UVB exposure activated both AP-1 and NFAT transactivation, arsenite alone at low concentration showed neither an observable

effect on transactivation of AP-1 and NFAT, nor a co-inductive effect with UVB on the activation of AP-1 and NFAT. This was not consistent with their effects on COX-2 expression, suggesting that AP-1 and NFAT might not be involved in this COX-2 co-induction by arsenite and UVB treatment. This led us to test the potential contribution of NF- κ B to COX-2 co-induction by arsenite and UVB. The results indicated that UVB and arsenite at low concentration did result in co-induction of NF- κ B transactivation in the same Cl41 cells under the same experimental conditions. We further found that knockout of either the NF- κ B component p65 or p50 resulted in a dramatic reduction of COX-2 expression, demonstrating that NF- κ B activation was a major pathway responsible for the mediation of COX-2 co-induction due to arsenite and UVB exposure in Cl41 cells. Taken together with the important role of COX-2 in skin carcinogenesis, we anticipated that co-induction of COX-2 by arsenite and UVB radiation, and its role in the inhibition of UVB-induced cell apoptosis, was at least one of mechanisms underlying the reported co-carcinogenic activities of arsenite in UVR-induced skin carcinogenesis. These studies provided additional insights into understanding the molecular mechanisms that underline the co-carcinogenic effect of arsenite with UV radiation.

Acknowledgments

GRANT SUPPORT

This work was supported in part by grants from NIH/NCI CA112557 (CH), and from NIH/NIEHS ES010344 (CH and MC) and ES-000260 (CH and MC)

We appreciate Dr. John Hodgkiss for his critical reading of the manuscript.

ABBREVIATIONS

AP-1	activator protein-1
Bcl-2	B-cell lymphoma-2
Bcl-xL	B-cell lymphoma-extra large
BEAS-2B	bronchial epithelial cell line
COX	cyclooxygenase
DMEM	Dulbecco's Modified Eagle's Medium
FBS	fetal bovine serum
HO-1	heme oxygenase 1
MEFs	mouse embryonic fibroblasts
MEM	minimal essential medium
MTT	3-[4,5-dimethylthiazol-2-yl]-2,5-diphenyltetrazolium bromide
NADPH	nicotinamide adenine dinucleotide phosphate
NFAT	nuclear factor of activated T cells
NF-κB	nuclear factor- κ B
UVB	ultraviolet-B

References

1. Ball P. Arsenic-free water still a pipedream. *Nature*. 2005; 436:313. [PubMed: 16034383]
2. Nriagu, JO. *Environmental Chemistry of Arsenic*. Dekker; New York: 2002.

3. Chen CJ, Wang CJ. Ecological correlation between arsenic level in well water and age-adjusted mortality from malignant neoplasms. *Cancer Res.* 1990; 50:5470–5474. [PubMed: 2386951]
4. Fawell J. Physiological factors and environmental carcinogenesis. Multiple risk factors associated with arsenic-induced skin cancer: effects of chronic liver disease and malnutritional status. *Hum Exp Toxicol.* 1995; 14:464–465. [PubMed: 7612313]
5. Cabrera HN, Gomez ML. Skin cancer induced by arsenic in the water. *J Cutan Med Surg.* 2003; 7:106–111. [PubMed: 12447618]
6. El Ghissassi F, Baan R, Straif K, Grosse Y, Secretan B, Bouvard V, Benbrahim-Tallaa L, Guha N, Freeman C, Galichet L, Coglian V. A review of human carcinogens--part D: radiation. *Lancet Oncol.* 2009; 10:751–752. [PubMed: 19655431]
7. Longstreth J, de Gruijl FR, Kripke ML, Abseck S, Arnold F, Slaper HI, Velders G, Takizawa Y, van der Leun JC. Health risks. *J Photochem Photobiol B.* 1998; 46:20–39. [PubMed: 9894351]
8. Meghaug I, Olseth JA, Reuder J. UV radiation and skin cancer in Norway. *J Photochem Photobiol B.* 2009; 96:232–241. [PubMed: 19628404]
9. Burns FJ, Uddin AN, Wu F, Nadas A, Rossman TG. Arsenic-induced enhancement of ultraviolet radiation carcinogenesis in mouse skin: a dose-response study. *Environ Health Perspect.* 2004; 112:599–603. [PubMed: 15064167]
10. Ouyang W, Zhang D, Ma Q, Li J, Huang C. Cyclooxygenase-2 induction by arsenite through the IKKbeta/NFkappaB pathway exerts an antiapoptotic effect in mouse epidermal Cl41 cells. *Environ Health Perspect.* 2007; 115:513–518. [PubMed: 17450217]
11. Smith WL, Garavito RM, DeWitt DL. Prostaglandin endoperoxide H synthases (cyclooxygenases)-1 and -2. *J Biol Chem.* 1996; 271:33157–33160. [PubMed: 8969167]
12. Matsuzuka T, Miller K, Pickel L, Doi C, Ayuzawa R, Tamura M. The synergistic induction of cyclooxygenase-2 in lung fibroblasts by angiotensin II and pro-inflammatory cytokines. *Mol Cell Biochem.* 2009; 320:163–171. [PubMed: 18827978]
13. Xu K, Chang CM, Gao H, Shu HK. Epidermal growth factor-dependent cyclooxygenase-2 induction in gliomas requires protein kinase C-delta. *Oncogene.* 2009; 28:1410–1420. [PubMed: 19169273]
14. Tiano HF, Loftin CD, Akunda J, Lee CA, Spalding J, Sessoms A, Dunson DB, Rogan EG, Morham SG, Smart RC, Langenbach R. Deficiency of either cyclooxygenase (COX)-1 or COX-2 alters epidermal differentiation and reduces mouse skin tumorigenesis. *Cancer Res.* 2002; 62:3395–3401. [PubMed: 12067981]
15. Fischer SM, Lo HH, Gordon GB, Seibert K, Kelloff G, Lubet RA, Conti CJ. Chemopreventive activity of celecoxib, a specific cyclooxygenase-2 inhibitor, and indomethacin against ultraviolet light-induced skin carcinogenesis. *Mol Carcinog.* 1999; 25:231–240. [PubMed: 10449029]
16. Ding J, Li J, Xue C, Wu K, Ouyang W, Zhang D, Yan Y, Huang C. Cyclooxygenase-2 induction by arsenite is through a nuclear factor of activated T-cell-dependent pathway and plays an antiapoptotic role in Beas-2B cells. *J Biol Chem.* 2006; 281:24405–24413. [PubMed: 16809336]
17. Samaha HS, Hamid R, ElBayoumy K, Rao CV, Reddy BS. The role of apoptosis in the modulation of colon carcinogenesis by dietary fat and by the organoselenium compound 1,4-phenylenebis(methylene) selenocyanate. *Cancer Epidem Biomar.* 1997; 6:699–704.
18. Tsujii M, DuBois RN. Alterations in cellular adhesion and apoptosis in epithelial cells overexpressing prostaglandin endoperoxide synthase 2. *Cell.* 1995; 83:493–501. [PubMed: 8521479]
19. Ding J, Wu K, Zhang D, Luo W, Li J, Ouyang W, Song L, Huang C. Activation of both nuclear factor of activated T cells and inhibitor of nuclear factor-kappa B kinase beta-subunit/nuclear factor-kappa B is critical for cyclooxygenase-2 induction by benzo[a]pyrene in human bronchial epithelial cells. *Cancer Sci.* 2007; 98:1323–1329. [PubMed: 17640307]
20. Huang C, Zhang D, Li J, Tong Q, Stoner GD. Differential inhibition of UV-induced activation of NF kappa B and AP-1 by extracts from black raspberries strawberries blueberries. *Nutr Cancer.* 2007; 58:205–212. [PubMed: 17640167]
21. Huang C, Ma WY, Li J, Dong Z. Arsenic induces apoptosis through a c-Jun NH2-terminal kinase-dependent, p53-independent pathway. *Cancer Res.* 1999; 59:3053–3058. [PubMed: 10397243]

22. Ouyang W, Ma Q, Li J, Zhang D, Liu ZG, Rustgi AK, Huang C. Cyclin D1 induction through I κ B kinase beta/nuclear factor- κ B pathway is responsible for arsenite-induced increased cell cycle G1-S phase transition in human keratinocytes. *Cancer Res.* 2005; 65:9287–9293. [PubMed: 16230390]
23. Huang C, Huang Y, Li J, Hu W, Aziz R, Tang MS, Sun N, Cassady J, Stoner GD. Inhibition of benzo(a)pyrene diol-epoxide-induced transactivation of activated protein 1 and nuclear factor κ B by black raspberry extracts. *Cancer Res.* 2002; 62:6857–6863. [PubMed: 12460899]
24. Guo W, Yang Z, Xia Q, Liu J, Yu Y, Li J, Zuo Z, Zhang D, Li X, Shi X, Huang C. Arsenite stabilizes HIF-1 protein through p85-mediated up-regulation of inducible Hsp70 protein expression. *Cell Mol Life Sci.* 2011; 68:475–488. [PubMed: 20835880]
25. Mammone T, Gan D, Collins D, Lockshin RA, Marenus K, Maes D. Successful separation of apoptosis and necrosis pathways in HaCaT keratinocyte cells induced by UVB irradiation. *Cell Biol Toxicol.* 2000; 16:293–302. [PubMed: 11201053]
26. Liu XJ, Shi ST, Ye JL, Liu LQ, Sun M, Wang CB. Effect of polypeptide from *Chlamys farreri* on UVB-induced ROS/NF- κ B/COX-2 activation and apoptosis in HaCaT cells. *J Photochem Photobiol B.* 2009; 96:109–116. [PubMed: 19481953]
27. Pauloin T, Dutot M, Joly F, Warnet JM, Rat P. High molecular weight hyaluronan decreases UVB-induced apoptosis and inflammation in human epithelial corneal cells. *Mol Vision.* 2009; 15:577–583.
28. Kapahi P, Takahashi T, Natoli G, Adams SR, Chen Y, Tsien RY, Karin M. Inhibition of NF- κ B activation by arsenite through reaction with a critical cysteine in the activation loop of I κ B kinase. *J Biol Chem.* 2000; 275:36062–36066. [PubMed: 10967126]
29. Song L, Li JX, Zhang DY, Liu ZG, Ye JP, Zhan QM, Huang CS. IKK beta programs to turn on the GADD45 alpha-MKK4-JNK apoptotic cascade specifically *via* p50 NF- κ B in arsenite response. *J Cell Biol.* 2006; 175:607–617. [PubMed: 17116751]
30. Ouyang WM, Luo WJ, Zhang DY, Jian JL, Ma Q, Li JX, Huang CS. PI-3K/Akt pathway-dependent cyclin D1 expression is responsible for arsenite-induced human keratinocyte transformation. *Environ Health Persp.* 2008; 116:1–6.
31. Pi J, He Y, Bortner C, Huang J, Liu J, Zhou T, Qu W, North SL, Kasprzak KS, Diwan BA, Chignell CF, Waalkes MP. Low level, long-term inorganic arsenite exposure causes generalized resistance to apoptosis in cultured human keratinocytes: potential role in skin co-carcinogenesis. *Int J Cancer.* 2005; 116:20–26. [PubMed: 15756686]
32. Buckman SY, Gresham A, Hale P, Hruza G, Anast J, Masferrer J, Pentland AP. COX-2 expression is induced by UVB exposure in human skin: implications for the development of skin cancer. *Carcinogenesis.* 1998; 19:723–729. [PubMed: 9635856]
33. Shi B, Isseroff RR. Arsenite pre-conditioning reduces UVB-induced apoptosis in corneal epithelial cells through the anti-apoptotic activity of 27 kDa heat shock protein (HSP27). *J Cell Physiol.* 2006; 206:301–308. [PubMed: 16114012]
34. Kundu JK, Chang EJ, Fujii H, Sun B, Surh YJ. Oligonol inhibits UVB-induced COX-2 expression in HR-1 hairless mouse skin--AP-1 and C/EBP as potential upstream targets. *Photochem Photobiol.* 2008; 84:399–406. [PubMed: 18221453]
35. Tang X, Kim AL, Kopelovich L, Bickers DR, Athar M. Cyclooxygenase-2 inhibitor nimesulide blocks ultraviolet B-induced photocarcinogenesis in SKH-1 hairless mice. *Photochem Photobiol.* 2008; 84:522–527. [PubMed: 18266822]
36. Cho JW, Park K, Kweon GR, Jang BC, Baek WK, Suh MH, Kim CW, Lee KS, Suh SI. Curcumin inhibits the expression of COX-2 in UVB-irradiated human keratinocytes (HaCaT) by inhibiting activation of AP-1: p38 MAP kinase and JNK as potential upstream targets. *Exp Mol Med.* 2005; 37:186–192. [PubMed: 16000872]
37. Ouyang W, Ma Q, Li J, Zhang D, Ding J, Huang Y, Xing MM, Huang C. Benzo[a]pyrene diol-epoxide (B[a]PDE) upregulates COX-2 expression through MAPKs/AP-1 and IKKbeta/NF- κ B in mouse epidermal C141 cells. *Mol Carcinog.* 2007; 46:32–41. [PubMed: 16921490]
38. Song L, Li JX, Hu MR, Huang CS. Both IKK alpha and IKK beta are implicated in the arsenite-induced AP-1 transactivation correlating with cell apoptosis through NF- κ B activity-independent manner. *Exp Cell Res.* 2008; 314:2187–2198. [PubMed: 18508047]

39. Chan PC, Huff J. Arsenic carcinogenesis in animals and in humans: mechanistic, experimental, and epidemiological evidence. *Environ Carcinog Ecotoxicol Rev.* 1997; C15:83–122.
40. Gomez-Camirero, A.; Howe, P.; Hughes, M.; Kenyon, E.; Lewis, DR.; Moore, M.; Ng, J.; Aitio, A.; Becking, G. Arsenic and Arsenic Compounds. World Health Organization; Geneva: 2001.
41. Tseng WP. Effects and dose--response relationships of skin cancer and blackfoot disease with arsenic. *Environ Health Perspect.* 1971; 19:109–119. [PubMed: 908285]
42. Pi J, Yamauchi H, Kumagai Y, Sun G, Yoshida T, Aikawa H, Hopenhayn-Rich C, Shimojo N. Evidence for induction of oxidative stress caused by chronic exposure of Chinese residents to arsenic contained in drinking water. *Environ Health Perspect.* 2002; 110:331–336. [PubMed: 11940449]
43. Ng, JC. Ph D Thesis. University of Queensland; Australia: 1999. Bioavailability and Toxicology of Arsenic in the Environment.
44. Simeonova PP, Wang S, Toriuma W, Kommineni V, Matheson J, Unimye N, Kayama F, Harki D, Ding M, Vallyathan V, Luster MI. Arsenic mediates cell proliferation and gene expression in the bladder epithelium: Association with activating protein-1 transactivation. *Cancer Res.* 2000; 60:3445–3453. [PubMed: 10910055]
45. Straub AC, Clark KA, Ross MA, Chandra AG, Li S, Gao X, Pagano PJ, Stolz DB, Barchowsky A. Arsenic-stimulated liver sinusoidal capillarization in mice requires NADPH oxidase-generated superoxide. *Clin Invest.* 2008; 118:3980–3989.
46. Andrew AS, Karagas MR, Hamilton JW. Decreased DNA repair gene expression among individuals exposed to arsenic in United States drinking water. *Int J Cancer.* 2003; 104:263–268. [PubMed: 12569548]
47. Hartwig A, Groblichoff UD, Beyersmann D, Natarajan AT, Filon R, Mullenders LH. Interaction of arsenic(III) with nucleotide excision repair in UV-irradiated human fibroblasts. *Carcinogenesis.* 1997; 18:399–405. [PubMed: 9054635]
48. Danaee H, Nelson HH, Liber H, Little JB, Kelsey KT. Low dose exposure to sodium arsenite synergistically interacts with UV radiation to induce mutations and alter DNA repair in human cells. *Mutagenesis.* 2004; 19:143–148. [PubMed: 14981161]
49. Sykora P, Snow ET. Modulation of DNA polymerase beta-dependent base excision repair in cultured human cells after low dose exposure to arsenite. *Toxicol Appl Pharmacol.* 2008; 228:385–394. [PubMed: 18252256]
50. Ding W, Liu W, Cooper KL, Qin XJ, de Souza Bergo PL, Hudson LG, Liu KJ. Inhibition of poly(ADP-ribose) polymerase-1 by arsenite interferes with repair of oxidative DNA damage. *J Biol Chem.* 2009; 284:6809–6817. [PubMed: 19056730]
51. Hanahan D, Weinberg RA. The hallmarks of cancer. *Cell.* 2000; 100:57–70. [PubMed: 10647931]
52. Taketo MM. Cyclooxygenase-2 inhibitors in tumorigenesis (Part II). *J Natl Cancer Inst.* 1998; 90:1609–1620. [PubMed: 9811310]
53. Xu L, Zhang L, Yi Y, Kang HK, Datta SK. Human lupus T cells resist inactivation and escape death by upregulating COX-2. *Nat Med.* 2004; 10:411–415. [PubMed: 14991050]
54. Dannenberg AJ, Lippman SM, Mann JR, Subbaramaiah K, DuBois RN. Cyclooxygenase-2 and epidermal growth factor receptor: pharmacologic targets for chemoprevention. *J Clin Oncol.* 2005; 23:254–266. [PubMed: 15637389]
55. Tjiu JW, Liao YH, Lin SJ, Huang YL, Tsai WL, Chu CY, Kuo ML, Jee S. H. Cyclooxygenase-2 overexpression in human basal cell carcinoma cell line increases antiapoptosis, angiogenesis, and tumorigenesis. *J Invest Dermatol.* 2006; 126:1143–1151. [PubMed: 16528365]
56. Yan Y, Li J, Ouyang W, Ma Q, Hu Y, Zhang D, Ding J, Qu Q, Subbaramaiah K, Huang C. NFAT3 is specifically required for TNF-alpha-induced cyclooxygenase-2 (COX-2) expression and transformation of C141 cells. *J Cell Sci.* 2006; 119:2985–2994. [PubMed: 16803872]
57. Huang C, Bode AM, Chen NY, Ma WY, Li J, Nomura M, Dong Z. Transactivation of AP-1 in AP-1-luciferase reporter transgenic mice by arsenite and arsenate. *Anticancer Res.* 2001; 21:261–267. [PubMed: 11299744]
58. Zuo Z, Cai T, Li J, Zhang D, Yu Y, Huang C. Hexavalent chromium Cr(VI) up-regulates COX-2 expression through an NF B/c-Jun/AP-1-dependent pathway. *Environ Health Perspect.* 2012; 120:547–553. [PubMed: 22472290]

59. Cai T, Li X, Ding J, Luo W, Li J, Huang C. A cross-talk between NFAT and NF- κ B pathways is crucial for nickel-induced COX-2 expression in Beas-2B cells. *Curr Cancer Drug Targets*. 2011; 11:548–559. [PubMed: 21486220]
60. Chai CY, Huang YC, Hung WC, Kang WY, Chen WT. Arsenic salt-induced DNA damage and expression of mutant p53 and COX-2 proteins in SV-40 immortalized human uroepithelial cells. *Mutagenesis*. 2007; 22:403–408. [PubMed: 17906315]
61. Kitchin KT, Conolly R. Arsenic-induced carcinogenesis-oxidative stress as a possible mode of action and future research needs for more biologically based risk assessment. *Chem Res Toxicol*. 2010; 23(2):327–335. [PubMed: 20035570]
62. Ma Y, Wang JZ, Liu LX, Zhu HQ, Chen XN, Pan SH, Sun XY, Jiang HC. Genistein potentiates the effect of arsenic trioxide against human hepatocellular carcinoma: Role of Akt and nuclear factor-kappa B. *Cancer Lett*. 2011; 301:75–84. [PubMed: 21078540]
63. Flockhart RJ, Diffey BL, Farr PM, Lloyd J, Reynolds NJ. NFAT regulates induction of COX-2 and apoptosis of keratinocytes in response to ultraviolet radiation exposure. *FASEB J*. 2008; 22:4218–4227. [PubMed: 18708588]

Fig. 1. Arsenite at low concentration rendered C141 cells resistant to pro-apoptosis effects upon UVB radiation

C141 cells (2×10^5) were seeded into each well of 6-well plates. After being cultured at 37°C overnight, the cells were treated with UVB (1KJ/m²), or arsenite (5 μM) for 48 h, or C141 cells were pretreated with arsenite (5 μM) for 30 min and then exposed to UVB (1KJ/m²). The cells were incubated for an additional 48 h and cell death was imaged under microscopy (A), or monitored with MTT assay (B). Each bar indicates the mean $\pm$ SD of three replicate assay wells. The symbol (*) indicates that the cell survival rate significantly decreased from the control medium ($p < 0.05$). The symbol () indicates that the cell survival rate significantly increased from UVB radiation alone ($p < 0.05$). Apoptotic cells were analyzed by flow cytometry with PI (propidium iodide) staining (C) or by the detection of DNA fragments in agarose gel electrophoresis (D) and assay for cleavage of caspase 3 (E).

Fig. 2. Arsenite and/or UVB induced COX-2 expression in C141 cells

(A–C) C141 cells (2×10^5) were seeded into each well of 6-well plates. After being cultured at 37°C overnight, the cells were treated with UVB (1KJ/m²) then cultured for 6 h or 12h (A), or various concentrations of arsenite for 12 h (B), or pretreatment of C141 cells with arsenite for 30 min then treated with UVB at 1KJ/m², then cultured for 12 h (C). The cells were then extracted with SDS-sample buffer, and cell extracts were subjected to Western blotting with specific antibodies against either COX-2 or β -Actin. Quantification of COX-2 expression was normalized to β -Actin using the Image J program. (D) COX-2 luciferase reporter stable transfectant was seeded into 96-well plates. The cells were exposed to arsenite for 6 h, or pretreated with arsenite for 30 min then treated with UVB at 1KJ/m², then cultured for 6 h. The cells were then extracted with 50 μ L lysis buffer for luciferase activity assay, as described in our previous studies. The results are expressed as COX-2 induction relative to the control medium (relative COX-2 induction). Each bar indicates the mean $\pm$ SD of three replicate assay wells. The symbol (*) indicates a significant increase from the control medium ($p < 0.05$); the symbol (♥) indicates a significant increase from UVB radiation alone ($p < 0.05$);

Fig. 3. NF-κB, not AP-1 or NFAT, mediated the COX-2 co-induction due to arsenite and UVB exposure

(A–D), AP-1 luciferase reporter stable transfectants (A), NFAT luciferase reporter stable transfectants (B), or NF-κB luciferase reporter stable transfectants (C), or IKK-κM stable transfectants harboring COX-2 promoter luciferase (D) were seeded into 96-well plates and cultured in 10% FBS-DMEM at 37°C until the cell density reached 75–80%. The cells were exposed to arsenite for 6 h, or pretreated with arsenite for 30 min, then treated with UVB at 1KJ/m², then cultured for 6 h, and then extracted with 50μL lysis buffer for luciferase activity assay. The results are expressed as AP-1, NFAT or NF-κB activity, or COX-2 promoter activity induction relative to the control medium. Each bar indicates the mean ± SD of three replicate assay wells. The symbol (*) indicates a significant increase from the control medium ($p < 0.05$). The symbol (Δ) indicates a significant increase from UVB radiation alone ($p < 0.05$); The symbol (◇) indicates a significant increase from that in transfectants expressing Mock vector ($p < 0.05$) (E), mouse embryonic fibroblasts (MEFs), with either p65 deficient (p65^{-/-}) or p50 deficient (p50^{-/-}), were cultured in 10% FBS DMEM until 75–80% confluence. The cells were exposed to arsenite for 12 h, or pretreated with arsenite for 30 min then treated with UVB (1KJ/m²), then cultured for 12 h, and then extracted with SDS sample buffer. The cell extracts were subjected to Western blotting with specific antibodies against COX-2, P65, P50 or β-Actin. Quantification of COX-2 expression was normalized to β-Actin using the Image J program.

Fig. 4. COX-2 was the mediator for arsenite protection of UVB-exposed C141 cells from apoptosis

COX-2 constitutive expression (**A**) and knockdown (**C**) were identified with Western blotting as indicated; **B** and **D**, C141 cells (2×10^5) were seeded into each well of 6-well plates. After being cultured at 37°C overnight, the cells were treated with UVB (1KJ/m²), or pretreated with 5 μ M of arsenite for 30 min and then exposed to UVB (1KJ/m²) and cultured for an additional 36h. The cellular apoptosis was analyzed using flow cytometry with PI staining. The experiments were repeated three times.