P. Velankar, M.D.

A CASE OF REVERSE TRANSIENT ISCHEMIC DILATATION

Pradnya Velankar, M.D.;^a Venkateshwar Polsani, M.D.;^b Erasmo de la Pena-Almaguer, M.D.;^c Su Min Chang, M.D.^b

^aHouston Methodist Hospital, Houston, Texas; ^bHouston Methodist DeBakey Heart & Vascular Center, Houston Methodist Hospital, Houston, Texas; ^cInstituto de Cardiología y Medicina Vascular, Hospital San José Tec de Monterrey, Monterrey, Mexico

Introduction

Transient ischemic dilatation (TID) is a phenomenon that occurs when there is subendocardial ischemia and the post-stress tomograms show a larger left ventricular cavity size compared to the rest images. We present a rare case of reverse TID in which the opposite findings are seen.

Case Report

A 65-year-old female with a past medical history of hypertension was admitted for left-sided chest pain. An electrocardiogram and three sets of cardiac enzymes were negative. Single-photon emission computed tomography (SPECT) with regadenoson was negative for ischemia. The patient was in atrial fibrillation at 131 bpm during stress imaging (Figure 1), while on rest imaging she was in sinus bradycardia at 59 bpm (Figure 2). Stress images showed a smaller left ventricular cavity (ungated volume of 32 mL) and thicker wall compared to the rest images (ungated volume of 56 mL) (Figures 3, 4). TID ratio was 0.57. Reverse TID occurs with tachyarrhythmias during stress imaging, in which a reduction in diastolic filling time and higher counts obtained during systole lead to a smaller LV size, and bradycardia during rest imaging, where longer diastolic filling time and higher counts obtained during diastole lead to a larger LV size.^{1,2} The clinical significance is unknown.

Figure 1. Rhythm strip showing atrial fibrillation during stress (heart rate = 131 bpm).

Figure 2. Rhythm strip showing sinus bradycardia during rest portion of nuclear stress test (heart rate = 59 bpm).

Conflict of Interest Disclosure: The authors have completed and submitted the *Methodist DeBakey Cardiovascular Journal* Conflict of Interest Statement and none were reported.

Funding/Support: The authors have no funding disclosures.

Keywords: transient ischemic dilatation, reverse transient ischemic dilatation, SPECT

Figure 3. Stress images show a smaller and thicker left ventricle compared to the resting images.

Figure 4. Stress image on left shows smaller left ventricular size (ungated volume of 32 mL) compared to the rest image on right (ungated volume of 56 mL).

References

- Gadiraju R, Bommireddipalli S, Ahuja A, Klein E, DePuey G. Sustained supraventricular tachycardia resulting in a pattern of apparent “reverse transient ischemic dilatation” and an underestimated ejection fraction. *J Nucl Cardiol*. 2010 Jan-Feb;17(1):153-7.
- Hardebeck C. The influence of heart rate on gated SPECT images. *J Nucl Cardiol*. 2007 Jan;14(1):e1-5.