Published in final edited form as:

Conserv Genet Resour. 2012 December ; 4(4): 979–981. doi:10.1007/s12686-012-9687-5.

17 novel polymorphic microsatellite markers for the giant water bug, *Abedus herberti* (Belostomatidae)

T. S. Daly-Engel,

Entomology Department, University of Arizona, Tucson, AZ 85721, USA

R. L. Smith,

Entomology Department, University of Arizona, Tucson, AZ 85721, USA

D. S. Finn,

Department of Zoology, Oregon State University, Corvallis, OR 97331, USA

M. E. Knoderbane,

Department of Zoology, Oregon State University, Corvallis, OR 97331, USA

I. C. Phillipsen, and

Department of Zoology, Oregon State University, Corvallis, OR 97331, USA

D. A. Lytle

Department of Zoology, Oregon State University, Corvallis, OR 97331, USA

Abstract

The giant water bug (*Abedus herberti*) is a large flightless insect that is a keystone predator in aridland aquatic habitats. Extended droughts, possibly due to climate change and groundwater pumping, are causing once-perennial aquatic habitats to dry, resulting in serious conservation concern for some populations. *A. herberti* also exhibits exclusive male parental care, which has made it a model organism for studying mating systems evolution. Here we describe 17 novel polymorphic microsatellite loci developed for *A. herberti*. Number of alleles per locus ranged from 2 to 15, and average observed and expected heterozygosities were 0.579 and 0.697, respectively. These loci can successfully resolve both population genetic structure among sites separated by 3–100 km ($F_{ST} = 0.08–0.21$, $P < 0.0001$), and divergent mating strategies within local populations, making them highly useful for conservation genetics studies of this vulnerable species.

Keywords

Hemiptera; Stream ecology; Mating systems; Polygyny; Population genetics

The giant water bug *Abedus herberti* (Belostomatidae) is a keystone aquatic invertebrate of conservation concern. This long-lived, flightless predator occupies perennial aquatic habitats in aridlands of Arizona, USA and northern Mexico. Extirpation of populations, possibly due

to climate change and water withdrawals by humans, is ongoing and leads to significant ecosystem-level changes (Bogan and Lytle 2011). Its naturally fragmented distribution and conservation status have made *A. herberti* the subject of recent studies on phylogeography (Finn et al. 2007), population stability (Finn et al. 2009), and behavioral adaptations (Lytle et al. 2008). Belostomatids are also interesting because males provide exclusive parental care, and *A. herberti* has become a model system for studying sperm competition (Smith 1979) and mating systems evolution (Smith 1997). Although mitochondrial DNA is useful for resolving broad-scale patterns of population structure (Finn et al. 2007; Daly-Engel et al. 2012), highly polymorphic markers such as microsatellites are needed to understand how genetic diversity varies across complex landscapes in the face of ongoing environmental change (Phillipsen and Lytle in review), and to answer questions at the intrapopulation scale, e.g., assigning parentage and kinship (Daly-Engel et al. 2010). Here, we present 17 polymorphic microsatellite loci that are useful for both purposes.

Tissue samples from two *A. herberti* were used by Genetic Identification Services (GIS; Chatsworth, CA) to construct four enriched microsatellite libraries for CA-, ATG-, CATC- and TAGA-nucleotide repeat motifs using standard cloning protocol (see Murray et al. 2008). GIS used DesignerPCR v.1.03 (Research Genetics, Inc) to design primers for 37 microsatellites that we then optimized in 34–77 unrelated *A. herberti* collected from the Santa Rita Mountains of Arizona. Leg clips from adult bugs were stored in 20 % DMSO-saturated salt buffer (Seutin et al. 1991) and DNA was extracted using Qiagen DNeasy Blood & Tissue kits (Valencia, CA). Loci were electrophoresed for optimal annealing temperature and allelic diversity using a series of 3 % agarose gels run at 80v for approximately 2 h. All forward primers were labeled with proprietary dyes obtained from Applied Bio-systems (Foster City, California; Table 1) except for AHA003, where the reverse primer was labeled for guanine excess. PCR reactions consisted of 0.1 U Biolase *Taq* DNA polymerase (Bioline; Randolph, Massachusetts), 1x *Taq* buffer, 0.50–0.05 μ m of each primer (Table 1), 200 μ m each dNTP, and 2.0 mm MgCl₂. PCR amplification consisted of an initial denaturation at 95 °C for 4 min, followed by 35 cycles of 1 min at 95 °C, 30 s at 52 °C, and 30 s at 72 °C, followed by a final extension at 72 °C for 20 min. PCR products were resolved with an ABI 3730 automated sequencer and visualized using ABI PRISM GENEMAPPER Software 3.0 (Applied Biosystems).

Of the 37 loci provided by GIS, 17 (46 %) were successfully genotyped (Table 1). We estimated heterozygosity and tested for deviation from Hardy–Weinberg Equilibrium (HWE) using GENETOP 3.4 (Raymond and Rousset 1995). One locus, AHC011, only amplified in 16 individuals, but we included these data in hopes that they will be useful to future investigators. ARLEQUIN 3.11 (Excoffier et al. 2005) indicated that no loci were positive for linkage disequilibrium after Bonferroni correction. 4 loci were out of HWE because of heterozygote deficiency, which MICROCHECKER (van Oosterhout et al. 2004) inferred was due to the presence of null alleles.

Data from two studies demonstrate the utility of these loci in identifying fine-scale genetic structure and determining mating system in wild populations of *A. herberti*. A subset of ten loci was used to resolve population genetic structure among 19 collection sites (median sample size = 27) from four Arizona mountain ranges (Phillipsen and Lytle in review). Euclidean distances separating the sites ranged from 3 to 100 km. Loci were amplified in three multiplex PCR sets using Qiagen's Multiplex PCR kits: Set A (AHA010, AHA114, AHB009, AHD006), Set B (AHA103, AHA112, AHA106), and Set C (AHA111, AHA002, AHB107). Exact tests performed in FSTAT with 1000 replications (Goudet 1995) showed significant genetic differentiation across loci, with F_{ST} values ranging from 0.08 to 0.21 ($P < 0.0001$; Phillipsen and Lytle in review). A different subset of 12 loci was used to test for heritability in a single male *A. herberti* and 32 of his offspring. Loci were amplified in three

different multiplex PCR sets: Set A (AHA005, AHA112, AHA106, AHA103), Set B (AHA114, AHA010, AHA03a, AHB009), and Set C (AHA003, AHA002, AHB107, AHC011). For multiplexing, primers were mixed in three pools at optimal concentration (Table 1) and added to the PCR at 1/5 total reaction volume. Paternal alleles were inherited in expected 50:50 ratios, and four maternal alleles were detected among the brood, indicating a polygynous mating system.

This suite of polymorphic markers is the first developed for a member of the genus *Abedus*, of which the species *A. herberti* has become a model organism in the study of arthropod evolution and aridland conservation ecology. The successful resolution of both broad- and fine-scale population genetic structure and identification of individual mating systems in *A. herberti* demonstrate the utility of these loci for conservation genetics research.

Acknowledgments

We thank Brad Freese, Jeff Eble, Saul Macias, Chris Goforth, Mike Bogan, Mike Blouin, and Arlo Pelegrin for help with genetic analyses and collecting specimens. Funding was provided by National Science Foundation grant DEB-0445366 to D.A.L., a Howard Hughes Medical Institute scholarship to M.E.K., grant No. 2 K12 GM000708 to the PERT Program at the University of Arizona on behalf of T.D.E. from the National Institutes of Health, and Oregon State University (URISC Program).

References

- Bogan MT, Lytle DA. Severe drought drives novel community trajectories in desert stream pools. *Freshw Biol.* 2011; 56:2070–2081.
- Daly-Engel TS, Grubbs RD, Feldheim KA, Bowen BW, Toonen RJ. Is multiple mating beneficial or unavoidable? Low multiple paternity and genetic diversity in the shortspine spurdog (*Squalus mitsukurii*). *Mar Ecol Prog Ser.* 2010; 403:255–267.
- Daly-Engel TS, Randall JE, Bowen BW. Is the Great Barracuda (*Sphyrna barracuda*) a reef fish or a pelagic fish? The phylogeographic perspective. *Mar Biol.* 2012; 159:975–985.
- Excoffier L, Laval LG, Schneider S. Arlequin ver. 3.0: an integrated software package for population genetics data analysis. *Evolut Bioinform Online.* 2005; 1:47–50.
- Finn DS, Blouin MS, Lytle DA. Population genetic structure reveals terrestrial affinities for a headwater stream insect. *Freshw Biol.* 2007; 52:1881–1897.
- Finn DS, Bogan MT, Lytle DA. Demographic stability metrics for conservation prioritization of isolated populations. *Conserv Biol.* 2009; 23(5):1185–1194. [PubMed: 19774708]
- Goudet J. FSTAT (version 1.2): a computer program to calculate F-statistics. *J Hered.* 1995; 86:485–486.
- Lytle DA, Bogan MT, Finn DS. Evolution of aquatic insect behaviours across a gradient of disturbance predictability. *Proc Royal Soci B.* 2008; 275:453–462.
- Murray KL, Fleming TH, Gaines MS, Williams DA. Characterization of polymorphic microsatellite loci for two species of phyllostomid bats from the Greater Antilles (*Erophylla sezekorni* and *Macrotus waterhousii*). *Mol Ecol Resour.* 2008; 8:596–598. [PubMed: 21585843]
- Raymond M, Rousset F. GENEPOP Version 1.2. Population genetics software for exact tests and ecumenicism. *J Hered.* 1995; 86:248–249.
- Seutin G, White BN, Boag PT. Preservation of avian blood and tissue samples for DNA analyses. *Can J Zool.* 1991; 69(1):82–90.
- Smith RL. Repeated copulation and sperm precedence: paternity assurance for a male brooding water bug. *Science.* 1979; 205:1029–1031. [PubMed: 17795564]
- Smith, RL. Evolution of paternal care in the giant water bugs (Heteroptera: Belostomatidae).. In: Choe, J.; Crespi, B., editors. The evolution of social behavior in insects and arachnids. Cambridge University Press; UK: 1997. p. 116-149.
- Phillipsen IC, Lytle DA. Aquatic insects in a sea of desert: landscape genetics reveals connections between species biology and population structure. *Ecography.* (in review) (submitted).

van Oosterhout CV, Hutchinson W, Wills D, Shipley P. Micro-checker: software for identifying and correcting genotype errors in microsatellite data. *Mol Ecol Notes*. 2004; 4:535–538.

Table 1

Microsatellite loci isolated from *Abedus herberti*

Locus	Primer sequence (5'–3')	Dye	T _a (°C)	[p] (mM)	Repeat Motif	Size range	N	A	H _o	H _e	P
AHA002	F: GCGAGGGTGTCTCACTTGG R: GCCGATAGAGTCGTTGTCG	NED	52	0.20	di-	195–215	70	6	0.786	0.781	0.174
AHA003	F: GGGGGGCTCAAAAAATTGA R: AGATTGTAGTTGCCACACACA	FAM	52	0.20	di-	190–198	73	5	0.671	0.716	0.842
AHA03a	F: GTGTGGCAACTACAATCTG R: GCTCCAGGCTTATGCTAATC	FAM	52	0.25	di-	106–114	66	5	0.712	0.710	0.760
AHA005	F: TTTACTGGTCCGGATATAGAC R: CAAAATGAAGATAGGGGAAGAA	VIC	52	0.15	di-	147–217	71	11	0.775	0.809	0.189
AHA010	F: GTTCCGCATATAACAAACAATC R: AATCGAGCCCTACAAAGTTAATC	NED	52	0.30	di-	149–251	43	10	0.814	0.827	0.228
AHA102	F: AGACCCCTTCATCCACTACTACC R: CTACCCCTCTACACTCCAC	NED	52	–	di-	257–315	35	10	0.600	0.854	<0.001*
AHA103	F: ACGATTTCGGTGAGATAG R: AAAAGCCTCCCTTTGAAGTCC	NED	52	0.20	di-	197–241	44	15	0.886	0.879	0.135
AHA104	F: CCTTCATTGCGAATCCTC R: TTCCTTCATCCTCAGTAGTGC	FAM	52	–	di-	258–308	41	9	0.220	0.674	<0.001*
AHA106	F: GCATTATTGGCGTCTGCTA R: GCTATCTGTGCTGCTCTCATTC	FAM	52	0.25	di-	258–280	34	4	0.471	0.619	0.148
AHA111	F: TCGTCGTACAAATTCGCAAAAC R: CCCCCGTGTCCCACTTAA	VIC	52	–	di-	190–204	40	6	0.625	0.661	0.068
AHA112	F: GGGTTGCGAATGTTATCTC R: CGGGCTTTTCCTTATGTAG	PET	52	0.15	di-	148–176	72	10	0.806	0.837	0.498
AHA114	F: AATAATTTTCTCCCCCTATCA R: TTGGCTACTTGACGCATATAGA	PET	52	0.50	di-	251–327	58	15	0.707	0.793	0.239
AHB009	F: GCCTTCTGTATCGCCCAATAC R: TCAGGTAGAGGAAAAACATTGTG	VIC	52	0.15	tri-	231–264	75	7	0.360	0.362	0.202
AHB107	F: GCCCCATTATCCTCATAGTC R: GTGGCTGTGAAAGCGTAAC	VIC	52	0.05	tri-	228–240	77	5	0.701	0.719	0.566

Locus	Primer sequence (5' -3')	Dye	T _a (°C)	[p] (mM)	Repeat Motif	Size range	N	A	H _o	H _e	P
AHC011	F: GAGCCAGCAAGCGTAACA R: GCCGCAAGCAGTTCAGTA	PET	52	0.30	tet-	243-277	16	5	0.625	0.746	0.621
AHD003	F: CTCCTCCCTCTCACTTATTG R: CTCCTGTCGTATTGTTCT	PET	52	-	tet-	100-104	34	2	0.147	0.409	<0.001*
AHD006	F: CTCCTGAGGCTCTATTTTATCA R: GCCCAAAACAGATTGGAATC	PET	52	-	tet-	100-108	34	3	0.249	0.453	<0.001*

T_a annealing temperature, [p] primer concentration in multiplex pools, Repeat Motif di-, tri-, and tetra-nucleotide repeats, N number of individuals genotyped, A number of alleles, H_o observed heterozygosity, H_e expected heterozygosity, P P value for deviation from HW

* statistically significant