CORRESPONDENCE

Fentanyl for the Treatment of Tumor-Related Breakthrough Pain

by Dr. med. univ. et scient. med. Helmar Bornemann-Cimenti,
Dr. med. univ. Mischa Wejbora, Mag. rer. nat. Dr. scient. med. Istvan S. Szilagyi,
Prof. Dr. med. univ. Andreas Sandner-Kiesling in volume 16/2013

Lack of Precision in Colloquial Language Results in Incorrect Therapeutic Recommendation

In terms of numbers, the most common pain in tumor patients is incident pain, more rarely “end of dose” pain, as well as mixed forms and overlap phenomena. By contrast, genuine breakthrough pain—described in palliative medicine as an independent type—is rare. The authors define breakthrough pain correctly as a sudden-onset, transient exacerbation of generally well-controlled background pain (1). Additionally, we wish to mention that the frequency of genuine breakthrough pain cannot be influenced by increasing patients’ opiate-based medication.

In routine clinical practice, the term of breakthrough pain has unfortunately undergone a tendency to be used in a generalized and universal manner; often it is used for all additional pain states in tumor patients.

The pain exacerbation that occurs most often is incident pain, which primarily does not require therapy with rapidly effective fentanyl preparations. Affected patients can easily be informed instructed as to how they should deal with looming pain. The available short-acting morphine and hydromorphone preparations are appropriate and very effective.

The increased risk of opioid misuse and dependency should not be underestimated for the new, rapidly effective fentanyl preparations, especially if a patient has already shown signs of addictive behavior (alcohol, nicotine, etc). Patients receiving rapid-release fentanyl preparations are not fit to drive, which is in marked contrast to patients regularly receiving slow-release morphine.

It is not only for reasons of cost that the indication for “transmucosal” fentanyl should be strictly defined: to be used in “genuine” breakthrough pain and after therapeutic failure of conventional, fast-acting opiates.

DOI: 10.3238/arztebl.2013.0598a

REFERENCES

1. Bornemann-Cimenti H, Wejbora M, Szilagyi IS, Sandner-Kiesling A: Fentanyl for the treatment of tumor-related breakthrough pain. *Dtsch Arztebl Int* 2013; 110(16): 271–7.

Willi Ketterer

Palliativstation am DRK Krankenhaus Neuwied
willi.ketterer@drk-kh-neuwied.de

Conflict of interest statement

The author declares that no conflict of interest exists.

In Reply

Ketterer explained certain important aspects of our article (1) in an even more poignant and lucid way. He adds that the definition of breakthrough pain is often wrongly used in vernacular language. As early as in 2002, Mercadante et al discussed in a consensus paper from the European Association for Palliative Care that the English term “breakthrough pain” is used inconsistently and suggested using the term “episodic pain,” which, however, has not become widely accepted and adopted (2). Furthermore, they said that translation into different languages—as into German with the word “Durchbruchschmerz” [breakthrough pain]—only served to increase the semantic confusion.

With regard to predictable episodes of pain, such as the above-mentioned incident pain, we agree with our colleague’s explanations. Prevention, whose first step is surely providing the patient with information, is always the best therapeutic option. Where this is not enough, a differentiated use of medication options available should be selected, including that of non-fentanyl-preparations.

We actually did discuss the risk of addiction in our article (1). The scarce evidence should not be misinterpreted as a low risk. We would assess the topic of fitness to drive in a similar way. Even though currently no scientific studies of the topic exist, the conclusion should not be that the substances under discussion give no cause for concern in this respect.

We therefore also reach the conclusion that the use of rapid-onset fentanyl requires a strict and restricted indication and—as we said in our review article—needs to be handled responsibly by both doctors and patients (1).

DOI: 10.3238/arztebl.2013.0598b

REFERENCES

1. Bornemann-Cimenti H, Wejbora M, Szilagyi IS, Sandner-Kiesling A: Fentanyl for the treatment of tumor-related breakthrough pain. *Dtsch Arztebl Int* 2013; 110(16): 271–7.
2. Mercadante S, Radbruch L, Caraceni A, et al.: Episodic (breakthrough) pain: consensus conference of an expert working group of the European Association for Palliative Care. *Cancer* 2002; 94: 832–9.

Dr. med. univ. et scient. med. Helmar Bornemann-Cimenti

Dr. med. univ. Mischa Wejbora
Mag. rer. nat. Dr. scient. med. Istvan S. Szilagyi
Prof. Dr. med. univ. Andreas Sandner-Kiesling
Universitätsklinik für Anästhesiologie und Intensivmedizin
Medizinische Universität Graz
helmar.bornemann@medunigraz.at

Conflict of interest statement

Professor Sandner-Kiesling has had conference attendance fees and travel expenses reimbursed by Pfizer, Mundipharma, Cephalon, Grünenthal, and Fresenius. He has received lecture fees from Cephalon and Grünenthal. The other authors declare that no conflict of interest exists.