CORRESPONDENCE

Chronic Tinnitus: an Interdisciplinary Challenge

by Dr. med. Peter M. Kreuzer, Dr. med. Veronika Vielsmeier, PD Dr. med. Berthold Langguth in volume 16/2013

Remember Self-Help

The article (1) unfortunately omits any mention of self-help. The charitable self-help organization *Deutsche Tinnitus Liga* (DTL, the German tinnitus association) makes a valuable contribution to support affected persons in this context (2). However, the authors deserve credit for explaining the topic of tinnitus so comprehensively. At the same time it should be clear that we are mostly talking about management, not cure. For patients with severe tinnitus, even just reducing the burden is a realistic therapeutic objective.

Scientific evaluations have shown that of the 14 000 DTL members, 16% have grade 4 tinnitus (mini tinnitus questionnaire–12, *Mini-TF12*) (3). In order to confirm the effectiveness of self-help in qualified self-help groups, the DTL has started a large, prospective 3-year study in collaboration with the University Medical Center Hamburg-Eppendorf (2, 4).

In our experience there are better grading scales for the severity of tinnitus than the external assessment that the authors list in the *Box*. The DTL on its home page provides a tinnitus test for interested parties that is based on the scientifically evaluated mini-tinnitus questionnaire 12 and is being undertaken by some 20 000 affected persons every year. 16% of these are most severely affected (3).

The article mentions “tinnitus patients” in several places, sometimes for all those affected, but elsewhere for those who are actually suffering, which may well be termed “patients.” If suffering caused by tinnitus is the main symptom then tinnitus should not be subsumed under the symptom category H 93.1 in the ICD-10 but should be integrated as a disease under a new category that is yet to be established. The DTL—supported by the Association of the Scientific Medical Societies in Germany (AWMF)—has taken this up with the German Institute of Medical Documentation and Information (*Deutsches Institut für Medizinische Dokumentation und Information*, DIMDI) for the purposes of the ICD-10-GM 2014.

DOI: 10.3238/arztebl.2013.0599a

REFERENCES

1. Kreuzer PM, Vielsmeier V, Langguth B: Chronic tinnitus: an interdisciplinary challenge. *Dtsch Arztebl Int* 2013; 110(16): 278–84.
2. Goebel G: Was ist Tinnitus-Selbsthilfe und welche Bedeutung hat sie? In: Hesse G (ed.): *Tinnitus*. Stuttgart: Thieme 2008: 175–77.
3. Goebel G: Psychische Komorbidität bei Tinnitus. *Psychiatr Psychother* Up2date 2010; 4: 389–408.

4. Kofahl C, Werner S, Goebel G: Evaluation der Wirksamkeit von Selbsthilfegruppenarbeit in der Deutschen Tinnitus-Liga e. V. (DTL). *Tinnitus-Forum* 2012; 3: 26–7.

Volker Albert

Michael Bergmann

Deutsche Tinnitus-Liga e. V., Wuppertal
m.bergmann@tinnitus-liga.de

Conflict of interest statement

Mr Albert is the honorary chair of the Deutsche Tinnitus Liga e.V. (the German Tinnitus Association, reg assoc).

Mr Bergmann is the managing director of the Deutsche Tinnitus Liga e.V. (the German Tinnitus Association, reg assoc).

The Therapeutic Objective: Reducing the Stress Caused by Tinnitus

The article laudably provided an overview of the topic from an interdisciplinary perspective (1). One cannot but agree with the authors in objecting to the still prevalent therapeutic nihilism. At the same time it should be emphasized that management of tinnitus is the rule, rather than its actual cure. Especially in severely affected patients, the realistic therapeutic objective is a significant reduction in the stress caused by tinnitus.

Cognitive behavioral therapy (CBT) is the only genuinely evidence based therapy, but this goes almost unnoticed in the article owing to the very short explanation and cautious wording (“the best-evaluated treatment for tinnitus”).

In order to assess the indication for treatment (and therapeutic success) it is vital to determine the extent of the stress caused by tinnitus. Relevant validated questionnaires are not only available, as was mentioned almost as an aside, but they should be an integral component of the diagnostic evaluation. The first one to mention in this context is the tinnitus questionnaire ([2] or the mini-tinnitus questionnaire-12 [3]).

Psychologically oriented tinnitus therapy, which ranges from one-off counseling to multimodal tinnitus management therapy including CBT, should be used in a differentiated manner, depending on the stress caused by the tinnitus and the psychological comorbidity (4). Psychological comorbidity, which often also includes somatoform disorders, correlates with the severity of the tinnitus. Especially highly stressed tinnitus patients should undergo targeted screening for psychological disorders (3, 4). Psychological comorbidities are not only a consequence of tinnitus; rather, pre-existing mental disorders are often a form of adaptive processing of the tinnitus and as such constitute a crucial risk factor for its chronification and decompensation (4). Treating psychological comorbidities should not, as shown in *Figure 2*, precede (behavioral therapeutic) tinnitus therapy but be treated alongside the tinnitus.

DOI: 10.3238/arztebl.2013.0599b

REFERENCES

1. Kreuzer PM, Vielsmeier V, Langguth B: Chronic tinnitus: an interdisciplinary challenge. *Dtsch Arztebl Int* 2013; 110(16): 278–84.
2. Goebel G, Hiller W: Tinnitus-Fragebogen (TF). Ein Instrument zur Erfassung von Belastung und Schweregrad bei Tinnitus. Göttingen, Bern, Toronto, Seattle: Hogrefe Verlag für Psychologie 1998.
3. Hiller W, Goebel G: Rapid assessment of tinnitus-related psychological distress using the Mini-TQ. *International Journal of Audiology* 2004; 43: 600–4.
4. Goebel G: Psychische Komorbidität bei Tinnitus. *Psychiatr Psychother Up2date* 2010; 4: 389–408.

Dr. med. Ulrich Stappert

Prof. Dr. med. Gerhard Goebel

Prof. Dr. med. Ulrich Voderholzer

Schön Klinik Roseneck, Prien a. Chiemsee
grempert@schoen-kliniken.de

Conflict of interest statement

Professor Goebel has received royalties for the tinnitus questionnaire from Hogrefe publishers. The other two authors declare that no conflict of interest exists.

Research Is Needed

The authors are to be thanked for their valuable review article (1). We wish to underline the importance of continuous evolution of the therapeutic relationship, and point out the urgent need for qualitative research of complex disorders.

In all experience, the otherwise harmless condition that is tinnitus usually exacerbates when the affected person's own life is "internally inconsistent." A developmental step is imminent, and stress is a sign of exhausted coping strategies. Complex mental functions such as self-perception and world view, identity and self-consistency are affected; variables that are complex but can be researched (2).

The cited superior multiprofessional treatment (3) includes, as mentioned by the authors, psychotherapy and counseling, as well as sociotherapy, physical therapy, exercise therapy, and speech therapy, according to the complexity of the disorder and the patient. It is also our experience that in addition to verbal psychotherapy, it is interventions oriented towards the body and the senses that make multimodal therapy effective. The location of the symptom is the auditory system as the "hub," where the external world is transformed by the sensory stimulus into internal, psychological, experience. And it is there where, in case of increasing tension and discrepancy between the external and internal worlds, the tormenting "warning alarm signal" tinnitus is experienced.

It was surprising to read in the summary that, in addition to behavioral therapy, the main focus in research is on apparatus-based solutions. This gives rise to hopes for simple solutions, but it also implies a reductionist vision of humans and disease. In spite of causing constant impaired hearing, tinnitus fluctuates substantially depending on the subject's psychological state, something that many studies in (animal) models have not been able to reproduce.

The subject as the actual location of suffering, with dimensions such as development, maturity, self-image,

sense, authenticity, and consistency must not be lost. This is only indirectly accessible to drug and apparatus-based treatment, but it plays the crucial part in the sustained recovery from subjective tinnitus.

DOI: 10.3238/arztebl.2013.0600a

REFERENCES

1. Kreuzer PM, Vielsmeier V, Langguth B: Chronic tinnitus: an interdisciplinary challenge. *Dtsch Arztebl Int* 2013; 110(16): 278–84.
2. Fonagy P, Gergely G, Jurist E, Target M: Affektregulierung, Mentalisierung und die Entwicklung des Selbst. Stuttgart: Klett-Cotta 2002.
3. Cima RF, Maes IH, Joore MA, et al.: Specialised treatment based on cognitive behaviour therapy versus usual care for tinnitus: a randomised controlled trial. *Lancet* 2012; 379: 1951–9.

Dr. med. Tobias Roeckl

HNO-Klinik Dr. Gaertner, München

t.roeckl@gaertnerklinik.de

Conflict of interest statement

The author declares that no conflict of interest exists.

Psychiatrists Are not the First Port of Call

In their review article (1), our psychiatrist colleagues state that patients with tinnitus can be treated meaningfully. In this setting, the treatment approaches are similarly multifarious as the affected patients are different. In our long years of experience, the following are helpful:

- A fundamental neuro-otological competence that includes the concrete auditory perception
- An understanding of psychosomatic approaches
- And, above all else, empathy for the symptom and the affected person.

As a rule, disorder-specific treatment that is audiotically based and considers the possible causes as well as the sustaining factors—psychosomatic or somatopsychic—will be successful (2, 3). Psychiatric help may be indicated if relevant psychiatric disorders determine the tinnitus (3)—but psychiatrists are certainly not the first port of call in tinnitus therapy.

In cases where all outpatient therapies have been exhausted, large effect sizes have been found for inpatient psychotherapy, which is quite common in Germany—even though its efficacy, as the authors noted, has thus far not been evaluated in randomized controlled studies.

In contrast to studies of apparatus-based approaches or psychotropic drugs it is difficult to imagine how tinnitus patients who require treatment and have been admitted as inpatients would be divided into a psychotherapeutically treated group and a placebo group. On examining results from suitable psychosomatic hospitals—which are accessible to psychiatrists—with regard to diagnostic evaluation that is related to symptoms as well as psychological, one will find not only notable and sustained improvements of symptoms (4) but also therapeutically very effective reductions in the accompanying symptoms of anxiety and depression, with large effect sizes (3).

By contrast, therapeutic approaches (such as neuro-modulation) that promise suffering patients the complete extinction of their tinnitus with the help of medical apparatus, will mostly lead to disappointment, which in turn is also a serious adverse effect.

DOI: 10.3238/arztebl.2013.0600b

REFERENCES

1. Kreuzer PM, Vielsmeier V, Langguth B: Chronic tinnitus: an interdisciplinary challenge. *Dtsch Arztebl Int* 2013; 110(16): 278–84.
2. Hesse G: Die Neurootologisch psychosomatische Habituationstherapie. *HNO* 2009; 56: 686–93.
3. Schaaf H, Eichenberg C, Hesse G: Tinnitus und das Leiden am Tinnitus. Ein Plädoyer für eine störungsspezifische Betrachtungsweise und ein differenziertes Vorgehen anhand 287 ambulanter und 128 stationär behandelter Tinnitus Patienten. *Der Psychotherapeut* 2010; 55: 225–32.
4. Goebel G, Kahl M, Arnold W, Fichter M: 15-year prospective follow-up study of behavioral therapy in a large sample of inpatients with chronic tinnitus. *Acta Otolaryngol Suppl* 2006; 556: 70–9.

Dr. med. Helmut Schaaf

Prof. Dr. med. habil. Gerhard Hesse

Tinnitus-Klinik Dr. Hesse und Gleichgewichtssambuanz der Klinik Dr. Hesse im Stadtkrankenhaus Bad Arolsen
hschaaf@tinnitus-klinik.net

Conflict of interest statement

The authors declare that no conflict of interest exists.

In Reply:

We are pleased to note how much of the correspondence in response to our article emphasized the importance of forging a continuous therapeutic relationship and using an interdisciplinary treatment approach (1), including the willingness to mutually look beyond one's own perspective. We wish to comment briefly on some of the criticisms:

The representatives from the Deutsche Tinnitus Liga (German tinnitus association) and those from a specialized psychosomatic hospital (Dr Stattrop, Professor Goebel, and Professor Voderholzer) correctly remind us, that the currently available therapeutic options are mainly dealing with tinnitus management, not cure. We agree with this statement, but we think that the availability of treatment options for tinnitus management should not prevent us from optimizing these approaches further and from working on techniques aiming for a cure.

We also wish to focus more closely on the statement that cognitive behavioral therapy is the only “genuinely” evidence based therapy. In our article we presented, for all therapeutic approaches, the available evidence according to generally used criteria. In our opinion, the argument that certain interventions cannot be studied in randomized controlled trials is as unrewarding as the wholesale rejection of apparatus-based approaches. It should also be borne in mind that the questionnaires commonly used in clinical studies are suitable for reflecting the degree of stress due to tinnitus over the course of the treatment, but that the

perception of the tinnitus is not incorporated in any way. In the reality of clinical care, many of those affected desire not only help in managing their tinnitus but also an actual reduction in their impression of the noise. We leave it to our readers to decide what counts as “only genuinely evidence based” on this background.

Regarding the categorization of tinnitus according to Biesinger, criticized by Albert and Bergmann in their reader's letter: we aimed to provide general practitioners and otorhinolaryngologists working in Germany's statutory health insurance system with a pragmatic decision aid, to enable them to distinguish quickly between those affected and those more severely impaired by tinnitus. In our experience, the patient's own comments in the personal encounter are much more reliable for a clinical assessment than the total score in a self-assessment questionnaire, which may result in a distorted picture. This in no way diminishes the importance of questionnaires for scientific purposes and for quality assurance. Because of space limitations we were not able to discuss all the different available self-rating scales.

Space was also the primary reason for not being able to discuss self-help services in any detail. A recent review article (published in July 2013) (2), which was not yet available at the time our article went to print, found that specific self-help interventions offer significant benefits compared with mere information provision or discussion groups.

With regard to the comments of Dr Schaaf and Professor Hesse we wish to add that we did not postulate any categorical distinction between psychiatric and psychosomatic approaches anywhere in our article. Without wanting to go into detail about health political contexts, we are of the opinion that the therapist's individual experience and willingness to enter into an interdisciplinary dialogue provide the key to successful treatment. Usually a general practitioner or otorhinolaryngologist would be the first port of call for persons affected by tinnitus, not a psychiatrist or psychosomatic specialist (3). In contrast to what Stattrop et al. say in their letter, we are convinced that counseling should not be subsumed as an advisory and information-providing talk under “psychologically oriented tinnitus therapy” but as an integral component of any tinnitus therapy. It was our central aim to provide doctors working within the statutory health insurance system with an overview of the currently available, evidence-based therapeutic options as an orientation for clinical decision making—in addition to emphasizing the reasonableness of an interdisciplinary therapeutic approach.

DOI: 10.3238/arztebl.2013.0601

REFERENCES

1. Cima RF, Maes IH, Joore MA, et al.: Specialised treatment based on cognitive behaviour therapy versus usual care for tinnitus: a randomised controlled trial. *Lancet* 2012; 379: 1951–9.
2. Nyenhuis N, Golm D, Kroner-Herwig B: A systematic review and meta-analysis on the efficacy of self-help interventions in tinnitus. *Cogn Behav Ther* 2013; 42: 159–69.

3. Hall DA, Lainez MJ, Newman CW, et al.: Treatment options for subjective tinnitus: self reports from a sample of general practitioners and ENT physicians within Europe and the USA. BMC Health Serv Res 2011; 11: 302.
4. Kreuzer PM, Vielsmeier V, Langguth B: Chronic tinnitus: an interdisciplinary challenge. Dtsch Arztebl Int 2013; 110(16): 278–84.

Dr. med. Peter M. Kreuzer
Dr. med. Veronika Vielsmeier
PD Dr. med. Berthold Langguth
 Universität Regensburg
 peter.kreuzer@medbo.de

Conflict of interest statement

Dr. Kreuzer has received third-party funding for study projects from Cerbomed GmbH. He has received reimbursement of congress attendance fees and travel costs from the Tinnitus Research Initiative. In addition, travel and training costs have been paid on his behalf by Servier, Pfizer, AstraZeneca, Lilly, Bristol Myers Squibb, and Lundbeck.

Dr. Vielsmeier has received reimbursement of congress attendance fees and travel costs from the Tinnitus Research Initiative.

Dr. Langguth has received consulting fees from ANM, Autifony, Merz, Novartis, and Sanofi, and lecture fees from Merz and ANM. He holds patents for the method of neuronavigational positioning of the TMS coil for treatment of tinnitus and of treatment with cyclobenzaprine and naltrexone. Furthermore, he has received revenue from sales of the "Textbook of Tinnitus".