Published in final edited form as:

Biochemistry. 2013 August 13; 52(32): 5415–5420. doi:10.1021/bi400790j.

Phosphatidylinositol is crucial for the mechanosensitivity of *Mycobacterium tuberculosis* MscL

Dalian Zhong¹ and Paul Blount^{1,*}

¹From the Department of Physiology, UT Southwestern Medical center at Dallas, 5323 Harry Hines Blvd, Dallas TX 75390-9040

Abstract

The bacterial mechanosensitive channel of large conductance (MscL) directly senses and responds to membrane tension. It serves as an “emergency release valve” upon acute decreases in the osmotic environment, thus preventing cell lysis. It is one of the best studied mechanosensitive channels and serves as a paradigm of how a channel senses and responds to its membrane environment. The MscL protein is highly conserved, found throughout the bacterial kingdom, and has been shown to encode a functional mechanosensitive channel in all species where it has been studied. However, channels from different species have shown some functional variance; an extreme example is the *Mycobacterium tuberculosis* MscL, which when heterologously expressed in *Escherichia coli* requires significantly more membrane tension for gating than the endogenous *E. coli* MscL. We previously speculated that the membrane environment or factors not found in *E. coli* promoted the proper gating of the *M. tuberculosis* MscL channel in its native environment. Here, by reconstituting the *M. tuberculosis* and *E. coli* MscL channels in various lipids, we demonstrate that inclusion of phosphatidylinositol, a lipid found in *M. tuberculosis* but not *E. coli*, is sufficient for gating of the *M. tuberculosis* MscL channel within a physiological range of membrane tension.

MscL is one of the best studied mechanosensitive channels from any species^{1–4}. It was initially isolated from *E. coli*, where many of the initial studies were performed. However, it was a homologue, *Mycobacterium tuberculosis* MscL (Mt-MscL), which was found to be amenable to crystallization. The crystal structure of Mt-MscL, solved at 3.6 Å, revealed that the protein forms a homopentameric complex (Figure 1 A, right two panels)⁵; although there have been controversies on the oligomeric structure, specifically for a *Staphylococcus aureus* orthologue⁶, the pentameric nature of the complex *in vivo* has been recently confirmed^{7, 8}. Each subunit of the complex contains an N-terminal helix that runs along the membrane, two transmembrane domains, TM1 and TM2, which are connected by a periplasmic loop, and a C-terminal helix that form a helical bundle from the five subunits^{5, 9, 10} (Figure. 1A, left panel). Several studies have led to proposals for the function of each domain, as well as a cohesive model for conformational changes that occur upon channel gating^{2, 4, 11}. The function of the protein appears to be preserved among many bacterial species; a previous study had found that the *mscL* genes from a variety of bacterial species encoded a mechanosensitive channel activity when expressed in *E. coli*; however, Mt-MscL was not among the initial orthologues studied¹². Thus, after the crystal structure had been published, we sought to determine if the Mt-MscL was functional, i.e., could we definitively state that Mt-MscL was an orthologue. We found that although MscL-like

*Corresponding author: Paul Blount, Department of Physiology, UT Southwestern Medical center at Dallas, 5323 Harry Hines Blvd, Dallas, TX, USA, Tel: (214) 645-6014; Paul.Blount@utsouthwestern.edu.

The authors declare no competing financial interest.

channel activities could be recorded, significantly more membrane tension was required to gate the channel when expressed heterologously in *E. coli*¹³. In addition, unlike other orthologues, wild type Mt-MscL was unable to rescue MscS/MscL double null *E. coli* cells from an acute osmotic down shock¹³. These properties of the Mt-MscL channel make the acquisition of recordings extremely difficult; thus, in spite of the advantage of an available crystal structure, most MscL studies have continued to be performed on the *E. coli* MscL (Ec-MscL) orthologue.

It is well accepted that specific lipids can play an important role in the function of membrane proteins, including ion channels. For example, anionic lipid components are crucial to channel function of the bacterial inward rectifier potassium channel KcsA¹⁴. In the original functional characterization of Mt-MscL, we postulated that there was an environmental factor missing from *E. coli* that would allow normal gating in its native environment¹³. It seems reasonable that a membrane lipid component could be this factor. The inner membrane of *E. coli* is composed of 75–85% of phosphatidylethanolamine (PE), 10–20% of the anionic phosphatidylglycerol (PG) and about 5–15% of cardiolipin (CL)¹⁵. However, while the *M. tuberculosis* membrane is also dominated by PE lipids and contains a small percentage of CL, the second major component is the anionic lipid phosphatidylinositol (PI)^{16, 17}, which can account for up to a quarter of total phospholipids in *Mycobacteria tuberculosis* membranes¹⁶. Hence, the major difference in the lipid composition is the primary anionic lipid. Here we have studied the effects of different lipid components on the Ec- and Mt-MscL channels and found that inclusion of PI is sufficient for normal Mt-channel function.

MATERIALS AND METHODS

Strains and protein expression-Wild type *E. coli* MscL and *M. tuberculosis* MscL with C-terminal six-his-tags were inserted into pET21a expression construct and transformed into *E. coli* strain PB116 cells for protein expression^{18, 19}. Wild type *E. coli* MscS with a six-his-tag was subcloned into pB10b expression vector and expressed in PB104 (mscL:Cm) cells^{18, 19}. Cultures were grown routinely in Lennox broth medium (Fisher Scientific, Pittsburg, PA, USA) containing 100 ug/ml ampicillin. Expression was induced by addition of 1 mM IPTG at 37°C. Cells were harvested after 3 hours of induction.

Protein purification- Protein purification was performed as previously described²⁰. Briefly, cells were resuspended in base buffer containing 10 mM Potassium Phosphate Buffer at pH 8.0 (KPi), 300 mM NaCl, plus 0.5 ug/ml Dnase, 1 mg/ml Lysozyme, 1 tablet protease inhibitor cocktail (Roche, Indianapolis, IN, USA), and subjected to two passages through a French pressure cell at 16,000 psi to complete the lysis at 4°C. The lysate was subjected to extraction by addition of 15 mM Imidazole and 2% Triton. MscLH6 or MscSH6 was purified from the membrane by using Ni-NTA resin (Thermo Scientific, Rockford, IL, USA). The Ni-NTA was added to the cleared lysate and incubated at room temperature for 1 hour. Then the resin was washed once with extraction buffer (10 mM KPi, pH 8.0, 300 mM NaCl, 15 mM Imidazole, and 2% Triton), twice with wash buffer (10 mM KPi, pH 8.0, 300 mM NaCl, 20 mM Imidazole, and 1% Triton) and transferred to a column. The column was washed with 30 ml wash buffer and eluted with elution buffer (10 mM KPi, pH 8.0, 300 mM NaCl, 250 mM Imidazole, and 0.2% Triton). The elution fractions were subjected to SDS-PAGE and protein was pooled for concentrating and BCA assay (Pierce, Rockford, IL, USA).

Co-reconstitution- Lipid vesicles for reconstitution were prepared as previously described²⁰. Lipids, Azolectin, also known as Soy Total Lipid Extract (Soy PC 20% Cat# 541601), cardiolipin (CL), 1,2-dioleoyl-*sn*-glycero-3-XX (DOXX), where XX is one of the following

headgroups: phosphatidylcholine (PC), phosphatidylglycerol (PG), phosphatidylinositol (PI), or phosphatidylethanolamine (PE), were utilized (Avanti polar lipids, Inc, Alabaster, AL, USA). Briefly, lipids were dissolved in chloroform, proper amount of different lipids were added to a glass tube, mixed and dried under argon. The lipid film was further dried under vacuum for another 2 hours and resuspended to 20 mg/ml buffer (10 mM Tris, pH 7.5, 1mM EDTA, and 1mM EGTA) and rehydrated at 45°C for at least 2 hours. The lipid vesicles were formed by bath sonication. MscLH6 (Mt-MscLH6 or Ec-MscLH6) and MscSH6 proteins were combined with lipid vesicles to a 1:500 protein-to-lipid mass ratio. Detergent was removed by dialysis with 3x 1 000 ml of buffer (5 mM Tris, pH 7.5, 100 mM NaCl, 0.2 mM EDTA, and 0.02% NaN₃) containing Bio-beads (Bio-Rad, Hercules, CA, USA). The resultant liposomes were collected by a 20-minute centrifugation at 30 psi in an airfuge. The pellet was resuspended to 1 mg/ 5ul in buffer (10 mM MOPS, pH 7.5, 5% ethylene glycerol), and desiccated overnight under vacuum at 4 °C.

Electrophysiology- Desiccated liposomes were rehydrated in buffer (0.25 mM Hepes, pH 7.5, 10 mM KCl, and 2 mM MgCl₂ plus 320 mM sucrose) at a lipid concentration of 90 mg/ml as described previously²¹. Excised patches were studied at room temperature under symmetrical patch buffer conditions (5mM Hepes, pH 7.5, 200 mM KCl, 40 mM MgCl₂). Data were acquired at -20 mV at a sampling rate of 20 kHz filter, using an AxoPatch 200B amplifier in conjunction with Axoscope software (Axon Instruments, Union City, CA, USA). A piezoelectric pressure transducer (World Precision Instruments, Sarasota, FL, USA) was used to monitor the pressure introduced to membrane by suction throughout the experiments. For neutral liposomes, the presence of 200 mM KCl inhibited the formation of unilamellar blisters; thus patch buffer containing no KCl was used in the chamber, and KCl was added to a final concentration to 200 mM after the seal formation. The channels were gated by applying negative pressure to the patch. Trace acquisition and analyses were performed using Clamfit10 from pCLAMP10 (Axon instruments).

Mechanosensitivity was determined as previously described^{18, 22}. Briefly, MscS is used as an internal standard within the patch, the ratio of gating pressure for MscL divided by the pressure of MscS, expressing as pL/pS ratio, is determined. For neutral liposomes, the channel threshold pressure was used instead of pL/pS ratio, due to the reversal of MscS/ MscL sensitivities, which made analysis impractical. All data were obtained from at least three independent recordings performed in at least two independent protein purifications and reconstitutions.

Statistics- Data were analyzed for statistical significance using the Student's *t* test.

RESULTS

We sought to determine the functional properties of the Ec- and Mt-MscL channels in different anionic lipid environments. We have previously utilized another mechanosensitive channel endogenously expressed in *E. coli*, MscS (mechanosensitive channel of small conductance), as an internal standard control^{18, 22}. This allows for a more accurate determination of differences in sensitivity to membrane tension, or for simplicity what we will call mechanosensitivity. This approach is easily performed when the channels are expressed in giant cells derived from *E. coli*. The mechanosensitivity of the channels are then expressed as the pressure threshold of MscL gating divided by pressure threshold of MscS (pL/pS). This convention was utilized in our previous study, where we found that the Mt-MscL required a much greater pL/pS ratio (2.8) relative to the Ec-MscL channel (pL/pS of 1.4) when heterologously expressed in giant *E. coli* cells¹³; or put another way, the Mt-MscL channel was far less sensitive to membrane tension when expressed in *E. coli* native membranes.

We wanted to use MscS as a standard in reconstitution studies as reported by Battle and coworkers²³, where we tested the effect of different lipid compositions on channel function. Thus, we co-reconstituted and assayed by patch clamp purified MscL and MscS proteins into liposomes composed of synthetic lipids that would largely reproduce either the *E. coli* inner membrane (85%DOPE: 13%DOPG: 2%CL) or that of *M. tuberculosis* (85%DOPE: 13%DOPI: 2%CL). Figure 2A shows typical current traces of the channels with the pressure threshold (mmHg) under each trace. Figure 2B summarizes the mechanosensitivity threshold (pL/pS) of Mt-MscL and Ec-MscL in the different lipid conditions. Consistent with our results from the giant *E. coli* cells, in the PG containing liposomes the activation threshold of Mt-MscL was significantly higher than the threshold required for Ec-MscL channel activation. In contrast, in PI containing *M. tuberculosis*-like (Mt-like) lipids, the Mt-MscL became more mechanosensitive, with a ratio approaching that observed for the Ec-MscL, thus indicating the mechanosensitivity of Mt-MscL channel is within the normal range when 13% PI is present.

To further verify this finding, we co-reconstituted MscS and Mt-MscL in azolectin lipids, which contain approximately 12% PI. Consistent with the results from lipids containing PI, we found that Mt-MscL was more mechanosensitive in azolectin when compared to PG containing lipids. These results support the observation that the presence of PI is sufficient for an increased mechanosensitivity of Mt-MscL. While it appears that in both cases PI greatly enhances the mechanosensitivity of Mt-MscL, it does not quite fully achieve the mechanosensitivity of Ec-MscL; this could indicate yet another subtle factor within the native environment of *M. tuberculosis* that is missing, or simply be a slight species variance.

Even though a fully open state of Mt-MscL was observed on many occasions when reconstituted in synthetic lipid membranes containing anionic lipids, the channel activities appeared heterogeneous. In a subset of patches, a channel showed a behavior in which it obtained very short open dwell times, beyond equipment and computational resolution. Figure 2C shows a typical trace with such channel behavior in synthetic lipids containing PI. The mechanisms underlying this heterogeneity are currently unknown. Nevertheless, channels showing different kinetic properties had no difference in their mechanosensitivity.

The data suggest that PI interacts with the Mt-MscL to allow it to function at more physiological levels of membrane tension. However, from the above data we could not formally rule out an alternative possibility: that PG inhibits Mt-MscL channel gating. To test this, we removed all negatively charged lipid headgroups and reconstituted the channels into neutral lipids containing 50% DOPE and 50% DOPC. Unfortunately, both MscL channels opened at tensions lower than MscS, making it difficult to obtain an accurate threshold ratio. Therefore, we were confined to assaying absolute pressure rather than threshold ratios. The results, however, were clear. Figure 3A shows typical traces of Mt-MscL and Ec-MscL in 50% DOPE: 50% DOPC liposomes. As shown in Figure. 3B, we still needed over twice the pressure to activate Mt-MscL relative to Ec-MscL. Thus, it appears that PG within the *E. coli* membrane does not inhibit Mt-MscL channel function; instead, PI increases the sensitivity of the Mt-MscL channel.

DISCUSSION

While it has been known for a number of years that the Mt-MscL was not functional when expressed in *E. coli* cells, the underlying reasons were not clear. Here we find that it is because PI, a lipid not expressed in *E. coli* is crucial for its normal function. Our results largely explain the data obtained in the original characterization of Mt-MscL, except for one observation. In the previous study, the Mt-MscL channel activity was characterized in several systems including azolectin lipids, which contains approximately 12% PI; in contrast

to the present study, the previous results suggested that the Mt-MscL required more stimuli to gate¹³. However, there are complications to the interpretation of the previous study. The reconstituted protein was N-terminally multi-his tagged, and modification at this end of the protein has been shown to disrupt channel function^{22,24,25}; here we have used a C-terminally tagged protein. In addition, the experiments could not be well controlled because at the time MscS had not yet been observed in a reconstitution system and was therefore not used as an internal control. Hence, in this study we repeated this experiment and utilized our co-reconstitution approach to allow us to better interpret the results. The results we report here support the observation that the presence of PI is crucial and sufficient for an increased mechanosensitivity of Mt-MscL.

Lipids, especially negatively charged lipids, have been shown to modulate many different channel types^{26–28}. However, this has not been true for MscL. Although one study suggested anionic lipids helped in the reconstitution of functional Ec-MscL channels²⁹, this was performed using flux assays; the best evidence using patch clamp as an assay suggests anionic lipids play little if any role in normal Ec-MscL function²⁰ (also see¹¹ for a detailed discussion). The relative lack of influence of lipid composition on Ec-MscL channel function seemed surprising, given the function of the protein: directly sensing biophysical changes in the lipid membrane. Thus, the finding that PI is sufficient for normal Mt-MscL activity demonstrates not only that lipids can influence MscL channel activity, it also indicates that lipid interactions may be dependent upon the orthologue being studied.

The mechanism for how PI influences the gating of Mt-MscL would be of prime interest. There have been mutations that influence mechanosensitivity, for instance E104Q and V15C gate at lower tension than wild type Mt-MscL³⁰; however, these studies were performed in *E. coli* spheroplasts and therefore do not reflect mutations that influence the protein interactions with PI. On another front, Powl and co-workers have identified the C-terminal charged cluster, specifically Arg-98, Lys-99, and Lys-100, as a hot spot for anionic lipid binding in Mt-MscL³¹; here again, however, the studies were performed in a reconstitution system where the liposomes did not contain PI. Hence, the mechanisms of how and where the Mt-MscL channel interacts specifically with the PI headgroup remains a mystery that will require further study.

One of the surprising findings was that MscL opened at lower tension than MscS when co-reconstituted in neutral lipids. At first glance, this seems to contradict observations by Momura and co-workers, who reported the activation ratios of pL/pS in 70% PE: 30% PC liposome about 1.7, indicating that MscL opened at a higher pressure threshold than MscS³². However, the lipid composition used here is slightly different and is composed of less PE, having a ratio of 50% PE: 50% PC. It is currently unclear why this reversal occurs; however, previous studies have shown localization of MscS at the cell poles was CL-dependent, thus CL may influence the function of MscS³³. Another possibility is that the increased PE plays a role; it has been shown that PE makes MscL less sensitive to membrane tension, presumably because of the small headgroup size^{20,34}; thus, it is possible that this is exaggerated for MscS, causing the observed reversal in channel opening. Regardless, the pressure required for Mt-MscL to open was significantly higher than that for Ec-MscL in neutral lipids. Our results demonstrate that PG doesn't inhibit Mt-MscL channel gating.

Together the data strongly suggest that Mt-MscL is indeed an orthologue, serving the function of an osmotic emergency release valve when expressed in its native environment. The reason for its apparent inability to function at physiological membrane tensions in previous studies was because PI, which is expressed in the *M. tuberculosis* but not *E. coli* membranes, increases the sensitivity of the Mt-MscL channel so it is within a physiological range.

Acknowledgments

The work was supported by Grant I-1420 of the Welch Foundation, Grant RP100146 from the Cancer Prevention & Research Institute of Texas (CPRIT; <http://www.cprit.state.tx.us/>), and Grant GM061028 from the National Institutes of Health. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

The authors thank Christina Eaton and Robin Wray for technical assistance, Dr. Limin Yang for her helpful advice, and Christina Eaton and Drs. Irene Iscla, Limin Yang and Hannah Malcolm for useful discussions and critical reading of the manuscript.

ABBREVIATIONS

MscL	mechanosensitive channel of large conductance
MscS	mechanosensitive channel of small conductance
Ec-MscL	<i>Escherichia coli</i> MscL
Mt-MscL	<i>Mycobacterium tuberculosis</i> MscL
Ec-like	<i>Escherichia coli</i> -like
Mt-like	<i>Mycobacterium tuberculosis</i> -like
CL	cardiolipin
DOPC	1,2-dioleoyl- <i>sn</i> -glycero-3-phosphatidylcholine
DOPG	1,2-dioleoyl- <i>sn</i> -glycero-3-phosphatidylglycerol
DOPI	1,2-dioleoyl- <i>sn</i> -glycero-3-phosphatidylinositol
DOPE	1,2-dioleoyl- <i>sn</i> -glycero-3-phosphatidylethanolamine

References

1. Blount, P.; Iscla, I.; Li, Y. Mechanosensitive channels and sensing osmotic stimuli in bacteria. In: Martinac, B., editor. Sensing with ion channels. Springer-Verlag Press; Berlin: 2008. p. 25-47.
2. Blount, P.; Iscla, I.; Moe, PC.; Li, Y. MscL: The bacterial mechanosensitive channel of large conductance. In: Hamill, OP., editor. Mechanosensitive Ion Channels (a volume in the Current Topics in Membranes series). Elsevier Press; St. Louis, MO: 2007. p. 202-233.
3. Blount, P.; Li, Y.; Moe, PC.; Iscla, I. Mechanosensitive channels gated by membrane tension: Bacteria and beyond. In: Kamkin, A.; Kiseleva, I., editors. Mechanosensitive ion channels (a volume in the Mechanosensitivity in Cells and Tissues, Moscow Academia series). Springer Press; New York: 2008. p. 71-101.
4. Booth IR, Blount P. Microbial Emergency Release Valves: the MscS and MscL Families of Mechanosensitive Channels. J Bacteriol. 2012; 194:4802–4809. [PubMed: 22685280]
5. Chang G, Spencer RH, Lee AT, Barclay MT, Rees DC. Structure of the MscL homolog from *Mycobacterium tuberculosis*: A gated mechanosensitive ion channel. Science. 1998; 282:2220–2226. [PubMed: 9856938]
6. Liu Z, Gandhi CS, Rees DC. Structure of a tetrameric MscL in an expanded intermediate state. Nature. 2009; 461:120–124. [PubMed: 19701184]
7. Dorwart MR, Wray R, Brautigam CA, Jiang Y, Blount P. S. aureus MscL is a pentamer in vivo but of variable stoichiometries in vitro: implications for detergent-solubilized membrane proteins. PLoS Biol. 2010; 8:e1000555. [PubMed: 21151884]
8. Iscla I, Wray R, Blount P. The oligomeric state of the truncated mechanosensitive channel of large conductance shows no variance in vivo. Protein Sci. 2011; 20:1638–1642. [PubMed: 21739498]
9. Yang LM, Wray R, Parker J, Wilson D, Duran RS, Blount P. Three Routes To Modulate the Pore Size of the MscL Channel/Nanovalve. ACS Nano. 2012; 6:1134–1141. [PubMed: 22206349]

10. Steinbacher, S.; Bass, R.; Strop, P.; Rees, DC. Structures of the prokaryotic mechanosensitive channels MscL and MscS. In: Hamill, OP., editor. *Mechanosensitive Ion Channels* (a volume in the *Current Topics in Membranes* series). Elsevier Press; St. Louis, MO: 2007. p. 1-20.
11. Iscla I, Blount P. Sensing and Responding to Membrane Tension: The Bacterial MscL Channel as a Model System. *Biophys J*. 2012; 103:169–174. [PubMed: 22853893]
12. Moe PC, Blount P, Kung C. Functional and structural conservation in the mechanosensitive channel MscL implicates elements crucial for mechanosensation. *Mol Microbiol*. 1998; 28:583–592. [PubMed: 9632260]
13. Moe PC, Levin G, Blount P. Correlating a protein structure with function of a bacterial mechanosensitive channel. *J Biol Chem*. 2000; 275:31121–31127. [PubMed: 10846181]
14. Iwamoto M, Oiki S. Amphipathic antenna of an inward rectifier K⁺ channel responds to changes in the inner membrane leaflet. *Proc Natl Acad Sci U S A*. 2013; 110:749–754. [PubMed: 23267068]
15. Raetz CR. Enzymology, genetics, and regulation of membrane phospholipid synthesis in *Escherichia coli*. *Microbiol Rev*. 1978; 42:614–659. [PubMed: 362151]
16. Jackson M, Crick DC, Brennan PJ. Phosphatidylinositol is an essential phospholipid of mycobacteria. *J Biol Chem*. 2000; 275:30092–30099. [PubMed: 10889206]
17. Haites RE, Morita YS, McConville MJ, Billman-Jacobe H. Function of phosphatidylinositol in mycobacteria. *J Biol Chem*. 2005; 280:10981–10987. [PubMed: 15634688]
18. Blount P, Sukharev SI, Moe PC, Schroeder MJ, Guy HR, Kung C. Membrane topology and multimeric structure of a mechanosensitive channel protein of *Escherichia coli*. *EMBO J*. 1996; 15:4798–4805. [PubMed: 8890153]
19. Ou X, Blount P, Hoffman RJ, Kung C. One face of a transmembrane helix is crucial in mechanosensitive channel gating. *Proc Natl Acad Sci U S A*. 1998; 95:11471–11475. [PubMed: 9736761]
20. Moe P, Blount P. Assessment of potential stimuli for mechano-dependent gating of MscL: effects of pressure, tension, and lipid headgroups. *Biochemistry*. 2005; 44:12239–12244. [PubMed: 16142922]
21. Folgering JH, Kuiper JM, de Vries AH, Engberts JB, Poolman B. Lipid-mediated light activation of a mechanosensitive channel of large conductance. *Langmuir*. 2004; 20:6985–6987. [PubMed: 15301476]
22. Blount P, Sukharev SI, Schroeder MJ, Nagle SK, Kung C. Single residue substitutions that change the gating properties of a mechanosensitive channel in *Escherichia coli*. *Proc Natl Acad Sci U S A*. 1996; 93:11652–11657. [PubMed: 8876191]
23. Battle AR, Petrov E, Pal P, Martinac B. Rapid and improved reconstitution of bacterial mechanosensitive ion channel proteins MscS and MscL into liposomes using a modified sucrose method. *FEBS Lett*. 2009; 583:407–412. [PubMed: 19111548]
24. Iscla I, Wray R, Blount P. On the structure of the N-terminal domain of the MscL channel: helical bundle or membrane interface. *Biophys J*. 2008; 95:2283–2291. [PubMed: 18515388]
25. Häse CC, Ledain AC, Martinac B. Molecular dissection of the large mechanosensitive ion channel (MscL) of *E. coli* - mutants with altered channel gating and pressure sensitivity. *J Membr Biol*. 1997; 157:17–25. [PubMed: 9141355]
26. D'Avanzo N, McCusker EC, Powl AM, Miles AJ, Nichols CG, Wallace BA. Differential lipid dependence of the function of bacterial sodium channels. *PLoS One*. 2013; 8:e61216. [PubMed: 23579615]
27. Roberts-Crowley ML, Mitra-Ganguli T, Liu L, Rittenhouse AR. Regulation of voltage-gated Ca²⁺ channels by lipids. *Cell Calcium*. 2009; 45:589–601. [PubMed: 19419761]
28. Valiyaveetil F, Hermolin J, Fillingame RH. pH dependent inactivation of solubilized F1F0 ATP synthase by dicyclohexylcarbodiimide: pK(a) of detergent unmasked aspartyl-61 in *Escherichia coli* subunit c. *Biochimica et biophysica acta*. 2002; 1553:296–301. [PubMed: 11997138]
29. Powl AM, East JM, Lee AG. Anionic phospholipids affect the rate and extent of flux through the mechanosensitive channel of large conductance MscL. *Biochemistry*. 2008; 47:4317–4328. [PubMed: 18341289]

30. Shapovalov G, Bass R, Rees DC, Lester HA. Open-state disulfide crosslinking between *Mycobacterium tuberculosis* mechanosensitive channel subunits. *Biophys J*. 2003; 84:2357–2365. [PubMed: 12668444]
31. Powl AM, East JM, Lee AG. Heterogeneity in the Binding of Lipid Molecules to the Surface of a Membrane Protein: Hot Spots for Anionic Lipids on the Mechanosensitive Channel of Large Conductance MscL and Effects on Conformation. *Biochemistry*. 2005; 44:5873–5883. [PubMed: 15823046]
32. Nomura T, Cranfield CG, Deplazes E, Owen DM, Macmillan A, Battle AR, Constantine M, Sokabe M, Martinac B. Differential effects of lipids and lyso-lipids on the mechanosensitivity of the mechanosensitive channels MscL and MscS. *Proc Natl Acad Sci U S A*. 2012; 109:8770–8775. [PubMed: 22586095]
33. Romantsov T, Battle AR, Hendel JL, Martinac B, Wood JM. Protein localization in *Escherichia coli* cells: comparison of the cytoplasmic membrane proteins ProP, LacY, ProW, AqpZ, MscS, and MscL. *J Bacteriol*. 2010; 192:912–924. [PubMed: 20008071]
34. Perozo E, Kloda A, Cortes DM, Martinac B. Physical principles underlying the transduction of bilayer deformation forces during mechanosensitive channel gating. *Nat Struct Biol*. 2002; 9:696–703. [PubMed: 12172537]

FIGURE 1.

The crystal structure of Mt-MscL and alignment of MscL homologs. A. Crystal structure of Mt-MscL, a homopentamer with each subunit containing two transmembrane domains (TM1 and TM2), an N-terminal S1-helix, C-terminal helix and linkers between helices. B. Amino acid sequence alignment of MscL homologs from *M. tuberculosis* (Mt-MscL) and *E. coli* (Ec-MscL). Identical and similar residues are shown in boxes. The functional regions of MscL protein, the S1-helix, TM1, Periplasmic loop, TM2, and C-terminal helix are shown above the sequences.

FIGURE 2.

Mechanosensitivity of Mt-MscL and Ec-MscL in different membrane lipid conditions. A. Patch clamp recording of Mt-MscL and Ec-MscL in *E. coli*-like (Ec-like) lipids and Mt-like lipids. Upper and lower rows show the typical current traces of Mt-MscL (left) and Ec-MscL (right) along with the Ec-MscS in PG containing Ec-like lipids (85%DOPE:13% DOPG: 2%CL) and PI containing Mt-like lipids (85%DOPE:13%DOPI:2% CL), respectively. Pressure thresholds are shown under each trace. “O” and “C” refer to channel open and closed states. The scale bar is shown at the top right. B. Pressure sensitivity analysis of Mt-MscL and Ec-MscL in Ec-like lipids, Mt-like lipids, and Azolectin. * $p < 0.05$, **** $p = 0.0001$ (two tailed un-paired student t test). C. A short open dwell behavior observed in a subset of

Mt-MscL channels in Mt-like lipids. “O” and “C” refer to channel open and closed states. The scale bar is shown at the top right.

FIGURE 3.

Pressure threshold of Mt-MscL and Ec-MscL in neutral lipids. A. Patch clamp recording of Mt-MscL and Ec-MscL in lipids composed of 50%DOPE:50%DOPC. "O" and "C" refer to channel open and closed states. The pressures for MscL open are shown under each trace. The scale bar is shown at the top right. B. Pressure threshold analysis of Mt-MscL and Ec-MscL in lipids composed of 50%DOPE:50%DOPC. *** $p=0.0006$ (two tailed unpaired student t test).