Published in final edited form as:

Transplantation. 2013 November 15; 96(9): . doi:10.1097/TP.0b013e318298fa61.

The Decline in Living Kidney Donation in the United States: Random Variation or Cause for Concern?

James R. Rodrigue, PhD¹, Jesse D. Schold, PhD², and Didier A. Mandelbrot, MD¹

¹Center for Transplant Outcomes and Quality Improvement. The Transplant Institute, Beth Israel Deaconess Medical Center. Harvard Medical School, Boston, Massachusetts, USA

²Quantitative Health Sciences, Cleveland Clinic, Cleveland, Ohio, USA

Abstract

The annual number of living kidney donors in the United States peaked at 6,647 in 2004. The preceding decade saw a 120% increase in living kidney donation. However, since 2004, living kidney donation has declined in all but one year, resulting in a 13% decline in the annual number of living kidney donors from 2004 to 2011. The proportional decline in living kidney donation has been more pronounced among men, blacks, younger adults, siblings, and parents. In this paper, we explore several possible explanations for the decline in living kidney donation, including an increase in medical unsuitability, an aging transplant patient population, financial disincentives, public policies, and shifting practice patterns, among others. We conclude that the decline in living donation is not merely reflective of random variation, but one that warrants action by transplant centers, the broader transplant community, and state and national governments.

Keywords

living donation; live donor kidney transplantation

Introduction

Live donor kidney transplantation (LDKT) accounts for one-third of kidney transplants performed in the United States and continues to offer superior outcomes compared to maintenance dialysis and deceased donor kidney transplantation for individuals with end-stage kidney disease (1). Figure 1 shows a remarkable 265% increase in the annual number of living donors from 1988 to 2004. In each year during this time period, there was an increase in living donation relative to the previous year. This increase in living donation may be attributed to several factors, including advances in histocompatibility testing, new laparoscopic surgical techniques, findings of comparable graft survival outcomes with genetically related and unrelated living donors, and greater public awareness about the need for organ donors.

However, since the peak of 6,647 donors in 2004, the annual number of living donors has declined from the previous year in all but one year (2009), despite the emergence of novel programs to expand kidney donation (2–8). This observed trend differs from recent increases in living donation seen in some regions of the world (e.g., United Kingdom, Japan,

Corresponding author: James R. Rodrigue, Ph.D., The Transplant Institute, Beth Israel Deaconess Medical Center, 110 Francis Street, 7th Floor, Boston, MA 02215, (617) 632-9821 (phone), (617) 632-9820 (fax), jrodrig@bidmc.harvard.edu.

DISCLOSURE

The authors of this manuscript have no conflicts of interest to disclose.

Netherlands, Mexico, and Australia) (11). This recent living donation trend in the United States has not received the scrutiny one might expect in light of its implications for transplant candidates. In this paper, we examine the recent decline in living donation, including whether it is specific to certain subgroups or populations, factors that might explain it, and what the transplant community should do, if anything, to attenuate it. Our intent is to stimulate further discussion and scientific inquiry into living donation trends and patterns in the United States.

Is the decline in living donation specific to particular subgroups or populations?

We extracted data from the Organ Procurement and Transplantation Network (OPTN) website to examine trends in living donation within subgroups (1). To facilitate our examination of pattern shifts before and after the peak number of living donors observed in 2004, we divided living donation into two eras reflecting identical time periods: Era 1 (1998 to 2004) and Era 2 (2005 to 2011). The proportions of living donors for each category in the two eras were compared using *t*-tests for proportions.

Donor Gender

Women represent 57% to 62% of all living donors in each of the last 15 years (Figure 2). There was a decline in male donors as a proportion of total donors from 41.9% in Era 1 to 39.8% in Era 2 ($p=0.006$). The number of male donors peaked in Era 1 (2004) and declined consistently throughout Era 2. In contrast, the three highest annual totals for female donors occurred during Era 2 (2005, 2009, 2010).

Donor Race/Ethnicity

Non-Hispanic whites represent the majority (70%) of living donors over the last 15 years. Living donation among non-Hispanic whites and blacks peaked in Era 1 (both in 2004). The number of non-Hispanic white ($n=4,045$) and black ($n=680$) donors in 2011 are the lowest annual totals for both races in more than a decade. Black donors as a proportion of total donors dropped significantly from 13.4% in Era 1 to 12.2% in Era 2 ($p=0.002$). In contrast, the annual number of both Hispanic and Asian living donors has increased over time, and there was a significant increase in the proportion of total donors from Era 1 to Era 2 for Hispanics and Asians (11.9% to 13.6%, $p<0.001$ and 2.9% to 3.4%, $p=0.010$, respectively).

Donor Age

There has been a steady decline in the annual number of living donors <50 years old since 2004 (Figure 3). Living donation among adults aged 18 to 34 years and 35 to 49 years peaked in Era 1 (2004) and Era 2 (2005), respectively, and the number of 2011 donors for both age groups was the lowest in 12 years. Living donors in the 18 to 34 and 35 to 49 age groups as a proportion of total donors dropped significantly from Era 1 to Era 2 (33.5% to 30.6%, $p<0.001$ and 46.9% to 44.0%, $p=0.002$, respectively). In contrast, the number of annual living donors aged 50 to 64 years increased steadily since 1998. Their proportion of total donors increased from 18.6% in Era 1 to 23.9% in Era 2 ($p<0.001$). Compared to younger donors, the number of annual donors aged 50 to 64 years and 65+ years peaked only more recently in Era 2 (both in 2010).

Donor Relation

Sibling donors, historically the most common type, peaked in Era 1 (2001) and have declined substantially since then. For the first time, in 2010 another type of living donor (unrelated non-spouses) exceeded the annual number of sibling donors. Parent donors,

excluding pediatric transplant recipients, also peaked in Era 1 (2001) and have decreased over time, although not as precipitously as sibling donors. The number of sibling (n=1,185) and parent (n=323) donors in 2011 reflects the lowest annual totals for both groups in more than two decades. Both siblings and parents as a proportion of total donors dropped significantly from Era 1 to Era2 (33.1% to 24.1%, $p<0.001$ and 13.9% to 9.4%, $p=0.001$, respectively). Both unrelated non-spouse and child donors increased sharply during Era 1, peaked early in Era 2 (2006 and 2005, respectively) and declined in recent years.

Region

All 11 organ procurement regions in the United States experienced an increase in the annual number of living kidney donors from 1998 to 2004 (1). However, the annual number of living donors peaked in Era 1 (Regions 1, 2, 5, and 6) or early in Era 2 (Regions 3, 7, 10, and 11) in the majority of regions. In contrast, more recent peaks (2009, 2010) in annual living donors have been observed in Regions 4 (Oklahoma, Texas), 8 (Colorado, Iowa, Kansas, Missouri, Nebraska, Wyoming), and 9 (New York, Western Vermont); indeed, these regions have seen an increase in living donation during Era 2.

Overall, while the annual number of living donors has declined since the peak in 2004, the decline appears more pronounced and sustained among men, blacks, younger adults, siblings, and parents. All of these subgroups reached an annual peak in Era 1, have been on a persistent decline in Era 2, and their respective 2011 totals were the lowest observed in a decade or more.

What are possible hypotheses to explain the decline in living kidney donation?

Change in deceased donor allocation

In October 2005, the United Network for Organ Sharing (UNOS) implemented a new policy that prioritized the allocation of kidneys from deceased donors less than 35 years old to pediatric candidates less than 18 years old (10). A commonly heard refrain within the transplant community is that the decline in living donation is attributable to this allocation change. The central hypothesis is that preferential access to young deceased donor kidneys results in substantially shorter waiting time and, consequently, lower LDKT rates. There is some empirical support for this supposition. Agarwal et al.(11) found that the mean waiting time for pediatric patients who received a deceased donor kidney declined significantly from pre to post policy implementation, as did the mean number of LDKTs to pediatric recipients.

However, there are factors that argue against attributing the decline in living donation solely to this allocation policy change. For instance, if the policy change caused the decline in total number of living donors, there would be a decrease in the number of living donors for pediatric candidates and for the total candidate population, but not for adult candidates. Instead, we observe that the number of living donors for adult candidates has declined in all but one year following policy implementation and there were 11% fewer living donors for adult candidates in 2011 (n = 5,468) compared to 2005 (n = 6,148). Moreover, while there is good evidence that the policy change led to a reduction in LDKT rates for pediatric candidates (11), living donors for pediatric patients had already peaked in 2002 and was in decline before the policy change in 2005. We conclude, therefore, that the decline in the total number of living donors is partially, but not entirely, due to the allocation policy change in 2005.

Increased emphasis on performance oversight

The Scientific Registry of Transplant Recipients (SRTR) has published outcomes for transplant centers in the United States for over a decade (12). These data contain information on center volume, donor and recipient characteristics as well as risk adjusted post-transplant graft and patient survival. In 2007 (Era 2), the Center for Medicare and Medicaid Services (CMS) published Conditions of Participation (COP) which explicitly aligns transplant center outcomes with potential loss of public funding using very similar outcomes as the SRTR (13). One concern of increased regulatory oversight is a suppression of innovation, a decline in the use of novel therapies, and an increase in more conservative policies and practices due to the potential impact on center performance reports (14,15). Schold et al.(16) recently found that transplant programs receiving a low performance evaluation between 2007 and 2010 had an average reduction of 22 transplants over the study period compared to an average increase of 8 transplants among centers that did not receive a low performance evaluation ($p=0.001$). This difference was significant across donor types, including an average net decrease of 15 LDKTs among centers with a low performance evaluation as compared to an average decrease of only 3 LDKTs among centers without a low performance evaluation.

Largely in response to widely publicized deaths of living liver and kidney donors, the Secretary of Health and Human Services directed the OPTN in 2006 (Era 2) to develop policies regarding living organ donors and organ donor recipients and stipulated that program noncompliance with these policies would result in consequences similar to those for noncompliance with OPTN deceased donor transplantation. The OPTN subsequently developed policy regarding the medical and psychosocial evaluation of potential living donors, the informed consent process, the inclusion of independent living donor advocates in the evaluation process, and minimum requirements for the submission of living donor follow-up data (OPTN Policy 12.0, Living Donation, February 1, 2013).

An important question is whether the increased regulatory oversight and scrutiny of center outcomes will have an appreciable impact on growth of transplantation including living donation. Centers may have more reluctance to select “marginal” candidates or adults for living donation due to the potential impact on center report cards. Also, if centers with performance concerns or insufficient resources to meet new OPTN policy requirements are more reluctant to sustain or expand their LDKT programs, this may contribute to a decline in living donation. Finally, the degree to which increased media attention of living donor deaths and increased regulatory oversight in Era 2 impacted the willingness to the general public to pursue living kidney donation is unknown, but these factors should be considered when evaluating contributors to the decline in living donation.

Changes in donor selection criteria

In the United States, the expansion of living donor eligibility criteria is well documented (17–19). This includes not only expanding the types of donors who are acceptable candidates (e.g., unrelated, anonymous non-directed, older donors), but also the medical and psychosocial eligibility criteria. Many more programs have shifted practices and are now willing to consider donors with well-controlled hypertension, mild obesity, and stable psychiatric disorders. It is likely that the increase in living donation observed in Era 1 is attributable, in part, to these shifting donor eligibility criteria.

While programs may be more willing to evaluate a wider range of potential donors, there may be some retraction in other aspects of the donor evaluation (e.g., moderate obesity). Also, some programs now apply different eligibility criteria across different groups of donor candidates (e.g., race, obesity), based on outcomes published mostly in Era 2 (20–24). For

example, young blacks with risk factors for metabolic syndrome may be excluded due to studies demonstrating higher rates of CKD in this population compared to whites (20,25). Finally, the availability and evolution of radiological and histocompatibility techniques have led to more refined evaluation of the renovascular anatomy and immuno-compatibility. For example, increased measurement of donor specific antibodies may rule out some donor candidates. While more detailed assessment and testing is warranted in light of expanding eligibility criteria, widespread adoption of more refined testing may be contributing to a higher incidence of donor exclusion in the last decade. It could also be argued that advancements in immunological and histocompatibility screening have led to the notable growth of kidney exchange and desensitization protocols that have enabled adults – previously considered ineligible – to become living donors (26–28). While the total number of donors participating in kidney exchanges surged from 65 in Era 1 to 1,593 in Era 2, it is too early to determine whether growth in these types of programs can offset the observed decline in living donation rates.

A changing transplant population

Increasingly, older adults comprise the majority of new additions to the kidney transplant waiting list (1,29,30). Adults 50 years and older comprised 45% (8,437/18,942) of all adults listed for kidney transplantation in 1998, but this representation increased dramatically to 60% (19,540/32,650) in 2011 (Figure 4). Of particular note is the increase in adults 65+ years on the waiting list, from 8% (1,574/18,942) in 1998 to 17% (5,431/32,650) in 2011. One possible implication of the increasingly older transplant population is a correspondingly smaller network of potential living donors for these patients. As adults age, their peripheral social networks contract due to employment loss, retirement, and more restrictions on leisure activities (31). Chronic kidney disease, dialysis treatment, and quality of life limitations may further increase social isolation, which may reduce the transplant candidate's overall pool of potential donors. Moreover, while older living donors may yield favorable transplant outcomes (32,33), the risk of medical exclusion likely increases with age. Siblings of older transplant candidates increasingly become ineligible for donation due to declining kidney function and comorbidities, and parents are almost always excluded from donation by these factors. The decline in mean household size in the United States during the last 50 years (3.47 in 1950 to 2.63 in 2009) may also be contributing to the notable retraction in the number of sibling donors. Overall, older transplant candidates simply may not have as many potential donors as younger transplant candidates, which may explain the lower LDKT rate in older adults. In 2011, 50% of transplant recipients 18 to 34 years old received a LDKT, compared to only 30% and 34% of recipients 50 to 64 years old and 65+ years, respectively (1).

Transplant waiting time

Shorter waiting times for deceased donor kidney transplantation might impact center practices (e.g., more stringent donor selection criteria, less likely to implement new programs to increase LDKT rates) as well as the willingness of patients and family members to consider living donation (34). Indeed, there is some evidence that LDKT rates are lower in centers with shorter transplant waiting times (35,36). However, these studies are based largely on data from Era 1. In light of median waiting times exceeding 5 years in some regions, it cannot yet be determined whether the relationship between waiting time and LDKT rates has been maintained throughout Era 2.

Health status of the general population

Obesity, pre-diabetes, diabetes, and hypertension are alarmingly prevalent in the United States (37–40). These health conditions contribute to the incidence of chronic kidney disease and may also account for a smaller pool of potential living donors, particularly in subgroups

with a higher incidence of these risk factors (e.g., Blacks, older adults). Diabetes is an absolute contraindication to living donation, while obesity and hypertension are strong relative contraindications in many instances. The percentage of adults with diabetes in the United States increased from 6.3% in Era 1 to 8.1% in Era 2 (41). Similarly, the percentage of obese adults (BMI>30) increased from 22.7% in Era 1 to 27.4% in Era 2 (42). Collectively, these risk factors are the most common medical reasons for exclusion from living donation, particularly among minorities (43–45).

Financial disincentives in an economic crisis

The economic downturn in the United States is well documented (46). Since 2007, unemployment claims have risen, median household income has declined, and housing foreclosures have increased (46). These economic pressures on individuals and families may have negatively impacted willingness to pursue living donation, especially among those with low or moderate incomes and concerns about employment security. Indeed, Gill et al. (47) showed that the decline in living donation in the United States is most pronounced among adults with lower household income, particularly among blacks.

The indirect and direct costs associated with living donation have not been systematically captured. However, typical costs include those for travel, lodging, meals, childcare, and other service related costs while the donor is recovering (47). Lost wages are common for living donors, even for those with short-term disability or limited medical leave benefits. Out-of-pocket expenses are reported to range from \$0 to \$20,000, with an average of about \$5,000 when all direct and indirect costs are considered (48–50). In times of economic uncertainty, these costs may be a deterrent to living donation for otherwise willing potential donors. There may be heightened concerns about employment security or stability during a recession and less willingness to absorb lost wages, which is greater than one month's household income for the majority of living donors in the United States (51). While the U.S. Department of Health and Human Services provides some financial assistance for low-income donors (i.e., National Living Donor Assistance Center), these reimbursements are limited largely to travel and lodging. Reimbursements for lost wages are not permitted as part of this program. Recipients or family members who otherwise might have helped the living donor meet their financial obligations during recovery may be limited by the same financial hardships as the donor. Most transplant recipients have similar incomes to their living donors and, therefore, may not be able to compensate donors for out-of-pocket expenses (51). Overall, adults may be less willing to assume the financial and occupational risks of living donation during an economic downturn than in more prosperous times, and this may be contributing to the observed decline in living donation in the United States.

What can be done to attenuate the decline in living kidney donation?

We propose a multi-layered approach to attenuate the decline in living kidney donation, one that engages key stakeholders – transplant centers, the broader transplant community, and state and national governments. Transplant centers could examine trends in their LDKT rate since 2004 and use existing quality improvement processes to identify, implement, and evaluate strategies to address any downward pressures on LDKT. For instance, some centers have successfully employed patient navigators (6), home-based education (5), dialysis center-based interventions (8), and enhanced educational processes in the transplant center (7,52,53) to increase LDKT willingness and rates. These quality improvement initiatives and findings could then be disseminated to the broader transplant community for further evaluation and replication. The broader transplant community, using existing organizational structures (e.g., UNOS Living Donor Committee, American Society of Transplantation's Living Donor Community of Practice) could identify and disseminate “best practices” for removing barriers to living donation and LDKT as well as develop policies to reduce

financial disincentives to living donation. At the level of state and national governments, legislation could be developed to expand tax reduction incentives for living donor and prohibit discrimination against living donors by insurers. Together, the transplant community and government agencies could modify existing legislation and increase funding to allow existing financial assistance programs to reimburse living donors for lost wages. Additionally, transplant regulations and performance report cards must be carefully evaluated for their potential adverse impact on LDKT and living donation accessibility.

The decline in living donation over the last several years is indisputable. It is tempting to attribute this decline to random variation and to assume the number of living donors will return to levels observed in 2004. However, despite novel programs to help patients find suitable living donors, national financial programs and state tax incentives for living donors, and heightened general public awareness of living kidney donation, there is considerable downward pressure on living donation rates. Furthermore, the recent decline in living donation appears most prominent among men, blacks, and younger and lower income adults, which may further exacerbate existing LDKT disparities (54–59). We believe that this persistent downturn in living donation should be cause for action within the transplant community, as it potentially leads to longer waiting times for transplantation, greater dialysis exposure, higher death rates on the waiting list, lower graft and patient survival for recipients, and higher overall healthcare costs for the care of patients with end-stage renal disease. Programmatic, scientific, policy, and legislative efforts to identify and remove barriers to living donation are needed to ensure that LDKT remains a viable option for patients with chronic kidney disease.

Acknowledgments

The project described is supported by Award Number R01DK079665 (JR) and R01DK085185 (JR, DM, JS) from the National Institute of Diabetes and Digestive and Kidney Diseases. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institute of Diabetes and Digestive and Kidney Diseases or the National Institutes of Health. Preparation of this manuscript was also supported, in part, by the Julie Henry Research Fund and the Center for Transplant Outcomes and Quality Improvement, The Transplant Institute, Beth Israel Deaconess Medical Center, Boston, MA.

Abbreviations

BMI	Body mass index
CMS	Center for Medicare and Medicaid Services
COP	Conditions of Participation
ESRD	End-stage renal disease
LDKT	Live donor kidney transplantation
OPTN	Organ Procurement and Transplantation Network
SRTR	Scientific Registry of Transplant Recipients
UNOS	United Network for Organ Sharing

References

1. Organ Procurement and Transplantation Network (OPTN) and Scientific Registry of Transplant Recipients (SRTR). OPTN/SRTR 2011 Annual Data Report. Rockville, MD: Department of Health and Human Services, Health Resources and Services Administration, Healthcare Systems Bureau, Division of Transplantation; 2012. p. 11-46.

2. Melcher ML, Leese DB, Gritsch HA, Milner J, Kapur S, Busque S, et al. Chain transplantation: initial experience of a large multicenter program. *Am J Transplant*. 2012; 12:2429–36. [PubMed: 22812922]
3. Segev DL, Gentry SE, Warren DS, Reeb B, Montgomery RA. Kidney paired donation and optimizing the use of live donor organs. *JAMA*. 2005; 293:1883–90. [PubMed: 15840863]
4. Roth AE, Sonmez T, Unver MU, Delmonico FL, Saidman SL. Utilizing list exchange and nondirected donation through ‘chain’ paired kidney donations. *Am J Transplant*. 2006; 6:2694–705. [PubMed: 16981911]
5. Rodrigue JR, Cornell DL, Lin JK, Kaplan B, Howard RJ. Increasing live donor kidney transplantation: a randomized controlled trial of a home-based educational intervention. *Am J Transplant*. 2007; 7:394–401. [PubMed: 17173659]
6. Sullivan C, Leon JB, Sayre SS, Marbury M, Ivers M, Pencak JA, Bodziak KA, Hricik DE, Morrison EJ, Albert JM, Navaneethan SD, Reyes CM, Sehgal AR. Impact of navigators on completion of steps in the kidney transplant process: a randomized, controlled trial. *Clin J Am Soc Nephrol*. 2012; 7:1639–45. [PubMed: 22798540]
7. Hays R, Waterman AD. Improving preemptive transplant education to increase living donation rates: reaching patients earlier in their disease adjustment process. *Prog Transplant*. 2008; 18:251–6. [PubMed: 19186577]
8. Pradel FG, Suwannaprom P, Mullins CD, Sadler J, Bartlett ST. Short-term impact of an educational program promoting live donor kidney transplantation in dialysis centers. *Prog Transplant*. 2008; 18:263–72. [PubMed: 19186579]
9. Horvat LD, Shariff SZ, Garg AX. Donor Nephrectomy Outcomes Research (DONOR) Network. Global trends in the rates of living kidney donation. *Kidney Int*. 2009; 75:1088–98. [PubMed: 19225540]
10. Organ Procurement and Transplantation Network. [Accessed November 14, 2012] Policy 3.5. Organ Distribution: Allocation of Deceased Kidneys. Available from: <http://optn.transplant.hrsa.gov/policiesAndBylaws/policies.asp>
11. Agarwal S, Oak N, Siddique J, Harland RC, Abbo ED. Changes in pediatric renal transplantation after implementation of the revised deceased donor kidney allocation policy. *Am J Transplant*. 2009; 9:1237–42. [PubMed: 19422349]
12. Dickinson DM, Shearon TH, O’Keefe J, Wong HH, Berg CL, Rosendale JD, Delmonico FL, Webb RL, Wolfe RA. SRTR center-specific reporting tools: Posttransplant outcomes. *Am J Transplant*. 2006; 6:1198–211. [PubMed: 16613596]
13. Department of Health and Human Services. Centers for Medicare and Medicaid Services. 42 CFR Parts 405, 482, 488, and 498 Medicare Program; Hospital Conditions of Participation: Requirements for Approval and Re-Approval of Transplant Centers To Perform Organ Transplants; Final Rule. *Federal Register*. 2007; 72(61):15198–280.
14. Abecassis MM, Burke R, Klintmalm GB, Matas AJ, Merion RM, Millman D, Olthoff K, Roberts JP. American Society of Transplant Surgeons. American Society of Transplant Surgeons transplant center outcomes requirements--a threat to innovation. *Am J Transplant*. 2009; 9:1279–86. [PubMed: 19392984]
15. Schold JD, Arrington CJ, Levine G. Significant alterations in reported clinical practice associated with increased oversight of organ transplant center performance. *Prog Transplant*. 2010; 20:279–87. [PubMed: 20929114]
16. Schold JD, Buccini LD, Srinivas TR, Srinivas RT, Poggio ED, Flechner SM, Soria C, Segev DL, Fung J, Goldfarb DA. The association of center performance evaluations and kidney transplant volume in the United States. *Am J Transplant*. In Press.
17. Rodrigue JR, Pavlakis M, Danovitch GM, Johnson SR, Karp SJ, Khwaja K, Hanto DW, Mandelbrot DA. Evaluating living kidney donors: relationship types, psychosocial criteria, and consent processes at US transplant programs. *Am J Transplant*. 2007; 7:2326–32. [PubMed: 17845566]
18. Davis CL, Cooper M. The state of U.S. living kidney donors. *Clin J Am Soc Nephrol*. 2010; 5:1873–80. [PubMed: 20634322]

19. Mandelbrot DA, Pavlakis M, Danovitch GM, Johnson SR, Karp SJ, Khwaja K, Hanto DW, Rodrigue JR. The medical evaluation of living kidney donors: a survey of US transplant centers. *Am J Transplant.* 2007; 7:2333–43. [PubMed: 17845567]
20. Lentine KL, Schnitzler MA, Xiao H, Saab G, Salvalaggio PR, Axelrod D, Davis CL, Abbott KC, Brennan DC. Racial variation in medical outcomes among living kidney donors. *N Engl J Med.* 2010; 363:724–32. [PubMed: 20818874]
21. Lentine KL, Patel A. Risks and outcomes of living donation. *Adv Chronic Kidney Dis.* 2012; 19:220–8. [PubMed: 22732041]
22. Ommen ES, Winston JA, Murphy B. Medical risks in living kidney donors: absence of proof is not proof of absence. *Clin J Am Soc Nephrol.* 2006; 1:885–95. [PubMed: 17699301]
23. Heimbach JK, Taler SJ, Prieto M, Cosio FG, Textor SC, Kudva YC, Chow GK, Ishitani MB, Larson TS, Stegall MD. Obesity in living kidney donors: clinical characteristics and outcomes in the era of laparoscopic donor nephrectomy. *Am J Transplant.* 2005; 5:1057–64. [PubMed: 15816886]
24. Young A, Storsley L, Garg AX, Treleaven D, Ngan CY, Cuerden MS, Karpinski M. Health outcomes for living kidney donors with isolated medical abnormalities: a systematic review. *Am J Transplant.* 2008; 8:1878–90. [PubMed: 18671676]
25. Steiner RW. ‘Normal for now’ or ‘at future risk’: a double standard for selecting young and older living kidney donors. *Am J Transplant.* 2010; 10:737–41. [PubMed: 20199512]
26. Saidman SL, Roth AE, Sönmez T, Unver MU, Delmonico FL. Increasing the opportunity of live kidney donation by matching for two- and three-way exchanges. *Transplantation.* 2006; 81:773–82. [PubMed: 16534482]
27. Melancon JK, Cummings LS, Graham J, Rosen-Bronson S, Light J, Desai CS, Girlanda R, Ghasemian S, Africa J, Johnson LB. Paired kidney donor exchanges and antibody reduction therapy: novel methods to ameliorate disparate access to living donor kidney transplantation in ethnic minorities. *J Am Coll Surg.* 2011; 212:740–5. [PubMed: 21463825]
28. Wilpert J, Fischer KG, Pisarski P, Wiech T, Daskalakis M, Ziegler A, Neumann-Haefelin E, Drognitz O, Emmerich F, Walz G, Geyer M. Long-term outcome of ABO-incompatible living donor kidney transplantation based on antigen-specific desensitization. An observational comparative analysis. *Nephrol Dial Transplant.* 2010; 25:3778–86. [PubMed: 20466677]
29. Abecassis M, Bridges ND, Clancy CJ, Dew MA, Eldadah B, Englesbe MJ, et al. Solid-organ transplantation in older adults: current status and future research. *Am J Transplant.* 2012; 12:2608–22. [PubMed: 22958872]
30. Danovitch G, Savransky E. Challenges in the counseling and management of older kidney transplant candidates. *Am J Kidney Dis.* 2006; 47:S86–97. [PubMed: 16567243]
31. Fung HH, Carstensen LL, Lang FR. Age-related patterns in social networks among European Americans and African Americans: implications for socioemotional selectivity across the life span. *Int J Aging Hum Dev.* 2001; 52:185–206. [PubMed: 11407486]
32. Young A, Kim SJ, Speechley MR, Huang A, Knoll GA, Prasad GV, Treleaven D, Diamant M, Garg AX. Donor Nephrectomy Outcomes Research (DONOR) Network. Accepting kidneys from older living donors: impact on transplant recipient outcomes. *Am J Transplant.* 2011; 11:743–50. [PubMed: 21401866]
33. Gill J, Bunnapradist S, Danovitch GM, Gjertson D, Gill JS, Cecka M. Outcomes of kidney transplantation from older living donors to older recipients. *Am J Kidney Dis.* 2008; 52:541–52. [PubMed: 18653267]
34. Rodrigue JR, Ladin K, Pavlakis M, Mandelbrot DA. Disclosing recipient information to potential living donors: preferences of donors and recipients, before and after surgery. *Am J Transplant.* 2011; 11:1270–8. [PubMed: 21645257]
35. Segev DL, Gentry SE, Montgomery RA. Association between waiting times for kidney transplantation and rates of live donation. *Am J Transplant.* 2007; 7:2406–13. [PubMed: 17845574]
36. Reese PP, Feldman HI, Bloom RD, Abt PL, Thomasson A, Shults J, et al. Assessment of variation in live donor kidney transplantation across transplant centers in the United States. *Transplantation.* 2011; 91:1357–63. [PubMed: 21562451]

37. Crawford AG, Cote C, Couto J, Daskiran M, Gunnarsson C, Haas K, Haas S, Nigam SC, Schuette R. Prevalence of obesity, type II diabetes mellitus, hyperlipidemia, and hypertension in the United States: findings from the GE Centricity Electronic Medical Record database. *Popul Health Manag.* 2010; 13:151–61. [PubMed: 20521902]
38. Sentell TL, He G, Gregg EW, Schillinger D. Racial/ethnic variation in prevalence estimates for United States prediabetes under alternative 2010 American Diabetes Association criteria: 1988–2008. *Ethn Dis.* 2012; 22:451–8. [PubMed: 23140076]
39. Ogden CL, Carroll MD, Kit BK, Flegal KM. Prevalence of obesity in the United States, 2009–2010. *NCHS Data Brief.* 2012; 82:1–8. [PubMed: 22617494]
40. Guo F, He D, Zhang W, Walton RG. Trends in prevalence, awareness, management, and control of hypertension among United States adults, 1999 to 2010. *J Am Coll Cardiol.* 2012; 60:599–606. [PubMed: 22796254]
41. Centers for Disease Control and Prevention (CDC). Crude and Age-Adjusted Incidence of Diagnosed Diabetes per 1,000 Population Aged 18–79 Years, United States, 1980–2011. National Center for Health Statistics, Division of Health Interview Statistics; <http://www.cdc.gov/diabetes/statistics/incidence/fig2.htm> [Accessed December 5, 2012]
42. Centers for Disease Control and Prevention (CDC). Prevalence of obesity among adults aged 20 years and over: United States, 1997–June 2011. National Center for Health Statistics, Division of Health Interview Statistics; http://www.cdc.gov/nchs/data/nhis/earlyrelease/201112_06.pdf [Accessed December 5, 2012]
43. Moore DR, Feurer ID, Zaydfudim V, Hoy H, Zavala EY, Shaffer D, Schaefer H, Moore DE. Evaluation of living kidney donors: variables that affect donation. *Prog Transplant.* 2012; 22:385–92. [PubMed: 23187057]
44. Weng FL, Dhillon N, Lin Y, Mulgaonkar S, Patel AM. Racial differences in outcomes of the evaluation of potential live kidney donors: a retrospective cohort study. *Am J Nephrol.* 2012; 35:409–15. [PubMed: 22517188]
45. Norman SP, Song PX, Hu Y, Ojo AO. Transition from donor candidates to live kidney donors: the impact of race and undiagnosed medical disease states. *Clin Transplant.* 2011; 25:136–45. [PubMed: 20047616]
46. United States Bureau of Labor Statistics. [Accessed November 10, 2012] The recession of 2007 – 2009. BLS spotlight on statistics. Feb. 2012 Available from: http://www.bls.gov/spotlight/2012/recession/pdf/recession_bls_spotlight.pdf
47. Gill J, Dong J, Rose CL, Gill JS. The retraction of living donation is most marked among the poor. *J Am Soc Nephrol.* 2010; 21:103A. [PubMed: 19959717]
48. Clarke KS, Klarenbach S, Vlaicu S, Yang RC, Garg AX. The direct and indirect economic costs incurred by living kidney donors: a systematic review. *Nephrol Dial Transplant.* 2006; 21:1952–60. [PubMed: 16554329]
49. Gaston RS, Danovitch GM, Epstein RA, Kahn JP, Matas AJ, Schnitzler MA. Limiting financial disincentives in live organ donation: A rational solution to the kidney shortage. *Am J Transplant.* 2006; 6:2548–55. [PubMed: 16889608]
50. Rodrigue JR, Cross NJ, Newman RC, Widows MR, Guenther RT, Kaplan B, Morgan MA, Howard RJ. Patient-reported outcomes for open versus laparoscopic living donor nephrectomy. *Prog Transplant.* 2006; 16:162–9. [PubMed: 16789708]
51. Gill JS, Gill J, Barnieh L, Dong J, Rose C, Johnston O, Tonelli M, Klarenbach S. Income of living kidney donors and the income difference between living kidney donors and their recipients in the United States. *Am J Transplant.* 2012; 12:3111–8. [PubMed: 22882723]
52. Schweitzer EJ, Yoon S, Hart J, Anderson L, Barnes R, Evans D, et al. Increased living donor volunteer rates with a formal recipient family education program. *Am J Kidney Dis.* 1997; 29:739–45. [PubMed: 9159309]
53. Boulware LE, Hill-Briggs F, Kraus ES, Melancon JK, Falcone B, Ephraim PL, et al. Effectiveness of educational and social worker interventions to activate patients' discussion and pursuit of preemptive living donor kidney transplantation: a randomized controlled trial. *Am J Kidney Dis.* 2013; 61:476–86. [PubMed: 23089512]

54. Waterman AD, Rodrigue JR, Purnell TS, Ladin K, Boulware LE. Addressing racial and ethnic disparities in live donor kidney transplantation: priorities for research and intervention. *Semin Nephrol.* 2010; 30:90–8. [PubMed: 20116653]
55. Purnell TS, Hall YN, Boulware LE. Understanding and overcoming barriers to living kidney donation among racial and ethnic minorities in the United States. *Adv Chronic Kidney Dis.* 2012; 19:244–51. [PubMed: 22732044]
56. Hall EC, James NT, Garonzik Wang JM, Berger JC, Montgomery RA, Dagher NN, Desai NM, Segev DL. Center-level factors and racial disparities in living donor kidney transplantation. *Am J Kidney Dis.* 2012; 59:849–57. [PubMed: 22370021]
57. Ladin K, Rodrigue JR, Hanto DW. Framing disparities along the continuum of care from chronic kidney disease to transplantation: barriers and interventions. *Am J Transplant.* 2009; 9:669–74. [PubMed: 19344460]
58. Stolzmann KL, Bautista LE, Gangnon RE, McElroy JA, Becker BN, Remington PL. Trends in kidney transplantation rates and disparities. *J Natl Med Assoc.* 2007; 99:923–32. [PubMed: 17722672]
59. Jindal RM, Ryan JJ, Sajjad I, Murthy MH, Baines LS. Kidney transplantation and gender disparity. *Am J Nephrol.* 2005; 25:474–83. [PubMed: 16127268]

Figure 1.
Number of living kidney donors, United States, 1988 – 2012.

Figure 2.
Number of living kidney donors by gender and era, United States, 1998 – 2011.

Figure 3.
Number of living kidney donors by age group, United States, 1998 – 2011.

Figure 4.
Number of adults added to the kidney transplant waiting list by age group, United States, 1998 – 2011.