Published in final edited form as:

Synth Commun. 2013 ; 43(16): 2171–2177. doi:10.1080/00397911.2012.693239.

SUPERACID-PROMOTED HYDROXYALKYLATION OF 1,2-INDANDIONES

Adam F. Tracy, Matthew P. Abbott, and Douglas A. Klumpp

Department of Chemistry and Biochemistry, Northern Illinois University, DeKalb, IL 60115

Abstract

1,2-Indandione reacts efficiently with arenes to give 2,2-diaryl-1-indanones by the hydroxyalkylation reaction. The Brønsted superacid $\text{CF}_3\text{SO}_3\text{H}$ (triflic acid) is an effective catalyst for these condensation reactions. The requisite 1,2-indandiones were prepared from the 1-indanones.

INTRODUCTION

The hydroxyalkylation reaction is an acid-catalyzed condensation of aldehydes and ketones with arenes.¹ It is a common industrial reaction, as it is used to prepare feedstock chemicals, aromatic monomers, and condensation polymers. In 1995, Shudo and coworkers described a superacid-promoted hydroxyalkylation reaction of 2,3-butanedione (**1**) to give the condensation product **3** in excellent yield (eq 1).² The diprotonated, superelectrophile (**2**) was proposed as an intermediate in the reaction.³ Several other recent manuscripts have described superacid-promoted hydroxyalkylation reactions with 1,2-dicarbonyl compounds.⁴ For example, products **5–7** are prepared from the respective condensation reactions of acenaphthenequinone, isatin, and parabanic acid, with benzene and triflic acid.

(1)

These types of superacid-promoted reactions have also recently been shown to be useful in polymerization chemistry.⁵ In the following Communication, we describe superacid-promoted hydroxyalkylation reactions of 1,2-indandione and substituted derivatives. The conversions lead to 2,2-diaryl-1-indanone products and a mechanism is proposed involving a deprotonated superelectrophilic intermediate.

RESULTS AND DISCUSSION

Although 1,2-indandione (**7**) is available from commercial suppliers, substituted derivatives may be prepared from the 1-indanones.⁶ Thus, 4-bromoindanone was converted to the oxime **8** by reaction with isoamyl nitrite and hydrolysis liberates the substituted 1,2-indandione **9** (eq 2). Similar reactions were done with 5-methoxyindanone and 4-methoxyindanone to prepare the respective 1,2-diketones. We also sought to compare the reactions of 2,2-dichloro-1-indanones (**10**) and 2,2-dimethoxy-1-indanones (**11**) as possible substrates in the synthesis.

(2)

(3)

(4)

The desired compounds were prepared by chlorination of the 1-indanone (eq 3) and by acetalization of 1,2-indandione (eq 4).^{7,6}

With 1,2-indandione, the condensation product (**12**) is formed in 81% yield by reaction at the 2-position (Scheme 1). The identity of **12** was verified by comparison with an authentic sample of **12** (prepared from a different approach⁸) and it was clear that condensation occurs exclusively at the 2-carbonyl group. The regiochemistry of condensation was also confirmed by X-ray crystallographic analysis. During the course of our studies, we found that some 1,2-indandiones were prone to decomposition during storage. This prompted us to explore the possibility of protecting the diketone as the dimethoxy acetal. Although the acetal substrates were easily prepared and indefinitely stable, the condensation reaction from **11** provided compound **12** in a disappointing 43% yield. As an alternative strategy from the dichloride, compound **10** gave product **12** in a fair yield of 61%.

Using substituted arenes, the condensation reactions provide the products in good yields. Thus, reaction of **7** with triflic acid and fluorobenzene or toluene gives the respective products (**13-14**) in 85% and 67% yields (Scheme 2). The electrophilic aromatic substitution

occurs with excellent regioselectivity, as GC and NMR analysis shows at least a 20:1 *para:ortho* ratio. This is a trend previously observed with other 1,2-dicarbonyl systems – very good regioselectivity is seen in the condensation reactions with alkylbenzenes and monohalogenated benzenes.⁴ We also obtained the condensation product from 1,3-dichlorobenzene (**15**) although the yield is considerably lower. In some cases, minor amounts of the intermediate product inden-1-ones (i.e., **16**) could also be detected by GCMS analysis. Finally, substituted 1,2-indandiones were also prepared and reacted with CF₃SO₃H and benzene. 6-Methoxy and 5-methoxy-substituted 1,2-indandiones provide condensation products **17** and **18**, respectively. Similarly, 4-bromo-1,2-indandione gives the condensation product **19** in excellent yield from CF₃SO₃H.

Hydroxyalkylation reactions generally occur via carboxonium intermediates by protonation of aldehyde or ketone carbonyl groups.¹ In the case of 1,2-indandione (**7**), initial protonation could occur at either the 1- or 2-carbonyl group (Scheme 3). DFT calculations (B3LYP 6-311G (d,p) level) indicate that protonation at the 1-carbonyl group is significantly favored.⁹ However, condensation occurs at the 2-carbonyl. Benzene (and other arenes) are weak nucleophiles, and consequently, direct attack of benzene at the 2-carbonyl group of **21** is not considered a likely reaction step. Two plausible mechanisms can be suggested. A second protonation of **21** leads to superelectrophile **22** and reaction with benzene then occurs to provide adduct **23**. This mechanistic path could also involve only a partially diprotonated species where the 2-carbonyl group is forming a strong hydrogen bond to the superacid. Alternatively, reaction through the monoprotinated species (**20**) could directly give **23** and further reaction steps then lead to product **12**.

CONCLUSION

Using the Brønsted superacid CF₃SO₃H, 1,2-indandiones form condensation products with benzene and substituted arenes. The chemistry provides easy access to aryl-functionalized 1-indanones. Moreover, the condensation occurs with good positional selectivity in reactions with substituted benzenes.

EXPERIMENTAL

General Methods

Solvents and reagents were purchased from commercial suppliers and used as received. The triflic acid was distilled from an Ar atmosphere prior to its use. Compounds were purified using standard flash chromatography grade silica gel. Low-resolution mass spectra were obtained from a commercial GC instrument equipped with a mass selective detector. High-resolution mass spectra were obtained by an external analytical laboratory. All NMR spectra were obtained from a Bruker Avance DRX-500 instrument.

General Experimental Procedures

Preparation of condensation products 12-15 and 17-19—The 1,2-indandione (1 mmol) is dissolved in CH₂Cl₂ (5 mL) and the aromatic compound (1 mL) is added. The vessel is purged with Ar and then triflic acid (1 mL, 11 mmol) is slowly added to the solution with stirring. After 3 hr., the mixture is poured over ice. The resulting mixture is

extracted twice with chloroform and the organic extracts are combined. The organic solution is washed with water (2x) then brine (2x), and dried over anhydrous sodium sulfate. Filtration and removal of the solvent provides the crude condensation product. In some cases, the resulting product is greater than 95% pure (GC analysis) and requires no further purification. However, mixtures were purified by flash chromatography (hexanes:ether).

Alternative condensation procedure for 12—1,2-Indandione (0.15g, 1 mmol) is dissolved in benzene (5 mL) and anhydrous aluminum chloride is added (0.67 g, 5 mmol) is added. The solution is refluxed under Ar for 4 hr and then poured over ice. The resulting mixture is extracted twice with chloroform and the organic extracts are combined. The organic solution is washed with water (2x) then brine (2x), and dried over anhydrous sodium sulfate. Filtration and removal of the solvent provides the crude condensation product. Compound **12** is then isolated by flash chromatography (hexanes:ether).

Acknowledgments

Financial support from the NIH-National Institute of General Medical Sciences (GM085736-01A1) is gratefully acknowledged.

References

1. (a) Hofmann, JE.; Schriesheim, A. Friedel-Crafts and Related Reaction. Olah, GA., editor. Vol. 2. Wiley; New York, NY: 1964. p. 597-640.(b) March, J. Advanced Organic Chemistry. 4. Wiley; New York, NY: 1992. p. 548-549.
2. Yamazaki T, Saito S, Ohwada T, Shudo K. Tetrahedron Lett. 1995; 36:5749.
3. For a review of superelectrophiles, see: Olah GA, Klumpp DA. Superelectrophiles and Their Chemistry. Wiley & Sons New York 2008
4. (a) Klumpp DA, Zhang Y, Do D, Kartika R. Appl Catal A. 2008; 336:128.(b) Klumpp DA, Yeung KY, Prakash GKS, Olah GAJ. Org Chem. 1998; 63:4481.(c) Klumpp DA, Yeung KY, Prakash GKS, Olah GA. SynLett. 1998:918.
5. (a) Zolotukhin MG, Hernandez MCG, Lopez AM, Fomina L, Cedillo G, Nogales A, Ezquerro T, Rueda D, Colquhoun HM, Fromm KM, Ruiz-Trevino A, Ree M. Macromolecules. 2006; 39:4696. (b) Colquhoun HM, Zolotukhin MG, Khalilov LM, Dzhemilev UM. Macromolecules. 2001; 34:1122.
6. Petrovskaia O, Taylor BM, Hauze DB, Carroll PJ, Joullie MM. J Org Chem. 2001; 66:7666. [PubMed: 11701019]
7. Quintiliano SAP, Silva LF Jr. J Braz Chem Soc. 2007; 18:1281.
8. Klumpp DA, Garza M, Lau S, Shick B, Kantardjieff K. J Org Chem. 1999; 64:7635.
9. Frisch, MJ.; Trucks, GW.; Schlegel, HB.; Scuseria, GE.; Robb, MA.; Cheeseman, JR.; Scalmani, G.; Barone, V.; Mennucci, B.; Petersson, GA.; Nakatsuji, H.; Caricato, M.; Li, X.; Hratchian, HP.; Izmaylov, AF.; Bloino, J.; Zheng, G.; Sonnenberg, JL.; Hada, M.; Ehara, M.; Toyota, K.; Fukuda, R.; Hasegawa, J.; Ishida, M.; Nakajima, T.; Honda, Y.; Kitao, O.; Nakai, H.; Vreven, T.; Montgomery, JA., Jr; Peralta, JE.; Ogliaro, F.; Bearpark, M.; Heyd, JJ.; Brothers, E.; Kudin, KN.; Staroverov, VN.; Kobayashi, R.; Normand, J.; Raghavachari, K.; Rendell, A.; Burant, JC.; Iyengar, SS.; Tomasi, J.; Cossi, M.; Rega, N.; Millam, NJ.; Klene, M.; Knox, JE.; Cross, JB.; Bakken, V.; Adamo, C.; Jaramillo, J.; Gomperts, R.; Stratmann, RE.; Yazyev, O.; Austin, AJ.; Cammi, R.; Pomelli, C.; Ochterski, JW.; Martin, RL.; Morokuma, K.; Zakrzewski, VG.; Voth, GA.; Salvador, P.; Dannenberg, JJ.; Dapprich, S.; Daniels, AD.; Farkas, Ö.; Foresman, JB.; Ortiz, JV.; Cioslowski, J.; Fox, DJ. Gaussian 09, Revision B.01. Gaussian, Inc; Wallingford CT: 2009.

Scheme 1.
Synthetic approaches to 2,2-diphenyl-1-indanone (**12**).

Scheme 2.

Products (**13-15**) from the reaction of 1,2-indandione.

Scheme 3.

Proposed reaction mechanism with DFT calculated relative energies of monocations **20** and **21**.

Table 1

Analytical data for condensation products **13-15** and **17-19**.

	MP	¹ H NMR	¹³ C NMR	Mass spectrum (low res.; EI)	Mass spectrum (high res.; EI)
13	143–145°C	CDCl ₃ , : 3.91 (s, 2H), 6.98 (m, 4H), 7.25–7.31 (m, 4H), 7.44–7.47 (m, 1H), 7.56 (d, 1H, <i>J</i> =7.3 Hz), 7.68–7.71 (m, 1H), 7.84 (d, 1H, <i>J</i> =7.6 Hz)	CDCl ₃ , : 44.8, 61.5, 115.3 (d, <i>J</i> _{C-F} =21.2 Hz), 125.3, 125.9, 128.2, 129.7 (d, <i>J</i> _{C-F} =7.9 Hz), 135.4, 135.6, 139.1, 151.6, 161.7 (d, <i>J</i> _{C-F} =245 Hz), 204.9	320 (M+), 270, 225, 196, 190	Calcd for C ₂₁ H ₁₄ OFe ₂ , 320.10128, found, 320.10207
14	108–111°C	CDCl ₃ , : 2.34 (s, 6H), 3.92 (s, 2H), 7.11 (d, 4H, <i>J</i> =8.2 Hz), 7.20 (d, 4H, <i>J</i> =8.2 Hz), 7.41–7.44 (m, 1H), 7.53 (d, 1H, <i>J</i> =7.6 Hz), 7.65–7.68 (m, 1H), 7.83 (d, 1H, <i>J</i> =7.6 Hz)	CDCl ₃ , : 20.9, 44.9, 62.2, 125.2, 125.8, 127.8, 128.0, 129.1, 135.2, 135.8, 136.3, 140.6, 152.1, 205.6	312 (M+), 269, 253, 220, 178	Calcd for C ₂₃ H ₂₀ O, 312.15142, found, 312.15193
15	194–196°C	CDCl ₃ , : 4.21 (s, 2H), 7.08 (d, 2H, <i>J</i> =8.4 Hz), 7.18 (dd, 2H, <i>J</i> =8.6, 2.2 Hz), 7.45–7.48 (m, 6H), &.66–7.69 (m, 1H), 7.92 (d, 1H, <i>J</i> =7.4 Hz)	CDCl ₃ , : 43.8, 62.9, 124.5, 126.2, 127.1, 128.0, 131.4, 131.8, 134.1, 134.9, 135.8, 136.3, 137.1, 152.0, 204.8	424/422/420 (M+), 387/385, 286, 252/250, 143, 125	Calcd for C ₂₁ H ₁₂ OCl ₄ , 419.96422, found, 419.96345
17	151–154°C	CDCl ₃ , 300MHz, : 3.86 (s, 3H), 3.87 (s, 2H), 7.25–7.33 (m, 12H), 7.44 (d, 1H, <i>J</i> = 9.3).	CDCl ₃ , 300MHz, : 44.3, 55.6, 63.6, 106.2, 124.8, 126.6, 126.7, 128.2, 129.4, 136.9, 143.6, 144.9, 159.8, 205.3	314 (M+), 253, 237, 223, 165.	Calcd for C ₂₂ H ₁₈ O ₂ , 314.13068, found, 314.13160
18	135–137°C	CDCl ₃ , 300MHz, : 3.86 (s, 3H), 3.87 (s, 2H), 7.25–7.33 (m, 12H), 7.439 (d, 1H, <i>J</i> = 9.3 Hz)	CDCl ₃ , 300MHz, : 44.9, 55.7, 62.9, 109.11, 115.9, 126.7, 126.9, 128.2, 128.4, 128.9, 143.9, 155.0, 165.9, 203.5	314 (M+), 285, 237, 223, 165	Calcd for C ₂₂ H ₁₈ O ₂ , 314.13068, found, 314.13000
19	114–117°C	CDCl ₃ , 300MHz, : 3.90 (s, 1H), 7.24–7.38 (m, 11H), 7.79 (d, 1H, <i>J</i> = 7.5 Hz), 7.84 (dd, 1H, <i>J</i> = 7.8, 8.7 Hz).	CDCl ₃ , 300MHz, : 45.6, 62.9, 121.4, 124.0, 127.0, 128.6, 129.8, 137.6, 138.0, 142.9, 151.7, 204.6	364/362 (M+), 284, 265, 252, 178	Calcd for C ₂₁ H ₁₅ OBr, 362.03063, found, 362.03151