Published in final edited form as:

Anesth Analg. 2013 October ; 117(4): 1010–1016. doi:10.1213/ANE.0b013e3182a1bf1c.

Neuraxial Anesthesia Decreases Post-operative Systemic Infection Risk Compared to General Anesthesia in Knee Arthroplasty

Jiabin Liu, MD, PhD[§], Chenjuan Ma, PhD[¶], Nabil Elkassabany, MD, MSCE[§], Lee A. Fleisher, MD[§], and Mark D. Neuman, MD, M.Sc[§]

[§]Department of Anesthesiology and Critical Care, the Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA, USA 19104.

[¶]Center for Health Outcomes and Policy Research, University of Pennsylvania, School of Nursing, Philadelphia, PA, USA 19104.

Abstract

Background—Surgical stress has been previously shown to result in immune disturbance. Neuraxial anesthesia (NA) has been long hypothesized to blunt undesired surgical insults, and thus limit immune compromise and improve surgical outcomes. We hypothesized that neuraxial anesthesia would decrease postoperative infectious complications compared to general anesthesia among knee arthroplasty patients.

Methods—We studied the American College of Surgeons National Surgical Quality Improvement Program (NSQIP) database from 2005 to 2010. There were total of 16,555 patients included in our final cohort, with 9,167 patients receiving general anesthesia (GA) and 7,388 patients receiving spinal or epidural anesthesia (neuraxial anesthesia). Outcomes of interest included infection-related 30-day postoperative complications, including surgical site related infections, pneumonia, urinary tract infection, sepsis, septic shock, and a composite endpoint of any systemic infection. Multivariable logistic regression was performed to test for effect of anesthesia type while adjusting for the influence of pre-existing comorbidities.

Results—The overall mortality was 0.24% and 0.15% among NA and GA subjects respectively ($P=0.214$). NA subjects had lower unadjusted incidences of pneumonia ($P=0.035$) and composite systemic infection ($P=0.006$). After risk adjustment for pre-existing co-morbidities, NA was associated with lower odds of pneumonia (OR=0.51, [95% CI 0.29–0.90]), and lower odds of composite systemic infection (OR=0.77, 95% CI [0.64–0.92]).

Conclusions—Our study suggested that neuraxial anesthesia was associated with lower adjusted odds of both pneumonia and a composite outcome of any systemic infectious complication within 30 days of surgery compared to general anesthesia.

Introduction

There are total over 500,000 total knee arthroplasty procedures performed annually in the United States, and the total number of cases is increasing every year ¹. Most of these knee arthroplasty candidates are elderly and some have extensive co-existing cardiac, pulmonary, and other systemic diseases ².

Surgical stress and anesthesia result in various metabolic and endocrine disturbances, and may lead to generalized immune suppression ^{3–6}. Neuraxial anesthesia (NA), including spinal and/or epidural anesthesia, reduces the surgical stress response ^{7,8}. It has been long

1. Author: Jiabin Liu, MD, PhD
- Title: Assistant Professor of Anesthesiology and Critical Care
 - Affiliation: Department of Anesthesiology and Critical Care, the Perelman School of Medicine, the University of Pennsylvania, Philadelphia, PA, USA 19104.
 - hypothesized that neuraxial anesthesia may reduce complications and improve surgical
 - Jiabin.Liu@uphs.upenn.edu
 - Contribution: <This author helped design the study, conduct the study, data analysis, and prepare the manuscript.>
 - Attestation: < Liu, Jiabin approved the final manuscript. Liu, Jiabin attests to the integrity of the original data and the analysis reported in this manuscript. Liu, Jiabin is the archival author.>
 - Conflicts of Interest: <“None”>
2. Author: Chenjuan Ma, PhD
- Title: Research Assistant
 - Affiliation: Center for Health Outcomes and Policy Research, the University of Pennsylvania, School of Nursing, Philadelphia, PA, USA 19104.
 - chenjuan@nursing.upenn.edu
 - Contribution: < This author helped conduct the study and data analysis >
 - Attestation: < Ma, Chenjuan approved the final manuscript. Ma, Chenjuan attests to the integrity of the original data and the analysis reported in this manuscript.>
 - Conflicts of Interest: <“None”>
3. Author: Nabil Elkassabany, MD, MSCE
- Title: Assistant Professor of Anesthesiology and Critical Care
 - Affiliation: Department of Anesthesiology and Critical Care, the Perelman School of Medicine, the University of Pennsylvania, Philadelphia, PA, USA 19104.
 - Nabil.Elkassabany@uphs.upenn.edu
 - Contribution: <This author helped design the study and prepare the manuscript.>
 - Attestation: < Nabil Elkassabany approved the final manuscript.>
 - Conflicts of Interest: <“None”>
4. Author: Name, Lee A. Fleisher, MD
- Title: Rober Dunning Dripps Professor of Anesthesiology and Critical Care
 - Affiliation: Department of Anesthesiology and Critical Care, the Perelman School of Medicine, the University of Pennsylvania, Philadelphia, PA, USA 19104.
 - Lee.Fleisher@uphs.upenn.edu
 - Contribution: <This author helped design the study and prepare the manuscript.>
 - Attestation: < Lee A. Fleisher approved the final manuscript.>
 - Conflicts of Interest: <“None”>
5. Author: Mark D. Neuman, MD, M.Sc
- Title: Assistant Professor of Anesthesiology and Critical Care
 - Affiliation: Department of Anesthesiology and Critical Care, the Perelman School of Medicine, the University of Pennsylvania, Philadelphia, PA, USA 19104.
 - Mark.Neuman@uphs.upenn.edu
 - Contribution: <This author helped design the study, data analysis, and prepare the manuscript.>
 - Attestation: < Mark D. Neuman approved the final manuscript. Mark D. Neuman attests to the integrity of the original data and the analysis reported in this manuscript.>
 - Conflicts of Interest: <“None”>

Name of Department(s) and Institution(s):

Department of Anesthesiology and Critical Care, the Perelman School of Medicine, the University of Pennsylvania, Philadelphia, PA, USA 19104.

Institution Review Board, the University of Pennsylvania, 3624 Market Street, Suite 301S, Philadelphia, PA 19104. Phone: 1-215-573-2540

outcomes by blocking noxious afferent inputs and decreasing surgery-related disturbance^{9,10}. These effects of neuraxial anesthesia, particularly those on the immune response, might have a profound effect on the risk of infectious complications. Nonetheless, while prior studies have suggested that neuraxial anesthesia is beneficial among selected patient groups^{11–15}, the small sample sizes preclude firm conclusion regarding the association of anesthesia type and infectious complications.

In this study, we focused on knee arthroplasty, which is one of the most common and painful procedures regularly performed with general and/or neuraxial anesthesia worldwide. We hypothesized that neuraxial anesthesia (NA) would decrease postoperative infectious complications compared to general anesthesia (GA) among knee arthroplasty patients. Drawing on a prospectively collected and validated clinical database, we compared rates of pneumonia, urinary tract infection, sepsis, septic shock, surgical site infections, and a composite outcome of any systemic infectious events according to anesthesia type.

Methods

Data source

This study was exempted by the institutional review board of the University of Pennsylvania (Philadelphia, Pennsylvania, USA). We acquired the American College of Surgeons National Surgical Quality Improvement Program (NSQIP) database from 2005 to 2010 (<http://site.acsnsqip.org>). NSQIP is a nationally validated, risk-adjusted, outcome-based program to measure surgical outcomes. NSQIP collects data from patient's medical chart, as opposed to many large database studies, which utilize administrative and claims data from insurance claim forms. NSQIP data is nationally benchmarked to be comparable across various hospitals. The data are collected prospectively, validated, and submitted from multiple states across hundreds of hospitals, and each hospital submission is guided by randomization process to minimize sampling bias. There are 136 variables collected prospectively. The data include demographic information, pre-operative risk factors, intra-operative variables, and post-operative events for 30 days after their operation. All reported data were validated by vigorous quality checks by NSQIP. The full list of information collected is available at NSQIP (http://site.acsnsqip.org/wp-content/uploads/2012/03/2010-User-Guide_FINAL.pdf). The NSQIP participant use data files included 152,490 subjects in 2005 and 2006, 211,407 subjects in 2007, 271,368 subjects in 2008, 336,190 subjects in 2009, and 363,431 subjects in 2010.

Study Sample

To define our study cohort, we included patients with the Current Procedural Terminology (CPT) code for either partial or total knee arthroplasty as their principal procedure (CPT codes 27447, 27446, 27445, 27438, 27487, and 27486). There were total of 18137 entries with the listed CPT codes as principal procedure. We excluded patients with bilateral knee arthroplasty as defined by a relevant concurrent CPT code since patients undergoing bilateral knee arthroplasty have been reported to have a significantly increased risks of peri-operative complications¹⁶. We also excluded patients who received an anesthesia type other than general, spinal, or epidural. Next, we excluded patients who had pre-existing infections or infectious conditions as documented in the NSQIP database including pneumonia, systemic inflammatory response syndrome (SIRS), sepsis, septic shock, or contaminated/infected/dirty wound classification. Last, we excluded ventilator dependent patients.

Exposure variable

The Participant Use Data File coded anesthesia type into the following categories: general, epidural, spinal, regional, monitored anesthesia care, local, other, none, or unknown. In

cases where general anesthesia was used concurrently with another type of anesthesia, patients were coded as receiving general anesthesia. For the present study, we grouped patients receiving spinal and epidural anesthesia together in a single category as “neuraxial anesthesia”. All patients undergoing knee arthroplasty with anesthesia type other than general, epidural, or spinal anesthesia were excluded from the study, including patients with primary anesthesia type as regional/peripheral nerve block.

Study Variables

The NSQIP dataset contains demographic information (age, gender, height, weight, race), type of anesthesia, American Society of Anesthesiologists (ASA) Physical Status Classification, level of functional dependence prior to surgery in activities of daily living (independent, partially dependent, or totally dependent), wound classification (clean, clean/contaminated, contaminated, or dirty/infected), and comorbidities. For this study, we created variables corresponding to the following comorbidities: severe chronic obstructive pulmonary disease, congestive heart failure, coronary artery disease (defined as history of myocardial infarction, prior percutaneous coronary intervention, previous cardiac surgery, or history of angina in one month before surgery), peripheral vascular disease (defined as history of revascularization/amputation for peripheral vascular disease, or rest pain/gangrene), hypertension requiring medications, diabetes mellitus with or without insulin treatment, end stage liver disease (defined as presence of ascites and esophageal varices), kidney failure (defined as acute renal failure or currently on dialysis), central nervous system (CNS) disease (defined as impaired sensorium, coma >24 hours, history of transient ischemic attack, cerebrovascular accident/stroke with or without neurological deficit, or tumor involving CNS), spinal cord injury (defined as hemiplegia, paraplegia, or quadriplegia), and active malignancy (defined as disseminated cancer, chemotherapy or radiotherapy for malignancy).

Outcome Variables

We obtained data on eight postoperative infectious complications. These included superficial wound infection, deep incisional wound infection, organ space surgical site infection, surgical wound disruption, pneumonia, urinary tract infection, sepsis, and septic shock. Details on variable definitions are listed in table 1 as defined in NSQIP user guide. In brief, all eight variables were defined either as diagnosed by surgeon or attending physician, or on the basis of pre-defined clinical and laboratory criteria. We additionally defined a ninth composite outcome of any systemic infectious complications as the occurrence of any infection within 30 days, including all eight infection related variables as listed above.

Statistical Analysis

All data analysis was executed in STATA 12.1 statistical software (StataCorp LP, College Station, TX, USA). Initial analyses used Fisher’s exact test and chisquare test for categorical data, and Student T-test and Wilcoxon test for interval data, with statistic significance as $P < 0.05$. Our main focus was comparing infectious outcomes between NA and GA groups. For complications for which we observed statistically significant associations with anesthesia type, we next developed a multi-variable logistic regression model to control for potential confounding by pre-existing comorbidities and other patient factors. The independent variables included types of anesthesia, age, gender, race, BMI, and pre-existing comorbidities listed above. In addition, we included surgical time and year of surgery as independent variables. The surgical time was defined as total operation time in minutes. The surgical time was first log transformed to adjust the skewed distribution. We also included perioperative bleeding and transfusion as an independent variable. The perioperative bleeding and transfusion was defined as positive with either received transfusion of PRBC or whole blood from 72 hours before surgery to 72 hours postoperatively. The modeling

process used backward elimination with a threshold of $P < 0.2$ for variable inclusion. Adjusted odds ratio (OR), 95% confidence interval (CI), and P values were reported. We elected to use the word “risk” in replace of “odds” in the discussion session given the low underlying incidence rates. The adequacy of modeling was assessed by C-statistics. We used heteroskedasticity robust standard errors in all regression models¹⁷.

We also conducted regression analysis based on propensity score in order to compensate for influence from nonrandom selection of general anesthesia versus neuraxial anesthesia as previously described^{18,19}. Propensity score analysis is popular in analyzing low incident events to control for confounder²⁰. Propensity scores were calculated to reflect the probability of receiving neuraxial anesthesia with logistic regression model that included all study variables as listed above (including age, gender, BMI, race, ASA Status, level of functional status, and all comorbidities). The patient population was then stratified into five quintiles based on the propensity scores. For each significant finding from the multi-variable regression analysis, we conducted logistic regression analysis with the dependent variable, type of anesthesia, and propensity score quintiles to validate the significance.

Results

We identified 18,137 patients who underwent either partial or total knee arthroplasty. We excluded 222 (1.22%) entries with bilateral knee arthroplasty, 1095 patients with types of anesthesia other than neuraxial or general anesthesia, including regional anesthesia/peripheral nerve block (n=955), monitored anesthesia care (n=77), other (n=45), local (n=1), or unknown (n=17) anesthesia type. We further eliminated patients with pre-existing conditions, including current pneumonia (n=5), ventilator dependency (n=3), sepsis/septic shock/SIRS (n=88), and dirty/infected/contaminated wound (n=169). In the remaining cohort of 16555 patients, 9167 patients received general anesthesia (GA), 6875 patients received spinal anesthesia, and 513 patients received epidural anesthesia.

There were 7388 (44.63%) subjects in neuraxial anesthesia (NA) group and 9167 (55.37%) subjects in the general anesthesia (GA) group (Table 2). The NA group was older than the GA group. The gender distributions within each group were similar with more female patients than males. The NA group had lower BMI on average compared to GA group. Patients were predominantly white in both groups.

There were several significant differences among the prevalence of pre-existing comorbidities and physical status between GA and NA groups (Table 3). The GA group had higher incidence of pre-operative renal failure (0.35% vs 0.16%), bleeding disorders (3.41% vs 1.88%), and prior surgery within 30 days preoperatively (0.54% vs 0.22%) compared to the NA group. The NA group had more patients with partially dependent or completely dependent function status in activities of daily living prior to surgery (4.45% vs 3.69%) and more patients with hypertension requiring medication (70.36% vs 68.19%) than GA group.

The overall mortality was 0.24% and 0.15% among NA and GA subjects respectively ($P=0.214$). The NA group had statistically significant lower incidences of pneumonia (0.24% vs 0.45%, $P=0.035$) and composite systemic infection (2.95% vs 3.73%, $P=0.006$) (Table 4).

Logistic regression analysis with risk adjustment from pre-existing co-morbidities demonstrated that NA was associated with lower odds of pneumonia occurrence (OR=0.51 [95% CI 0.29–0.90]) (Table 5). The regression model for pneumonia risk included anesthesia type, age, gender, race, CNS disease, diabetes, dyspnea, and functional health status prior to surgery. The C-statistic of the regression model was 0.71. The pneumonia risk was further evaluated in logistic regression analysis with propensity score analysis, and

results were comparable (95% CI 0.29–0.92, Supplement Table 1). Logistic regression analysis of composite systemic infection also indicated that NA was associated with lower odds for any systemic infection than GA (OR=0.77 [95% CI 0.64–0.92]) (Table 6). The independent variables included in this model were anesthesia type, age, race, BMI, steroid use for chronic condition, diabetes, dyspnea, functional health status prior to surgery, ASA classification, and surgical time. The C-statistic of the regression model was 0.64. Similarly, we conducted logistic regression analysis on composite systemic infection risk with propensity quintiles, the results showed same significance (95% CI 0.64–0.92, Supplement Table 2).

Discussion

Our study using a large prospectively collected database of 16,555 knee arthroplasty patients showed that neuraxial anesthesia was associated with lower incidence of pneumonia and composite systemic infection relative to general anesthesia within 30 days post-operatively. The data suggested a 49% reduction in pneumonia risk ($P=0.021$) and 23% reduction of any systemic infectious complication risk ($P=0.004$).

Our work adds to the past literature on the comparative effectiveness of differing anesthesia types for preventing postoperative infectious complications. Rodgers et al. conducted a meta-analysis on post-operative infectious complications across a range of surgical procedures¹¹. Their analysis indicated a decreased risk of pneumonia in NA patients, with odds ratio of 0.61 (95% CI: 0.48–0.76) based on 28 clinical trials. A similar meta-analysis study by Nishimori et al.¹² also focused on pneumonia risk after abdominal aortic surgery. This analysis included seven studies and demonstrated a combined odds ratio of 0.64 (95% CI: 0.38–1.05) in favor of epidural anesthesia without statistical significance. Wu et al. conducted a retrospective study of Medicare patients and found the use of epidural analgesia may contribute to an increase in 30 days pneumonia risk with odds ratio of 1.91 (95% CI: 1.09–3.34)¹⁵. It is noteworthy that Wu et al included various surgical procedures, such as segmental excision of the lung, complete pneumonectomy, partial excision of large intestine, anastomosis of the esophagus, total knee replacement and revision, abdominal hysterectomy, pancreaticoduodenectomy, nephrectomy, cystectomy, hepatotomy/lobectomy of the liver, gastrectomy, and radical retropubic prostatectomy. The heterogeneous nature of the surgical procedures and the inclusion of both thoracic and lumbar epidural anesthesia among these procedures might partially explain the differences between the results of the current study and Wu et al¹⁵.

There have been very few past studies examining the association of anesthesia type with postoperative infectious complications among patients undergoing orthopedic procedures^{13,14,21}. A meta-analysis by Parker et al. of nine clinical trials with a total of 1125 patients did not show any reduction in pneumonia risk by NA in hip fracture surgery in adults¹³. The differences between our findings and those of Parker et al. may be explained in part by clinical differences between hip fracture patients and patients undergoing elective hip or knee arthroplasty. Another meta-analysis on total hip and knee replacement surgery had insufficient power with respect to infection with combined sample size of 204 subjects¹⁴. Chang et al. conducted a retrospective study on 3081 patients with total hip or knee replacement between 2002 and 2006, and concluded that general anesthesia was associated with a greater odds of surgical site infection compared to epidural or spinal anesthesia²¹. We examined four surgical-site related infection variables, including superficial surgical site infection, deep incisional surgical site infection, organ space surgical site infection, and surgical wound disruption. However, none differed significantly between general anesthesia and neuraxial anesthesia. Notably, the overall incidence of surgical site infections was lower in our study than in the study of Chang et al. It is unclear of whether

this is due to the fact that our study only included knee replacement, or that our patients are from more recent years than included in this earlier work. Our study suggested that spinal or epidural anesthesia decreased pneumonia risk among knee arthroplasty by about 49%. This observation could be attributed to less systemic immune disturbance in the NA group comparing to GA. In addition, the potential absence of airway instrumentation and mechanical ventilation in the NA group might also be an important confounder as GA is related to elevated risks of airway trauma and atelectasis.

We acknowledge that the retrospective observational nature is a limitation of this study. Future prospective randomized studies are needed to confirm our findings. Another limitation of the current study lies in the coding system of anesthesia type. In case of concurrent use of general anesthesia with neuraxial anesthesia, patients were coded as receiving general anesthesia. There is no detail information on general anesthesia, such as with endotracheal tube, LMA, or others, which could contribute to the differences in outcome complications. This study is also limited by the number of variables listed in the NSQIP database. There are other important risk factors that might be important for infectious risk, such as hypothermia, level of immediate postoperative mental status, and etc. However, there is no information in the NSQIP database. The current study included patients from 2005 to 2010, during which prophylactic antibiotic regime might change over time, which could affect the infectious outcomes. However, there is no information in the NSQIP database on prophylactic antibiotic regime. The hospital identification and surgeon identification information were also removed by NSQIP in order to comply with participation agreement between NSQIP and participating sites. Furthermore, information on postoperative analgesia techniques was missing from NSQIP dataset, which may affect the surgical outcomes.

Nonetheless, our study suggests that NA may offer advantages over GA for knee arthroplasty by reducing the likelihood of selected infectious complications, especially pneumonia. While this study confirmed our hypothesis clinically, it does not provide explanations regarding the mechanisms underlying the associations we observed here. Studies on immune response to various anesthesia techniques have been inconclusive due to the heterogeneity among subjects.

In conclusion, our study demonstrated that neuraxial anesthesia was associated with a lower incidence of both pneumonia and any systemic infectious complication within 30 days of surgery compared to general anesthesia in patients undergoing knee arthroplasty. While neuraxial anesthesia is advantageous from 30-day post-operative infection aspect, the underlying mechanism is still unknown. As such, further prospective investigations are needed to confirm our findings.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Funding:

Departmental career developmental fund to JL. Department of Anesthesiology and Critical Care, the Perelman School of Medicine, the University of Pennsylvania, Philadelphia, PA, USA 19104.

References

1. Cram P, Lu X, Kates SL, Singh JA, Li Y, Wolf BR. Total knee arthroplasty volume, utilization, and outcomes among Medicare beneficiaries, 1991–2010. *JAMA*. 2012; 308:1227–1236. [PubMed: 23011713]
2. Cram P, Vaughan-Sarrazin MS, Wolf B, Katz JN, Rosenthal GE. A comparison of total hip and knee replacement in specialty and general hospitals. *J Bone Joint Surg Am*. 2007; 89:1675–1684. [PubMed: 17671004]
3. Gottschalk A, Sharma S, Ford J, Durieux ME, Tiouririne M. Review article: the role of the perioperative period in recurrence after cancer surgery. *Anesth Analg*. 2010; 110:1636–1643. [PubMed: 20435944]
4. Kennedy BC, Hall GM. Neuroendocrine and inflammatory aspects of surgery: do they affect outcome? *Acta Anaesthesiol Belg*. 1999; 50:205–209. [PubMed: 10603996]
5. Rosendahl W, Schulz U, Teufel T, Irtel von Brenndorf C, Gupta D. Surgical stress and neuroendocrine responses in infants and children. *J Pediatr Endocrinol Metab*. 1995; 8:187–194. [PubMed: 8521193]
6. Wada H, Seki S, Takahashi T, Kawarabayashi N, Higuchi H, Habu Y, Sugahara S, Kazama T. Combined spinal and general anesthesia attenuates liver metastasis by preserving TH1/TH2 cytokine balance. *Anesthesiology*. 2007; 106:499–506. [PubMed: 17325508]
7. Liu SS, Block BM, Wu CL. Effects of perioperative central neuraxial analgesia on outcome after coronary artery bypass surgery: a meta-analysis. *Anesthesiology*. 2004; 101:153–161. [PubMed: 15220785]
8. Tonnesen E, Wahlgreen C. Influence of extradural and general anaesthesia on natural killer cell activity and lymphocyte subpopulations in patients undergoing hysterectomy. *Br J Anaesth*. 1988; 60:500–507. [PubMed: 3377925]
9. Tverskoy M, Cozakov C, Ayache M, Bradley EL Jr, Kissin I. Postoperative pain after inguinal herniorrhaphy with different types of anesthesia. *Anesth Analg*. 1990; 70:29–35. [PubMed: 2297102]
10. Woolf CJ, Salter MW. Neuronal plasticity: increasing the gain in pain. *Science*. 2000; 288:1765–1769. [PubMed: 10846153]
11. Rodgers A, Walker N, Schug S, McKee A, Kehlet H, van Zundert A, Sage D, Futter M, Saville G, Clark T, MacMahon S. Reduction of postoperative mortality and morbidity with epidural or spinal anaesthesia: results from overview of randomised trials. *BMJ*. 2000; 321:1493. [PubMed: 11118174]
12. Nishimori M, Ballantyne JC, Low JH. Epidural pain relief versus systemic opioid-based pain relief for abdominal aortic surgery. *Cochrane Database Syst Rev*. 2006:CD005059. [PubMed: 16856074]
13. Parker MJ, Handoll HH, Griffiths R. Anaesthesia for hip fracture surgery in adults. *Cochrane Database Syst Rev*. 2004:CD000521. [PubMed: 15494999]
14. Choi PT, Bhandari M, Scott J, Douketis J. Epidural analgesia for pain relief following hip or knee replacement. *Cochrane Database Syst Rev*. 2003:CD003071. [PubMed: 12917945]
15. Wu CL, Hurley RW, Anderson GF, Herbert R, Rowlingson AJ, Fleisher LA. Effect of postoperative epidural analgesia on morbidity and mortality following surgery in medicare patients. *Reg Anesth Pain Med*. 2004; 29:525–533. [PubMed: 15635510]
16. Stundner O, Chiu YL, Sun X, Mazumdar M, Fleischut P, Poultides L, Gerner P, Fritsch G, Memtsoudis SG. Comparative perioperative outcomes associated with neuraxial versus general anesthesia for simultaneous bilateral total knee arthroplasty. *Reg Anesth Pain Med*. 2012; 37:638–644. [PubMed: 23080348]
17. White H. A Heteroskedasticity-Consistent Covariance Matrix Estimator and a Direct Test for Heteroskedasticity. *Econometrica*. 1980; 48:817–838.
18. Rosenbaum PR, Rubin RD. The central role of the propensity score in observational studies for causal effects. *Biometrika*. 1983; 70:41–55.
19. Joffe MM, Rosenbaum PR. Invited commentary: propensity scores. *Am J Epidemiol*. 1999; 150:327–333. [PubMed: 10453808]

20. Cepeda MS, Boston R, Farrar JT, Strom BL. Comparison of logistic regression versus propensity score when the number of events is low and there are multiple confounders. *Am J Epidemiol*. 2003; 158:280–287. [PubMed: 12882951]
21. Chang CC, Lin HC, Lin HW, Lin HC. Anesthetic management and surgical site infections in total hip or knee replacement: a population-based study. *Anesthesiology*. 2010; 113:279–284. [PubMed: 20657202]

Table 1

NSQIP Definition of Eight Major Infection Outcome Variables Adapted from NSQIP User Guide.

Variable Label	Variable Definition
Superficial surgical site infection	Infection occurs within 30 days after the operation and the infection involves only skin or subcutaneous tissue of the incision and at least one of the following: purulent drainage, organisms isolated, or at least one of the following (pain or tenderness, localized swelling, redness, or heat)
Deep Incisional surgical site infection	Infection occurs within 30 days after the operation and the infection appears to be related to the operation involves deep soft tissues of the incision and at least one of the following: purulent drainage, spontaneously dehisces or is deliberately opened by a surgeon, an abscess or other evidence of infection.
Organ Space surgical site infection	Infection occurs within 30 days after the operation and the infection appears to be related to the operation and the infection involves any part of the anatomy (e.g., organs or spaces), other than the incision, which was opened or manipulated during an operation and at least one of the following: purulent drainage from a drain, organisms isolated, or abscess or other evidence of infection involving the organ/space.
Surgical Wound Disruption	Separation of the layers of a surgical wound, which may be partial or complete, with disruption of the fascia within 30 days of the operation.
Pneumonia	<p>Meet one of the following TWO criteria within 30 days of the operation:</p> <p>Criterion 1: Rales or dullness to percussion on physical examination of chest AND any of the following: new onset of purulent sputum, organism isolated from blood culture, tracheal, or bronchial specimen.</p> <p>Criterion 2: Chest radiography evidence AND any of the following: new onset of purulent sputum, organism isolated (from blood culture, tracheal, bronchial specimen, or in respiratory secretions), antibody titer increase for pathogen, or histopathologic evidence of pneumonia</p>
Urinary Tract Infection	<p>Meet one of the following TWO criteria within 30 days of the operation:</p> <ol style="list-style-type: none"> 1 One of the following: fever, urgency, frequency, dysuria, suprapubic tenderness AND positive urine culture. 2 Two of the following: fever, urgency, frequency, dysuria, suprapubic tenderness AND any of the following: dipstick test positive for leukocyte esterase and/or nitrate, pyuria, organisms seen on Gram stain of unspun urine.
Sepsis	Report this variable if the patient has two of the following clinical signs and symptoms: Temp >38 C or <36 C, HR >90 bpm, RR >20/min or PaCO ₂ <32 mmHg, WBC >12,000 (or <4000) cells/mm ³ , >10% immature forms, anion gap acidosis AND one of the following: positive blood culture, or clinical documentation of purulence or positive culture.
Septic Shock	Clinical signs and symptoms of SIRS or sepsis AND documented organ and/or circulatory dysfunction. Examples of organ dysfunction include: oliguria, acute alteration in mental status, acute respiratory distress. Examples of circulatory dysfunction include: hypotension, requirement of inotropic or vasopressor agents.

http://site.acsnsqip.org/wp-content/uploads/2012/03/2010-User-Guide_FINAL.pdf

Table 2

Patient Demographic Information Summary by Anesthesia Type.

	General Anesthesia		Neuraxial Anesthesia		P-value
	N/Mean	%/SD	N/Mean	%/SD	
Total Subjects (N)	9167	55.37%	7388	44.63%	
Age:	66.41 ±	10.61	67.51 ±	10.17	<0.0001
Gender: Male	3262	35.69%	2730	37.00%	0.084
Female	5878	64.31%	4649	63.00%	
Unknown	27		9		
BMI:	32.83 ±	7.97	32.36 ±	7.29	0.0001
Race: White	7311	82.82%	5691	82.85%	<0.001
Black	614	6.96%	572	8.33%	
Hispanic	727	8.24%	427	6.22%	
Others	176	1.99%	179	2.61%	
Unknown	339		519		

All values were reported as mean ± SD, or percentage. The unit of measure for BMI is kilogram/meter². "Other race" included patients listed as Native Hawaiian or Pacific Islander, Asian or Pacific Islander, Asian, and American Indian or Alaska Native in the NSQIP database.

Table 3

Prevalence of Pre-existing Comorbidities.

Comorbidity	General Anesthesia		Neuraxial Anesthesia		P-value
	N	%	N	%	
ASA classification					
4-Life Threat	179	1.96	128	1.74	0.291
3-Severe Disturbance	4449	48.67	3543	48.03	
2-Mild Disturbance	4340	47.48	3583	48.58	
1-No Disturbance	173	1.89	122	1.65	
Functional health status					
Prior to Surgery					
Dependent	6	0.07	11	0.15	0.020
Partially Dependent	332	3.62	318	4.30	
Independent	8828	96.31	7059	95.55	
Congestive heart failure (CHF) in 30 days before surgery	17	0.19	7	0.09	0.152
Coronary Artery Disease	1011	11.03	776	10.51	0.290
Peripheral Vascular Disease	61	0.67	48	0.65	0.923
HTN requiring medication	6251	68.19	5198	70.36	0.003
Dyspnea					
At Rest	26	0.28	24	0.32	0.350
Moderate Exertion	872	9.51	657	8.89	
History of severe COPD	338	3.69	252	3.41	0.354
End Stage Liver Disease	5	0.05	1	0.01	0.235
Renal Failure	32	0.35	12	0.16	0.022
Diabetes mellitus with oral agents or insulin	409	4.46	291	3.94	0.164
Insulin					
Oral/Non-insulin	1267	13.82	1064	14.40	
None	7491	81.72	6033	81.66	
Central Nervous System Disease	535	5.84	417	5.65	0.615
Spinal Cord Injury	42	0.46	23	0.31	0.169
Disseminated cancer, Chemotherapy/Radiotherapy	37				
Bleeding disorders	0.40	23	0.31	0.364	
	313	3.41	139	1.88	<0.001
Prior Operation within 30 days	49	0.54	16	0.22	0.001

ASA: American Society of Anesthesiologists; HTN: Hypertension; COPD: Chronic Obstructive Pulmonary Disease.

Table 4

Incidences of 30-days Post-operative Complications.

Variable Label	General Anesthesia		Neuraxial Anesthesia		P-value
	N	%	N	%	
Superficial Surgical Wound Infection	79	0.86	55	0.74	0.433
Deep Incisional Wound Infection	20	0.22	15	0.20	0.867
Organ/Space Surgical Site Infection	20	0.22	14	0.19	0.733
Surgical Wound Disruption	24	0.26	21	0.28	0.881
Pneumonia Occurrences	41	0.45	18	0.24	0.035
Urinary Tract Infection Occurrences	154	1.68	97	1.31	0.055
Sepsis Occurrences	47	0.51	24	0.32	0.073
Septic Shock Occurrences	11	0.12	8	0.11	1
Composite Systemic Infection	342	3.73	218	2.95	0.006

Table 5

Multivariable Regression Analysis, ORs, and 95% CI with Pneumonia Occurrence as Dependent Variable.

Independent Variable	Odds Ratio	95% CI	P-value
Anesthesia Type	0.51	(0.29 – 0.90)	0.021
Age	1.66	(1.31 – 2.11)	<0.001
Gender	0.53	(0.31 – 0.89)	0.016
Race – other	5.39	(2.14 – 13.57)	<0.001
Central Nervous System Disease	1.84	(0.89 – 3.82)	0.099
Dyspnea – Moderate Exertion/At Rest	1.98	(0.99 – 3.95)	0.052
Diabetes – Insulin Dependent	2.58	(1.14 – 5.85)	0.023
Functional health status Prior to Surgery – Partially Dependent/Dependent	2.00	(0.80 – 4.99)	0.139

Odds Ratio and 95% CI were reported to two digits, while P-values were reported to three digits. We also analyzed potential interactions between anesthesia type and all other independent variables via univariable regression analysis, and no significant interactions were identified (all P>0.13).

Table 6

Multivariable Regression Analysis, ORs, and 95% CI with Composite Systemic Infection as Dependent Variable.

Independent Variable	Odds Ratio	95% CI	P-value
Anesthesia Type	0.77	(0.64 – 0.92)	0.004
Age	1.23	(1.12 – 1.35)	<0.001
Race –Hispanic	0.70	(0.48 – 1.03)	0.072
Race –Other	1.82	(1.13 – 2.95)	0.014
Body Mass Index	1.02	(1.01 – 1.03)	0.001
Steroid Use for Chronic Condition	1.71	(1.11 – 2.63)	0.015
Diabetes – Insulin Dependent	1.54	(1.11 – 2.15)	0.011
Dyspnea – Moderate Exertion/At Rest	1.22	(0.94 – 1.58)	0.131
Functional health status Prior to Surgery – Partially Dependent/Dependent	1.35	(0.94 – 1.95)	0.106
ASA Classification: 3-Severe Disturbance	1.52	(1.25 – 1.85)	<0.001
ASA Classification: 4-Life Threatening Concurrent Disease	1.80	(1.09 – 2.99)	0.022
Operation Year	0.92	(0.84 – 1.01)	0.065
Surgical Time (log[minutes])	1.94	(1.52 – 2.48)	<0.001

Odds Ratio and 95% CI were reported to two digits, while P-values were reported to three digits. Odds ratio for Surgical Time is per unit of the transformed time (log[minutes]).