CORRESPONDENCE

Adverse Drug Events in Older Patients Admitted as an Emergency: The Role of Potentially Inappropriate Medication in Elderly People (PRISCUS)

by PD Dr. med. Harald Dormann, Anja Sonst, Dr. med. Fabian Müller, et al. in volume 13/2013

Study Design

The primary objective of the study—to clarify whether PRISCUS list medications are more commonly associated with adverse drug reactions (ADRs) than other medications—has been confirmed by the authors. But this seems not convincing due to the questionable primary goal, the preferred study design and some classifications being missing or misleading.

Regarding the primary goal of the study: If the intention had been to provide an answer to the primary question it would have sufficed to study the frequency and severity of ADRs after administration of PRISCUS drugs (PDs) versus non-PRISCUS drugs in patients older than 65 with identical indications. In addition to the over-65s (n=351) however, 401 subjects younger than 65 were included. If the intention was to find an answer to the question of whether PDs are less well tolerated by older patients than by younger patients, then this comparison would have been the correct one to confirm the statement that older patients react more sensitively to PDs than younger ones—the indication being the same, obviously.

Regarding the study design: The authors favored accidentally collected data from a cohort of patients in an emergency unit over a prospective, randomized controlled comparative study. The populations (under comparison) were of different sizes and inhomogeneous: in patients aged 65 or older, the most commonly observed disorders were circulatory disorders (35%) and gastrointestinal disorders (10.5%); In patients younger than 65, circulatory disorders were present only in 17.7% of cases, and psychological/behavioral disorders occurred in 10.2%. This means that medication therapy would be different in both compared groups, something that was not explained anywhere in the article. In total, only 149 patients with ADRs were evaluated.

Regarding the classification: Medication errors (caused by the physician or non-compliance) cannot be subsumed into ADRs. In order to classify ADRs, information on severity/risk is required, especially for the comparison between PRISCUS drugs and non-PRISCUS drugs. The category “unexpected ADRs” is lacking. Did such reactions not occur?

Of interest, albeit unmentioned, is the fact that in the patient population under study, medication errors occurred in more patients (n=166) than adverse drug reactions (n=149). A further surprising result (disconcerting aspect) is (in) the drastic increase in medication errors in 90 patients between the age of 70 and 89.

In conclusion: Medication therapy will have to be improved, especially for populations at risk. DOI: 10.3238/arztebl.2013.0674a

REFERENCES

1. Dormann H, Sonst A, Müller F, et al.: Adverse drug events in older patients admitted as an emergency—the role of potentially inappropriate medication in elderly people (PRISCUS). *Dtsch Arztebl Int* 2013; 110(13): 213–9.

Dr. med. Karla Lehmann
Kottmar OT Walddorf
Karla.Lehmann@t-online.de

In Reply:

The authors thank Dr Lehmann for her contribution.

Regarding the study question: The analysis we conducted of the risk potential of PRISCUS drugs is a subanalysis of a study that included more than 2200 patients in an emergency unit prospectively in an intensive drug safety test. The data on younger patients were reported only for completeness's sake. As Dr Lehmann rightly comments, the PRISCUS-independent comparison of younger and older patients shows an increase in medication errors and adverse drug reactions in older patients.

Regarding the study design: The design of a prospective, randomized, placebo-controlled study (RCT) was not feasible for our research question. By means of an intensified drug safety test conducted by two independent expert teams and an additional sample of the patient files that were primarily assessed to be normal, a new gold standard in the clinical evaluation of adverse drug reactions was defined. We wish to point out that it was not the aim of our study to evaluate the safety of the PRISCUS drugs compared with possible alternative preparations. An RCT would formally be better in that setting, albeit difficult to implement in practice.

Regarding the classification: Only 52 of the 166 medication errors manifested clinically. In our analyses we distinguish between many events on the basis of medication errors and adverse effects that occur even when the drug is taken as instructed.

In the context of the present study we established that the classifications of medication errors and adverse drug reactions as used up to now are coming up against their limitations, if several drugs contribute to an adverse reaction and medication errors are observed additionally (2).

The authors agree with Dr Lehmann that medication therapy requires particular attention especially in at-risk populations such as elderly and multimorbid patients. The focus should therefore not be exclusively on PRISCUS drugs. DOI: 10.3238/arztebl.2013.0674b

REFERENCES

1. Dormann H, Sonst A, Müller F, Vogler R, et al.: Adverse drug events in older patients admitted as an emergency—the role of potentially inappropriate medication in elderly people (PRISCUS.) *Dtsch Arztebl Int* 2013; 110(13): 213–9.
2. Bürkle T, Müller F, Patapovas A, et al.: A new approach to identify, classify and count drug-related events. *Br J Clin Pharmacol* 2013; 76 (Suppl S1): 56–68.

Prof. Dr. med. Harald Dormann
Klinikum Fürth
Harald.Dormann@klinikum-fuerth.de

Conflict of interest statement

All authors declare that no conflict of interest exists.