Published in final edited form as:

J Gastrointest Surg. 2008 May ; 12(5): 783–794. doi:10.1007/s11605-007-0391-9.

Quality of Life in Pancreatic Cancer: Analysis by Stage and Treatment

Stefano Crippa,

Wang Ambulatory Care Center 460, Department of Surgery, Massachusetts General Hospital, Harvard Medical School, 15 Parkman Street, Boston, MA 02114, USA

Ismael Domínguez,

Wang Ambulatory Care Center 460, Department of Surgery, Massachusetts General Hospital, Harvard Medical School, 15 Parkman Street, Boston, MA 02114, USA

J. Ruben Rodríguez,

Wang Ambulatory Care Center 460, Department of Surgery, Massachusetts General Hospital, Harvard Medical School, 15 Parkman Street, Boston, MA 02114, USA

Oswaldo Razo,

Wang Ambulatory Care Center 460, Department of Surgery, Massachusetts General Hospital, Harvard Medical School, 15 Parkman Street, Boston, MA 02114, USA

Sarah P. Thayer,

Wang Ambulatory Care Center 460, Department of Surgery, Massachusetts General Hospital, Harvard Medical School, 15 Parkman Street, Boston, MA 02114, USA

David P. Ryan,

Wang Ambulatory Care Center 460, Department of Oncology, Massachusetts General Hospital, Harvard Medical School, 15 Parkman Street, Boston, MA 02114, USA

Andrew L. Warshaw, and

Wang Ambulatory Care Center 460, Department of Surgery, Massachusetts General Hospital, Harvard Medical School, 15 Parkman Street, Boston, MA 02114, USA

Carlos Fernández-del Castillo

Wang Ambulatory Care Center 460, Department of Surgery, Massachusetts General Hospital, Harvard Medical School, 15 Parkman Street, Boston, MA 02114, USA

Carlos Fernández-del Castillo: cfernandez@partners.org

Abstract

In pancreatic cancer patients, survival and palliation of symptoms should be balanced with social and functional impairment, and for this reason, health-related quality of life measurements could play an important role in the decision-making process. The aim of this work was to evaluate the quality of life and survival in 92 patients with different stages of pancreatic adenocarcinoma who underwent surgical and/or medical interventions. Patients were evaluated with the Functional Assessment of Cancer Therapy questionnaires at diagnosis and follow-up (3 and 6 months). At diagnosis, 28 patients (30.5%) had localized disease (group 1) and underwent surgical resection, 34 (37%) had locally advanced (group 2), and 30 (32.5%) metastatic disease (Group 3).

Improvement in quality of life was found in group 1, while in group 3, it decreased at follow-up ($p = 0.03$). No changes in quality of life in group 2 were found. Chemotherapy/chemoradiation seems not to significantly modify quality of life in groups 2 and 3. Median survival time for the entire cohort was 9.8 months (range, 1–24). One-year survival was 74%, 30%, and 16% for groups 1, 2, and 3 respectively ($p = 0.001$). Pancreatic cancer prognosis is still dismal. In addition to long-term survival benefits, surgery impacts favorably quality of life.

Keywords

Pancreatic cancer; Quality of life; Meaningful important difference; Palliation; Survival

Introduction

Pancreatic cancer remains the fourth leading cause of cancer-related death in the USA. Of the 32,180 patients diagnosed with pancreatic adenocarcinoma in 2006, the great majority will die within 2 years from initial diagnosis.¹ Less than 20% of the patients are candidates for surgical resection, which remains the only treatment offering the possibility of long-term survival, even if this is only 10% to 25%.^{2–4} The remaining patients present with locally advanced or metastatic disease, and for them, chemotherapy or chemoradiation represent the current standard of care with the aim of improving survival.^{5–7}

During the course of disease, 70% to 80% of patients with pancreatic head tumors develop obstructive jaundice, and 10% to 20% duodenal obstruction.^{8–10} Many also will manifest pain, which is probably the most disturbing and incapacitating symptom in advanced pancreatic cancer.¹¹ Adequate palliation of biliary and duodenal obstruction, as well as of pain, is one of the goals in the treatment of these patients. However, optimal methods of palliation remain controversial in terms of patient benefit perception and durability.^{12–15} Many studies have compared different treatments and palliative procedures in different stages of the disease, but few have considered these issues among the entire spectrum of pancreatic cancer patients.^{4,16}

To assess the value of a treatment, physicians routinely use “physician-centered” objective outcomes, such as disease recurrence, complications, treatment toxicity, or survival, but infrequently consider patients’ perception and quality of life.¹⁷ Health-related quality of life (HQOL) seeks to measure the impact of disease process on physical, psychological, and social aspects of the person’s life and feeling of well-being,^{18–20} and recently, has become an important subject in pancreatic cancer care, with the aim of measuring the impact of different interventions on patients’ health and life.^{18,21–24} It has become clear that in a disease-like pancreatic cancer, in which patients have a short life expectancy, improvements in survival and treatment-related complications must be carefully balanced against HQOL outcomes to define better approaches while considering patients’ personal needs.

The aim of this prospective study is to analyze the effects of therapeutic and palliative treatments on health outcomes and HQOL in a contemporary cohort of patients with pancreatic cancer.

Patients and Methods

After obtaining Institutional Review Board approval, 105 patients with histologically proven ductal adenocarcinoma of the pancreas, treated at Massachusetts General Hospital between September 2004 and January 2006, were enrolled in this study after obtaining informed consent. Demographics, clinical presentation, laboratory and radiologic findings, type of surgery, postoperative morbidity and mortality, pathology, neoadjuvant or adjuvant

treatments, type of palliation, disease recurrence, and number of readmissions were recorded. Patients with adenocarcinoma arising in intraductal papillary mucinous neoplasm of the pancreas were excluded.

Patients were classified in three clinical stages: apparently localized cancer (group 1), locally advanced (group 2), and metastatic (group 3). Localized cancer permitted resection. Locally advanced cancer was defined as extension of the neoplasm outside the pancreas with major vascular encasement or other features precluding potentially curative resection. Local recurrence was defined as recurrent retroperitoneal mass or regional lymph nodes in patients who had undergone pancreatic resection with curative intent. Metastases were defined as a relapse of disease in the peritoneal cavity or at any distant site.

Follow-up data was obtained through direct contact with patients' oncologists, primary care physicians, and families.

Quality of Life Assessment

To assess health-related quality of life, the fourth version of the Functional Assessment of Cancer Therapy (FACT-G) and the hepatobiliary and pancreatic specific module (FACT-Hep) were used. The FACT-G is a 27-item self-report instrument that assesses four different dimensions of quality of life: physical (seven items), social (seven items), emotional (six items), and functional (seven items). The specific subscale for hepatobiliary and pancreatic diseases (FACT-Hep) has 18 additional items. Patients responded to HQOL questions on a five-point Likert-type scale ranging from 0 (not at all) to 4 (very much). FACT-G and FACT-Hep have been previously validated in cancer populations.^{25–27}

From these two questionnaires, three scores were obtained: the FACT-G score, which is the sum of the four subscales; the FACT-Hep, which is the sum of the FACT-G and the disease-specific module (FACT-Hep), and the Trial Outcome Index (TOI), which is the sum of the physical, functional and disease-specific module. The TOI has been demonstrated to be a sensitive indicator of clinical outcome.²⁷

Each questionnaire was applied at diagnosis (baseline) and sent by mail after 3 and 6 months from the initial diagnosis.

Data Analysis

Descriptive statistics were used to summarize demographic and clinical characteristics. We performed paired and unpaired *t* student and χ^2 tests when comparing nominal and categorical variables, respectively. In the case of a nonparametric distribution, a Mann–Whitney *U* was done. One-way ANOVA with post hoc analysis was performed for multiple comparisons with normal distribution, and the probability was adjusted by Tukey's correction. A 5% of significance was accepted. Survival curves were constructed with the Kaplan–Meier method.

HQOL scores were analyzed from a statistical and clinical significance viewpoint. For the statistical analysis, due to a lack of complete sets of questionnaires in some patients, a cross-sectional analysis was done, and all patients who provided HQOL questionnaires at baseline were compared with all those provided follow-up HQOL data. Thus, scores available at 3- and 6-month follow-up were grouped together as “follow-up scores.” For the purpose of the clinical analysis, scores available at 3- and 6-month were considered separately. HQOL scores were compared among the three groups at baseline and follow-up and within each group (baseline versus follow-up scores). Moreover, HQOL scores were compared in the entire cohort ($n = 92$) or in groups 2 and 3 patients ($n = 64$) to evaluate differences in the following independent variables: age, resection, CA 19.9 level, adjuvant/neoadjuvant

treatment in the entire cohort, and need for stents, chemotherapy/ chemoradiation, and celiac block in groups 2 and 3.

Clinical significance was determined using the meaningful important difference (MID). MID is defined as “the smallest difference in score in the outcome of interest that informed patients perceive as important, either beneficial or harmful, and that would lead the patient or clinician to consider a change in the management”.^{27,28} MIDs measure the clinical relevant changes in HQOL perception derived from a treatment or the disease itself. To estimate MID, the mean of all HQOL scores (FACT-G, FACT-Hep, TOI) were first assessed, and subsequently, differences between mean HQOL scores at baseline and at 3- and 6-months were evaluated. Differences in mean HQOL scores considered clinically significant (MIDs) were as follows: 8–9 for FACT-Hep, 7–8 for FTOI and 6–7 for FACT G.²⁷

Results

Of the 102 patients initially enrolled, ten with no complete HQOL questionnaires at baseline were excluded, and therefore, final analysis was performed in 92 patients (46 women and 46 men; mean age $\pm$ SD of 66 ± 10 years; range, 42–88).

At diagnosis, 28 patients (30.5%) had localized disease (group 1), 34 (37%) had locally advanced disease (group 2), and 30 (32.5%) had metastatic disease (group 3).

Demographics

Demographic characteristics of the three groups are shown in Table 1. There were no differences among the three groups with regard to age, sex, and tumor location. Jaundice was the most common presenting symptom in group 1 (86% of patients), while its frequency was lower in groups 2 and 3 (21% and 14%, respectively; $P = 0.0001$). Abdominal pain and weight loss were more likely to be associated with locally advanced or metastatic disease. As expected, median CA 19.9 was significantly higher in patients with metastatic disease.

Diagnostic and Palliative Procedures

Table 2 shows diagnostic and palliative procedures as well as data regarding hospital stay and readmissions. All the patients underwent at least one computed tomography (CT) scan. The mean number of CTs performed per patient was significantly greater in those with localized disease. The number of endoscopic ultrasounds (EUS), endoscopic retrograde cholangiopancreatography (ERCP) and endoscopically placed biliary stents was greater in group 2 patients, but the difference was significant only when comparing EUS between groups 2 and 3 (83% versus 53%, $P = 0.01$). Only three patients (10%) in groups 3 and two (6%) in group 2 required a percutaneous transhepatic biliary drainage during their clinical course. No patient in group 1 underwent percutaneous or intraoperative celiac block compared to 14 patients (41%) in group 2 and 6 (20%) in group 3. Enteral stents were used to palliate malignant duodenal obstruction in six patients (18%) in group 2 and in two patients (7%) in group 3. Four patients in group 2 and one in group 3 had both enteral and biliary stents.

The overall mean hospital stay and the number of readmissions were significantly shorter in patients with metastatic disease. Fifty-six percent of patients with locally advanced disease required two or more readmissions, compared to 36% in group 1 and 27% in group 3. There were no differences between groups 2 and 3 in the number of surgical palliative procedures. Five patients with locally advanced disease received intraoperative radiation therapy (IORT) in association with palliative surgery.

Treatment

Twenty-nine patients underwent pancreatic resection with curative intent. Of these, 28 patients had localized disease at diagnosis, and one patient had locally advanced disease and underwent resection after chemoradiation. Table 3 shows the intraoperative, postoperative, and pathologic features of these patients. Overall morbidity was 24%, mortality was nil, and no patient required surgical re-exploration.

Of the 29 resected patients, four underwent neoadjuvant chemoradiation (three patients of group 1 and one of group 2), and 21 underwent adjuvant chemoradiation. Of these patients, ten received gemcitabine, and the remaining, 5-fluorouracil. Four patients declined further treatment.

With regard to the 34 patients in group 2, 18 (53%) underwent chemoradiation (including the five who received also IORT), 11 (32%) chemotherapy alone, and five (15%) refused any treatment or were considered unsuitable for chemotherapy or chemoradiation because of major comorbidities. In group 3 ($n = 30$), 24 patients (80%) underwent chemotherapy, three (10%) chemoradiation, and three (10%) refused treatment. In patients who underwent chemotherapy alone, gemcitabine was administered as single agent in 14 cases, and in association with new agents, as part of clinical trials in the remaining ten patients.

Survival

All but two patients were followed until death or at least 12 months. The median survival for the entire cohort was 9.8 months (mean $\pm$ SD, 9.5 ± 6 ; range, 1–24). Figure 1a shows survival of the entire cohort, and Fig. 1b the survival of the three different groups. One- and 2-year survival for the entire cohort was 39% and 23%. One-year survival was 74%, 30%, and 16% for groups 1, 2, and 3, respectively ($P = 0.001$). The median survival for group 3 was 5.8 months, for group 2 8.6 months, and for group 1, it has not been reached. With regard to the 29 patients who underwent surgical resection, 12 of them developed tumor recurrence at a median time of 7 months from operation. Sites of recurrence were distant metastases in eight patients and locoregional in four. Ten patients died of disease while two remain alive with disease. The remaining 17 patients are alive with no evidence of disease at a median of 14 months (range, 12–23).

The mean survival for the five patients who underwent IORT was 10.6 ± 6.2 months. Two of these patients died for tumor progression after 8.8 and 8.6 months, while the remaining patients are alive with stable disease.

Quality of Life

Baseline questionnaires were completed in 92 patients (100%). The rate of completed questionnaires decreased to 56% at 3 months and to 48% at 6 months. Excluding dead patients, the rate of completed questionnaires was 63% at 3 months and 59% at 6 months.

Table 4 shows HQOL scores at baseline and during follow-up. At baseline, no statistical differences were found in the HQOL scores among the three groups, while during follow-up, patients in group 1 had higher HQOL scores compared to groups 2 and 3. Comparisons within each group showed an improvement of HQOL scores from baseline to follow-up in groups 1 and 2, and a worsening of all scores in group 3.

Clinically meaningful changes (MIDs) from baseline to 3- and 6-months were found in groups 1 and 3 but not in group 2 (Fig. 2). The MIDs in group 1 were toward improvement and in group 3 toward deterioration.

Since HQOL scores at baseline were homogeneous among the three groups, we performed statistical analysis of HQOL scores considering different variables in the entire cohort ($n = 92$; Table 5). Patients who underwent resection ($n = 29$) had significantly higher HQOL scores at follow-up compared to nonresected ones ($n = 63$; $P = 0.001$). Patients with CA 19-9 values above 200 U/l ($n = 46$) had lower HQOL scores both at baseline ($P = 0.01$) and during follow-up ($P = 0.04$). There were no statistically significant differences in HQOL scores regarding age (cut-off, 60 years) and chemotherapy/chemoradiation both at baseline and during follow-up.

Patients in groups 2 and 3 who required biliary or enteral stents, celiac block, and chemoradiation/chemotherapy alone were analyzed from a clinical viewpoint (Figs. 3, 4, and 5). Patients with locally advanced or metastatic disease who underwent stent placement had a decrease in HQOL scores at 3 months but a clinically significant recovery at 6 months, whereas patients without stents did not have significant changes in HQOL mean scores at 3 and 6 months (Fig. 3). Patients who underwent chemoradiation showed no significant differences in HQOL scores at 3 and 6 months, whereas those who received chemotherapy alone presented a significant decrease in HQOL score at 3 months with a nonsignificant improvement at 6 months (Fig. 4).

Patients who underwent celiac block had a decrease in HQOL score at 3 months and a significant recovery at 6 months. No changes in quality of life were found in patients without celiac block from baseline to follow-up (Fig. 5).

The statistical analysis of these variables in groups 2 and 3 patients was not significant.

Discussion

“The outlook in carcinoma of the pancreas continues to be grim.” With this peremptory sentence, Morrow and colleagues summarized the 1975–1980 experience of Memorial Sloan–Kettering Cancer Center in treating 231 patients with ductal adenocarcinoma of the pancreas.²⁹ They reported a resectability rate of 16.9% and a median survival of 18 months for patients who underwent resection, and of only 4 months for those who had surgical bypass. Over the last 25 years, many efforts have focused on improving outcomes in pancreatic cancer patients. Morbidity and mortality after pancreatic surgery have decreased markedly, chemotherapy and chemoradiation regimens have been developed, and 5-year survival rates of up to 25% have been reported in resected patients.^{2,16,30–35}

While many studies have evaluated specific stages of pancreatic cancer, few have reported data on the entire spectrum of disease, considering treatments, palliative procedures, and outcomes.¹⁶ The present study specifically addresses this point to give the reader a “snapshot” of pancreatic cancer treatment at the beginning of the twenty-first century. Unfortunately, the picture that emerges continues to be as grim as that described by Morrow 25 years ago. Our data shows that, despite chemotherapy and chemoradiation, the median survival is only 5.8 months for patients with metastatic disease and 8.6 months for those with locally advanced cancers, these two groups together constituting two thirds of the pancreatic cancer population seen in a cancer center. The one-year survival rate for the entire cohort was only 39%, and 58% of these are patients who underwent resection. Our survival rates did not significantly differ from those generally reported in the literature.^{4–7,30–35} Moreover, we considered a consecutive series of patients newly diagnosed with pancreatic cancer, not a selected population with favorable prognostic factors in which better survival rates can be achieved.^{16,24,33}

Surgery remains the only possibility of long-term survival for patients with pancreatic cancer, but the great majority of them are not amenable to resection even after neoadjuvant

treatments.^{16,36} In this series, 53% of group 2 patients underwent chemoradiation, but only in one case was sufficient “downstaging” obtained to allow subsequent surgical resection. These data underscore that palliation rather than curative treatment still remains the most relevant goal in the great majority of patients with pancreatic cancer.

In addition to evaluating treatment and survival, our study assessed longitudinal changes in HQOL, using validated instruments administered at diagnosis and during follow-up. Physicians aiming to keep patients comfortable and free of symptoms must evaluate the impact of these interventions on patients’ quality of life.¹⁷ To the best of our knowledge, this is the first prospective study that considers HQOL in patients with the full spectrum of localized, locally advanced, and metastatic pancreatic cancer.

The instruments used for HQOL evaluation in pancreatobiliary diseases range from visual analogue scales to generic HQOL (FACT-G or EORTC QLQ-C30) or disease-specific (FACT-Hep) questionnaires.^{18,23,24,37} We used both the FACT-G and FACT-Hep questionnaires, which have an excellent test–retest reliability, and high internal consistency are easy to complete and have been validated for patients with pancreatobiliary cancers.^{23–26} In addition to the general and disease-specific scores we also evaluated the F-TOI, a functional index that is a sensitive indicator of clinical outcome.²⁷

In the present study, data were analyzed to look for both statistical and clinical significance, which account for two different perspectives in HQOL interpretation.^{28,38} The purpose of statistical analysis is to quantify the importance of differences in a cohort of patients (population), and the sample sizes inevitably affect statistical power. However, statistically significant changes in a general population setting may not be meaningful in the context of single or few individuals. Clinical significance focuses upon detecting changes that are important in the patient’s perspective and therefore relevant in the management of individual patients. MIDs were used to define clinical significance in this study.²⁷

Interestingly, our data shows no statistical or clinical significant differences in HQOL scores at baseline among the three groups, which were also homogeneous for age, sex, and site of tumor. Only patients with localized disease who underwent surgical resection (group 1) had a subsequent improvement in quality of life: Scores on almost all HQOL scales improved during follow-up after surgical resection. In contrast, a decrease of all the scores was evident in group 3, while a slight increase was found in group 2 (Table 4).

It is difficult to compare HQOL results among different studies because of differences in instruments, methodology, and patient population.^{37,39} In some, various periampullary tumors or pancreatic diseases were considered,^{18,21} or HQOL was not assessed longitudinally but only after the therapeutic intervention.²³ Not surprising therefore, our results differ from those previously reported in the literature. Schniewind et al.³⁷ in a prospective study evaluating HQOL in a group of 91 patients who underwent pancreaticoduodenectomy for pancreatic cancer, showed a large decrease in most HQOL scales after surgery, followed by a slow recovery to preoperative levels. A similar trend was found by Nieveen van Dijkum et al.²¹ who compared pancreaticoduodenectomy versus biliary and duodenal bypass for pancreatic and periampullary carcinomas. Farnell et al.²⁴ comparing pancreaticoduodenectomy with or without extended lymphadenectomy in pancreatic cancer, found a decrease in most of HQOL scores from baseline to 4 months after surgery in both groups.

In contrast, we found an improvement of HQOL scores from baseline to follow-up only after surgical resection. This difference was clinically but not statistically significant despite pancreatic surgical resections being extensive procedures associated with potential major complications. The difference between group 1 versus groups 2 and 3 certainly depend upon

the differences in cancer stage and are influenced by the curative potential of surgical resection and the rapid evolution of the disease in unresected patients. It is nonetheless remarkable that HQOL scores improved during follow-up in group 1 patients, even though 73% of them underwent adjuvant chemoradiation.

A common methodological problem of HQOL studies is missing data, which may lead to bias.^{21,23,37} Specifically, missing data can result in overestimation of HQOL since very sick or dying patients are less likely to complete the questionnaires. In our study, this defect might be more prevalent in groups 2 and 3, but it is unlikely that overestimation affected group 1 because the majority of patients were alive with no recurrent disease 3 and 6 months after surgery.

In patients with advanced disease, we evaluated the impact of different palliative procedures and treatment on HQOL. In patients who develop jaundice and/or gastrointestinal obstruction but who are judged to be unresectable, endoscopic procedures are our first choice for palliation, while surgical bypass is generally performed for tumors found to be unresectable at laparotomy. Several studies in assessing the feasibility and efficacy of endoscopic biliary and enteral stenting as an alternative to surgical bypasses have shown that stenting is associated with lower costs and better quality of life when compared to surgical bypass.^{10,12,13} In our cohort, patients who underwent stent placement had a decrease in HQOL at 3 months but a clinically significant improvement at 6 months, while there were no significant changes in patients who did not require or receive stents (Fig. 3).

Pain control is a major issue in patients with advanced pancreatic cancer.^{11,15} Recently, Yan et al. and Wong et al. showed that,^{14,40} compared with standard analgesia, celiac block is associated with a significant but limited reduction of pain but does not improve either quality of life or survival. They concluded that celiac block should not replace standard pain control measures but should be used selectively as an adjunct. Pain relief based on systemic analgesics was successfully obtained in 59% of group 2 and 80% of group 3 patients. Patients who underwent celiac block had a decrease in HQOL at 3 months but a clinical significant improvement at 6 months, while no changes in HQOL were detected in the remaining patients (Fig. 5).

Chemotherapy and chemoradiation did not seem to impact HQOL differently in patients with locally advanced and metastatic disease. At 3 months, worse scores were found in the chemotherapy group, although this can be explained by the fact that chemotherapy was preferentially performed in patients with metastatic pancreatic cancer (Fig. 4).

Conclusion

In conclusion, because the rate of cure for pancreatic cancer continues to be very low, palliation of symptoms remains the more attainable goal for most cases. This study shows that there is a good overall impact of surgical and medical interventions on the quality of life in patients with pancreatic cancer. Despite potential perioperative and long-term complications, pancreatic resection improves quality of life of those with localized disease. Chemoradiation and chemotherapy do not negatively impact the quality of life in patients with locally advanced disease, but chemotherapy in patients with metastatic disease is associated with a significant decrease in quality of life during follow-up, due either to chemotherapy, the progression of the cancer, or both.

Acknowledgments

Dr. Crippa was supported by the International Hepato-Pancreato-Biliary Association (2007 Warren Fellowship Award). Dr. Domínguez was supported by the Fundación México en Harvard A.C.

References

1. Jemal A, Murray T, Ward E, Samuels A, Tiwari RC, Ghafoor A, Feuer EJ, Thun MJ. Cancer Statistics, 2005 CA. *Cancer J Clin.* 2005; 55:10–30.
2. Yeo CJ, Cameron JL, Lillemoe KD, Sohn TA, Campbell KA, Sauter PK, Coleman J, Abrams RA, Hruban RH. Pancreaticoduodenectomy with or without distal gastrectomy and extended retroperitoneal lymphadenectomy for periaampullary adenocarcinoma, part 2: randomized controlled trial evaluating survival, morbidity and mortality. *Ann Surg.* 2002; 236:355–366. [PubMed: 12192322]
3. Ferrone CR, Finkelstein DM, Thayer SP, Muzikansky A, Fernandez-del Castillo C, Warshaw AL. Perioperative CA19-9 levels can predict stage and survival in patients with resectable pancreatic adenocarcinoma. *J Clin Oncol.* 2006; 24:2897–2902. [PubMed: 16782929]
4. Sener SF, Fremgen A, Menck HR, Winchester DP. Pancreatic cancer: a report of treatment and survival trends for 100,313 patients diagnosed from 1985–1995, using the national cancer database. *J Am Coll Surg.* 1999; 189:1–7. [PubMed: 10401733]
5. Talamonti MS, Small W, Mulcahy MF, Wayne JD, Attaluri V, Coletti LM, Zalupski MM, Hoffman JP, Freedman GM, Kinsella TJ, Philip PA, McGrinn CJ. A multi-institutional phase II trial of preoperative full-dose gemcitabine and concurrent radiation for patients with potentially resectable pancreatic carcinoma. *Ann Surg Oncol.* 2006; 13:150–158. [PubMed: 16418882]
6. Yip D, Karapetis C, Strickland A, Steer CB, Goldstein D. Chemotherapy and radiotherapy for inoperable advanced pancreatic cancer. *Cochrane Database Syst Rev.* 2006; 19:CD00093.
7. Ko AH, Dito E, Schillinger B, Venook AP, Bergsland EK, Tempero MA. Phase III study of fixed dose rate gemcitabine with cisplatin for metastatic adenocarcinoma of the pancreas. *J Clin Oncol.* 2006; 24:379–385. [PubMed: 16344320]
8. Lillemoe KD, Pitt HA. Palliation. *Surgical and otherwise Cancer.* 1996; 78:605–614.
9. Lillemoe KD, Cameron JL, Hardacre JM, Sohn TA, Sauter PK, Coleman J, Pitt HA, Yeo CJ. Is prophylactic gastrojejunostomy indicated for unresectable periaampullary cancer? A prospective randomized trial. *Ann Surg.* 1999; 230:322–328. [PubMed: 10493479]
10. Mortenson MM, Ho HS, Bold RJ. An analysis of cost and clinical outcome in palliation for advanced pancreatic cancer. *Am J Surg.* 2005; 190:406–411. [PubMed: 16105527]
11. Latori KJ, Hjermstad MJ, Wester T, Buanes T, Loge JH. Symptom profiles and palliative care in advanced pancreatic cancer: a prospective study. *Support Care Cancer.* 2006; 14:1126–1133. [PubMed: 16601947]
12. Artifon EL, Sakai P, Cunha JE, Dupont A, Filho FM, Hondo FY, Ishioka S, Raju GS. Surgery or endoscopy for palliation of biliary obstruction due to metastatic pancreatic cancer. *Am J Gastroenterol.* 2006; 101:2031–2037. [PubMed: 16968509]
13. Maire F, Hammel P, Ponsot P, Aubert A, O'Toole D, Hentic O, Levy P, Ruszniewski P. Long-term outcome of biliary and duodenal stents in palliative treatment of patients with unresectable adenocarcinoma of the head of the pancreas. *Am J Gastroenterol.* 2006; 101:735–742. [PubMed: 16635221]
14. Yan BM, Myers RP. Neurolytic celiac plexus block for pain control in unresectable pancreatic cancer. *Am J Gastroenterol.* 2007; 102:430–438. [PubMed: 17100960]
15. Lillemoe KD, Cameron JL, Kaufman HS, Yeo CJ, Pitt HA, Sauter PK. Chemical splanchnicectomy in patients with unresectable pancreatic cancer. A prospective randomized trial. *Ann Surg.* 1993; 217:447–455. [PubMed: 7683868]
16. Kuhlmann KF, de Castro SM, Wesseling JG, ten Kate FJ, Offerhaus GJ, Busch OR, van Gulik TM, Obertop H, Gouma DJ. Surgical treatment of pancreatic adenocarcinoma; actual survival and prognostic factors in 343 patients. *Eur J Cancer.* 2004; 40:549–558. [PubMed: 14962722]
17. Velanovich V. Using quality-of-life measurements in clinical practice. *Surgery.* 2007; 141:127–133. [PubMed: 17263966]
18. Huang JJ, Yeo CJ, Sohn TA, Lillemoe KD, Sauter PK, Coleman J, Hruban RH, Cameron JL. Quality of life and outcomes after pancreaticoduodenectomy. *Ann Surg.* 2000; 231:890–898. [PubMed: 10816633]

19. Velanovich V. Using quality-of-life instruments to assess surgical outcomes. *Surgery*. 1999; 126:1–4. [PubMed: 10418584]
20. Testa MA, Simonson D. Assessment of quality of life outcomes. *N Engl J Med*. 1996; 334:835–840. [PubMed: 8596551]
21. Nieveen van Dijkum EJ, Kuhlmann KF, Terwee CB, Obertop H, de Haes JC, Gouma DJ. Quality of life after curative or palliative surgical treatment of pancreatic and periampullary carcinoma. *Br J Surg*. 2005; 92:471–477. [PubMed: 15672431]
22. Billings BJ, Christein JD, Harmsen WS, Harrington JR, Chari ST, Que FG, Farnell MB, Nagorney DM, Sarr MG. Quality-of-life after total pancreatectomy: is it really that bad on long-term follow-up? *J Gastrointest Surg*. 2005; 9:1059–1066. [PubMed: 16269376]
23. Nguyen TC, Sohn TA, Cameron JL, Lillemoe KD, Campbell KA, Coleman J, Sauter K, Abrams RA, Hruban RH, Yeo CJ. Standard vs. radical pancreaticoduodenectomy for periampullary adenocarcinoma: a prospective, randomized trial evaluating quality of life in pancreaticoduodenectomy survivors. *J Gastrointest Surg*. 2003; 7:1–9. [PubMed: 12559179]
24. Farnell MB, Pearson RK, Sarr MG, DiMaggio EP, Burgart LJ, Dahl TR, Foster N, Sargent DJ. Pancreas Cancer Working Group. A prospective randomized trial comparing standard pancreaticoduodenectomy with pancreaticoduodenectomy with extended lymphadenectomy in resectable pancreatic head adenocarcinoma. *Surgery*. 2005; 138:618–628. [PubMed: 16269290]
25. Cella DF, Tulsky DS, Gray G, Sarafian B, Linn E, Bonomi A, Silberman M, Yellen SB, Winicour P, Brannon J. The functional assessment of cancer therapy scale: development and validation of the general measure. *J Clin Oncol*. 1993; 11:570–579. [PubMed: 8445433]
26. Heffernan N, Cella DF, Webster K, Odom L, Martone M, Passik S, Bookbinder M, Fong Y, Jarnagin W, Blumgart L. Measuring health-related quality of life in patients with hepatobiliary cancers: the functional assessment of cancer therapy—hepatobiliary questionnaire. *J Clin Oncol*. 2002; 20:2229–2239. [PubMed: 11980994]
27. Steel JL, Eton DT, Cella D, Olek MC, Carr BI. Clinically meaningful differences in health-related quality of life in patients diagnosed with hepatobiliary carcinoma. *Ann Oncol*. 2006; 17:304–312. [PubMed: 16357021]
28. Bro3ek JL, Guyatt HG, Schünemann HJ. How a well-grounded minimal important difference can enhance transparency of labeling claims and improve interpretation of a patient reported outcome measure. *Health Qual Life Outcomes*. 2006; 4:1–7. [PubMed: 16393335]
29. Morrow M, Hilaris B, Brennan MF. Comparison of conventional surgical resection, radioactive implantation, and bypass procedure for exocrine carcinoma of the pancreas 1975–1980. *Ann Surg*. 1984; 199:1–5. [PubMed: 6419687]
30. Trede M, Schwall G, Saeger HD. Survival after pancreaticoduodenectomy: 118 consecutive resections without mortality. *Ann Surg*. 1990; 211:430–438.
31. Yeo CJ, Cameron JL, Sohn TA, Lillemoe KD, Pitt HA, Talamini MA, Hruban RH, Ord SE, Sauter PK, Coleman J, Zahurak ML, Grochow LB, Abrams RA. Six hundred fifty consecutive pancreaticoduodenectomies in the 1990s: pathology, complications, outcomes. *Ann Surg*. 1997; 226:248–260. [PubMed: 9339931]
32. Warshaw AL, Fernandez-del Castillo C. Pancreatic carcinoma. *N Engl J Med*. 1992; 326:455–465. [PubMed: 1732772]
33. Nitecki SS, Sarr MG, Colby TV, van Heerden JA. Long-term survival after resection for ductal adenocarcinoma of the pancreas. Is it really improving? *Ann Surg*. 1995; 221:59–66. [PubMed: 7826162]
34. Sohn TA, Yeo CJ, Cameron JL, Koniaris L, Kaushal S, Abrams RA, Sauter PK, Coleman J, Hruban RH, Lillemoe KD. Resected adenocarcinoma of the pancreas—616 patients: results, outcomes, and prognostic indicators. *J Gastroint Surg*. 2000; 4:567–579.
35. Riall TS, Nealon WH, Goodwin JS, Zhang D, Kuo YF, Townsend CM Jr, Freeman JL. Pancreatic cancer in the general population: improvements in survival over the last decade. *J Gastrointest Surg*. 2006; 10:1212–1224. [PubMed: 17114008]
36. Masucco P, Capussotti L, Magnino A, Sperti E, Gatti M, Muratore A, Sgotto E, Gabriele P, Aglietta M. Pancreatic resections after chemoradiotherapy for locally advanced ductal

- adenocarcinoma: analysis of perioperative outcome and survival. *Ann Surg Oncol*. 2006; 13:1201–1208. [PubMed: 16955382]
37. Schniewind B, Bestmann B, Henne-Bruns D, Faendrich F, Kremer B, Kuechler T. Quality of life after pancreaticoduodenectomy for ductal adenocarcinoma of the pancreatic head. *Br J Surg*. 2006; 93:1099–1107. [PubMed: 16779883]
38. Revicki D. FDA draft guidance and health-outcomes research. *Lancet*. 2007; 369:540–542. [PubMed: 17307086]
39. Holzner B, Bode RK, Hahn EA, Cella D, Kopp M, Sperner-Unterweger B, Kemmler G. Equating EORTC QLQ-C30 and FACT-G scores and its use in oncological research. *Eur J Cancer*. 2006; 42:3169–3177. [PubMed: 17045472]
40. Wong GY, Schroeder DR, Carns PE, Wilson JL, Martin DP, Kinney MO, Mantilla CB, Warner DO. Effect of neurolytic celiac plexus block on pain relief, quality of life, and survival in patients with unresectable pancreatic cancer: a randomized controlled trial. *JAMA*. 2004; 291:1092–1099. [PubMed: 14996778]

Figure 1.

a Kaplan–Meier overall survival curve in 92 patients with pancreatic cancer. One- and 2-year survival was 39 and 23 months. **b** Kaplan–Meier survival curves in patients with localized (group 1, $n = 28$), locally advanced (group 2, $n = 34$), and metastatic pancreatic cancer (group 3, $n = 30$) at initial presentation. One-year survival was 74%, 30%, and 16% for groups 1, 2, and 3, respectively ($P = 0.001$).

Figure 2. Clinical changes in health-related quality of life (*HQOL*) of patients with local (group 1), locally advanced (group 2), and metastatic (group 3) pancreatic cancer. *Square and triangle* Meaningful important differences (MID).

Figure 3. Clinical changes in HQOL in patients affected by locally advanced and metastatic pancreatic cancer with or without biliary/enteral stents. *Square and triangle* Meaningful important differences (MID).

Figure 4. Clinical changes in HQOL of patients with locally advanced and metastatic pancreatic cancer who underwent chemotherapy or chemoradiation. *Square* Meaningful important differences (MID).

Figure 5. Clinical changes in HQOL of patients with locally advanced and metastatic pancreatic cancer who underwent celiac block or did not. *Square, triangle; and circle* Meaningful Important Differences (MID).

Table 1

Demographics, Presenting Symptoms, and Tumor Site in 92 Patients with Pancreatic Cancer Classified According to the Clinical Stage (Group 1, Localized Disease; Group 2, Locally Advanced Disease; Group 3, Metastatic Disease)

	Group 1 (28 Patients), n (%)	Group 2 (34 Patients), n (%)	Group 3 (30 Patients), n (%)	1 vs. 2 (P value)	2 vs. 3 (P value)	1 vs. 3 (P value)
Age (years), mean $\pm$ SD	65 $\pm$ 10	66 $\pm$ 11	65 $\pm$ 11	1	1	1
Sex						
Male	14 (50)	16 (47)	16 (53)	0.5	0.4	0.5
Female	14 (50)	18 (53)	14 (47)			
Presenting symptoms						
Jaundice	24 (86)	7 (21)	4 (14)	0.0001	0.3	0.0001
Abdominal pain	3 (11)	13 (38)	9 (30)	0.1	0.6	0.2
Weight loss	1 (3)	14 (41)	17 (56)	0.0001	0.1	0.0001
Tumor site						
Proximal	21 (75)	25 (73.5)	19 (63)	0.5	0.2	0.3
Distal	7 (25)	9 (26.5)	11 (37)			
Median CA 19.9 (U/L)	60.5	264	1536	0.2	0.02	0.003

Table 2
Diagnostic and Palliative Procedures in 92 Patients with Pancreatic Cancer Classified According to the Clinical Stage (Group 1, Localized Disease; Group 2, Locally Advanced Disease; Group 3, Metastatic Disease)

	Group 1 (28 Patients)	Group 2 (34 Patients)	Group 3 (30 Patients)	1 vs. 2 <i>P</i> Value	1 vs. 3 <i>P</i> Value	2 vs. 3 <i>P</i> Value
Abdominal CT						
Overall number	117	122	77			
Number of patients ^a (%)	28 (100)	34 (100)	30 (100)	–	–	–
Mean number/pt	4±2.6	3.6±2.7	2.6±2	0.9	0.03	0.3
EUS						
Overall number	18	31	19			
Number of patients ^a (%)	17 (61)	28 (83)	16 (53)	0.5	0.6	0.01
Mean number/pt	1±0.2	1±0.1	1.2±0.4	0.6	0.2	0.4
ERCP						
Overall number	23	38	22			
Number of patients ^a (%)	14 (50)	23 (68)	16 (53)	0.1	0.8	0.2
Mean number/pt	1.64±0.7	1.65±0.7	1.3±0.6	0.9	0.2	0.1
Diagnostic laparoscopy (%)	4 (14)	10 (30)	6 (20)	0.1	0.4	0.3
Biliary stents						
Overall number	20	33	19			
Number of patients ^a (%)	14 (50)	20 (62.5)	14 (47)	0.2	0.8	0.2
Mean number/pt	1.4±0.5	1.65±0.7	1.3±0.6	0.4	0.5	0.2
Hospital stay						
Two or more admissions (%)	10 (36)	19 (56)	8 (27)	0.1	0.4	0.01
Overall mean hospital stay ± SD	15±12	14±11.5	8±7	0.5	0.001	0.03
Celiac block (%)	0	14 (41)	6 (20)	–	–	0.6
Enteral stent (%)	0	6 (18)	2 (7)	–	–	0.1
Palliative surgery (%)	0	6 (17.5)	4 (13)	–	–	0.4

Palliative surgery consisted of gastrojejunostomy (*n* = 5) and gastrojejunostomy plus a hepaticojunostomy (*n* = 1) in group 2; gastrojejunostomy and hepaticojunostomy (*n* = 1), hepaticogastrostomy (*n* = 1) in group 3.

^aOverall number of patients who underwent at least one examination/procedure in each group

Table 3

Perioperative Findings and Postoperative Complications in 29 Patients Who Underwent Pancreatic Resection for Ductal Adenocarcinoma

	Number of Patients (%)
Type of resection	
Pancreaticoduodenectomy ^a	21 (73%)
Distal pancreatectomy and splenectomy	7 (24%)
Total pancreatectomy	1 (3%)
Mean blood loss (ml)	670±430
Mean operative time (min)	286±78
Patients requiring blood transfusions	3 (10)
Postoperative complications	
Overall morbidity	7 (24)
Pancreatic fistula	2 (7)
Abscess	2 (7)
Mean postoperative length of stay (days)	8±2
Surgical-related mortality ^b	0
Neoadjuvant chemoradiation ^c	4 (13)
Adjuvant chemoradiation	21 (73)
Mean pathologic tumor size	30±12
Positive lymph nodes	17 (58)
Positive resection margins	3 (10)
Tumor grading	
G1	1 (3)
G2	14 (48.5)
G3	14 (48.5)
Presence of perineural infiltration	23 (79)
Presence of microvascular infiltration	18 (62)

They include 28 patients with localized neoplasm (group 1) and one patient with a locally advanced neoplasm (group 2) who underwent neoadjuvant chemoradiation.

^aWhipple procedure with pancreatojejunostomy was performed in all the cases.

^bSurgical-related mortality was defined as in-hospital or 30-day postoperative mortality.

^cNeoadjuvant chemoradiation was performed in one patient affected by a locally advanced neoplasm with clear evidence of vascular infiltration (group 2) and in three patients with localized disease (group 1).

Table 4
Health Related Quality of Life Scores (FACT G, FACT Hep, FTOI) in the Different Groups

HQOL Scores	Group 1 (28 Patients), Mean $\pm$ SD	Group 2 (34 Patients), Mean $\pm$ SD	Group 3 (30 Patients), Mean $\pm$ SD	1 vs. 2, <i>P</i> Value	2 vs. 3, <i>P</i> Value	1 vs. 3, <i>P</i> Value
Baseline						
FACT Hep	124.7 $\pm$ 21.6	122.6 $\pm$ 23.7	122.7 $\pm$ 20.2	0.9	0.9	0.9
FTOI	83.7 $\pm$ 19.4	82.5 $\pm$ 19.1	83.2 $\pm$ 16.8	0.9	0.9	0.9
FACT G	76.3 $\pm$ 14.4	72.5 $\pm$ 17.5	69.5 $\pm$ 19.3	0.7	0.7	0.3
Follow-up						
FACT Hep	134 $\pm$ 17.9	125 $\pm$ 21.4	113.3 $\pm$ 34.7	0.4	0.3	0.03
FTOI	93.5 $\pm$ 16	84.2 $\pm$ 17	74.7 $\pm$ 14	0.2	0.3	0.02
FACT G	78.8 $\pm$ 11.6	74.7 $\pm$ 14	68.7 $\pm$ 20	0.6	0.4	0.1
<i>P</i> value ^a						
FACT Hep	0.1	0.6	0.2			
FTOI	0.06	0.7	0.09			
FACT G	0.5	0.6	0.8			

^a Comparison between baseline and follow up scores within each group

Table 5

Health-Related Quality of Life Scores at Baseline and Follow-up in the Whole Cohort ($n = 92$) Considering the Following Variables: Pancreatic Resection, CA 19.9 (cut-off, 200 U/l), Age (cut-off, 60 years), Chemotherapy Versus Chemoradiation During the Disease Course

Variable	Baseline		Follow-up		p
	FACT-Hep	FTOI	FACT-G	FACT-Hep	
Resected	125.7±21.6	84.5±19.5	77±14.5	93.6±15	0.001
Nonresected	122.2±22	82.5±18	70.8±18.2	80.4±22	
CA 19-9>200	118.3±22.5	78.4±18.6	69.8±19	80±20.5	0.04 ^a
CA 19-9<200	128.8±19.7	88.4±16.5	75.6±15	90.8±19	
Age>60	126.2±19.5	84.3±18	74.4±16.8	86.5±21	NS
Age<60	116.4±25.3	80±19	68±18.1	82.8±19	
Chemotherapy	123.6±20	83.3±17	72±15	79±25.2	NS
Chemoradiation	122.8±20.5	82.2±17.6	74.7±14.4	87.6±15	

Statistically significant mean differences in FACT-Hep, FTOI and FACT-G scores. No statistically significant differences were found when comparing baseline versus follow-up scores for all the variables.

^aStatistically significant mean differences in FACT-Hep and FTOI scores