Published in final edited form as:

J Med Chem. 2013 April 11; 56(7): . doi:10.1021/jm4000374.

Chemistry and Pharmacological Studies of 3-Alkoxy-2,5-Disubstituted-Pyridinyl Compounds as Novel Selective $\alpha 4\beta 2$ Nicotinic Acetylcholine Receptor Ligands That Reduces Alcohol Intake in Rats

Yong Liu¹, Janell Richardson², Thao Tran², Nour Al-Muhtasib², Teresa Xie², Venkata Mahidhar Yenugonda¹, Hannah G. Sexton³, Amir H. Rezvani³, Edward D. Levin³, Niaz Sahibzada², Kenneth J. Kellar², Milton L. Brown¹, Yingxian Xiao^{2,*}, and Mikell Paige^{1,*}

¹Center of Drug Discovery, Georgetown University Medical Center, 3970 Reservoir Road, NW, Washington, D.C. 20057

²Department of Pharmacology and Physiology, Georgetown University School of Medicine, 3900 Reservoir Road, NW, Washington, D.C. 20057

³Department of Psychiatry and Behavioral Sciences, Duke University Medical Center, Durham, NC 27710

Abstract

Neuronal acetylcholine receptors mediate the addictive effects of nicotine and may also be involved in alcohol addiction. Varenicline, an approved smoking cessation medication, showed clear efficacy in reducing alcohol consumption in heavy-drinking smokers. More recently, sazetidine-A, which selectively desensitizes $\alpha 4\beta 2$ nicotinic receptors, was shown to significantly reduce alcohol intake in a rat model. To develop novel therapeutics for treating alcohol use disorder, we designed and synthesized novel sazetidine-A analogs containing a methyl group at the 2-position of the pyridine ring. In vitro pharmacological studies revealed that some of the novel compounds showed similar overall pharmacological property profiles with that of sazetidine-A, but exhibited reduced agonist activity across all nicotinic receptor subtypes tested. In animal studies, compound (*S*)-**9** significantly reduced alcohol uptake in rats. More importantly, preliminary results from studies in a ferret model indicate that these novel nAChR ligands have an improved adverse side-effect profile in comparison with that of varenicline.

Keywords

nicotinic acetylcholine receptors; sazetidine-A; varenicline; desensitization; addiction; alcohol use disorders

INTRODUCTION

Tobacco use, mainly via cigarette smoking, and alcohol consumption are among the leading preventable risk factors for premature mortality in the world. In 2000, tobacco use was the

*Correspondence regarding the medicinal chemistry design and synthesis should be addressed to: Phone +1-202-687-5341; map65@georgetown.edu. . *Correspondence regarding the pharmacological studies should be addressed to Phone +1-202-687-5390; yxiao02@georgetown.edu. .

Author Contributions The manuscript was written through contributions of all authors. All authors have given approval to the final version of the manuscript.

number one risk factor for mortality in the United States, which resulted in 435,000 premature deaths. In that same year, alcohol consumption was the number three risk factor, which was responsible for 85,000 premature deaths.¹ In addition to being an immense medical and public health problem, tobacco use and alcohol consumption also impose a huge social and economic burden to society. Given the grave consequences of nicotine and alcohol addiction there is obviously a great need for significant improvement in existing therapies for treating these disorders.

Although alcohol and nicotine addictions are usually categorized as separate disorders, heavy alcohol consumption and smoking occur very often in the same individual.^{2, 3} It has been well established that there are potent interactions between alcohol and nicotinic systems and that alcohol potentiates nicotine reward and self-administration.⁴⁻⁷

Neuronal nicotinic acetylcholine receptors (nAChRs) are found throughout the central nervous system (CNS) and peripheral nervous system (PNS). The receptors mediate the pharmacological effects of nicotine and play a key role in regulating nicotine self-administration behavior.⁸⁻¹⁰ The predominant nAChR subtype in mammalian CNS, which contains both $\alpha 4$ and $\beta 2$ subunits, is essential for nicotine addiction.¹¹⁻¹⁸ Several lines of evidence show that nAChRs are involved in mediating alcohol consumption as well. The classical non-competitive antagonist of nAChRs, mecamylamine, significantly reduced alcohol consumption in rats and mice.^{6, 19-21} Several other nicotinic ligands were reported to attenuate alcohol seeking and consumption in rats.^{22, 23} Furthermore, genetic studies showed that nicotinic receptor subunits are directly involved in alcohol consumption in mouse models.²⁴⁻²⁶

Varenicline, an nAChR ligand developed by Pfizer, was approved by the US Food and Drug Administration (FDA) in 2006 as a smoking cessation medication.²⁷⁻³⁰ Interestingly, smokers treated with varenicline reported that the medication also reduced their alcohol consumption.³¹ In a double-blind, placebo-controlled clinical trial, varenicline significantly reduced alcohol self-administration in heavy-drinking smokers.³² Furthermore, several studies using animal models showed that varenicline effectively reduced alcohol consumption in rats and mice.^{24, 25, 31, 33} Although varenicline appears to be safe for most people, adverse side effects have been reported.^{28, 29, 34}

In 2006, we proposed a new approach in the design of novel nicotinic therapeutics, which is based on the ability of nicotinic ligands to desensitize nAChRs.³⁵ The actions of nicotine and other nicotinic agonists to activate and then desensitize nAChRs have been known for more than 100 years³⁶ and conceptualized for more than 50 years.³⁷ A fundamental question about nicotinic cholinergic signaling in the CNS is how each of these two opposite actions contributes to the overall pharmacological effects of nicotine. It has been well known that nicotine, as well as all other known nicotinic agonists, is more potent in desensitizing nAChRs than in activating them.³⁸⁻⁴⁵

Sazetidine-A is a highly selective ligand for nAChRs that contain the $\beta 2$ subunit, especially the $\alpha 4 \beta 2$ nAChR subtype.³⁵ Sazetidine-A was shown to potently and selectively desensitize $\alpha 4 \beta 2$ nAChRs (see Table 2). In studies using animal behavioral models, sazetidine-A reduced nicotine self-administration,^{46, 47} improved performance in tests of attention,^{48, 49} and showed antidepressant and/or anti-anxiety effects.⁵⁰⁻⁵² Interestingly, sazetidine-A also significantly decreased alcohol intake in rats.⁴⁷

Herein we report the design, synthesis, and biological characterization of a new class of sazetidine-A analogs that have a reduced agonist effect on several nAChR subtypes and a reduced adverse side-effect profile in ferrets as compared with varenicline. These novel compounds could lead to more effective treatments for alcohol addiction.

RESULTS

Chemistry

Previous studies have shown that O-substitution at the 3-position of pyridine with a 2(*S*)-azetidinylmethanol substituent and the presence of an aliphatic chain at the 5-position, such as found in sazetidine-A, are key structural components for the selective desensitization of $\alpha_4\beta_2$ nAChRs.³⁵ During our studies of sazetidine-A and related analogs, it became clear that the in vivo stability of these compounds should be an important consideration for clinical development. Recently, the Baran and the MacMillan groups independently reported that the innate susceptibility for heteroaromatic hydroxylation could be investigated under oxidative conditions in the presence of a trifluoromethylating reagent.^{53, 54} We subjected sazetidine-A (Figure 1) to the conditions reported by Baran and co-workers and observed that trifluoromethylation occurred at the 2-position of the pyridine ring, albeit in 10% yield. This result suggested that the 2-position of the pyridine moiety could be susceptible to P450-mediated oxidation. We hypothesized that blocking the 2-position of the pyridine ring of sazetidine-A with a CF₃ group could result in a compound with increased stability. Therefore, we synthesized compound (*S*)-**22**, a sazetidine-A analog that incorporates a CF₃ group at the 2-position of the pyridine moiety (Figure 1). However, this analog exhibited a tremendous decrease in binding affinity to the target receptors (Table 1). It is conceivable that the electronegativity of the CF₃ group, or the large volume of the group, is responsible for the significant alteration in the pharmacological properties for this structural class. To test this possibility, we then synthesized compound (*S*)-**9** (Figure 1), which contains a 2-methyl substituent on the pyridine moiety in place of the CF₃ group in compound (*S*)-**22**. The 2-methyl group of compound (*S*)-**9** provides an electron donating substituent at the 2-position in contrast to the CF₃ group of compound (*S*)-**22**, as well as a smaller volume than that of the CF₃ group. We were delighted that in the binding assays, compound (*S*)-**9** showed a high binding affinity ($K_i = 12$ nM) to $\alpha_4\beta_2$ nAChRs and high selectivity for the $\alpha_4\beta_2$ nAChR subtype over two other major subtypes, $\alpha_3\beta_4$ and $\alpha_7\beta_1$ (Table 1). Furthermore, comparison with varenicline and sazetidine-A, compound (*S*)-**9** exhibited significantly lower agonist activities at both the $\alpha_4\beta_2$ and $\alpha_3\beta_4$ nAChRs (Table 2). This discovery indicated a possibility for developing a series novel nicotinic ligands that selectively bind to $\alpha_4\beta_2$ nAChRs with high affinities and potentially desensitize the receptor subtype with very low agonist activities across all major nAChR sub-types. To further study this new class of nAChR ligands, we explored the importance of the ring size and the stereocenter of the cyclic amine. The effect of changing the terminal group of the alkynyl appendage was also investigated.

The new sazetidine-A analogs designed in our study that contain a 2-methyl group on the pyridine moiety were synthesized by an optimized three-step procedure as shown in Scheme 1. Each enantiomer of Boc-protected 2-azetidinylmethanol or 2-pyrrolidinylmethanol was reacted with 5-bromo-2-methylpyridin-3-ol via standard Mitsunobu coupling conditions to afford bromides **3** and **4**. The alkynyl appendage was incorporated by Sonogashira homologation in the presence of triphenylphosphine in triethylamine with DMSO as a co-solvent. The *N*-Boc protecting group was then removed by treatment with trifluoroacetic acid or hydrogen chloride to afford the analogs **9** through **12** as the corresponding salt.

Analog **16** and **17**, which contain a terminal halogen substituent, were synthesized as outline in Scheme 2. The hydroxide group of compound **12** was tosylated by treatment with *p*-toluenesulfonyl chloride in methylene chloride at 0 °C. The resulting product was then treated with potassium fluoride in the presence of Kryptofix 2.2.2 in anhydrous THF to afford fluoroalkane **15**.⁵⁵ Compound **16** was obtained as the hydrochloride salt after deprotection of the *N*-Boc group by treatment with hydrogen chloride in methanol. The

chloroalkane analog **17** was originally afforded by a side reaction during the tosylation step when compound **12** was treated with *p*-toluenesulfonyl chloride in pyridine at room temperature. We have optimized the synthesis of the chloroalkane analog by treating compound **12** with triphenylphosphine and carbon tetrachloride in a sealed tube at 80 °C for 48 hours.⁵⁶ The deprotection step of the chlorinated product **14** was accomplished by treatment with hydrogen chloride in methanol to afford the hydrochloride salt of compound **17**.

The 2-trifluoromethylpyridinyl analog **22** was synthesized as shown in Scheme 3. The methyl aryl ether **18** was treated with hydrogen bromide in acetic acid at elevated temperatures to afford phenol **19**.⁵⁷ Mitsunobu homologation of phenol **19** with alcohol **1** then gave aryl bromide **20**. Homologation of 1-hexyne to compound **20** was accomplished following the Sonogashiro protocol as outlined in Scheme 1. However, for this substrate the reaction was accomplished at lower temperature and shorter reaction time than for the 2-methylpyridinyl substituted substrates, presumably because the electron-withdrawing properties of the trifluoromethyl substituent assists in the initial oxidative insertion of the palladium between the carbon and bromine atoms. It is important to note that higher temperatures and longer reaction times resulted in diminished yields. Deprotection of the *N*-Boc group was then effected by treatment with hydrogen chloride in methanol to afford compound **22** as the hydrogen chloride salt.

In vitro Binding Affinities for nAChR Subtypes

The binding affinities of the novel 2,5-disubstituted sazetidine-A analogs for the defined rat nAChR subtypes as well as for native nAChRs of rat forebrain were examined in binding competition studies against [³H]-epibatidine. For comparison, binding affinities of (–)-nicotine, varenicline and sazetidine-A were obtained from parallel binding experiments. To determine the selectivities of these compounds in binding assays among the three predominant nAChR subtypes, $\alpha 3\beta 4$, $\alpha 4\beta 2$ and $\alpha 7\beta 4\gamma 2$, the ratios of the corresponding K_i values ($\alpha 3\beta 4/\alpha 4\beta 2$ and $\alpha 7\beta 4\gamma 2$) were determined.

As reported previously³⁵ and shown in Table 1, sazetidine-A bound to $\alpha 4\beta 2$ nAChRs with a very high affinity ($K_i = 0.062$ nM), which was 31,000 and 11,000 times higher than its affinities for the $\alpha 3\beta 4$ and $\alpha 7\beta 4\gamma 2$ subtypes, respectively. The data suggest that sazetidine-A bound to each $\alpha 2$ -containing nAChR subtype with a much higher affinity than to the corresponding $\alpha 4$ -containing counterparts having the same β subunit.

As shown in Scheme 1, the incorporation of a methyl group at 2-position of the pyridine moiety in sazetidine-A afforded analog (*S*)-**12**. The K_i value of this compound for $\alpha 4\beta 2$ nAChRs is 4.3 nM, representing a lower binding affinity than that of sazetidine-A (Table 1). However, the binding affinity of (*S*)-**12** for $\alpha 4\beta 2$ nAChRs is 2 fold higher than that of (–)-nicotine. More importantly, analog (*S*)-**12** retained the excellent selectivity pattern seen with sazetidine-A with a slightly more favorable selectivity for the $\alpha 4\beta 2$ nAChRs over the $\alpha 3\beta 4$ and $\alpha 7\beta 4\gamma 2$ subtypes.

We then synthesized four analogs of compound (*S*)-**12**. Replacing the hydroxyl group at the terminal end of the alkyl chain with a fluorine atom (analog (*S*)-**16**) or a chlorine atom (analog (*S*)-**17**) did not lead to significant changes in the binding property profiles (Table 1). Similarly, removing the hydroxyl group (analog (*S*)-**9**) or replacing the end of the side chain with a cyclopropyl ring (analog (*S*)-**11**) resulted in little change in the binding profiles for this class of compounds (Table 1).

All closely related *S* enantiomers, including (*S*)-**9**, (*S*)-**11**, (*S*)-**16** and (*S*)-**17**, showed binding property profiles similar to that of (*S*)-**12**. In contrast, compound (*R*)-**9** showed

significantly lower binding affinities and reduced nAChR subtype selectivities than those of compound (*S*)-**9** (Table 1). Consistent with this finding, all analogs containing an *R* stereocenter in this study showed lower binding affinities and less nAChR subtype selectivities than analogs containing the *S* stereo-chemistry.

It is important to note that compound (*S*)-**10**, which has a five-membered ring pyrrolidine group, showed much lower affinities and nAChR subtype selectivities than those of compound (*S*)-**9**, which has a four-member ring azetidiny group (Table 1). However, compound (*S*)-**10** still had binding affinities at 4–2 nAChRs in the nanomolar range and maintained a greater than 1,000-fold selectivity for the 4–2 nAChR subtype over the 3–4 or 7 subtypes. In line with our emerging SAR, the *R* enantiomer of compound **10** showed reduced binding affinity and nAChR subtype selectivities in comparison to the corresponding *S* enantiomer.

Compound **22**, which contains a trifluoromethyl group at the 2-position of the pyridine ring, showed minimal binding affinity to every nAChR subtype tested. This may indicate that reducing the electron density in the pyridine ring and/or increasing steric bulk at the 2-position of the pyridine moiety can lead to lower binding affinities.

In Vitro Effects on nAChR Function

Six new compounds showing high binding affinities for 4–2 nAChRs and high selectivities for this subtype over 3–4 and 7 receptors in binding assays were chosen for functional studies (Table 2). The agonist activities for these analogs were assessed by measuring stimulated $^{86}\text{Rb}^+$ efflux from stably transfected cells, either expressing human 4–2 nAChRs or rat 3–4 receptors. The ability of the analogs to desensitize the two nAChR subtypes were determined by measuring nicotine-stimulated $^{86}\text{Rb}^+$ efflux after cells were pre-incubated with the test compounds for 10 min. For comparison, nicotine, varenicline and sazetidine-A were included in these experiments.

As expected, nicotine showed full agonist activities at both human 4–2 and rat 3–4 nAChRs (Table 2). Consistent with previous reports,^{27, 30, 58} in comparison with (–)-nicotine our studies show that varenicline had 45% of the efficacy in stimulating efflux from cells expressing the 4–2 nAChRs and 90% of the efficacy from cells expressing the 3–4 nAChRs. Sazetidine-A showed lower efficacy at the 4–2 nAChRs than that of varenicline. In contrast, we found that sazetidine-A's agonist activity at the 3–4 nAChRs was lower than that of (–)-nicotine or varenicline. Consistent with the purposes of developing this line of novel ligands, all 6 novel compounds tested showed much lower agonist activities than those of sazetidine-A at both receptor subtypes. Compounds (*S*)-**9**, (*S*)-**11**, and (*S*)-**12** had less than 20% of the agonist efficacy at the human 4–2 nAChRs in comparison to sazetidine-A. Therefore, though these three new compounds showed partial agonist activities at the 4–2 nAChR subtype, their efficacies of activations were much lower than that of varenicline. Compounds (*S*)-**10**, (*S*)-**16**, and (*S*)-**17** showed no detectable agonist activity at the human 4–2 receptor subtype. Additionally, all 6 compounds showed no detectable agonist activity at the rat 3–4 nAChRs.

Consistent with our previous report,³⁵ (–)-nicotine, varenicline, and sazetidine-A potently and selectively desensitize 4–2 nAChRs with $\text{IC}_{50(10)}$ values in the nanomolar range (Table 2). To our delight, compounds (*S*)-**9**, (*S*)-**11**, (*S*)-**12**, (*S*)-**16**, and (*S*)-**17** selectively desensitize the 4–2 nAChRs. The $\text{IC}_{50(10)}$ values for these compounds to desensitize the 4–2 nAChRs ranged from 51 nM for analog (*S*)-**12** to 260 nM for analog (*S*)-**17**. In contrast, the $\text{IC}_{50(10)}$ values for these compounds in desensitizing 3–4 nAChRs were higher than 10,000 nM.

Analog (*S*)-**10** has the lowest potency in desensitizing the $\alpha 4 \beta 2$ nAChRs among all new compounds studied. Furthermore, the potency of analog (*S*)-**10** to desensitize $\alpha 4 \beta 2$ nAChRs is only slightly higher than that to desensitize $\alpha 3 \beta 4$ nAChRs. This observation confirmed the importance of the four-member ring azetidiny group in sazetidine-A, (*S*)-**9**, (*S*)-**11**, (*S*)-**12**, (*S*)-**16**, and (*S*)-**17** for selective desensitization of the $\alpha 4 \beta 2$ nAChRs.

In Vivo Effects on Alcohol Intake in Rats

In our initial behavioral studies in an animal model, effects of compound (*S*)-**9** on alcohol self-administration in selectively bred alcohol-preferring rats (P rats) were assessed (Figure 3). Three doses of compound (*S*)-**9** were tested: 0.33, 1 and 3 mg/kg (s.c.). Compared with control vehicle, compound (*S*)-**9** significantly reduced alcohol intake in a dose dependent manner. At 0.33 mg/kg, compound (*S*)-**9** did not show a significant effect at any time point. The medium dose of 1 mg/kg reduced alcohol intake significantly ($p < 0.05$) over the 24-h time point after administration. At 3 mg/kg, the compound had significant effects on alcohol intake at both the 6-h ($p < 0.05$) and 24-h time points ($p < 0.0005$) (Figure 3).

Emetic Effects of Compound (*S*)-**9** in Ferrets

The effects of compound (*S*)-**9** on causing nausea and emesis were assessed in a ferret model. Varenicline was used as a control. Administration of compound (*S*)-**9** (1 to 30 mg/kg; s.c.) did not elicit any emetic episodes or stereotypical behaviors that are indicative of nausea ($n = 6$). In general, all animals remained alert and active with periods of inactivity that mirrored their baseline behavior within the 1-h observation period.

Varenicline (0.5 mg/kg; s.c.; $n = 5$) was administered to ferrets to assess its effects on emesis or stereotypical behaviors associated with nausea. Whereas no emetic episodes were observed, drug-induced nausea as evidenced by gagging (retching; mean # of episodes 2.80 ± 1.63) were evident following the varenicline administration in the majority of the animals ($n = 4/5$) tested. In one animal, this behavior was accompanied by oral stereotype (licking), whereas in two animals it was interspersed by episodes of backward walking. On the average, these behaviors manifested around ~ 2.20 min. In general, the locomotor behavior of these animals was noticeably reduced at ~ 1.05 min post-injection and remained so for the duration of the observation period (~ 1 h). Occasionally, when animals tried to ambulate they would immediately cease their movement after each attempt. This would be followed by a momentary increase in breathing giving the impression that the animal was in distress.

DISCUSSION AND CONCLUSION

We have developed a series of sazetidine-A analogs that contain a methyl group at the 2-position of the pyridine ring. Similar to sazetidine-A, compounds (*S*)-**9**, (*S*)-**10**, (*S*)-**11**, (*S*)-**12**, (*S*)-**16**, and (*S*)-**17** are highly selective to the $\alpha 4 \beta 2$ nAChR subtype over the $\alpha 3 \beta 4$ and $\alpha 7$ subtypes in binding assays. In vitro pharmacological studies to assess their functional effects at nicotinic receptors showed that these compounds selectively desensitize $\alpha 4 \beta 2$ nAChRs. Particularly noteworthy, we found that these compounds exhibit much lower agonist activities at all nAChR subtypes tested in comparison with sazetidine-A. In rat modeling, compound (*S*)-**9** significantly reduced alcohol self-administration. It is important to note that preliminary studies in a ferret model show that compound (*S*)-**9** has a better adverse effects profile than that of varenicline.

In our previous studies, sazetidine-A exhibited much higher binding affinity for $\alpha 4 \beta 2$ nAChRs than for $\alpha 3 \beta 4$ or $\alpha 7$ receptors.^{35, 59} After cells expressing $\alpha 4 \beta 2$ nAChRs were exposed to sazetidine-A for 10 min, sazetidine-A potently inhibited nicotine-stimulated

function mediated by the $\alpha 4 \beta 2$ nAChRs. Consistent with its high selectivity for the $\alpha 4 \beta 2$ receptor, sazetidine-A at concentrations up to 10 μ M had no effect on the function of $\alpha 3 \beta 4$ or $\alpha 7$ nAChRs. These observations are consistent with our hypothesis that sazetidine-A selectively desensitizes $\alpha 4 \beta 2$ receptors. In addition to its ability to cause long lasting desensitization of $\alpha 4 \beta 2$ receptors, Zwart and co-workers reported that sazetidine-A also elicited agonist activity at the $\alpha 4 \beta 2$ receptors.⁶⁰ It is very interesting that this agonist effect is specific to the stoichiometry of the subunits. Of the two stoichiometric forms of the $\alpha 4 \beta 2$ receptors, sazetidine-A showed full agonist activity at the $(\alpha 4)_2(\beta 2)_3$ receptors but nearly no agonist activity at the $(\alpha 4)_3(\beta 2)_2$ receptors.⁶¹ Thus, sazetidine-A, like nicotine itself, has both agonist and desensitizer effects on $\alpha 4 \beta 2$ containing nAChRs, but in contrast to nicotine, its desensitizing effects appear to predominate and be long lasting.^{35, 59, 62} For example, initial activation of the $\alpha 4 \beta 2$ nAChRs *in vitro* by sazetidine-A is brief, lasting only seconds, whereas the receptor desensitization that immediately follows activation lasts more than an hour even with continual washing of the cells.

Many studies to further understand the *in vitro* pharmacological properties of sazetidine-A and its behavioral effects in animal models have been reported.^{47-50, 52, 63-69} The results from these studies are highly supportive of our proposal to develop novel nicotinic therapeutics based on their ability to selectively desensitize $\alpha 4 \beta 2$ nAChRs. In the present study, we investigated the agonist and desensitization properties of these new analogs at human $\alpha 4 \beta 2$ nAChRs and rat $\alpha 3 \beta 4$ nAChRs (Table 2). For the efficacies of agonist activities at $\alpha 4 \beta 2$ nAChRs in comparison to nicotine, compounds (*S*)-**9**, (*S*)-**11**, and (*S*)-**12** showed very low relative E_{max} values at the $\alpha 4 \beta 2$ nAChRs as follows: 15%, 17% and 20%, respectively. Compounds (*S*)-**10**, (*S*)-**16**, and (*S*)-**17** did not show any measureable agonist activities at this subtype, and none of the compounds tested exhibited agonist activities at rat $\alpha 3 \beta 4$ nAChRs. It is important to note that all the new analogs tested showed significantly lower agonist activities at both $\alpha 4 \beta 2$ and $\alpha 3 \beta 4$ receptor subtypes than those of varenicline or sazetidine-A. With regards to desensitization activities, compounds (*S*)-**9**, (*S*)-**11**, (*S*)-**12**, (*S*)-**16**, and (*S*)-**17** potently and selectively desensitized the $\alpha 4 \beta 2$ nAChRs after a 10-min incubation period (Table 2). These *in vitro* pharmacological functional properties indicate the possibility for these compounds to desensitize $\alpha 4 \beta 2$ nAChRs *in vivo*.

As shown in Figure 3, compound (*S*)-**9** significantly reduced alcohol intake in a dose-dependent manner in the animal efficacy model using selectively bred alcohol preferring rats. Based on the *in vitro* pharmacological property profile of compound (*S*)-**9**, we speculate that the ability of compound (*S*)-**9** in reducing alcohol intake resulted mainly from its $\alpha 4 \beta 2$ nAChR desensitization properties. In our previous post-deprivation studies,⁴⁷ sazetidine-A at 3 mg/kg significantly reduced alcohol intake at the 2-, 4-, and 6-hour time points but not at the 24-hour time point. Interestingly, (*S*)-**9** showed a different pattern of the inhibition, significantly reducing the alcohol intake at the 6- and 24-hour time points but not at the 2- and 4-hour points. It is important to note that though the binding affinity of (*S*)-**9** to $\alpha 4 \beta 2$ nAChRs is lower than that of sazetidine-A (Table 1), both compounds reduced alcohol intake at 3 mg/kg. It is conceivable that (*S*)-**9** might have reached a higher brain concentration than sazetidine-A at the same dose given systemically.

Varenicline is marketed in the U.S. and many other countries as a smoking cessation aid. Clinical trials before and after its approval as well as post-marketing reports demonstrated its superior efficacy compared with placebo or other marketed smoking cessation therapeutics.^{28, 29, 70-72} In addition to its use as a smoking cessation drug, the significant efficacy of varenicline in reducing alcohol consumption in heavy-drinking smokers indicated the possibility for it to be used to treat alcohol use disorders.³² However, notable adverse events related to varenicline have been reported,^{28, 29, 34} which resulted in the recent ruling by the FDA that varenicline carry a black box label for suicidal ideation and suicidal

behavior (FDA Drug Safety Newsletter 2009). The most common adverse events reported for varenicline are nausea and vomiting. For example, 30-40% of people taking varenicline reported nausea in randomized clinical trials, which is greater than 3 times that reported for the placebo group.⁷³ The adverse side effects of varenicline are thought to be mediated through its strong agonist activities at the $\alpha 3$ $\beta 4$ and $\alpha 7$ nAChR subtypes.⁵⁸ As shown in Table 2, compound (*S*)-**9** did not show any agonist activity at the rat $\alpha 3$ $\beta 4$ nAChRs. In addition, preliminary studies using whole cell current measurements showed that compound (*S*)-**9** did not elicit any detectable current in cells expressing rat $\alpha 7$ nAChRs (Sahibzada et al, data not shown). Based on a comparison of the in vitro pharmacological property profile of compound (*S*)-**9** with that of varenicline, it is conceivable that compound (*S*)-**9** might be much less likely to induce nausea and vomiting than varenicline. The animal studies in a ferret nausea and emetic model showed a clear difference between varenicline and compound (*S*)-**9** in their potential to induce nausea and vomiting. Varenicline at 0.5 mg/kg (s.c., n=5) caused evident signs related to nausea and vomiting in 4 ferrets. In contrast, compound (*S*)-**9** at doses up to 30 mg/kg (s.c.) did not elicit any emetic episodes or stereotypical behaviors that are indicative of nausea (n=6).

In summary, a series of novel sazetidine-A analogs were designed and synthesized. The in vitro pharmacological evaluations revealed that these compounds were selective $\alpha 4$ $\beta 2$ nAChR desensitizers. Analog (*S*)-**9**, our lead compound, showed significant efficacy in reducing alcohol intake in a rat model. More importantly, analog (*S*)-**9** did not induce nausea and vomiting up to 30 mg/kg in ferrets, exhibiting a potentially better adverse side effect profile than that of varenicline. Therefore, we concluded that this series of analogs should be further studied in the development of new treatments for alcohol use disorders.

EXPERIMENTAL

General Chemistry Methods

All solvents and reagents were used as obtained from commercial sources unless otherwise indicated. All reactions were performed under nitrogen atmosphere unless otherwise noted. The ^1H and ^{13}C NMR spectra were recorded on a Varian 400MR spectrometer operating at 400 MHz for ^1H and 100 MHz for ^{13}C . Deuterated chloroform or deuterated methanol was used as the solvent for NMR experiments. ^1H Chemical shifts value (δ) are referenced to the residual non-deuterated components of the NMR solvents (δ =7.26 ppm for CHCl_3 and δ =3.31 ppm for CHD_2OD). The ^{13}C chemical shifts (δ) are referenced to CDCl_3 (central peak, δ =77.0 ppm) or CD_3OD (δ =49.0 ppm) as the internal standard. Optical rotation was detected on ADP220 Automatic polarimeter from Bellingham & Stanley Limited. Mass spectra were measured in positive mode electrospray ionization (ESI). The HRMS data were obtained on a Waters Q-TOF Premier mass spectrometer. TLC was performed on silica gel 60 F₂₅₄ plastic sheets. Column chromatography was performed using silica gel (35-75 mesh). Combustion analyses were performed at Atlantic Microlabs, Inc. Norcross, Ga.

General procedure for the Mitsunobu reaction: To a mixture of the N-Boc protected alcohol (1.0 eq), the 5-halogen-3-pyridinol (1.0 eq), and Ph_3P (1.3 eq) in anhydrous THF (0.10 M) was added DEAD (1.3 eq) dropwise at 0 °C under nitrogen atmosphere. After stirring for 2 days at room temperature, the solvent was removed under reduced pressure. The residue was purified by column chromatography on silica gel using a gradient of hexane-ethyl acetate (10:1 to 5:1) as the eluent to give the product in 68-90% yield.

General Procedure for the Sonogashira coupling reaction: A mixture of the Mitsunobu adduct (1.0 eq), alkyne (4.0 eq), $\text{Pd}(\text{PPh}_3)_2\text{Cl}_2$ (5 mol %), CuI (10 mol %), PPh_3 (10 mol %) in $\text{Et}_3\text{N}/\text{DMSO}$ (10:1, 0.12 M) was heated to 95 °C under a nitrogen atmosphere in a sealed tube for 60 h. The cooled reaction mixture was taken up in ethyl acetate, and the organic

phase was washed with water, brine, and then dried over Na₂SO₄. The extract was concentrated under a reduced pressure, and the residue was purified by column chromatography on silica gel using a gradient of CH₂Cl₂-ethyl acetate (16:1 to 10:1) as the eluent to give the product in 88-98% yield.

General Procedure (method A) for deprotection of the N-Boc group: To a stirred solution of the Sonogashira adduct (1.0 eq) in dichloromethane (0.10 M) was added trifluoroacetic acid (32 eq) dropwise at 0 °C under a nitrogen atmosphere. The reaction mixture was stirred for 3 h at room temperature. The solvent and excess TFA was then removed under reduced pressure. To the residue was added 2-3ml of methanol was added followed by dropwise addition of 10% aqueous NaOH solution at 0 °C until the pH value of the mixture was 9-10. After the mixture was stirred at room temperature for 30 min, the solution was taken up in dichloromethane, and the organic phase was washed with brine, dried over Na₂SO₄. The extract was concentrated under reduced pressure. The residue was purified by column chromatography on silica gel using a gradient of CH₂Cl₂-methanol (20:1 to 10:1) as the eluent to give the product in 75-83% yield.

General Procedure (method B) for deprotection of the N-Boc group: To the Boc-protected compound (1.0 mmol), the solution of 2 M HCl in methanol (10 ml) was added at 0 °C under a nitrogen atmosphere. The reaction mixture was allowed to warm to room temperature and stirred for 3 hours. The reaction mixture was concentrated under a reduced pressure, and the residue was purified by column chromatography on silica gel using a gradient of CH₂Cl₂-methanol (10:1 to 5:1) as the eluent to give the product as white solid in 73-92% yield.

(S)-tert-Butyl 2-((5-Bromo-2-methylpyridin-3-yloxy)methyl)azetidine-1-carboxylate ((S)-3)—Yield: 69% (light red solid). ¹H NMR (CDCl₃, 400MHz): 8.15 (s, 1H), 7.26 (s, 1H), 4.52 (m, 1H), 4.33 (m, 1H), 4.05 (m, 1H), 3.90 (m, 2H), 2.45 (s, 3H), 2.34 (m, 2H), 1.41 (s, 9H).

(R)-tert-Butyl 2-((5-bromo-2-methylpyridin-3-yloxy)methyl)azetidine-1-carboxylate ((R)-3)—Yield: 77% (light red solid). ¹H NMR (CDCl₃, 400MHz): 8.16 (s, 1H), 7.26 (s, 1H), 4.53 (m, 1H), 4.34 (m, 1H), 4.06 (m, 1H), 3.91 (m, 2H), 2.45 (s, 3H), 2.35 (m, 2H), 1.42 (s, 9H).

(S)-tert-Butyl 2-((5 – bromo –2– methylpyridin –3 yloxy)methyl) pyrrolidine-1-carboxylate ((S)-4)—Yield: 68% (light red solid). ¹H NMR (CDCl₃, 400MHz): 8.13 (s, 1H), 7.26 (br s, 1H), 4.15 (m, 2H), 4.00 (m, 0.5H), 3.85 (m, 0.5H), 3.41 (m, 2H), 2.41 (s, 3H), 1.96 (m, 4H), 1.48 (s, 9H).

(R)-tert-Butyl 2-((5 – bromo –2– methylpyridin –3-yloxy)methyl) pyrrolidine-1-carboxylate ((R)-4)—Yield: 70% (light red solid). ¹H NMR (CDCl₃, 400MHz): 8.12 (s, 1H), 7.26 (br s, 1H), 4.14 (m, 2H), 3.99 (m, 0.5H), 3.84 (m, 0.5H), 3.41 (m, 2H), 2.40 (s, 3H), 1.95 (m, 4H), 1.47 (s, 9H).

(S)-tert-Butyl 2-((5-(hex-1-ynyl)-2-methylpyridin-3-yloxy)methyl) azetidine-1-carboxylate ((S)-5)—Yield: 95% (light yellow oil). ¹H NMR (CDCl₃, 400MHz): 8.12 (d, 1H, *J*=1.2 Hz), 7.10 (d, 1H, *J*=1.2 Hz), 4.52 (m, 1H), 4.32 (m, 1H), 4.05 (dd, 1H, *J*=10, 2.8 Hz), 3.90 (m, 2H), 2.48 (s, 3H), 2.41 (t, 2H, *J*=7.2 Hz), 2.33 (m, 2H), 1.59 (m, 2H), 1.48 (m, 2H), 1.41 (s, 9H), 0.95 (t, 3H, *J*=7.2 Hz).

(R)-tert-Butyl 2-((5-(hex-1-ynyl)-2-methylpyridin-3-yloxy)methyl) azetidine-1-carboxylate ((R)-5)—Method B was used. Yield: 93% (light yellow oil). ¹H NMR

(CDCl₃, 400MHz): 8.10 (d, 1H, *J*=1.2 Hz), 7.09 (d, 1H, *J*=1.2 Hz), 4.50 (m, 1H), 4.30 (m, 1H), 4.03 (dd, 1H, *J*=10, 2.8 Hz), 3.89 (m, 2H), 2.46 (s, 3H), 2.39 (t, 2H, *J*=7.2 Hz), 2.31 (m, 2H), 1.57 (m, 2H), 1.46 (m, 2H), 1.39 (s, 9H), 0.94 (t, 3H, *J*=7.2 Hz).

(S)-tert-Butyl 2-((5-(hex-1-ynyl)-2-methylpyridin-3-yloxy)methyl) pyrrolidine-1-carboxylate ((S)-6)—Yield: 98% (light yellow oil). ¹H NMR (CDCl₃, 400MHz): 8.10 (s, 1H), 7.11 (br s, 1H), 4.14 (m, 2H), 4.00 (m, 0.5H), 3.83 (m, 0.5H), 3.42 (m, 2H), 2.44 (s, 3H), 2.41 (t, 2H, *J*=7.2 Hz), 1.96 (m, 4H), 1.60 (m, 2H), 1.48 (m, 11H), 0.95 (t, 3H, *J*=7.2 Hz).

(R)-tert-Butyl 2-((5-(hex-1-ynyl)-2-methylpyridin-3-yloxy)methyl) pyrrolidine-1-carboxylate ((R)-6)—Method B was used. Yield: 92% (light yellow oil). ¹H NMR (CDCl₃, 400MHz): 8.09 (s, 1H), 7.10 (br s, 1H), 4.13 (m, 2H), 3.99 (m, 0.5H), 3.82 (m, 0.5H), 3.41 (m, 2H), 2.43 (s, 3H), 2.40 (t, 2H, *J*=7.2 Hz), 1.95 (m, 4H), 1.59 (m, 2H), 1.48 (m, 11H), 0.95 (t, 3H, *J*=7.2 Hz).

(S)-tert-Butyl 2-((5-(cyclopropylethynyl)-2-methylpyridin-3-yloxy)methyl)azetidine-1-carboxylate ((S)-7)—Yield: 97% (light yellow oil). ¹H NMR (CDCl₃, 400MHz): 8.09 (d, 1H, *J*=1.2 Hz), 7.08 (d, 1H, *J*=1.2 Hz), 4.51 (m, 1H), 4.29 (m, 1H), 4.03 (dd, 1H, *J*=10, 2.8 Hz), 3.89 (m, 2H), 2.46 (s, 3H), 2.32 (m, 2H), 1.42 (m, 1H), 1.40 (s, 9H), 0.87 (m, 2H), 0.80 (m, 2H).

(S)-tert-Butyl 2-((5-(6-hydroxyhex-1-ynyl) 2-methylpyridin-3-yloxy) methyl) azetidine-1-carboxylate ((S)-8)—Yield: 89% (yellow oil). ¹H NMR (CDCl₃, 400 MHz): 8.04 (br s, 1H), 7.06 (s, 1H), 4.46 (m, 1H), 4.25 (m, 1H), 3.98 (dd, 1H, *J*=10, 2.0 Hz), 3.84 (m, 2H), 3.62 (t, 2H, *J*=6.0 Hz), 2.91 (br s, 1H), 2.41 (s, 3H), 2.39 (t, 2H, *J*=6.0 Hz), 2.26 (m, 2H), 1.65 (m, 4H), 1.34 (s, 9H).

(S)-3-(Azetidin-2-ylmethoxy)-5-(hex-1-ynyl)-2-methylpyridine ((S)-9)—Method A was used. Yield: 83% (light red oil). ¹H NMR (CDCl₃, 400MHz): 8.08 (d, 1H, *J*=1.6 Hz), 7.04 (d, 1H, *J*=1.6 Hz), 4.25 (m, 1H), 3.96 (m, 2H), 3.67 (m, 1H), 3.47 (m, 1H), 2.42 (s, 3H), 2.38 (m, 4H), 2.24 (m, 1H), 1.57 (m, 2H), 1.46 (m, 2H), 0.92 (t, 3H, *J*=7.2 Hz). ¹³C NMR (CDCl₃, 100MHz): 152.3, 148.0, 143.1, 119.7, 118.8, 92.7, 77.4, 72.4, 57.1, 44.3, 30.6, 23.9, 22.0, 19.2, 19.1, 13.5. HRMS (ESI) *m/z* calcd for C₁₆H₂₂N₂O (M + H)⁺ 259.1810, found 259.1813; [*D*₂₄ = 4.3 (*c* 0.77, CHCl₃). Anal. Calcd for C₁₆H₂₂N₂O•0.625H₂O: C, 71.28; H, 8.69; N, 10.39; O, 9.64. Found: C, 71.25; H, 8.46; N, 10.13; O, 9.89.

(R)-3-(Azetidin-2-ylmethoxy)-5-(hex-1-ynyl)-2-methylpyridine ((R)-9)—Method A was used. Yield: 75% (light red oil). ¹H NMR (CDCl₃, 400MHz): 8.07 (d, 1H, *J*=1.6 Hz), 7.04 (d, 1H, *J*=1.6 Hz), 4.24 (m, 1H), 3.94 (m, 2H), 3.66 (m, 1H), 3.45 (m, 1H), 2.41 (s, 3H), 2.36 (m, 3H), 2.24 (m, 2H), 1.56 (m, 2H), 1.45 (m, 2H), 0.92 (t, 3H, *J*=7.2 Hz). ¹³C NMR (CDCl₃, 100MHz): 152.3, 148.0, 143.1, 119.7, 118.8, 92.7, 77.4, 72.5, 57.1, 44.3, 30.6, 24.0, 22.0, 19.2, 19.0, 13.5. HRMS (ESI) *m/z* calcd for C₁₆H₂₂N₂O (M + H)⁺ 259.1810, found 259.1816; [*D*₂₅ = +10.8 (*c* 0.62, CHCl₃). Anal. Calcd for C₁₆H₂₂N₂O•0.375H₂O: C, 72.49; H, 8.65; N, 10.57. Found: C, 72.72; H, 8.67; N, 10.43.

(S)-5-(Hex-1-ynyl)-2-methyl-3-(pyrrolidin-2-ylmethoxy)pyridine ((S)-10)—Method A was used. Yield: 80% (light red oil). ¹H NMR (CDCl₃, 400MHz): 8.07 (d, 1H, *J*=1.6 Hz), 7.02 (d, 1H, *J*=1.6 Hz), 3.84 (m, 2H), 3.52 (m, 1H), 2.98 (m, 2H), 2.42 (s, 3H), 2.38 (t, 2H, *J*=7.2 Hz), 2.11 (br s, 1H), 1.93 (m, 1H), 1.78 (m, 2H), 1.56 (m, 3H), 1.45 (m, 2H), 0.92 (t, 3H, *J*=7.2 Hz). ¹³C NMR (CDCl₃, 100MHz): 152.3, 147.8, 143.0, 119.6, 118.8, 92.6, 77.5, 71.6, 57.0, 46.7, 30.6, 28.0, 25.4, 22.0, 19.3, 19.0, 13.5. HRMS (ESI) *m/z* calcd for

$C_{17}H_{24}N_2O$ ($M + H$)⁺ 273.1967, found 273.1964; [δ]_D²⁵ = +6.4 (c 1.05, $CHCl_3$). Anal. Calcd for $C_{17}H_{24}N_2O \cdot 0.375H_2O$: C, 73.15; H, 8.94; N, 10.04. Found: C, 73.29; H, 8.83; N, 9.99.

(R)-5-(Hex-1-ynyl)-2-methyl-3-(pyrrolidin-2-ylmethoxy) pyridine ((R)-10)—

Method A was used. Yield: 83% (light red oil). ¹H NMR ($CDCl_3$, 400MHz): 8.06 (d, 1H, J =1.6 Hz), 7.02 (d, 1H, J =1.6 Hz), 3.84 (m, 2H), 3.52 (m, 1H), 2.97 (m, 2H), 2.66 (br s, 1H), 2.40 (s, 3H), 2.37 (t, 2H, J =7.2 Hz), 1.93 (m, 1H), 1.78 (m, 2H), 1.55 (m, 3H), 1.44 (m, 2H), 0.91 (t, 3H, J =7.2 Hz). ¹³C NMR ($CDCl_3$, 100MHz): 152.2, 147.8, 143.0, 119.6, 118.8, 92.6, 77.4, 71.4, 57.0, 46.6, 30.6, 27.9, 25.3, 21.9, 19.2, 19.0, 13.5. HRMS (ESI) m/z calcd for $C_{17}H_{24}N_2O$ ($M + H$)⁺ 273.1967, found 273.1960; [δ]_D²⁵ = -4.4 (c 0.85, $CHCl_3$). Anal. Calcd for $C_{17}H_{24}N_2O \cdot 0.75H_2O$: C, 71.42; H, 8.99; N, 9.80. Found: C, 71.58; H, 8.80; N, 9.51.

(S)-3-(Azetidin-2-ylmethoxy)-5-(cyclopropylethynyl)-2-methylpyridine dihydrochloride ((S)-11)—

Method B was used. Yield: 91% (off white solid). ¹H NMR (CD_3OD , 400MHz): 8.18 (s, 1H), 7.72 (s, 1H), 4.93 (m, 1H), 4.47 (m, 2H), 4.11 (m, 2H), 2.71 (m, 2H), 2.60 (s, 3H), 1.54 (m, 1H), 0.95 (m, 2H), 0.81 (m, 2H). ¹³C NMR (CD_3OD , 100MHz): 154.4, 147.3, 140.2, 125.7, 122.8, 100.2, 71.5, 68.9, 60.3, 45.0, 21.9, 17.5, 9.3 (2C), 0.7. HRMS (ESI) m/z calcd for $C_{15}H_{18}N_2O$ ($M + H$)⁺ 243.1497, found 243.1509. [δ]_D²⁶ = -8.1 (c 0.82, MeOH). Anal. Calcd for $C_{15}H_{18}N_2O \cdot 2HCl \cdot 2.5H_2O$: C, 50.01; H, 6.99; N, 7.78. Found: C, 49.64; H, 6.82; N, 8.12.

(S)-6-(5-(Azetidin-2-ylmethoxy)-6-methylpyridin-3-yl)hex-5-yn-1-ol dihydrochloride ((S)-12)—

Method B was used. Yield: 87% (white solid). ¹H NMR (CD_3OD , 400MHz): 8.37 (s, 1H), 8.12 (s, 1H), 4.97 (m, 1H), 4.59 (m, 2H), 4.13 (m, 2H), 3.61 (t, 2H, J =5.2 Hz), 2.71 (m, 5H), 2.56 (m, 2H), 1.71 (m, 4H). ¹³C NMR (CD_3OD , 100MHz): 155.6, 146.2, 136.2, 129.4, 124.4, 99.3, 75.5, 69.6, 62.3, 60.0, 44.9, 32.7, 25.8, 21.8, 19.9, 15.7. HRMS (ESI) m/z calcd for $C_{16}H_{22}N_2O_2$ ($M + H$)⁺ 275.1760, found 275.1761. [δ]_D²⁶ = -7.1 (c 0.94, MeOH). Anal. Calcd for $C_{16}H_{22}N_2O_2 \cdot 2HCl \cdot H_2O$: C, 52.61; H, 7.17; N, 7.67. Found: C, 52.54; H, 6.92; N, 7.92.

(S)-tert-Butyl 2-(2-methyl-5-(6-(tosyloxy) hex-1-ynyl) pyridine-3-yloxy) methyl azetidine-1-carboxylate ((S)-13)—

To a stirred solution of **8** (374 mg, 1 mmol) in dry dichloromethane (25ml) was added DMAP (12 mg, 0.1 mmol), Et₃N (0.34 ml, 2.4 mmol) and tosyl chloride (228 mg, 1.2 mmol) at 0 °C under a nitrogen atmosphere. The reaction mixture was stirred overnight at room temperature. The mixture was taken up in dichloromethane, and the organic phase was washed with saturated aqueous NaHCO₃ solution, brine, and then dried over Na₂SO₄. The extract was concentrated under reduced pressure, and then residue was purified by column chromatography on silica gel using a gradient of hexane-ethyl acetate (6:1 to 3:1) as the eluent to give the product (**13**) 422 mg as light yellow oil, Yield: 80%. ¹H NMR ($CDCl_3$, 400MHz): 8.05 (s, 1H), 7.75 (d, 2H, J =8.4 Hz), 7.30 (d, 2H, J =8.0 Hz), 7.07 (s, 1H), 4.49 (m, 1H), 4.29 (m, 1H), 4.06 (t, 2H, J =6.4 Hz), 4.01 (dd, 1H, J =10.4, 2.4 Hz), 3.87 (m, 2H), 2.44 (s, 3H), 2.40 (s, 3H), 2.36 (t, 2H, J =6.8 Hz), 2.28 (m, 2H), 1.80 (m, 2H), 1.61 (m, 2H), 1.37 (s, 9H).

(S)-tert-Butyl 2-((5-(6-chlorohex-1-ynyl)-2-methylpyridin-3-yloxy)methyl) azetidine-1-carboxylate ((S)-14)—

A mixture of the compound **8** (374 mg, 1 mmol), PPh₃ (288 mg, 1.1 mmol), NaHCO₃ (20 mg, 0.24mmol) in dry CCl₄ (15ml) was heated to 80 °C under a nitrogen atmosphere in a sealed tube for 48 h. The cooled reaction mixture was concentrated under a reduced pressure, and the residue was purified by column chromatography on silica gel using a gradient of hexane-ethyl acetate (6:1 to 3:1) as the eluent to give the product **14** 271 mg as light yellow oil, Yield: 69%. ¹H NMR ($CDCl_3$,

400MHz): 8.09 (d, 1H, $J=1.2$ Hz), 7.08 (d, 1H, $J=1.2$ Hz), 4.49 (m, 1H), 4.30 (m, 1H), 4.02 (dd, 1H, $J=10$, 2.8 Hz), 3.88 (m, 2H), 3.57 (t, 2H, $J=6.4$ Hz), 2.45 (s, 3H), 2.44 (t, 2H, $J=6.8$ Hz), 2.31 (m, 2H), 1.92 (m, 2H), 1.74 (m, 2H), 1.38 (s, 9H).

(S)-tert-Butyl 2-((5-(6-fluorohex-1-ynyl)-2-methylpyridin-3-yloxy)methyl)azetidine-1-carboxylate ((S)-15)—A mixture of **13** (67 mg, 0.13 mmol), Kryptofix 2.2.2 (62 mg, 0.16 mmol), and KF (10 mg, 0.17 mmol) in dry THF (3 mL) was heated to reflux overnight under a nitrogen atmosphere. The cooled reaction mixture was concentrated under reduced pressure. The residue was purified by column chromatography on silica gel using a gradient of hexane-ethyl acetate (4:1) as the eluent to give the product 40 mg as light yellow oil, Yield: 85%. ^1H NMR (CDCl_3 , 400 MHz): 8.11 (s, 1H), 7.10 (s, 1H), 4.56 (t, 1H, $J=6.0$ Hz), 4.51 (m, 1H), 4.44 (t, 1H, $J=6.0$ Hz), 4.31 (m, 1H), 4.05 (dd, 1H, $J=10$, 2.8 Hz), 3.90 (m, 2H), 2.48 (m, 5H), 2.33 (m, 2H), 1.90 (m, 1H), 1.83 (m, 1H), 1.74 (m, 2H), 1.41 (s, 9H).

(S)-3-(Azetidin-2-ylmethoxy)-5-(6-fluorohex-1-ynyl)-2-methylpyridine dihydrochloride ((S)-16)—Method B was used. Yield: 92% (white solid). ^1H NMR (CD_3OD , 400 MHz): 8.09 (s, 1H), 7.49 (s, 1H), 4.91 (m, 1H), 4.54 (t, 1H, $J=6.0$ Hz), 4.42 (t, 1H, $J=6.0$ Hz), 4.41 (m, 2H), 4.11 (m, 2H), 2.69 (m, 2H), 2.53 (m, 5H), 1.89 (m, 1H), 1.82 (m, 1H), 1.73 (m, 2H). ^{13}C NMR (CD_3OD , 100 MHz): 153.6, 148.6, 143.1, 123.0, 121.5, 94.6, 84.5 (d, $J=163.1$ Hz), 77.9, 68.4, 60.5, 45.0, 30.8 (d, $J=20.3$ Hz), 25.6 (d, $J=4.9$ Hz), 22.0, 19.8, 18.7. ^{19}F NMR (CD_3OD , 376 MHz): -220.6. HRMS (ESI) m/z calcd for $\text{C}_{16}\text{H}_{21}\text{FN}_2\text{O}$ ($\text{M} + \text{H}$) $^+$ 277.1716, found 277.1714. [D] $^{25} = -7.2$ (c 1.16, MeOH). Anal. Calcd for $\text{C}_{16}\text{H}_{21}\text{FN}_2\text{O} \cdot 2\text{HCl} \cdot 2.5\text{H}_2\text{O}$: C, 48.74; H, 7.16; N, 7.10. Found: C, 48.69; H, 6.84; N, 7.06.

(S)-3-(Azetidin-2-ylmethoxy)-5-(6-chlorohex-1-ynyl)-2-methylpyridine dihydrochloride ((S)-17)—Method B was used. Yield: 90% (white solid). ^1H NMR (CD_3OD , 400 MHz): 8.39 (s, 1H), 8.17 (s, 1H), 4.98 (m, 1H), 4.62 (m, 2H), 4.13 (m, 2H), 3.64 (t, 2H, $J=6.4$ Hz), 2.73 (m, 5H), 2.58 (t, 2H, $J=6.8$ Hz), 1.96 (m, 2H), 1.80 (m, 2H). ^{13}C NMR (CD_3OD , 100 MHz): 154.3, 144.8, 134.4, 128.4, 122.9, 97.5, 74.1, 68.2, 58.6, 43.8, 43.4, 31.4, 25.1, 20.3, 18.0, 14.1. HRMS (ESI) m/z calcd for $\text{C}_{16}\text{H}_{21}\text{ClN}_2\text{O}$ ($\text{M} + \text{H}$) $^+$ 293.1421, found 293.1430. [D] $^{25} = -4.6$ (c 0.72, MeOH). Anal. Calcd. for $\text{C}_{16}\text{H}_{21}\text{ClN}_2\text{O} \cdot 2\text{HCl} \cdot 0.75\text{H}_2\text{O}$: C, 50.67; H, 6.51; N, 7.39. Found: C, 50.31; H, 6.12; N, 7.59.

5-Bromo-2-(trifluoromethyl)pyridin-3-ol (19)—A solution of 5-bromo-3-methoxy-2-(trifluoromethyl)pyridine (**18**) (0.87 g, 3.4 mmol) in a mixture of acetic acid (6 mL) and 33% HBr in acetic acid (12 mL) was stirred at 110 °C in a sealed tube overnight. Then potassium sodium tartrate tetrahydrate (7.8 g) was added slowly at 0 °C. The mixture was taken up in ethyl acetate after stirring for 10 minutes, the organic phase was washed with brine, and dried over sodium sulfate. The extract was concentrated under reduced pressure. The residue was purified by column chromatography on silica gel using a gradient of hexanes-ethyl acetate (1:2) as the eluent to give 0.65 g of the product (**19**) as gray solid with a yield of 79%. ^1H NMR (CD_3OD , 400 MHz): 8.18 (s, 1H), 7.58 (s, 1H). ^{13}C NMR (CD_3OD , 100 MHz): 154.6, 141.4, 134.7 (q, $J=34$ Hz), 128.8, 125.2, 123.3 (q, $J=272$ Hz). ^{19}F NMR (CDCl_3 , 376 MHz): -68.0. HRMS (ESI) m/z calcd for $\text{C}_6\text{H}_3\text{BrF}_3\text{NO}$ ($\text{M} + \text{H}$) $^+$ 241.9428, found 241.9434.

(S)-tert-Butyl 2-((5-bromo-2-(trifluoromethyl)pyridin-3-yloxy)methyl)azetidine-1-carboxylate ((S)-20)—General Procedure for the Mitsunobu Reaction was

used. Yield: 88% (brown oil). ^1H NMR (CDCl_3 , 400 MHz): 8.31 (s, 1H), 7.62 (s, 1H), 4.55 (m, 2H), 4.11 (m, 1H), 3.88 (m, 2H), 2.36 (m, 2H), 1.41 (s, 9H).

(S)-tert-Butyl 2-((5-(hex-1-ynyl)-2-(trifluoromethyl)pyridin-3-yloxy)methyl)azetidine-1-carboxylate ((S)-21)—A mixture of compound (20) (290 mg, 0.7 mmol), 1-Hexyne (0.3 ml, 2.8 mmol), $\text{Pd}(\text{PPh}_3)_2\text{Cl}_2$ (25mg, 0.036mmol), CuI (13.5 mg, 0.071 mmol), PPh_3 (18.6 mg, 0.071 mmol) in $\text{Et}_3\text{N}/\text{DMSO}$ (5 ml/0.5 ml) was heated to 50 °C under a nitrogen atmosphere overnight. The cooled reaction mixture was taken up in ethyl acetate, and the organic phase was washed with water, brine, and then dried over Na_2SO_4 . The extract was concentrated under a reduced pressure, and the residue was purified by column chromatography on silica gel using a gradient of CH_2Cl_2 -ethyl acetate (50:1 to 20:1) as the eluent to give the product 265 mg as yellow oil, yield: 91%. ^1H NMR (CDCl_3 , 400 MHz): 8.22 (s, 1H), 7.40 (s, 1H), 4.52 (m, 2H), 4.10 (m, 1H), 3.89 (m, 2H), 2.44 (t, 2H, $J=7.2$ Hz), 2.36 (m, 2H), 1.59 (m, 2H), 1.48 (m, 2H), 1.41 (s, 9H), 0.96 (t, 3H, $J=7.2$ Hz).

(S)-3-(Azetidin-2-ylmethoxy)-5-(hex-1-ynyl)-2-(trifluoromethyl) pyridine hydrochloride ((S)-22)—Method B was used. Yield: 73% (white solid). ^1H NMR (CD_3OD , 400MHz): 8.25 (s, 1H), 7.74 (s, 1H), 4.81 (m, 1H), 4.46 (m, 2H), 4.02 (m, 2H), 2.72 (m, 1H), 2.60 (m, 1H), 2.51 (t, 2H, $J=7.2$ Hz), 1.63 (m, 2H), 1.52 (m, 2H), 0.98 (t, 3H, $J=7.2$ Hz). ^{13}C NMR (CD_3OD , 100MHz): 153.8, 144.3, 135.5 (q, $J=34$ Hz), 127.4, 125.2, 123.1 (q, $J=272$ Hz), 98.3, 77.1, 69.9, 59.7, 44.7, 31.6, 23.0, 22.1, 19.8, 13.9. ^{19}F NMR (CD_3OD , 376MHz): -67.3. HRMS (ESI) m/z calcd for $\text{C}_{16}\text{H}_{19}\text{F}_3\text{N}_2\text{O}$ ($\text{M} + \text{H}$) $^+$ 313.1528, found 313.1525. [D] $^{25}=-3.9$ (c 0.86, MeOH). Anal. Calcd for $\text{C}_{16}\text{H}_{19}\text{F}_3\text{N}_2\text{O}\cdot\text{HCl}$: C, 55.10; H, 5.78; N, 8.03. Found: C, 55.49; H, 5.65; N, 7.87.

Cell Lines and Cell Culture

The cell lines expressing defined rat nAChR subtypes were established previously by stably transfecting HEK 293 cells with rat nAChR subunit genes.^{74, 75} The cell line expressing human $\alpha 2$ nAChRs, YX 4 2H1, was established recently. These cell lines were maintained in minimum essential medium (MEM) supplemented with 10% fetal bovine serum, 100 units/ml penicillin G, 100 mg/ml streptomycin and selective antibiotics at 37°C with 5% CO_2 in a humidified incubator. Fetal bovine serum was provided by Gemini Bio-Products (Woodland, CA). Tissue culture medium and antibiotics were obtained from Invitrogen Corporation (Carlsbad, CA), unless otherwise stated.

[^3H] Epibatidine Radioligand Binding Assay

Membrane preparation procedures and binding assays were described previously.^{74, 75} Briefly, cultured cells at >80% confluence were removed from their flasks (80 cm^2) with a disposable cell scraper and placed in 10 ml of 50 mM Tris-HCl buffer (pH 7.4, 4 °C). The cell suspension was centrifuged at $10,000 \times g$ for 5 min and the pellet was collected. The cell pellet was then homogenized in 10 ml buffer with a polytron homogenizer and centrifuged at $36,000 g$ for 10 min at 4 °C. The membrane pellet was resuspended in fresh buffer, and aliquots of the membrane preparation were used for binding assays. The concentration of [^3H]-epibatidine used was ~500 pM for competition binding assays. Nonspecific binding was assessed in parallel incubations in the presence of 300 μM nicotine. Bound and free ligands were separated by vacuum filtration through Whatman GF/C filters treated with 0.5% polyethylenimine. The filter-retained radioactivity was measured by liquid scintillation counting. Specific binding was defined as the difference between total binding and nonspecific binding. Data from competition binding assays were analyzed using Prism 5 (GraphPad Software, San Diego, CA). The K_d values for [^3H]-epibatidine used for calculating K_i values of nAChR subtypes were 0.02 nM for $\alpha 2$, 0.08 nM for $\alpha 4$, 0.03

nM for 3 2, 0.3 nM for 3 4, 0.04 nM for 4 2, 0.09 nM for 4 4, 1.8 nM for 7 and 0.05 for rat forebrain.

⁸⁶Rb⁺ Efflux Assay

Functional properties of compounds at nAChRs expressed in the transfected cells were measured using ⁸⁶Rb⁺ efflux assays as described previously.^{35, 75} In brief, cells expressing human 4 2 or rat 3 4 nAChRs were plated into 24-well plates coated with poly-D-lysine. The plated cells were grown at 37 °C for 18 to 24 hour to reach 85-95% confluence. The cells were then incubated in growth medium (0.5 ml/well) containing ⁸⁶Rb⁺ (2 μCi/ml) for 4 hour at 37°C. The loading mixture was then aspirated, and the cells were washed four times with 1 ml buffer (15 mM HEPES, 140 mM NaCl, 2 mM KCl, 1 mM MgSO₄, 1.8 mM CaCl₂, 11 mM Glucose, pH 7.4). One ml of buffer with or without compounds to be tested was then added to each well. After incubation for 2 min, the assay buffer was collected for measurements of ⁸⁶Rb⁺ released from the cells. Cells were then lysed by adding 1 ml of 100 mM NaOH to each well, and the lysate was collected for determination of the amount of ⁸⁶Rb⁺ that was in the cells at the end of the efflux assay. Radioactivity of assay samples and lysates was measured by liquid scintillation counting. Total loading (cpm) was calculated as the sum of the assay sample and the lysate of each well. The amount of ⁸⁶Rb⁺ efflux was expressed as a percentage of ⁸⁶Rb⁺ loaded. Stimulated ⁸⁶Rb⁺ efflux was defined as the difference between efflux in the presence and absence of nicotine. For obtaining EC₅₀ and E_{max} values, stimulation curves were constructed in which 8 different concentrations of a ligand were included in the assay. For obtaining an IC₅₀₍₁₀₎ value, inhibition curves were constructed in which eight different concentrations of a compound were applied to cells for 10 min before 100 μM nicotine was applied to measure stimulated efflux. EC₅₀, E_{max} and IC₅₀₍₁₀₎ values were determined by nonlinear least-squares regression analyses (GraphPad, San Diego, CA).

General Procedures for Effect on Alcohol Intake in Rats

Adult male rats obtained from a colony of selectively bred alcohol-preferring rats (P rats) maintained at Indiana University School of Medicine. Rats were housed that were fitted with two 100 mL Richter tubes for the recording of water and alcohol intake. Animals were kept under a constant room temperature of 22 ± 1 °C and 12:12 light-dark cycle (7:00 a.m.–7:00 p.m. dark). Animals were fed 5001 Rodent Chow (Lab Diet, Brentwood, MO, USA). All procedures were approved by the IACUC at Duke University Medical Center.

After a week of handling and habituation, rats were given free access to water in a graduated Richter tube for 1 day. Next, they were given free access only to a solution of 10% (v/v) alcohol for 3 consecutive days. Thereafter, rats were given free access to water and a solution of 10% alcohol throughout the study. Water and alcohol intake were indexed by graduated Richter drinking tubes.

An acute study was conducted to determine a dose-response for compound (*S*)-9. After the establishment of a stable baseline for alcohol and water intake rats were injected subcutaneously with 0.33, 1 and 3 mg/kg of compound (*S*)-9 or the same volume of the vehicle (1 mg/kg). Alcohol and water intake were measured at, 2, 4, 6 and 24 hr after the drug administration. The preference for alcohol solution, [(alcohol volume) / (alcohol + water volume) × 100], was calculated for the same time points as alcohol and water intake⁴⁷. All animals (n=18) received all treatments following a crossover design with random assignment. The interval between injections was at least 3 days.

Solution of 10% (v/v) ethyl alcohol was prepared twice weekly from a solution of 200% ethanol mixed with tap water. Solutions of compound (*S*)-**9** was prepared weekly in 100 mM HCl solution and isotonic saline and was injected subcutaneously in a volume of 1 ml/kg.

The data were assessed by the analysis of variance with a between subjects factor of strain and a repeated measures factor of compound (*S*)-**9** dose. $P < 0.05$ (two-tailed) was used as the threshold for significance.

General Procedures for Emesis Studies in Ferrets

Studies were undertaken in male ferrets (*Mustela putorius furo*) weighing 0.9-1.5 kg (Marshall Farms, NY). They were housed in controlled conditions of room temperature (22°C) and light (12:12 h light-dark cycle) with free access to food and water. The technique that was employed to study emesis is one that was adapted from Osinski, et al.⁷⁶ Prior to each experiment, food was withheld overnight, whereas water was provided *ad libitum*. All tested compounds tested were constituted in 0.9% saline. Each ferret was tested with either varenicline (0.5 mg/kg) or compound (*S*)-**9** (1-30 mg/kg).

Emetic testing consisted of placing animals individually in a polycarbonate enclosure. Following a 30 min acclimation period, each animal randomly received a subcutaneous injection of varenicline or compound (*S*)-**9** in the subscapular region. A 60 min observation and video recording followed the drug administration. Following this period, animals that showed drug effects of either nausea and/or emesis were administered intraperitoneally a 1mg/kg combination of granisetron and dexamethasone and returned to their home enclosure.⁷⁷ All drugs were dissolved in sterile saline.

Video analysis of the incidence of emesis, stereotypical behaviors (such as licking, mouth-clawing, backward walking, gagging, and burying of the head in cage) and GI side effects (diarrhea) was done off-line. [Note: Stereotypical behaviors are thought to be reflective of a subjective sensation of nausea.]

Acknowledgments

We thank the Center of Drug Discovery, Georgetown University for support. We also acknowledge the Georgetown University Medical Center – Lombardi Comprehensive Cancer Center Proteomics & Metabolomics Shared Resource, which is partially supported by NIH/NCI grant P30-CA051008. Sazetidine-A was generously provided by National Institute on Drug Abuse through Research Triangle Institute (Research Triangle Park, NC). We thank Dr. Hans Rollema for providing varenicline through Global Compound Transfer Research, Pfizer (Groton, CT). We thank Dr. Lawrence Lumeng of the University of Indiana for providing P Rats (NIH- R24 AA015512).

Funding Sources This research was supported by NIH grants U19DA027990, R21DA032489 and R03DA025947.

ABBREVIATIONS

nAChR	nicotinic acetylcholine receptors
CNS	central nervous system
PNS	peripheral nervous system

REFERENCES

1. Mokdad AH, Marks JS, Stroup DF, Gerberding JL. Actual causes of death in the United States, 2000. JAMA. 2004; 291:1238–1245. [PubMed: 15010446]
2. Toneatto A, Sobell LC, Sobell MB, Kozlowski LT. Effect of cigarette smoking on alcohol treatment outcome. J. Subst. Abuse. 1995; 7:245–252. [PubMed: 7580233]

3. Moss HB, Chen CM, Yi HY. Subtypes of alcohol dependence in a nationally representative sample. *Drug Alcohol Depend.* 2007; 91:149–158. [PubMed: 17597309]
4. Rose JE, Brauer LH, Behm FM, Cramblett M, Calkins K, Lawhon D. Psychopharmacological interactions between nicotine and ethanol. *Nicotine Tob. Res.* 2004; 6:133–144. [PubMed: 14982697]
5. Mitchell SH, de Wit H, Zacny JP. Effects of varying ethanol dose on cigarette consumption in healthy normal volunteers. *Behav. Pharmacol.* 1995; 6:359–365. [PubMed: 11224344]
6. Le AD, Corrigan WA, Harding JW, Juzysch W, Li TK. Involvement of nicotinic receptors in alcohol self-administration. *Alcohol Clin. Exp. Res.* 2000; 24:155–163. [PubMed: 10698366]
7. Tizabi Y, Copeland RL Jr, Louis VA, Taylor RE. Effects of combined systemic alcohol and central nicotine administration into ventral tegmental area on dopamine release in the nucleus accumbens. *Alcohol Clin. Exp. Res.* 2002; 26:394–399. [PubMed: 11923594]
8. Lindstrom J. Nicotinic acetylcholine receptors in health and disease. *Mol. Neurobiol.* 1997; 15:193–222. [PubMed: 9396010]
9. Kellar, KJ.; Xiao, Y. Neuronal nicotinic receptors: one hundred years of progress. In: Sibley, DR.; Hanin, I.; Kuhar, M.; Skolnick, P., editors. *Handbook of Contemporary Neuropharmacology*. Vol. Vol. 1. John Wiley & Sons, Inc.; Hoboken: 2007. p. 107-145.
10. Changeux JP. Nicotinic receptors and nicotine addiction. *C. R. Biol.* 2009; 332:421–425. [PubMed: 19393973]
11. Flores CM, Rogers SW, Pabreza LA, Wolfe BB, Kellar KJ. A subtype of nicotinic cholinergic receptor in rat brain is composed of alpha 4 and beta 2 subunits and is up-regulated by chronic nicotine treatment. *Mol. pharmacol.* 1992; 41:31–37. [PubMed: 1732720]
12. Marks MJ, Burch JB, Collins AC. Effects of chronic nicotine infusion on tolerance development and nicotinic receptors. *J. Pharmacol. Exp. Ther.* 1983; 226:817–825. [PubMed: 6887012]
13. Schwartz RD, Kellar KJ. Nicotinic cholinergic receptor binding sites in the brain: regulation in vivo. *Science.* 1983; 220:214–216. [PubMed: 6828889]
14. Benwell ME, Balfour DJ, Anderson JM. Evidence that tobacco smoking increases the density of (–)-[3H]nicotine binding sites in human brain. *J. Neurochem.* 1988; 50:1243–1247. [PubMed: 3346676]
15. Breese CR, Marks MJ, Logel J, Adams CE, Sullivan B, Collins AC, Leonard S. Effect of smoking history on [3H]nicotine binding in human postmortem brain. *J. Pharmacol. Exp. Ther.* 1997; 282:7–13. [PubMed: 9223534]
16. Maskos U, Molles BE, Pons S, Besson M, Guiard BP, Guilloux JP, Evrard A, Cazala P, Cormier A, Mameli-Engvall M, Dufour N, Cloez-Tayarani I, Bemelmans AP, Mallet J, Gardier AM, David V, Faure P, Granon S, Changeux JP. Nicotine reinforcement and cognition restored by targeted expression of nicotinic receptors. *Nature.* 2005; 436:103–107. [PubMed: 16001069]
17. Picciotto MR, Zoli M, Rimondini R, Lena C, Marubio LM, Pich EM, Fuxe K, Changeux JP. Acetylcholine receptors containing the beta2 subunit are involved in the reinforcing properties of nicotine. *Nature.* 1998; 391:173–177. [PubMed: 9428762]
18. Tapper AR, McKinney SL, Nashmi R, Schwarz J, Deshpande P, Labarca C, Whiteaker P, Marks MJ, Collins AC, Lester HA. Nicotine activation of alpha4* receptors: sufficient for reward, tolerance, and sensitization. *Science.* 2004; 306:1029–1032. [PubMed: 15528443]
19. Blomqvist O, Ericson M, Johnson DH, Engel JA, Soderpalm B. Voluntary ethanol intake in the rat: effects of nicotinic acetylcholine receptor blockade or subchronic nicotine treatment. *Eur. J. Pharmacol.* 1996; 314:257–267. [PubMed: 8957244]
20. Blomqvist O, Ericson M, Engel JA, Soderpalm B. Accumbal dopamine overflow after ethanol: localization of the antagonizing effect of mecamylamine. *Eur. J. Pharmacol.* 1997; 334:149–156. [PubMed: 9369343]
21. Farook JM, Lewis B, Gaddis JG, Littleton JM, Barron S. Effects of mecamylamine on alcohol consumption and preference in male C57BL/6J mice. *Pharmacology.* 2009; 83:379–384. [PubMed: 19468256]
22. Farook JM, Lewis B, Gaddis JG, Littleton JM, Barron S. Lobeline, a nicotinic partial agonist attenuates alcohol consumption and preference in male C57BL/6J mice. *Physiol. Behav.* 2009; 97:503–506. [PubMed: 19268674]

23. Chatterjee S, Steensland P, Simms JA, Holgate J, Coe JW, Hurst RS, Shaffer CL, Lowe J, Rollema H, Bartlett SE. Partial agonists of the $\alpha 3\beta 4^*$ neuronal nicotinic acetylcholine receptor reduce ethanol consumption and seeking in rats. *Neuropsychopharmacology*. 2011; 36:603–615. [PubMed: 21048701]
24. Kamens HM, Andersen J, Picciotto MR. Modulation of ethanol consumption by genetic and pharmacological manipulation of nicotinic acetylcholine receptors in mice. *Psychopharmacology*. 2010; 208:613–626. [PubMed: 20072781]
25. Hendrickson LM, Zhao-Shea R, Pang X, Gardner PD, Tapper AR. Activation of $\alpha 4^*$ nAChRs is necessary and sufficient for varenicline-induced reduction of alcohol consumption. *J. Neurosci*. 2010; 30:10169–10176. [PubMed: 20668200]
26. Hendrickson LM, Gardner P, Tapper AR. Nicotinic acetylcholine receptors containing the $\alpha 4$ subunit are critical for the nicotine-induced reduction of acute voluntary ethanol consumption. *Channels*. 2011; 5:124–127. [PubMed: 21239887]
27. Coe JW, Brooks PR, Vetelino MG, Wirtz MC, Arnold EP, Huang J, Sands SB, Davis TI, Lebel LA, Fox CB, Shrikhande A, Heym JH, Schaeffer E, Rollema H, Lu Y, Mansbach RS, Chambers LK, Rovetti CC, Schulz DW, Tingley FD 3rd, O'Neill BT. Varenicline: an $\alpha 4\beta 2$ nicotinic receptor partial agonist for smoking cessation. *J. Med. Chem*. 2005; 48:3474–3477. [PubMed: 15887955]
28. Gonzales D, Rennard SI, Nides M, Oncken C, Azoulay S, Billing CB, Watsky EJ, Gong J, Williams KE, Reeves KR. Varenicline, an $\alpha 4\beta 2$ nicotinic acetylcholine receptor partial agonist, vs sustained-release bupropion and placebo for smoking cessation: a randomized controlled trial. *JAMA*. 2006; 296:47–55. [PubMed: 16820546]
29. Jorenby DE, Hays JT, Rigotti NA, Azoulay S, Watsky EJ, Williams KE, Billing CB, Gong J, Reeves KR. Efficacy of varenicline, an $\alpha 4\beta 2$ nicotinic acetylcholine receptor partial agonist, vs placebo or sustained-release bupropion for smoking cessation: a randomized controlled trial. *JAMA*. 2006; 296:56–63. [PubMed: 16820547]
30. Rollema H, Chambers LK, Coe JW, Glowa J, Hurst RS, Lebel LA, Lu Y, Mansbach RS, Mather RJ, Rovetti CC, Sands SB, Schaeffer E, Schulz DW, Tingley FD 3rd, Williams KE. Pharmacological profile of the $\alpha (4)\beta (2)$ nicotinic acetylcholine receptor partial agonist varenicline, an effective smoking cessation aid. *Neuropharmacology*. 2007; 52:985–994. [PubMed: 17157884]
31. Ericson M, Lof E, Stomberg R, Soderpalm B. The smoking cessation medication varenicline attenuates alcohol and nicotine interactions in the rat mesolimbic dopamine system. *J. Pharmacol. Exp. Ther*. 2009; 329:225–230. [PubMed: 19126781]
32. McKee SA, Harrison EL, O'Malley SS, Krishnan-Sarin S, Shi J, Tetraault JM, Picciotto MR, Petrakis IL, Estevez N, Balchunas E. Varenicline reduces alcohol self-administration in heavy-drinking smokers. *Biol. Psychiatry*. 2009; 66:185–190. [PubMed: 19249750]
33. Steensland P, Simms JA, Holgate J, Richards JK, Bartlett SE. Varenicline, an $\alpha 4\beta 2$ nicotinic acetylcholine receptor partial agonist, selectively decreases ethanol consumption and seeking. *Proc. Natl. Acad. Sci. U.S.A.* 2007; 104:12518–12523. [PubMed: 17626178]
34. Moore TJ, Furberg CD, Glenmullen J, Maltzberger JT, Singh S. Suicidal behavior and depression in smoking cessation treatments. *PLoS One*. 2011; 6:e27016. [PubMed: 22073240]
35. Xiao Y, Fan H, Musachio JL, Wei ZL, Chellappan SK, Kozikowski AP, Kellar KJ. Sazetidine-A, a novel ligand that desensitizes $\alpha 4\beta 2$ nicotinic acetylcholine receptors without activating them. *Mol. Pharmacol*. 2006; 70:1454–1460. [PubMed: 16857741]
36. Langley JN, Dickenson W. On the local paralysis of peripheral ganglia and on the connexion of different classes of nerve fibres with them. *Proc. R. Soc. Lond*. 1889; 46:423–431.
37. Katz B, Thesleff S. A study of the desensitization produced by acetylcholine at the motor end-plate. *J. Physiol*. 1957; 138:63–80. [PubMed: 13463799]
38. Hulihan-Giblin BA, Lumpkin MD, Kellar KJ. Acute effects of nicotine on prolactin release in the rat: agonist and antagonist effects of a single injection of nicotine. *J. Pharmacol. Exp. Ther*. 1990; 252:15–20. [PubMed: 2299587]
39. Grady SR, Marks MJ, Collins AC. Desensitization of nicotine-stimulated $[3H]$ dopamine release from mouse striatal synaptosomes. *J. Neurochem*. 1994; 62:1390–1398. [PubMed: 8133269]

40. Marks MJ, Grady SR, Yang JM, Lippiello PM, Collins AC. Desensitization of nicotine-stimulated 86Rb⁺ efflux from mouse brain synaptosomes. *J. Neurochem.* 1994; 63:2125–2135. [PubMed: 7964732]
41. Benwell ME, Balfour DJ, Birrell CE. Desensitization of the nicotine-induced mesolimbic dopamine responses during constant infusion with nicotine. *Br. J. Pharmacol.* 1995; 114:454–460. [PubMed: 7881744]
42. Pidoplichko VI, DeBiasi M, Williams JT, Dani JA. Nicotine activates and desensitizes midbrain dopamine neurons. *Nature.* 1997; 390:401–404. [PubMed: 9389479]
43. Quick MW, Lester RA. Desensitization of neuronal nicotinic receptors. *J. Neurobiol.* 2002; 53:457–478. [PubMed: 12436413]
44. Picciotto MR, Addy NA, Mineur YS, Brunzell DH. It is not “either/or”: activation and desensitization of nicotinic acetylcholine receptors both contribute to behaviors related to nicotine addiction and mood. *Prog. Neurobiol.* 2008; 84:329–342. [PubMed: 18242816]
45. Buccafusco JJ, Beach JW, Terry AV Jr. Desensitization of nicotinic acetylcholine receptors as a strategy for drug development. *J. Pharmacol. Exp. Ther.* 2009; 328:364–370. [PubMed: 19023041]
46. Levin ED, Rezvani AH, Xiao Y, Slade S, Cauley M, Wells C, Hampton D, Petro A, Rose JE, Brown ML, Paige MA, McDowell BE, Kellar KJ. Sazetidine-A, a selective $\alpha 4\beta 2$ nicotinic receptor desensitizing agent and partial agonist, reduces nicotine self administration in rats. *J. Pharmacol. Exp. Ther.* 2010; 332:933–939. [PubMed: 20007754]
47. Rezvani AH, Slade S, Wells C, Petro A, Lumeng L, Li TK, Xiao Y, Brown ML, Paige MA, McDowell BE, Rose JE, Kellar KJ, Levin ED. Effects of sazetidine-A, a selective $\alpha 4\beta 2$ nicotinic acetylcholine receptor desensitizing agent on alcohol and nicotine self administration in selectively bred alcohol-preferring (P) rats. *Psychopharmacology.* 2010; 211:161–174. [PubMed: 20535453]
48. Rezvani AH, Cauley M, Sexton H, Xiao Y, Brown ML, Paige MA, McDowell BE, Kellar KJ, Levin ED. Sazetidine-A, a selective $\alpha 4\beta 2$ nicotinic acetylcholine receptor ligand: effects on dizocilpine and scopolamine-induced attentional impairments in female Sprague-Dawley rats. *Psychopharmacology.* 2011; 215:621–630. [PubMed: 21274704]
49. Rezvani AH, Cauley M, Xiao Y, Kellar KJ, Levin ED. Effects of chronic sazetidine-A, a selective $\alpha 4\beta 2$ neuronal nicotinic acetylcholine receptors desensitizing agent on pharmacologically-induced impaired attention in rats. *Psychopharmacology.* 2012; 222:269–276. [PubMed: 22297831]
50. Kozikowski AP, Eaton JB, Bajjuri KM, Chellappan SK, Chen Y, Karadi S, He R, Caldarone B, Manzano M, Yuen PW, Lukas RJ. Chemistry and pharmacology of nicotinic ligands based on 6-[5-(azetidin-2-ylmethoxy)pyridin-3-yl]hex-5-yn-1-ol (AMOP-H-OH) for possible use in depression. *ChemMedChem.* 2009; 4:1279–1291. [PubMed: 19569163]
51. Turner JR, Castellano LM, Blendy JA. Nicotinic partial agonists varenicline and sazetidine-A have differential effects on affective behavior. *J. Pharmacol. Exp. Ther.* 2010; 334:665–672. [PubMed: 20435920]
52. Caldarone BJ, Wang D, Paterson NE, Manzano M, Fedolak A, Cavino K, Kwan M, Hanania T, Chellappan SK, Kozikowski AP, Olivier B, Picciotto MR, Ghavami A. Dissociation between duration of action in the forced swim test in mice and nicotinic acetylcholine receptor occupancy with sazetidine, varenicline, and 5-I-A85380. *Psychopharmacology.* 2011; 217:199–210. [PubMed: 21487659]
53. Ji Y, Brueckl T, Baxter RD, Fujiwara Y, Seiple IB, Su S, Blackmond DG, Baran PS. Innate C-H trifluoromethylation of heterocycles. *Proc. Natl. Acad. Sci. U.S.A.* 2011; 108:14411–14415. [PubMed: 21844378]
54. Nagib DA, MacMillan DW. Trifluoromethylation of arenes and heteroarenes by means of photoredox catalysis. *Nature.* 2011; 480:224–228. [PubMed: 22158245]
55. Ross TL, Honer M, Lam PY, Mindt TL, Groehn V, Schibli R, Schubiger PA, Ametamey SM. Fluorine-18F click radiosynthesis and preclinical evaluation of a new 18F-labeled folic acid derivative. *Bioconjug. Chem.* 2008; 19:2462–2470. [PubMed: 19053298]
56. Yadav JS, Deshpande PK, Sharma GVM. An Effective, Practical Method for the Synthesis of Chiral Propargyl Alcohols. *Tetrahedron.* 1990; 46:7033–7046.

57. Morgentin R, Jung F, Lamorlette M, Maudet M, Menard M, Ple P, Pasduet G, Renaud F. An efficient large-scale synthesis of alkyl 5-hydroxy pyridin- and pyrimidin-2-yl acetate. *Tetrahedron*. 2009; 65:757–764.
58. Mihalak KB, Carroll FI, Luetje CW. Varenicline is a partial agonist at $\alpha 4\beta 2$ and a full agonist at $\alpha 7$ neuronal nicotinic receptors. *Mol. Pharmacol.* 2006; 70:801–805. [PubMed: 16766716]
59. Xiao, Y.; Yasuda, RP.; Sahibzada, N.; Horton, L.; DiPietro, JR.; Iwueze, AF.; Paige, MA.; McDowell, BE.; Brown, ML.; Wolfe, BB.; Kellar, KJ. Pharmacological properties of sazetidine-A, a selective ligand of $\alpha 4\beta 2$ nicotinic acetylcholine receptors. Society for Neuroscience 38th Annual Meeting; Washington, DC. SFN; 2008. p. 31.12
60. Zwart R, Carbone AL, Moroni M, Bermudez I, Mogg AJ, Folly EA, Broad LM, Williams AC, Zhang D, Ding C, Heinz BA, Sher E. Sazetidine-A is a potent and selective agonist at native and recombinant $\alpha 4\beta 2$ nicotinic acetylcholine receptors. *Mol. Pharmacol.* 2008; 73:1838–1843. [PubMed: 18367540]
61. Carbone AL, Moroni M, Groot-Kormelink PJ, Bermudez I. Pentameric concatenated ($\alpha 4$)($\beta 2$)($\beta 3$) and ($\alpha 4$)($\beta 3$)($\beta 2$)($\beta 2$) nicotinic acetylcholine receptors: subunit arrangement determines functional expression. *Br. J. Pharmacol.* 2009; 156:970–981. [PubMed: 19366353]
62. Xiao, Y.; Yasuda, RP.; Sahibzada, N.; Wolfe, BB.; Horton, L.; Paige, M.; Brown, ML.; Kellar, KJ. Sazetidine-A selectively induces a long lasting desensitization of $\alpha 4\beta 2$ nicotinic acetylcholine receptors. 2010 Neuroscience Meeting, Society for Neuroscience; San Diego. 2010. p. 38.10
63. Alsharari SD, Carroll FI, McIntosh JM, Damaj MI. The antinociceptive effects of nicotinic partial agonists varenicline and sazetidine-a in murine acute and tonic pain models. *J. Pharmacol. Exp. Ther.* 2012; 342:742–749. [PubMed: 22678099]
64. Cucchiario G, Xiao Y, Gonzalez-Sulser A, Kellar KJ. Analgesic effects of Sazetidine-A, a new nicotinic cholinergic drug. *Anesthesiology*. 2008; 109:512–519. [PubMed: 18719450]
65. Hussmann GP, Turner JR, Lomazzo E, Venkatesh R, Cousins V, Xiao Y, Yasuda RP, Wolfe BB, Perry DC, Rezvani AH, Levin ED, Blendy JA, Kellar KJ. Chronic sazetidine-a at behaviorally active doses does not increase nicotinic cholinergic receptors in rodent brain. *J. Pharmacol. Exp. Ther.* 2012; 343:441–450. [PubMed: 22899752]
66. Johnson JE, Slade S, Wells C, Petro A, Sexton H, Rezvani AH, Brown ML, Paige MA, McDowell BE, Xiao Y, Kellar KJ, Levin ED. Assessing the effects of chronic sazetidine-A delivery on nicotine self-administration in both male and female rats. *Psychopharmacology*. 2012; 222:269–276. [PubMed: 22297831]
67. Levin ED, Hampton D, Xiao Y, Kellar KJ. Sazetidine-A, a nicotinic desensitizing agent, decrease nicotine self-administration in rats. *Biochem. Pharmacol.* 2007; 74:SMA-43.
68. Paterson NE, Min W, Hackett A, Lowe D, Hanania T, Caldarone B, Ghavami A. The high-affinity nAChR partial agonists varenicline and sazetidine-A exhibit reinforcing properties in rats. *Prog. Neuropsychopharmacol. Biol. Psychiatry*. 2010; 34:1455–1464. [PubMed: 20708056]
69. Zhang J, Xiao YD, Jordan KG, Hammond PS, Van Dyke KM, Mazurov AA, Speake JD, Lippiello PM, James JW, Letchworth SR, Bencherif M, Hauser TA. Analgesic effects mediated by neuronal nicotinic acetylcholine receptor agonists: Correlation with desensitization of $\alpha 4\beta 2$ (*) receptors. *Eur. J. Pharm. Sci.* 2012; 47:813–823. [PubMed: 23036283]
70. Nakamura M, Oshima A, Fujimoto Y, Maruyama N, Ishibashi T, Reeves KR. Efficacy and tolerability of varenicline, an $\alpha 4\beta 2$ nicotinic acetylcholine receptor partial agonist, in a 12-week, randomized, placebo-controlled, dose-response study with 40-week follow-up for smoking cessation in Japanese smokers. *Clin. Ther.* 2007; 29:1040–1056. [PubMed: 17692720]
71. Wang C, Xiao D, Chan KP, Pothirat C, Garza D, Davies S. Varenicline for smoking cessation: a placebo-controlled, randomized study. *Respirology*. 2009; 14:384–392. [PubMed: 19192221]
72. Tonstad S. Smoking cessation efficacy and safety of varenicline, an $\alpha 4\beta 2$ nicotinic receptor partial agonist. *J. Cardiovasc. Nurs.* 2006; 21:433–436. [PubMed: 17293731]
73. Cahill K, Stead L, Lancaster T. A preliminary benefit-risk assessment of varenicline in smoking cessation. *Drug Saf.* 2009; 32:119–135. [PubMed: 19236119]

74. Xiao Y, Kellar KJ. The comparative pharmacology and up regulation of rat neuronal nicotinic receptor subtype binding sites stably expressed in transfected mammalian cells. *J. Pharmacol. Exp. Ther.* 2004; 310:98–107. [PubMed: 15016836]
75. Xiao Y, Meyer EL, Thompson JM, Surin A, Wroblewski J, Kellar KJ. Rat alpha3/beta4 subtype of neuronal nicotinic acetylcholine receptor stably expressed in a transfected cell line: pharmacology of ligand binding and function. *Mol. Pharmacol.* 1998; 54:322–333. [PubMed: 9687574]
76. Osinski MA, Uchic ME, Seifert T, Shaughnessy TK, Miller LN, Nakane M, Cox BF, Brioni JD, Moreland RB. Dopamine D2, but not D4, receptor agonists are emetogenic in ferrets. *Pharmacol. Biochem. Behav.* 2005; 81:211–219. [PubMed: 15894081]
77. Fukunaka N, Sagae S, Kudo R, Endo T, Hirafuji M, Minami M. Effects of granisetron and its combination with dexamethasone on cisplatin-induced delayed emesis in the ferret. *Gen. Pharmacol.* 1998; 31:775–781. [PubMed: 9809477]

Figure 1.
Development of nicotinic desensitizers with reduced agonist activity.

Figure 3. Effect of acute administration of compound (S)-9 on alcohol intake in P-rats. Alcohol intakes (g/kg) were measured at 2, 4, 6 and 24 h after an injection of (S)-9. All values are expressed as mean $\pm$ SEM (n=18).

Scheme 1. Synthesis of Analogs 9-12

Reagents and condition: (a) DEAD, PPh_3 , THF, 0 °C-rt, 48 h, 68-77%; (b) 1-hexyne, $\text{Pd}(\text{PPh}_3)_2\text{Cl}_2$ (cat.), CuI (cat.), PPh_3 (cat.), Et_3N , DMSO, sealed tube, 95 °C, 60 h, 89-98%; (c) Method A: TFA, CH_2Cl_2 , 0 °C-rt, 3 h then 10% aqueous NaOH solution-methanol, 75-83%; Method B: HCl in methanol, 0 °C-rt, 3 h, 87-92%.

Scheme 2. Synthesis of Analogs 16-17

Reagents and condition: (a) p-TsCl, DMAP, Et₃N, DCM, 0 °C-rt, overnight, 80%; (b) Ph₃P, CCl₄, cat. NaHCO₃, sealed tube, 80 °C, 48 h, 69%; (c) Kryptofix 2.2.2, KF, THF, reflux, overnight, 85%; (d) HCl in methanol, 0 °C-rt, 3 h, 90%.

Scheme 3. Synthesis of Analog 22

Reagents and condition: (a) HBr, AcOH, 110 °C, sealed tube, overnight, 79%; (b) DEAD, PPh₃, THF, 0 °C-rt, 24 h, 88%; (c) 1-hexyne, Pd(PPh₃)₂Cl₂ (cat.), CuI (cat.), PPh₃ (cat.), Et₃N, DMSO, 50 °C, overnight, 91%; (d) HCl in methanol, 0 °C-rt, 3 h, 73%.

Table 1

Comparison of binding affinities of nicotine, varenicline and sazetidine-A for rat nAChR subtypes to those of the 3-alkoxy-2,5-substituted-pyridyl compounds.

Compound	K _i (nM) ^a										K _i Ratio					
	2	2	2	4	3	2	3	4	4	2	Fore-brain	7	4	2	7/ 4	2
Nicotine	12			110	47		440	10	40		517	12	44		520
Varenicline	0.48			94	2.5		390	0.12	28		37	1.1	3,300		310
Sazetidine-A	0.087			210	0.38		1,900	0.062	52		670	0.17	31,000		11,000
(R)-9	970			250,000	3,600		200,000	740	160,000		160,000	4,300	270		220
(S)-9	16			31,000	390		230,000	12	3,500		93,000	52	19,000		7,800
(R)-10	22,000			370,000	19,000		220,000	9,900	270,000		110,000	110,000	22		11
(S)-10	130			59,000	2,500		94,000	73	23,000		130,000	540	1,300		1,800	

Compound	K _i (nM) ^a										K _i Ratio				
	2	2	2	4	4	3	3	2	4	4	7	Fore-brain			
	2	2	2	4	4	3	3	2	4	4	7	3	4	4	2
(S)-11	13	9,600	340	150,000	9.6	750	150,000	31	16,000	16,000	16,000	16,000	16,000	16,000	16,000
(S)-12	5.3	26,000	220	300,000	4.3	3,300	170,000	11	70,000	40,000	40,000	40,000	40,000	40,000	40,000
(S)-16	8.7	24,000	230	150,000	5.1	3,000	150,000	15	29,000	29,000	29,000	29,000	29,000	29,000	29,000
(S)-17	8.6	29,000	300	92,000	5.3	3,700	67,000	32	17,000	13,000	13,000	13,000	13,000	13,000	13,000
(S)-22	8.6	29,000	300	92,000	5.3	3,700	67,000	32	17,000	13,000	13,000	13,000	13,000	13,000	13,000

^aK_i values of the compounds shown are the mean of 3 to 5 independent measurements (for clarity, SEM values were omitted, which were within 20% of the mean in most cases).

Table 2

Comparison of activation and desensitization of nAChR function by ligands.

Compound	4 2 nAChRs ^a			3 4 nAChRs ^a		
	EC ₅₀ ^b (nM)	E _{max} ^c (%)	IC ₅₀₍₁₀₎ ^d (nM)	EC ₅₀ (nM)	E _{max} (%)	IC ₅₀₍₁₀₎ (nM)
(-)-Nicotine	2,400	100	370	23,000	100	>10,000
varenicline	950	45	94	21,000	82	>10,000
sazetidine-A	24	40	11	30,000	12	>10,000
(S)-9	310	15	260	NA	NA	>10,000
(S)-10	NA	NA	1,600	NA	NA	6,400
(S)-11	160	17	110	NA	NA	>10,000
(S)-12	450	20	51	NA	NA	>10,000
(S)-16	NA	NA	130	NA	NA	>10,000
(S)-17	NA	NA	260	NA	NA	>10,000

^a Functional properties of each compounds were determined using stable cell lines expressing human 4 2 or rat 3 4 nAChRs.

^b EC₅₀ values show potencies of agonist activities.

^c E_{max} (%) values show relative efficacies of agonist activities, which were normalized to the E_{max} value of nicotine. ^d IC₅₀₍₁₀₎ values show potencies of desensitizer activities.

^d NA indicates no significant stimulated efflux was detected. All values shown are means of 3 to 9 independent experiments (for clarity, SEM values were omitted, which were within 20% of the mean in most cases).