Published in final edited form as:

Mov Disord. 2013 October ; 28(12): . doi:10.1002/mds.25588.

Sensorimotor adaptation of speech in Parkinson's disease

Fatemeh Mollaei^{1,2}, Douglas M. Shiller^{1,3,4}, and Vincent L. Gracco^{1,2,5}

¹Centre for Research on Brain, Language and Music, Montréal, Canada

²School of Communication Sciences and Disorders, McGill University, Montreal, Canada

³Centre de recherche CHU Sainte Justine, Montreal, Canada

⁴École d'orthophonie et d'audiologie, Université de Montréal, Canada

⁵Haskins Laboratories, New Haven, CT, USA

Abstract

Background—The basal ganglia (BG) are involved in establishing motor plans for a wide range of behaviors. Parkinson's disease (PD) is a manifestation of basal ganglia dysfunction associated with a deficit in sensorimotor integration and difficulty in acquiring new motor sequences, thereby affecting motor learning. Previous studies of sensorimotor integration and sensorimotor adaptation (SA) in PD have focused on limb movements using visual and force-field alterations.^{1–4}

Methods—Here we report the results from a SA experiment investigating the ability of PD patients to make speech motor adjustments to a constant and predictable auditory feedback manipulation. Participants produced speech while their auditory feedback was altered and maintained in a manner consistent with a change in tongue position. The degree of adaptation was associated with the severity of motor symptoms.

Results—The PD patients exhibited adaptation to the induced sensory error, however the degree of adaptation was reduced compared to healthy age-matched control participants.

Conclusions—The reduced capacity to adapt to a change in auditory feedback is consistent with reduced gain in the sensorimotor system for speech and with previous studies demonstrating limitations in the adaptation of limb movements following changes in visual feedback in PD patients.

Keywords

Sensorimotor adaptation; speech production; Parkinson's disease; speech motor learning; auditory feedback

INTRODUCTION

Although the cardinal symptoms associated with Parkinson's disease (PD) are predominantly associated with motor system impairment, sensory deficits are well

Corresponding author: Vincent L. Gracco, PhD, Centre for Research on Brain, Language & Music, McGill University, 3640 de la Montagne, Montreal, Quebec, H3G2A8, Canada, vincent.gracco@mcgill.ca.

Conflicts of interest/financial disclosures: Nothing to report.

Author Roles: F. Mollaei—Wrote first draft, executed of study including design; D. Shiller—major contribution to the design and statistical analysis, major contribution to the writing of the manuscript; V. Gracco—study conception, review and critique of statistics, major contribution to the writing of the manuscript.

Financial Disclosures of all authors: None.

known.⁵⁻⁸ Two of the major speech characteristics of PD are reduced vocal loudness and reduced respiratory control⁹ which appear to have a sensory contribution. For example, when asked to increase speech loudness PD patients often complain of a perception of being too loud.¹⁰ The misperception of self-generated loudness may be due to the apparent reduced sensitivity of laryngeal mucosal mechanoreceptors¹¹ generating a false sense of increased effort while speaking. In the auditory domain, when presented with a sudden change in auditory feedback of the pitch of their voice, PD subjects respond with a larger adjustment than non-affected control subjects.¹² Besides problems associated with speech loudness or pitch control. These speech symptoms are consistent with a more general problem of sensorimotor integration¹³⁻¹⁵.

One consequence of the sensory and motor impairments in PD patients may be reflected in the difficulty they exhibit in learning new motor skills and modifying existing motor behaviors.^{1,16,17} In speech, this deficit is observed in the extensive behavioral training needed to ameliorate their speech disorder.^{10,18,19} Until recently, behavioral treatment for PD resulted in minimal and short-term gains. In contrast, intensive, high effort training and a focus on improving sensorimotor processing has provided some of the most robust treatment effects with this population.²⁰ It appears that the speech symptoms in PD patients are not localized to the inability to adjust speech parameters such as loudness or articulation, but reflect a more general problem with sensory recalibration which may, by extension, reflect impair sensorimotor learning.

We addressed the question of sensorimotor integration and learning for speech through an experimental manipulation of auditory feedback in real-time during speech production. This manipulation introduces a predictable and constant feedback error associated with discrete acoustic properties of the speech signal which have predictable acoustic/motor mappings (e.g., vowel formant frequencies or consonant fricative energy). Over the course of a single experimental session these manipulations result in speech motor changes that adapt to the induced error.²¹⁻²⁷ For example, altering the first formant (F1) frequency of a vowel (an acoustic cue related to vertical tongue position) yields a change in speech motor output consistent with a compensatory change in tongue height.^{23,24,28,29} As a result, by evaluating the ability of subjects to engage a process of sensorimotor adaptation (SA) it is possible to evaluate whether and to what extent the direct sensorimotor (acoustic-articulatory) mapping is intact in PD subjects and to evaluate the ability to learn new sensorimotor mappings independent of vocal (pitch and loudness) or respiratory control.

We used a SA paradigm to assess the capacity of PD patients to alter their speech motor control in response to a perceived auditory-sensory error. We hypothesized that deficits in sensorimotor integration would limit the capacity for speech motor learning in PD patients --- a result that, if demonstrated, would have important implications for understanding the speech disorder of PD subjects as well as their ability to change their speech through behavioral intervention.

METHODS

The study was approved by the IRB at the Faculty of Medicine, McGill University. All participants gave written informed consent.

Participants

Participants included nine individuals with idiopathic PD (7 male, 2 female, mean age: 63.8 yrs) and nine healthy age-matched controls (4 male, 5 female, mean age: 60.3 yrs). The severity of PD motor symptoms, evaluated using the UPDRS, ranged from mild (15) to moderate (88), ($M = 33.2$, $SD = 13.5$). Cognitive functioning evaluated with the MMSE was

in the normal range for all PD patients (scores > 24). All patients were taking levodopa and other medication including dopaminergic and/or anticholinergics. Subjects were tested during their off state (6 hours off medication); one PD patient was tested with only two hours off medication due to disease severity.

Only two participants had a history of speech therapy; one could not remember the specifics of the treatment and one recalled treatment consisting of exercises focusing on increasing intelligibility and loudness.

For each PD participant, the Rainbow Passage was recorded for perceptual analysis. A licensed Speech-Language Pathologist rated participants' speech on 43 perceptual dimensions evaluating all the speech subsystems.³⁰ Overall, the severity of the PD participants was rated as moderate (1), mild (6) and within normal limits (2). We used the perceptual and UPDRS scores to assess any relationship between severity of speech and motor symptoms with the magnitude of adaptation.

Control subjects had no history of neurological impairment. All participants had binaural pure tone hearing thresholds of 40 dB HL or less at 250, 500, 1000, 2000 and 4000 Hz. None of the participants wore hearing-aids.

Audio recording

Subjects sat in front of a computer monitor and spoke into a microphone located 20cm from their mouth. The microphone signal was digitized at 44.1 kHz (16 bit). The microphone signal was presented back to the subjects (see below for details) through headphones (Beyerdynamic 880 Pro).

Speech was cued by presentation of the target word on the computer display. Each stimulus was presented for 2 seconds followed by a 1 second blank display. Subjects produced the target word at a comfortable rate and volume immediately following the visual presentation. Speech volume was maintained by providing the subject with a VU meter on the computer display.

Manipulation of auditory feedback

Subjects produced individual words under conditions of altered auditory feedback (see ^{24,28} for details). Briefly, the first formant frequency of the vowel / / in the target word "head" (/h d/) was manipulated with a digital signal processor (DSP) specialized for speech acoustic signals (VoiceOne, TC Helicon) in near-real time (11 ms delay). During maximal acoustic perturbation, the first formant frequency was increased by 30%, resulting in an acoustic signal that was closer to the vowel /æ/ (as in the word "had"). The altered speech signal was amplified and presented at a volume of approximately 70 dB. In order to reduce the subject's perception of their unmodified air- and bone-conducted feedback signal, the altered speech signal was mixed with pink noise (presented at approximately 60 dB).

Testing sequence

Subjects produced 200 repetitions of the word /h d/ ("head") in the following sequence: 30 trials produced under unaltered feedback condition (*baseline* phase), followed by the sudden introduction of the acoustic perturbation, and 100 trials under conditions of maximal acoustic perturbation (*hold* phase). Finally, auditory feedback was gradually returned to normal over 40 trials (*ramp-offset* phase) and then maintained for 30 trials to facilitate de-adaptation (*after-effect* phase). See Figure 1 for schematic.

Data Analysis

For each participant, the values of the first two formant frequencies (F1 and F2) were computed over a 30 ms window centered at the mid-point of the vowel / /. In addition we calculated the *difference* between F2 and F1 (F2-F1). Several factors motivated our use of this parameter. In typical productions of the vowels / / and /æ/, the contrast is associated with a change not only in F1, but in the difference between F2 and F1 --- /æ/ being associated with a smaller F2-F1 separation than / / (see, e.g., ^{31,32}). We were further motivated to include F2-F1 as a dependent measure because, in prior studies of speech adaptation (e.g., ²⁴), compensatory changes in speech output have been observed in both F2 and F1 in response to a manipulation of auditory feedback involving an increase in F1 alone.

In order to evaluate the changes in speech production across the 200 tokens in the speech adaptation task, individual F1, F2 and F2-F1 values were normalized and then averaged over successive blocks of 10 trials. The raw formant values (in Hz) were converted to reflect the proportional change in frequency relative to the baseline phase under normal (unaltered) feedback conditions. This eliminated the influence of baseline differences prior to averaging across subjects, in addition to representing the change in speech output in the same units as the feedback manipulation itself (a proportional change in F1).

For each dependent variable, a mixed-factorial 2-way ANOVA was carried out to examine differences between the PD vs. Control groups (GROUP, the between-subjects effect) at four time-points of interest during the adaptation protocol (PHASE, the within-subjects effect): 1) the beginning of the *hold* phase, characterizing the degree of adaptation immediately following the sudden onset of the feedback perturbation, 2) the end of the *hold* phase, characterizing the degree of adaptation at the end of 100 trials of practice under conditions of maximum feedback alteration, 3) the beginning of the *after-effect* phase reflecting vowel production after the gradual restoration of unaltered feedback and 4) at the end of the *after-effect* phase (at the end of 30 trials under normal auditory feedback conditions).

RESULTS

Baseline productions of the vowel / /, taken over trials 16–30 (under conditions of unaltered auditory feedback), were comparable between the PD and control groups. Mean and variability measures (SD) values of F1 and F2 frequency (Hz) are shown in Figure 2. No statistically reliable differences were observed between the groups for any of the measures (F1 mean: $t[16]=-0.12$, $p > 0.05$; F2 mean: $t[16]=0.23$, $p > 0.05$; F1 variability: $t[16]=-0.13$, $p > 0.05$; F2 variability: $t[16]=-0.32$, $p > 0.05$). Average RMS values were obtained from each subject's vowel recordings during the last 15 trials of the baseline and hold phases to evaluate changes in speech loudness over the course of the experiment. No statistically reliable differences were found for either group (PD means[SD]: baseline = 0.1819 [0.0175]; hold = 0.1844 [0.0143]; $t[8] = 2.1435$, $p = 0.07$) (Control means[SD]: baseline = 0.1660 [0.0157]; hold = 0.1824 [0.0136]; $t[8] = 0.1519$, $p = 0.88$).

Changes in F1 and F2 values throughout the speech adaptation procedure are shown in Figure 3. For the controls (top panel), compensation to the auditory feedback manipulation (decrease in F1 and increase in F2 relative to baseline) can be observed, with the changes beginning soon after the sudden introduction of the feedback manipulation (beginning in block 6) and maintained throughout the *hold* phase. The gradual restoration of normal auditory feedback during the *ramp-offset* phase, yielded a nearly complete de-adaptation in F1 and F2. By the final block of *after-effect* trials, vowel productions resembled those observed during the *null* phase. The PD group (Figure 3, bottom panel) also exhibited compensatory change in vowel formants beginning immediately after the sudden

introduction of the auditory feedback manipulation and continuing throughout the *hold* phase. However, the maximum change in F1 and F2 for the PD group at the end of the *hold* phase was notably reduced to that of the control group. The proportional change in F1 and F2 for the control group averaged -0.90 (-9.0%) and 0.044 (4.4%) respectively, while the proportional change in F1 and F2 for the PD group averaged -0.058 (-5.8%) and 0.014 (1.4%). Both groups exhibit a de-adaptation effect in F1 and F2 during the *ramp-offset* phase.

The effects of GROUP and PHASE on F1, F2 and F2-F1 were evaluated using a 2-way ANOVA. Mean formant values at four epochs are shown in Figure 4. For F1, a reliable main effect of PHASE ($F [2.17, 34.81] = 14.19, p < 0.01$) was found with no reliable effect of GROUP ($F [1,16] = 0.659, p > 0.05$) nor 2-way interaction ($F [2.17, 34.81] = 1.20, p > 0.05$). For F2 a significant main effect of PHASE ($F [3, 48] = 7.96, p < 0.01$) was found, however in this case the main effect of GROUP was also significant ($F [1,16] = 6.96, p < 0.01$). The 2-way interaction in F2 was not significant ($F [3, 48] = 0.485, p > 0.05$).

For the combined F2-F1 measure a significant main effect of PHASE ($F [2.25, 36.12] = 16.90, p < 0.01$) and a significant main effect of GROUP ($F [1, 16] = 9.12, p < 0.01$) was noted with no reliable interaction ($F [2.25, 36.12] = 0.932, p > 0.05$). Hence, while both groups exhibited a change in speech output in response to the different phases of the auditory feedback manipulation (owing to the reliable effect of PHASE), the overall magnitude of the compensatory change in vowel acoustics was found to be smaller in the PD group compared to controls.

An additional analysis was carried out to confirm whether the overall adaptation effect (F2-F1) at the end of the *Hold* phase reliably changed from baseline for both groups. A one-sample *t*-test comparing the results from each group against a hypothesized mean of zero confirmed that for each group, the adaptation effect was reliably different from zero (Controls: $t[8] = 5.34, p < 0.01$; PD Group: $t[8] = 4.39, p < 0.01$).

A final analysis focused on the relationship between the degree of adaptation in F1 or F2-F1, and the degree of severity of either the perceptual or UPDRS clinical rating scores. No significant correlations were found for any of the perceptual speech scores (by subsystem or overall) but a significant negative correlation was found between the degree of F1 adaptation and the severity of motor symptoms as indexed by the UPDRS scores ($r = -0.74, p = 0.011$).

DISCUSSION

During speech production, sensory feedback is used for both real-time control and for sensorimotor learning. Here we examined the manner in which PD, secondary to basal ganglia dysfunction, affected the process of sensorimotor integration and adaptation and by extension speech motor learning. The results of the present study illustrate that PD patients are capable of adapting their speech articulation in response to a change in auditory feedback related to vowel formants, however the result also showed a clear reduction in the magnitude of sensorimotor adaptation relative to control subjects.

The findings of the present study highlight a number of issues related to the speech production capacity of PD individuals, the role of sensory information in speech and the functions of the basal ganglia for speech.

Sensorimotor interactions for speech in PD

Overall, PD subjects were able to adjust their motor output in a consistent and predictable way to offset the specific auditory manipulation. The manipulation in F1 frequency was

accompanied in both groups by a reciprocal adjustment in F1 and F2 to partially offset the induced error. Thus, for both the control group and the PD group, the acoustic manipulation was mapped onto the appropriate articulatory/motor adjustment. Similar to results previously reported for both loudness and pitch perturbations¹² and delayed and frequency altered auditory feedback,³³ PD individuals maintain the functional ability to integrate sensory feedback with speech motor output. The change resulting from the disease, rather, appears to be related to the degree to which sensory feedback modulates or adjusts speech motor output.

Sensorimotor adaptation in PD

The sensorimotor adaptation paradigm involves the introduction of a constant and predictable change in re-afferent feedback to assess short-term sensorimotor learning. The current study is the first to evaluate motor learning in the speech domain in PD subjects. The reduction in adaptation for the PD group is consistent with the role of the BG in sequential motor learning³⁴ and previous observations of dysfunction in learning new motor sequences in PD.^{35,36} The present result is also consistent with prior findings of impaired sensorimotor learning in visually-guided reaching movements in PD patients relative to healthy controls.¹⁻³ Interestingly, the ability of the PD participants to adjust to the feedback alteration was related to their overall motor system involvement as measured by the UPDRS.

The reduced response to changes in auditory feedback associated with the control of the oral articulators stands in apparent contrast to a recent study demonstrating an *enhanced* laryngeal response following a perturbation in feedback related to loudness and fundamental frequency.¹² In that study, loudness and pitch feedback was altered suddenly and unexpectedly for brief periods during the production of sustained vowels in PD patients and age-matched controls. Following the manipulation, PD patients were observed to exhibit larger compensatory changes in loudness and fundamental frequency than the healthy controls.

There are two possible explanations for the different findings. One is that the auditory feedback mechanisms contributing to loudness, vocal pitch and articulatory control may be differentially impaired in PD. However, it is well known that increasing loudness is accompanied by changes in articulation and this relationship is a hallmark of one of the more effective behavioral treatments for PD (the Lee Silverman Voice Treatment or LSVT)^{37,38}. What is more likely is that all these parameters (loudness, pitch, articulation) are all affected in some complex way rather than all being independently controlled.

A perhaps more parsimonious explanation is that the tasks used by Liu *et al.* (2012) and our adaptation task tap into different sensorimotor processes. Liu *et al.* studied the immediate response to an unexpected and intermittent auditory feedback change during the production of a sustained vowel. Such a manipulation can be used to probe the control system (the operation of the subject's "internal model"), without examining changes to the model (i.e., no learning). In contrast, the present study employed an approach well established in the literature for studying speech motor learning.^{21,22,29,39-44} In this approach, a change in sensory feedback related to speech production is introduced and then maintained for a period of time, allowing subjects to gradually alter, with repeated practice, their speech motor plans in order to reduce the perceived sensory error. This approach, therefore, focuses more on the role of auditory feedback in speech motor learning (updating and maintaining the accuracy of the internal model) rather than on the use of sensory input for the immediate correction of motor output.

A recent study provided some empirical evidence and arguments in support of separate cortical regions used for learning but not for control.²⁸ These two related but independent processes, sensorimotor control and sensorimotor learning, may be differentially affected in PD. For normal speech production, BG damage may amplify reafferent feedback which creates the false perception in PD subjects of excessive loudness or effort. Under conditions in which speech motor patterns are modified, during therapy for example, BG damage may increase the threshold for reafferent feedback, reducing the overall system gain, and increasing the need for intensive and extensive training.

Acknowledgments

Funding agencies: This work was supported by grants from the Natural Sciences and Engineering Research Council of Canada, the Canadian Institutes of Health Research and the National Institutes of Health (NIDCD—R01DC012502.).

References

1. Contreras-Vidal JL, Buch ER. Effects of Parkinson's disease on visuomotor adaptation. *Exp Brain Res.* 2003; 150:25–32. [PubMed: 12698213]
2. Paquet F, Bedard MA, Levesque M, et al. Sensorimotor adaptation in Parkinson's disease: evidence for a dopamine dependent remapping disturbance. *Exp Brain Res.* 2008; 185:227–236. [PubMed: 17957363]
3. Messier J, Adamovich S, Jack D, Hening W, Sage J, Poizner H. Visuomotor learning in immersive 3D virtual reality in Parkinson's disease and in aging. *Exp Brain Res.* 2007; 179:457–474. [PubMed: 17146644]
4. Stern Y, Mayeux R, Hermann A, Rosen J. Prism adaptation in Parkinson's disease. *J Neurol Neurosurg Psychiatry.* 1988; 51:1584–1587. [PubMed: 3221228]
5. Snider SR, Fahn S, Isgreen WP, Cote LJ. Primary sensory symptoms in parkinsonism. *Neurology.* 1976; 26:423–9. [PubMed: 944393]
6. Koller WC. Sensory symptoms in Parkinson's disease. *Neurology.* 1984; 34:957–959. [PubMed: 6539873]
7. Hillen ME, Sage JI. Nonmotor fluctuations in patients with Parkinson's disease. *Neurology.* 1996; 47:1180–1183. [PubMed: 8909426]
8. Sailer A, Molnar GF, Paradiso G, Gunraj CA, Lang AE, Chen R. Short and long latency afferent inhibition in Parkinson's disease. *Brain.* 2003; 126:1883–1894. [PubMed: 12805105]
9. Duffy, JR. Motor speech disorders: Substrates, differential diagnosis, and management. St. Louis, Mo: Elsevier Mosby; 2005. p. 198
10. Liotti M, Ramig L, Willemse SC, Horstink MWIM, Maassen BAM. Improvement of voicing in patients with Parkinson's disease by speech therapy. *Neurology.* 2003; 61:1316–1317. [PubMed: 14610158]
11. Hammer MJ, Barlow SM. Laryngeal somatosensory deficits in Parkinson's disease: Implications for speech respiratory and phonatory control. *Exp Brain Res.* 2012; 201:401–409. [PubMed: 20012947]
12. Liu H, Wang EQ, Metman LV, Larson CR. Vocal responses to perturbations in voice auditory feedback in individuals with Parkinson's disease. *PLOS One.* 2012; 7:e33629. [PubMed: 22448258]
13. Caligiuri M, Abbs JH. Response properties of the perioral reflex in Parkinson's disease. *Exp Neurol.* 1987; 98:563–572. [PubMed: 3678433]
14. Diamond SG, Schneider JS, Markham CH. Oral sensorimotor defects in people with Parkinson's disease. *Adv Neurol.* 1987; 45:335–338. [PubMed: 3825706]
15. Schneider JS, Diamond SG, Markham CH. Deficits in orofacial sensorimotor function in Parkinson's disease. *Ann Neurol.* 1986; 19:275–282. [PubMed: 3963772]
16. Jobst EE, Melnick ME, Byl NN, Dowling GA, Aminoff MJ. Sensory perception in Parkinson disease. *Arch Neurol.* 1997; 54:450–4. [PubMed: 9109747]

17. Lewis GN, Byblow WD. Altered sensorimotor integration in Parkinson's disease. *Brain*. 2002; 125:2089–2099. [PubMed: 12183354]
18. Sarno MT. Speech impairment in Parkinson's disease. *Arch Phys Med Rehabil*. 1968; 49:269–275. [PubMed: 5651680]
19. Sapir, S.; Ramig, LO.; Fox, C. Voice, speech, and swallowing disorders. In: Pahna, R.; Lyons, KE., editors. *Handbook of Parkinson's disease*. New York: Informa healthcare; 2007.
20. Fox C, Ebersbach G, Ramig L, Sapir S. LSVT LOUD and LSVT BIG: Behavioral Treatment Programs for Speech and Body Movement in Parkinson Disease. *Parkinson's Disease*. 2012;10.1155/2012/391946
21. Houde JF, Jordan MI. Sensorimotor adaptation in speech production. *Science*. 1998; 279:1213–1216. [PubMed: 9469813]
22. Houde JF, Jordan MI. Sensorimotor adaptation of speech I: Compensation and adaptation. *J Speech Lang Hear Res*. 2002; 45:295–310. [PubMed: 12003512]
23. Purcell DW, Munhall KG. Adaptive control of vowel formant frequency: evidence from real-time formant manipulation. *J Acoust Soc Am*. 2006b; 120:966–977. [PubMed: 16938984]
24. Rochet-Capellan A, Ostry DJ. Simultaneous acquisition of multiple auditory-motor transformations in speech. *J Neurosci*. 2011; 31:2657–2662. [PubMed: 21325534]
25. Shiller DM, Sato M, Gracco VL, Baum SR. Perceptual recalibration of speech sounds following speech motor learning. *J Acoustic Soc Am*. 2009; 125:1103–1113.
26. Shiller DM, Gracco VL, Rvachew S. Auditory-Motor Learning during Speech Production in 9–11-Year-Old Children. *PLOS One*. 2012; 5:e12975. [PubMed: 20886033]
27. Villacorta VM, Perkell JS, Guenther FH. Sensorimotor adaptation to feedback perturbations of vowel acoustics and its relation to perception. *J Acoust Soc Am*. 2007; 122:2306–2319. [PubMed: 17902866]
28. Shum M, Shiller DM, Baum SR, Gracco VL. Sensorimotor integration for speech motor learning involves the inferior parietal cortex. *Eur J Neurosci*. 2011; 34:1817–22. [PubMed: 22098364]
29. Feng Y, Gracco VL, Max L. Integration of auditory and somatosensory error signals in the neural control of speech movements. *J Neurophysiol*. 2011; 106:667–679. [PubMed: 21562187]
30. Darley, FL.; Aronson, AE.; Brown, JR., editors. *Motor speech disorders*. Philadelphia: W. B. Saunders; 1975.
31. Peterson GE, Barley HL. Control methods used in a study of the vowels. *J Acoust Soci Am*. 1952; 24:175–184.
32. Kent, RD.; Read, C. *Acoustic analysis of speech*. 2. San Diego, CA: Singular Press; 2001.
33. Brendel B, Lowit A, Howell P. The effects of delayed and frequency shifted feedback on speakers with Parkinson's disease. Special issue: speech motor control conference proceedings. *J Med Speech Lang Pathol*. 2004; 12:131–138. [PubMed: 19330038]
34. Graybiel AM. The basal ganglia: learning new tricks and loving it. *Curr Opin Neurobiol*. 2005; 15:638–644. [PubMed: 16271465]
35. Frith CD, Bloxham CA, Carpenter KN. Impairments in the learning and performance of a new manual skill in patients with Parkinson's disease. *J Neurol Neurosurg Psychiatry*. 1986; 49:661–668. [PubMed: 3734823]
36. Swinnen SP, Steyvers M, Van DBL, Stelmach GE. Motor learning and Parkinson's disease: refinement of within-limb and between-limb coordination as a result of practice. *Behav Brain Res*. 2000; 111:1–2. [PubMed: 10840127]
37. Sapir S, Ramig LO, Fox CM. Intensive voice treatment in Parkinson's disease: Lee Silverman Voice Treatment. *Expert Reviews of Neurotherapeutics*. 2011; 11(6):815–830.
38. Ramig, LO.; Fox, CN.; Sapir, S. Speech and Voice Disorders in Parkinson's disease. In: Olanow, CW.; Stocchi, F.; Lang, AE., editors. *Parkinson's Disease: Non-Motor and Non-Dopaminergic Features*. 2011. p. 348–362.
39. Baum SR, McFarland DH. The development of speech adaptation to an artificial palate. *J Acoust Soc Am*. 1997; 102:2353–2359. [PubMed: 9348694]
40. Jones JA, Munhall KG. Perceptual calibration of F0 production: evidence from feedback perturbation. *J Acoust Soc Am*. 2000; 108:1246–1251. [PubMed: 11008824]

41. Jones JA, Munhall KG. Learning to produce speech with an altered vocal tract: the role of auditory feedback. *J Acoust Soc Am*. 2003; 113:532–543. [PubMed: 12558289]
42. Tremblay S, Shiller DM, Ostry DJ. Somatosensory basis of speech production. *Nature*. 2003; 423:866–869. [PubMed: 12815431]
43. Nasir SM, Ostry DJ. Somatosensory precision in speech production. *Curr Biol*. 2006; 16:1918–1923. [PubMed: 17027488]
44. MacDonald EN, Johnson EK, Forsythe J, Plante P, Munhall KG. Children’s development of self-regulation in speech production. *Curr Biol*. 2012; 22:113–117. [PubMed: 22197241]

Figure 1.

Testing sequence illustrating the scheme for introducing the auditory feedback manipulation. During the baseline trials subjects read words off a computer screen in the presence of unaltered auditory feedback (50 trials). During the Hold phase the feedback alteration was introduced suddenly and maintained for 100 trials followed by a period of gradual restoration of auditory feedback (40 trials) and another 30 trials with normal auditory feedback (After-effect). The feedback manipulation used was an increase in the first formant (F1) frequency of the subjects auditory feedback which is associated inversely with the height of the tongue in the oral cavity.

Figure 2. The mean first and second formant (F1 and F2) frequencies (left) and standard deviation of the mean of F1 and F2 (right) in Hertz for the baseline productions of the target vowel / / for the PD and Control subjects.

Figure 3.

Change in the acoustic vowel characteristics (F1 and F2) for the target word throughout the speech adaptation testing sequence. Shown are blocks of 10 trials with onset of the feedback alteration beginning at trial block 6 followed by 100 trials (10 blocks) under maximum feedback alteration (*hold* phase), and then gradual (RAMP-OFF) and subsequent complete restoration of unaltered feedback (A-E).

Figure 4.

Mean normalized formant values at the four time-points of interest during the speech adaptation procedure. See text for details.