Suspension of doctors

Reports highlight lack of human resources and legal expertise in NHS trusts

EDITOR—Although I agree with Empey that the recent direction from the Department of Health on the suspension of doctors represents progress—major progress with the mandatory need for referral to the National Clinical Assessment Authority (NCAA)—he has glossed over the fact that it has taken nine years for it to be produced.¹

He also failed to highlight the risk to patients' safety that is caused by suspensions and threats of suspensions, as exemplified by case 8 in the National Audit Office's report, in which a trust's culture was such that consultants feared bullying and victimisation if they raised any concerns about safety issues.²

Badly planned and executed suspensions compromise care for patients and put them at risk; in view of the requirements of the October 2002 Code of Conduct for Managers to give top priority to protecting patients from risk the rash, rushed, arbitrary suspension of a doctor should not be happening these days. If it does then the responsible manager should be held to account by either the trust board or even the NHS chief executive under the terms of the code. If those pathways of accountability fail then MPs on the Committee of Public Accounts can hold the Department of Health to account on behalf of the public.

The procedure and processes around suspensions of both senior managers and consultants are complex and require specialist human resources and legal skills and good, non-biased judgment to avoid unfair dismissal and inappropriate suspensions. A case could be made for establishing a central unit. Alternatively a more loose network of appropriate legal and human resources experts could be established to be called on by any trust to help in these difficult cases so that the proper processes are followed.

Nigel Dudley *consultant in elderly medicine*
St James's University Hospital, Leeds LS9 7TF
nigel.dudley@leedsth.nhs.uk

Competing interests: None declared.

GMC may be ultimate sacrifice

EDITOR—In his letter above and on bmj.com in response to Empey's editorial, Dudley says that a case could be made for establishing a unit that has the expertise to handle suspensions.^{1 2} I think a case should be made for re-establishing a unit to handle suspensions. This was a function that was carried out by the old regions and ceased when they were dissolved.

Ten years ago a regional medical officer commented that this power would devolve to local trusts.³ Each medical director would face the problem for the first time, and there would be no corporate knowledge. Each trust would make its own mistakes and would learn nothing, only for another trust to repeat its learning curve. All this repeated learning about disciplinary procedures would always be done at the expense of someone else's career. The regional medical officer was the later knighted Liam Donaldson—what better man to undertake the resolution of the problem he predicted all those years ago, as he is now chief medical officer?

I pointed out the conjunction of trusts lying about waiting lists and other trusts making less than frank referrals to the General Medical Council a year or so ago.⁴ Trusts have lied about waiting lists and are not so picky either about what they allege to the GMC. Trusts are not registered with and answerable to the GMC as medical practitioners are, and so in effect they can say what they like. That is why I suggested to the GMC that they put observers into hospital inquiries—so that they could see the goings-on first hand. The GMC declined the invitation.⁵

I now regard the GMC, packed as it is with Whitehall appointees, as both deaf and

blind, something it has in common with the BBC's board of governors. Both institutions have not aged well, are unfit for their purpose, and should be pensioned off.

Oliver R Dearlove *consultant anaesthetist*
Royal Manchester Children's Hospital, Manchester M27 4HA
o.dearlove@man.ac.uk

Competing interests: ORD is a councillor of the Royal College of Anaesthetists and is a full time NHS consultant. These views are his own and are not shared by the Royal College of Anaesthetists or his employer.

- 1 Dudley N. NAO reports on inappropriate waiting list adjustments and doctor suspensions highlight lack of NHS trust HR and legal expertise. Electronic response to: Suspension of doctors. *bmj.com* 2004. <http://bmj.com/cgi/eletters/328/7433/181#48242> (accessed 18 Feb 2004).
- 2 Empey D. Suspension of doctors. *BMJ* 2004;328:181-2. (24 January.)
- 3 Donaldson LJ. Doctors with problems in an NHS workforce *BMJ* 1994;308:1277-82.
- 4 Dearlove O. Liar! Liar! Electronic response to: Three trusts deliberately misreported data, says Audit Commission. *bmj.com* 2003. <http://bmj.com/cgi/eletters/326/7388/517#30282> (accessed 14 Feb 2004).
- 5 Tupper J. Re: GMC—the usual unaccountable mess continues. Electronic response to: GMC admits failings that left Shipman's patients unprotected. *bmj.com* 2003. <http://bmj.com/cgi/eletters/327/7426/1248#45662> (accessed 18 Feb 2004).

Medical suspensions may have ideological nature

EDITOR—Like Empey, I welcome the new guidance and directions from the Department of Health on the suspension of doctors.¹ I was suspended for six months by my NHS trust, from October 2000, before the National Clinical Assessment Authority (NCAA) was established, and I had to undergo a period of supervision for a year following a recommendation from the Royal College of Psychiatrists.

According to the National Audit Office I am in the minority of suspended doctors who have returned to work (40%).² The length of my suspension was below average (47 week average), not least because I went along with the recommendation from the college, even though I did not accept it. As noted by the National Audit Office, concerns exist about the thoroughness and rigour of external assessments.

My problem is the potential ideological nature of suspensions. Medicine, perhaps particularly psychiatry, is not an exact science. Attitudes and approaches vary from the biomedical to the psychosocial, and biomedical bias is reinforced by defensiveness about the personal nature of medicine.³ Psychiatry especially has been infused by conflict, perhaps best reflected in the debate about so called anti-psychiatry.⁴

1 Empey D. Suspension of doctors. *BMJ* 2004;328:181-2. (24 January.)

2 National Audit Office. *The management of suspensions of clinical staff in NHS hospital and ambulance trusts in England*. London: Stationery Office, 2003. (HC 1143 Session 2002-2003.)

I was told I needed retraining and would be sent for further education in organic psychiatry. The college assessors indicated that, if I did not accept their recommendation, my philosophy about psychiatry would need to be examined and my scepticism about the use of medication challenged. Although none of these things happened, it is difficult to think they were not motivating factors in my suspension.

I am not sure if critical psychiatry, which is the position I represent, has been strengthened by my experience. At least, information about critical psychiatry is now available on the Royal College of Psychiatrists' website.⁵

D B Double *consultant psychiatrist*
Norfolk Mental Health Care NHS Trust, Norfolk
NR6 5BE
dbdouble@dbdouble.co.uk

Competing interests: None declared.

- 1 Empey D. Suspension of doctors. *BMJ* 2004;328:181-2. (24 January.)
- 2 National Audit Office. *The management of suspensions of clinical staff in NHS hospital and ambulance trusts in England*. London: Stationery Office, 2003. (HC 1143 Session 2002-2003.)
- 3 Stewart M, Brown JB, Weston WW, McWhinney IR, McWilliam CL, Freeman TR. *Patient-centred medicine. Transforming the clinical method*. 2nd ed. Abingdon: Radcliffe Medical Press, 2003.
- 4 Double DB. The limits of psychiatry. *BMJ* 2002;324:900-4.
- 5 Royal College of Psychiatrists webguide. *Critical psychiatry*. www.rcpsych.ac.uk/info/webguide/critical.htm (accessed 14 Feb 2004).

Be careful with suspensions

EDITOR—With reference to Empey's editorial on the suspension of doctors,¹ suspensions that are badly performed cause human, psychological costs that cannot be measured easily and that can be widely different and sometimes devastating for doctors, their health, their families, and their subsequent lifestyles and working style.

This happens mainly when the authority is doing nothing positive while the doctor's skills and confidence are allowed to decline.

Italy has a blame culture; most failures in standards of care are caused by weaknesses in systems rather than individuals. Also Italy doesn't have a team practices' system, so no procedure is managed in the practice itself. Everything is decided by the local medical council, which could be "cleverly" reflexive or follow the suspension culture, often resulting in reactive and apparently arbitrary decisions being made.

Quick and objective investigations of allegations, adherence to reasonable time scales, and full consideration of alternatives to exclusion are needed. Often doctors lack knowledge, competencies, and experience to help them act with empathy, which is fundamental for real improvements in practice.

Francesco Carelli *national representative*
EURACT Council, Italian College of General Practitioners, 20123 Milan, Italy
carfra@tin.it

Competing interests: None declared.

- 1 Empey D. Suspension of doctors. *BMJ* 2004;328:181-2. (24 January.)

Ethics review in research

Research governance also delays research

EDITOR—The study by Hearnshaw and the accompanying editorial by Glasziou and Chalmers on ethics review highlight that ethics approval in the United Kingdom is a time consuming and costly process that may delay and potentially prevent ethical studies from taking place.^{1,2} We are aware of a very large multinational trial being conducted across Europe and North America in which the absence of a UK centre has been attributed to the perceived difficulties of obtaining ethics approval in the United Kingdom.

Yet delays in initiating research in the United Kingdom are now attributable not only to ethics review but also to the process of gaining research governance approval from health trusts.³ In our recent experience of organising multicentred, randomised controlled trials, the multitude of different application processes that have been implemented mean that research governance presents another large barrier to research.

The combined effect of current ethics and research governance requirements in the United Kingdom will have serious consequences for the number of trials that are carried out here in the future. In turn, this will damage patients' future health care.

David J Torgerson *director*
djt6@york.ac.uk

Jo C Dumville *research fellow*
York Trials Unit, Department of Health Sciences,
York University, York YO10 5DD

Competing interests: None declared.

- 1 Hearnshaw H. Comparison of requirements of research ethics committees in 11 European countries for a non-invasive interventional study. *BMJ* 2004;328:140-1. (17 January.)
- 2 Glasziou P, Chalmers I. Ethics review roulette: what can we learn? *BMJ* 2004;328:121-2. (17 January.)
- 3 Dumville JC, Watson J, Raynor P, Torgerson DJ. Research governance: a barrier to ethical research. *Q Rev Med* 2004 (in press).

Role of ethics committee review is interpreted widely

EDITOR—Hearnshaw and Glasziou and Chalmers highlight the inconsistencies and arbitrary decision making of UK research ethics committees and show that this may delay and suppress clinical research in the United Kingdom.^{1,2} I recently submitted a study that required six months, countless emails, and submission of 1443 sheets of A4 paper to gain approval to study 20 standard diagnostic biopsy specimens. This seems excessive as the subjects had already given consent for the use of surplus clinical material for research or teaching.

To examine whether application of ethics committee review is often so irregular, I reviewed all studies published in *Gut* and *Alimentary Pharmacology and Therapeutics* during 2003. Of 202 papers that required some type of research specific intervention (therapeutic or investigational), 169 (84%) explicitly stated that ethics committee review and informed consent had been obtained. Most of the omissions were questionnaires.

In 124 studies using surplus diagnostic clinical material (usually tissue biopsy specimens) only 59 (47%) confirmed ethics committee approval and 31 (25%) informed consent from the patients ($P < 0.001$ for both, compared with interventional studies).

Authors may have simply omitted this information. More likely, is that interpretation of the role of ethics committees varies widely, both in the United Kingdom and worldwide. It is imperative that a more uniform practice is applied both to protect patients and prevent any group of researchers being disadvantaged and research unavoidably being delayed or even not attempted. Referees and editors should become more proactive in ensuring such standards are uniform.

Ian L P Beales *clinical senior lecturer*
School of Medicine, Health Policy, and Practice,
University of East Anglia, Norfolk NR31 6TY
i.beales@uea.ac.uk

Competing interests: ILPB is actively engaged in clinical investigational research. All such studies are reviewed and approved by separate local ethics and research committees.

- 1 Hearnshaw H. Comparison of requirements of research ethics committees in 11 European countries for a non-invasive interventional study. *BMJ* 2004;328:140-1. (17 January.)
- 2 Glasziou P, Chalmers I. Ethics review roulette: what can we learn? *BMJ* 2004;328:121-2. (17 January.)

Ethics committees are risk averse

EDITOR—Hearnshaw and Glasziou and Chalmers have made points I would have made myself had my time not been taken with writing ethics applications.^{1,2}

I am currently coordinating a qualitative investigation into the attitudes of Asian families to child and adolescent mental health problems and services. Because the research involves service users, non-patient members of the community, children, and adults I have been required to make full submissions to three ethics committees, one university and two NHS. Despite the fact that the research had received very positive peer review from one of the top medical sociologists in the United Kingdom, each committee criticised the design or quality of the methods and required fairly substantial clarifications and alterations to patient information. For example, one committee required that I lengthen the children's information sheet to bring it into line with the guidelines of the Central Office for Research Committees (COREC) (www.corec.org.uk), while another said it was "too wordy" and required me to shorten it.

Meanwhile, our local health board is carrying out a similar focus group study without need for ethics approval because it is seen as "user involvement," is not hypothesis driven, and will not be published in the research literature.

Although I can see the need for a research governance framework, the current guidelines seem to have produced a climate of risk aversion without any weighing of possible benefits to patients. In my area, research by busy clinicians and by students on time limited MSc programmes is all but

grinding to a halt because the ethics barrier is seen as insurmountable. This is yet another example of clinical academia being strangled at its roots.

Helen J Minnis senior lecturer in child and adolescent psychiatry
Child Psychiatry, Section of Psychological Medicine, University of Glasgow, Glasgow G3 8SJ
h.minnis@clinmed.gla.ac.uk

Competing interests: None declared.

- 1 Hearnshaw H. Comparison of requirements of research ethics committees in 11 European countries for a non-invasive interventional study. *BMJ* 2004;328:140-1. (17 January.)
- 2 Glasziou P, Chalmers I. Ethics review roulette: what can we learn? *BMJ* 2004;328:121-2. (17 January.)

Time has come for reassessment

EDITOR—Glasziou and Chalmers are to be congratulated for their stand on the important issue of ethics review.¹ For many years MSc students in this school undertook projects considering policy questions facing local trusts and health authorities. These typically entailed a literature review and interviews with managers, clinicians, and other relevant stakeholders. These interviewees are now deemed research subjects, thus requiring each student to obtain ethics approval.

The lengthy process of getting approval has forced us to advise students to abandon this model of research, concentrating instead on desk based studies. The links forged with the key stakeholders meant that many of these projects led to improvements in the care provided, and it is far from clear, at least to me, that this development has increased the total sum of human welfare.

Martin McKee professor of European public health
London School of Hygiene and Tropical Medicine, London WC1E 7HT
martin.mckee@lshtm.ac.uk

Competing interests: None declared.

- 1 Glasziou P, Chalmers I. Ethics review roulette: what can we learn? *BMJ* 2004;328:121-2. (17 January.)

Exercise training in patients with chronic heart failure

More information is needed on outcomes in individual studies

EDITOR—The ExTraMATCH collaborative has conducted a meta-analysis for which there is a clear need.¹ The small size of studies of exercise training for patients with heart failure means that only a meta-analysis can adequately address the question of whether exercise training reduces mortality. Having read their report, I am left wanting to know more.

Is the omission of outcome data for the individual studies intended? The number of deaths in the training and control groups would be interesting for each study in turn. I have consulted the cited papers for

each study, but the mortality data are not reported in several of them, or are reported for a shorter period than indicated in table 1 of the meta-analysis.

A forest plot of the hazard ratios (with 95% confidence intervals) would be interesting for the identified studies. Figure 2 of the meta-analysis shows that the exercise regimen in one of the larger studies achieves a very large reduction in mortality, but the studies are not identified in that figure. Is the exercise intervention in this study doing something different from that in the others?

Also, did the authors consider including, in the proportional hazards regression, a covariate identifying the study in which a patient participated? If the average severity of disease in patients varied across studies, and such a covariate was not included in the regression equation, this may cause the observed treatment effect to decrease as follow up proceeded. The test of proportional hazards indicates that this is not a major problem, but this test is not always as sensitive as desired.

Chris Metcalfe lecturer in medical statistics
Department of Social Medicine, University of Bristol, Bristol BS8 2PR
chris.metcalfe@bristol.ac.uk

Competing interests: None declared.

- 1 Exercise training meta-analysis of trials in patients with chronic heart failure (ExTraMATCH). *BMJ* 2004;328:189-92. (24 January.)

Authors' reply

EDITOR—The figure shows the forest plots of the hazard ratios and 95% confidence intervals. The forest plots were removed during revision for reasons of space.

As Metcalfe observes, in one fairly large study the exercise regimen seemed to be achieving a greater beneficial effect.¹ In this, the training type (supervised cycling, 60 minutes three days a week) was similar to that in the other studies, at a slight lower intensity (60% of peak VO₂), but the factor that we suspect made the difference was the length of the programme which (at 420 days) was almost double the average of the others (213 (SD 135) days). This is in keeping with the non-significant trend observed in our subgroup analysis.

Although pathophysiological grounds exist to suspect that longer periods of training might be more likely to be effective, even meta-analysis of all available data cannot resolve the question clearly. It is possible that in a future study with more events such differences may become detectable.

We thank Metcalfe for suggesting insertion of a covariate identifying the study into the Cox survival regression. The beneficial

effect of exercise remains evident, both for mortality (odds ratio 0.67, 95% confidence interval 0.47 to 0.95) and for the combined

Hazard ratios and 95% confidence intervals for individual studies for effect of exercise training on risk of death and on risk of death and admission to hospital (full data available on bmj.com)

end point of mortality or admission (0.67, 0.52 to 0.87).

The benefits for health and survival of supervised exercise training (instead of general advice on healthy living) in heart failure are now established beyond reasonable doubt. We predict, however, that implementation will be far slower for this than for treatments such as drugs and implantable devices—both of which have a natural large industry lobby behind them.

Massimo F Piepoli member, ExTraMATCH collaborative
m.piepoli@imperial.ac.uk

Darrel P Francis member, ExTraMATCH collaborative
Clinical Cardiology, Royal Brompton Hospital, Imperial College of Science, Technology and Medicine, London SW3 6LY

Competing interests: None declared.

- 1 Belardinelli R, Georgiou D, Cianci D, Purcaro A. Randomized, controlled trial of long term moderate exercise training in chronic heart failure: effects on functional capacity, quality of life, and clinical outcome. *Circulation* 1999;99:1173-82.

Data for figures are available on bmj.com

Increase in non-evidence based use of antidepressants in children is cause for concern

EDITOR—The use of antidepressants in children is increasing, although the evidence for the effectiveness and safety of pharmaco-

logical treatment of depressive disorder in children and adolescents is scant¹ and widely debated²—particularly for selective serotonin reuptake inhibitors (SSRIs).^{3,4}

We used the ARNO database (<http://sanita.cineca.it/public/arno/arnoeng.htm>) to analyse drug prescriptions written during 2002 to 568 770 Italian patients under 18. A total of 1600 young people (2.8 per 1000) received at least one antidepressant, 1200 of them an SSRI and 297 a tricyclic antidepressant. Two thirds of prescriptions (8284/11 369) were for adolescents (age 14–17), mostly girls (4583/5872; rate 10.6 per 1000 v 5.9 per 1000 for boys).

These prescribing rates are lower than those reported for the United States (1–2%) and the Netherlands (4.4 per 1000), but nevertheless about 28 000 youths are exposed to treatment with antidepressants, 21 000 of them receiving SSRIs. We found a 4.5-fold increase in the rate of prescriptions for SSRIs (from 0.47 per 1000 to 2.11 per 1000) between the 2000 and 2002 datasets, which is worrying, even though it agrees with data from other reports.⁵

All the SSRIs prescribed (most commonly paroxetine) are unlicensed paediatric drugs, with the exception of sertraline for obsessive compulsive disorder in children under 6 years. The occurrence of depressive syndrome and obsessive compulsive disorder in children must be determined, along with their therapeutic needs. Appropriate, independently funded studies should be planned to guarantee effective and safe evidence based therapeutic approaches for children, adolescents, and their families.

Antonio Clavenna senior research fellow
clavenna@marionegri.it

Maurizio Bonati head
Laboratory for Mother and Child Health, Mario Negri Institute for Pharmacological Research, 20157 Milan, Italy

Elisa Rossi statistician

Marisa De Rosa senior researcher
Interuniversity Consortium, Bologna, Italy

Competing interests: None declared.

1 Hazell P. Depression in children and adolescents. *Clin Evid* 2003;10:375–85.

2 Ramchandani P. Treatment of major depressive disorder in children and adolescents. *BMJ* 2004;328:3–4. (3 January.)

3 Moynihan R. FDA advisory panel calls for suicide warnings over new antidepressants. *BMJ* 2004;328:303. (7 February.)

4 Committee on Safety of Medicines. Use of selective serotonin reuptake inhibitors (SSRIs) in children and adolescents with major depressive disorder (MDD). <http://www.mhra.gov.uk> (accessed 9 February 2004).

5 Magno Zito J, Safer DJ, dosReis S, Gardner JF, Soeken K, Boles M, et al. Rising prevalence of antidepressants among US youths. *Pediatrics* 2002;109:721–2.

Top up fees and medicine

Waive repayments completely if working life is spent in the NHS

EDITOR—If, as Mayor reports,¹ the BMA is concerned that one in four medical graduates is considering not practising medicine and that the introduction of top up fees might encourage more graduates into lucrative private work instead of public service, then why cap the fees at £3000 a year for medical students?

Why not instead charge them nearer the full costs but waive their repayments while they are earning less than £15 000 or working exclusively as a medical practitioner for the NHS or in another public sector?

Medical students who worked until retirement for the state would then never have to repay a penny of the costs of their expensive state funded education. At the very least, this should help applicants to medical schools think a little more carefully about their future careers before applying. Should they wish to work in private medicine or in another career unrelated to health that is similarly lucrative, they could at least help, through their repayments, to fund the training of the extra medical students needed to replace their lost efforts. Waiving top up fees for people who work for the state could be of wider benefit in encouraging people to use their education wisely.

According to the 2001 census,² there are already over 26 000 qualified medical doctors in England and Wales in employment but not working as medical practitioners. Only 5000 of these seem to have been working in non-medical public sector jobs.

Danny Dorling professor of geography
Department of Geography, University of Sheffield, Sheffield S10 2TN
Danny.Dorling@sheffield.ac.uk

Competing interests: DD went to university in Britain when access was free.

1 Mayor S. University fee changes may deter poor students from studying medicine. *BMJ* 2004;328:128. (17 January.)

2 National Statistics. *Census 2001: National Report for England and Wales*. Standard table 116. Available at www.statistics.gov.uk/StatBase/Product.asp?vlnk=10441 (accessed 15 Mar 2004).

Ability to commit to unspecified debt may take precedence over ability

EDITOR—The introduction of top up fees is highly emotive.¹ The selection of medical students has a markedly skewed socioeconomic distribution: seven out of 10 medical students come from a professional or managerial background. This does not reflect the make up of society as a whole.

The medical course is unique by virtue of its length, unavoidable expenditures, and the lack of opportunity for students to supplement their income with part time work. The government's proposals do not take into account the unique circumstances that medical students find themselves in. In fact, the student income and expenditure figures on which the government has based its reforms exclude medical students.

One of the major determinants of application to university is debt aversion; this is more marked in families who come from lower socioeconomic groups. The new proposals will increase the levels of debt for all medical students and cannot, therefore, be seen to encourage participation from all socioeconomic groups.

I am concerned that the government's reforms of student support will lead to a system where the willingness to commit to

unknown debt will take precedence over ability.

Daniel C Gibbons joint deputy chairman
Medical Students Committee, BMA, London
WC1H 9JP
d.gibbons@ucl.ac.uk

Competing interests: None declared.

1 Mayor S. University fee changes may deter poor students from studying medicine. *BMJ* 2004;328:128. (17 January.)

Regularly imbibing Dickens may inspire healthfulness

EDITOR—Anaesthetist Nick Jefferies's announcements on happiness unhappily contain an incorrect reference.¹ The one from Dickens he wrongly attributes to the *Pickwick Papers*. We must cordially insist, therefore, that he now read all the novels, to locate its proper source. This is no condign punishment, rather the reverse. In such perusal, particularly of Dickens's first and stupendously joyful novel, *Pickwick*, Jefferies may find, like millions of others since publication of the first instalment in 1836, an entrancing happiness beyond his wildest dreams.

Mr Micawber famously defined happiness

For reasons quite mystifying, regular imbibers of such literary delectations only seldom, and then only briefly, become patients. It is tempting to deduce that such items are restorative, even sometimes curative. However, such anecdotal observation and deduction have no place in modern, scientific, medical thought. Yet they are still mightily striking.

William G Pickering medical practitioner
7 Moor Place, Newcastle upon Tyne NE3 4AL
wgpi@hotmail.com

Competing interests: None declared.

1 Jefferies N. Happiness is ... *BMJ* 2004;328:453. (21 February.)