High-Resolution Mapping of Complex Traits with a Four-Parent Advanced Intercross Yeast Population

Francisco A. Cubillos,^{*,†,‡,1} Leopold Parts,^{§,***,1} Francisco Salinas,^{††} Anders Bergström,^{††} Eugenio Scovacricchi,^{*} Amin Zia,^{**} Christopher J. R. Illingworth,[§] Ville Mustonen,[§] Sebastian Ibstedt,^{§§} Jonas Warringer,^{§§} Edward J. Louis,^{*,***} Richard Durbin,[§] and Gianni Liti^{††,2}

^{*}Centre for Genetics and Genomics, Queen's Medical Centre, University of Nottingham, Nottingham, NG7 2UH, United Kingdom,

[†]Departamento de Ciencia y Tecnología de los Alimentos and [‡]Centro de Estudios en Ciencia y Tecnología de Alimentos, Universidad de Santiago de Chile, Santiago 9170201, Chile, [§]The Wellcome Trust Sanger Institute, Hinxton, CB10 1HH, United Kingdom, ^{**}Donnelly Centre for Cellular and Biomolecular Research and ^{††}Department of Cell and Systems Biology, University of

Toronto, Toronto, Ontario M5S 3B2, Canada, ^{††}Institute for Research on Cancer and Ageing of Nice, Centre National de la Recherche Scientifique UMR 7284–Institut National de la Santé et de la Recherche Médicale U1081–Université de Nice Sophia

Antipolis, Nice 06107, France, ^{§§}Department of Chemistry and Molecular Biology, University of Gothenburg, Gothenburg 40530, Sweden, and ^{***}Centre of Genetic Architecture of Complex Traits, University of Leicester, Leicester LE1 7RH, United Kingdom

ABSTRACT A large fraction of human complex trait heritability is due to a high number of variants with small marginal effects and their interactions with genotype and environment. Such alleles are more easily studied in model organisms, where environment, genetic makeup, and allele frequencies can be controlled. Here, we examine the effect of natural genetic variation on heritable traits in a very large pool of baker's yeast from a multiparent 12th generation intercross. We selected four representative founder strains to produce the *Saccharomyces* Genome Resequencing Project (SGRP)-4X mapping population and sequenced 192 segregants to generate an accurate genetic map. Using these individuals, we mapped 25 loci linked to growth traits under heat stress, arsenite, and paraquat, the majority of which were best explained by a diverging phenotype caused by a single allele in one condition. By sequencing pooled DNA from millions of segregants grown under heat stress, we further identified 34 and 39 regions selected in haploid and diploid pools, respectively, with most of the selection against a single allele. While the most parsimonious model for the majority of loci mapped using either approach was the effect of an allele private to one founder, we could validate examples of pleiotropic effects and complex allelic series at a locus. SGRP-4X is a deeply characterized resource that provides a framework for powerful and high-resolution genetic analysis of yeast phenotypes and serves as a test bed for testing avenues to attack human complex traits.

THE strong tendency for progeny to closely resemble their parents has turned out to be difficult to understand in detail. Nearly all traits, including lifetime risk for many common diseases, have a complex genetic basis that is determined by multiple quantitative trait loci (QTL) (Donnelly 2008; Manolio *et al.* 2009). The first step toward accurate models of trait variability, and a prerequisite for predicting and modulating them, is characterization of the underlying

genetic factors in the context of rest of the genome and their external environment. Research in model systems has led the way in this effort and produced powerful experimental and computational approaches for genetic mapping (Nordborg and Weigel 2008; Flint and Mackay 2009; Mackay *et al.* 2009).

A traditional, well-controlled approach for finding the QTL underlying natural phenotypic variation is to analyze a large number of progenies from two-parent crosses (Brem *et al.* 2002; Simon *et al.* 2008). Studies using this design have improved our understanding of complex traits and provided concrete evidence of natural segregating variants (Mackay *et al.* 2009), but have been limited in their scope with regard to the extent of genetic variation between the two parents. Mapping populations of popular model organisms ranging from fruit flies (King *et al.* 2012) and mice

Copyright © 2013 by the Genetics Society of America

doi: 10.1534/genetics.113.155515

Manuscript received July 25, 2013; accepted for publication September 8, 2013

Supporting information is available online at <http://www.genetics.org/lookup/suppl/doi:10.1534/genetics.113.155515/-/DC1>.

¹These authors contributed equally to this work.

²Corresponding author: Institute for Research on Cancer and Ageing of Nice, Centre National de la Recherche Scientifique UMR 7284–Institut National de la Santé et de la Recherche Médicale U1081, Faculté de Médecine, Université de Nice Sophia Antipolis, 28 Ave. de Valombrose, 06107 Nice Cedex 2, France. E-mail: gianni.liti@unice.fr

(Churchill *et al.* 2004; Durrant *et al.* 2011) to plants (Kover *et al.* 2009; Gan *et al.* 2011; Huang *et al.* 2011) has recently expanded the genetic and phenotypic diversity available to study by incorporating a wider repertoire of founder lines. These panels are the forefront of complex trait research and bear close resemblance to natural populations in multiple ways. Beyond increased variation, multiple founders introduce the possibility of more than two independent alleles at a locus and a larger space of potential epistatic interactions (Huang *et al.* 2012), while additional outcrossing rounds break linkage to further mix alleles. Nevertheless, lack of complete genotype information and the extent of remaining linkage in these recombinant lines have limited the power to detect small-effect QTL and identify the causative loci.

A multiparent mapping population has notably been missing in the budding yeast *Saccharomyces cerevisiae*, perhaps the most powerful eukaryotic model organism (Liti and Louis 2012). Yeast is a powerhouse of quantitative genetics due to its small genome size and very high recombination rate and the potential for accurate quantitative phenotyping, ease of obtaining and maintaining large mapping populations, and the ability to manipulate the genome at a single-base resolution. So far, nearly all yeast recombinant panels have been constructed by crossing the reference laboratory strain S288c (or one of its derivatives) with a wild isolate. However, laboratory strains poorly recapitulate the properties of natural populations, often represent phenotypic outliers when compared to the rest of the species (Warringer *et al.* 2011), and contain artificial auxotrophic markers that confound mapping experiments (Perlstein *et al.* 2007). To overcome this problem, we previously picked four natural yeast isolates sequenced in the *Saccharomyces* Genome Resequencing Project (SGRP) (Liti *et al.* 2009a) and released first-generation recombinant lines for each of the six pairwise crosses (Cubillos *et al.* 2011).

Here, we present SGRP-4X, a yeast mapping population obtained from outcrossing four wild founders representative of the main *S. cerevisiae* lineages for 12 generations. SGRP-4X contains >10 million segregants with fine-grained mosaic genomes and greatly reduced linkage, while retaining the phenotypic diversity of the parental strains. We demonstrate the power and resolution of QTL mapping in this population by both traditional linkage analysis on 179 genotyped and phenotyped individuals and a recently developed approach of sequencing the entire population under selection. SGRP-4X is a powerful, deeply characterized resource for high-resolution mapping of complex traits.

Materials and Methods

Generating the SGRP-4X advanced intercross lines

Two biological replicates of the intercrossed population were made. For replicate 1, parental strains YPS128 [North American (“NA”): *MATa*, *ho::HygMX*, *ura3::KanMX*] and DBVPG6044 [West African (“WA”): *MATα*, *ho::HygMX*,

ura3::KanMX] were crossed and grown overnight in complete media (YPDA) to generate diploid F₁ hybrids. In parallel, strains Y12 [Sake (“SA”): *MATa*, *ho::HygMX*, *ura3::KanMX*, *lys2::URA3*] and DBVPG6765 [Wine/European (“WE”): *MATα*, *ho::HygMX*, *ura3::KanMX*, *lys2::URA3*] were similarly crossed. To confirm successful crosses, we isolated individual colonies and performed mating tests, using tester strains Y55-2369 (*MATa*, *hoΔ*, *ura2-1*, *tyr1-1*) and Y55-2370 (*MATa*, *hoΔ*, *ura2-1*, *tyr1-1*) as well as diagnostic PCR for the *MAT* locus (Huxley *et al.* 1990). For replicate 2, we repeated the procedure described above with inverted combination of mating types (NA *MATα* × WA *MATa* and SA *MATα* × WE *MATa*). F₁ hybrids and haploid segregants were treated and generated as previously described (Parts *et al.* 2011). Briefly, F₁ hybrids were grown overnight and full plates were replicated onto KAc at 23° for sporulation during 10 days. Cells were collected in water, treated with an equal amount of ether, and vortexed for 10 min to kill unsporulated cells. After the cells were washed in water, they were resuspended in 900 ml of sterile water and treated with 100 ml of Zymolase (10 mg/ml) to remove the ascus. Cell mixtures were vortexed for 5 min to increase spore dispersion. In both replicates, haploid cells derived from both F₁ hybrids (NA × WA and SA × WE) were mixed in equal amounts, vortexed for 5 min, plated onto complete YPDA media, and grown overnight. Full plates were replica plated onto minimal media (MIN) to select for diploid F₂ hybrids containing genetic contributions from all four founders followed by a replica plating on YPDA. This procedure was repeated 11 times to create the F₁₂ population, which we term SGRP-4X. Finally, F₁₂ diploid hybrids were sporulated for 10 days in KAc at 23° and tetrads were dissected as previously described (Naumov *et al.* 1994). Viable spores with correct 2:2 segregations for the *MAT* locus and *ura3* and *lys2* auxotrophies were selected. We picked a total of 192 segregants (some from the same tetrad) from replicate 1 and stored them in glycerol stocks at −80°. We estimate that the pool goes through 10–15 mitotic generations during the high cell density replating cycles (YPDA/MIN/YPDA), between each sexual generation. This gives a point estimate of 150 cell divisions (12.5 generations × 12 intercross cycles) for SGRP-4X individuals starting from the founder strains, with a range from 120 to 180.

Sequencing and read mapping

The four founder strains, 192 isolated segregants, and large F₁₂ segregant pools (see below) were sequenced using the Illumina HiSeq and Illumina GAII platforms, with 2 × 108-bp paired end libraries prepared according to standard protocol (Kozarewa *et al.* 2009; Quail *et al.* 2012). We used the Sanger Centre sequencing pipelines for base calling and alignment. BWA version 0.5.8c (r1536) (Li and Durbin 2009) was used to map reads to the *S. cerevisiae* reference genome with options “aln -q 15 -t 2”, and we further filtered mappings to have quality scores of at least 30. Parental genome assembly and SNP calling will be reported in a separate article

(A. Bergström, J. T. Simpson, F. Salinas, L. Parts, A. Zia, A. N. Nguyen Ba, A. M. Moses, E. J. Louis, V. Mustonen, J. Warringer, R. Durbin, and G. Liti, unpublished results) and can be obtained from the web resource <http://www.moseslab.csb.utoronto.ca/sgrp/>.

Calling segregant genotypes

We called genotypes at each segregating site for each segregant, using a simple four-state hidden Markov model (HMM). The hidden state of the model for sample s and locus l is a 1-of-4 random variable $x_{s,l} = (x_{s,l,1}, \dots, x_{s,l,4})$ that emits base i corresponding to parent p with probability $e_{s,l,i} \wedge (1 - x_{s,l,p})(1 - e_{s,l,i}) \wedge x_{s,l,p}$, given the error probability of the base call $e_{s,l,i}$ calculated from the base quality. The transition probabilities are derived from approximate recombination rates assuming uniformly distributed events. We used $p(x_{s,l} = x_{s,l+1}) = \exp(-r d_{l,l+1})$, where r is the per-base recombination rate fixed at 1.2×10^{-5} , and $d_{l,l+1}$ is the distance between the two consecutive loci. We used standard methods to fit the HMM. In practice, as individual base calls were confident and the sequencing was of sufficiently high coverage, this approach made little difference compared to using maximum-likelihood estimate calls for private alleles and joining consecutive calls into haplotype blocks.

Phenotyping segregants and reciprocal hemizygous strains

The individual sequenced segregants and reciprocal hemizygous strains were subjected to precise growth phenotyping under three conditions (40° heat, 1.5 mM arsenite, and 100 µg/ml paraquat) in two technical replicates, using high-resolution microcultivation instruments (Bioscreen C; Oy Growth Curves, Raisio, Finland) for quantitative growth as previously described (Liti *et al.* 2009a; Cubillos *et al.* 2011; Warringer *et al.* 2011). Briefly, growth curves were dissected into three fitness variables, growth rate, growth efficiency, and growth lag, which were extracted using an automated procedure (Warringer and Blomberg 2003). Growth rate was calculated as population doubling time (hours) from the slope of the exponential phase, growth lag (hours) as the time until the start of detectable exponential phase, and growth efficiency (optical density units) as the optical density of cultures in stationary phase. The phenotypic variance for each F₁ cross and for F₁₂ outbred crosses was estimated as the sum of squares of deviations of single measurements from the mean. The number of transgressive segregants was calculated as previously described (Marullo *et al.* 2006), considering the fittest and weakest founders as phenotypic value boundaries.

Linkage mapping

We discarded segregants with evidence for diploid karyotype. For each phenotyped trait, we calculated the mean across technical replicates of the three fitness variables calculated as described above. We separately fitted the effect of each parental allele at every segregating site for each fitness trait

in a standard linear model. We used the genotype at the mating-type locus and auxotrophic markers *LYS2* and *URA3* as covariates, as we found these to have a detectable effect on the growth profiles. We considered only sites where genotypes were called for at least 100 segregants and each parental allele was observed at least four times. The linkage LOD scores (base 10) were used to call peaks with strongest signal of at least LOD 4 and their support regions within 3 LOD units of it. We repeated this procedure 1000 times with permuted genotypes, keeping the correlation structure in the phenotypes due to the covariates. The nominal P -values for the peaks were calculated as the average number of peaks detected in permutations with the same or a stronger maximum linkage signal. We retained peaks with empirical false discovery rate (FDR) < 0.5 and also calculated their q -values as the expected fraction of false positive calls when using the peak P -value as cutoff. While the cutoff of 0.5 is lenient, it has good reproducibility properties, and we have validated several of these moderate signals.

Pleiotropy calculation

We calculated linkage LOD scores as described above and performed permutations in the same way, recording for each peak the strongest linkage LOD score for the other traits of any parental allele for any site in a 40-kb window centered on the peak. We calculated the nominal P -value of a pleiotropic effect as the average number of peaks per permutation with a LOD score of at least 4 and a secondary LOD score corresponding to another phenotype stronger than the observed peak's best secondary LOD score in the same window. We called a pleiotropic signal if the nominal P -value was < 0.1, which corresponds to an FDR < 30% (9 of 25 pleiotropic QTL, 2.5 expected).

Reproducibility calculation

We took the regions previously mapped for the same growth traits in F₁ segregants (Cubillos *et al.* 2011) and calculated the strongest LOD score for the corresponding trait in any parental background in a 100-kb window centered on the locus. We calculated the nominal P -value as the fraction of all 100-kb windows in the genome that had an equal or stronger signal and calculated the P -values and their q -values in a standard way.

Pooled selection

Pools of millions of segregants were subjected to selection for high-temperature growth and nonselective media growth in duplicates as previously described (Parts *et al.* 2011). Briefly, two replicate pools of 10–100 million cells were collected from sporulation media for haploid and diploid F₁₂ populations and treated with ether and zymolase. Spores were plated on YPDA and incubated at 40° until full growth was obtained. Each plate was incubated for 48 hr and then resuspended in distilled water. Ten percent of the cells were used for the next replating and the rest for DNA extraction. In total, 10 time points (T₀, ..., T₁₀, corresponding to plates

0–10) for both control and heat-stress conditions were sampled, from which T0, T4, T8, and T10 were sequenced to an average genome coverage of 210× (Supporting Information, Table S6). We have previously argued that contributions of genetic drift and *de novo* mutations are negligible for detectable allele frequency changes in these experiments, due to the limited number of generations during the selection process (Parts *et al.* 2011; Illingworth *et al.* 2012).

Estimating allele frequencies

As a first pass, we estimated posterior allele frequencies for each parent separately. We considered all sites private to parent p . For each locus l , we inferred the posterior distribution of allele frequency $f_{c,r,t,l,p} \sim \text{Beta}(a_{c,r,t,l,p}, b_{c,r,t,l,p})$ for experimental condition c , replicate r , and time-point t following Parts *et al.* (2011). In short, $\text{prob}(f_{c,r,t,l,p} | D) \sim \prod_{l'} f_{c,r,t,l,p}^{x_{c,r,t,l,p}} (1 - f_{c,r,t,l,p})^{x_{c,r,t,l,p} - 1} r_{l,l'}$, where $x_{c,r,t,l,p}$ and $x_{c,r,t,l,\sim p}$ are the number of alleles of parent p and alleles not of parent p observed in condition c , replica r , time t , and locus l , and $r_{l,l'}$ is the recombination rate between loci l and l' . Loci l' were chosen to be at least 100 bp apart to ensure the sampled alleles come from independent reads. $r_{l,l'}$ was set to 0 if it was < 0.9 , so that only strongly linked sites were used in the allele frequency calculation. After the initial pass, we filtered out sites for which the posterior allele frequency mean $a_{c,r,t,l,p} / (a_{c,r,t,l,p} + b_{c,r,t,l,p})$ was at least 0.2 away from the empirical mean $x_{c,r,t,l,p} / (x_{c,r,t,l,p} + x_{c,r,t,l,\sim p})$ at the site, as in the examined cases these outliers were due to alignment issues. We then also included sites with a 2:2 segregation between parents and recalculated the posterior allele frequency estimates from the first round of posterior estimates with an expectation-maximization-like approach. To compare allele frequencies in the sequenced segregants to those of sequenced pool, we split the genome into nonoverlapping windows of 500 segregating sites, calculated Pearson's r for each parental allele frequency in each window separately, and used the median of all the calculated r 's as a summary statistic.

Locating QTL in pools

To identify selected regions, we estimated allele frequencies for each time point as outlined above, quantified their changes between sampled time points, compared the changes during selection to changes during a control experiment (pool grown under optimal conditions at 23°), and combined the comparisons across time points and replicates. For replica r of selection condition c , and every measured time point $t > 0$ (T4, T8, and T10), we calculated the posterior distribution of the difference in allele frequency $d_{c,r,t,l,p}$ at each locus l and parent p . We approximated the allele frequency posteriors with a normal distribution with mean $m_{c,r,t,l,p}$ and variance $v_{c,r,t,l,p}$, which gives a normal distribution for the difference distribution, $d_{c,r,t,l,p} \sim N(m_{c,r,t,l,p} - m_{c,r,t-1,l,p}, v_{c,r,t,l,p} + v_{c,r,t-1,l,p})$. We then calculated P -values of increasing and decreasing allele frequency for each time point at each locus and any ploidy, by numerically estimating $P_{c,r,t,l,p+} = \text{prob}(d_{c,r,t,l,p} > 0)$ and $P_{c,r,t,l,p-} = \text{prob}(d_{c,r,t,l,p} < 0)$. We combined these P -values

across time points, using Fisher's method, calculating $P_{c,r,l,p,s} = \chi^2(-2 \log(\text{prod}_t(P_{c,r,t,l,p,s})); 2^*T_r)$, where T_r is the number of time points sequenced for replicate r , and s is one of “−” for decrease and “+” for increase in allele frequency. We then calculated q -values from the P -values by considering the null distribution of all control experiment ($c = 0$) P -values $\{P_{0,r,l,p,s}\}_{r,l,p}$ calculated as above, according to the method of Storey and Tibshirani (2003). We expect allele frequencies not to be subject to selection during the control experiment, and thus any changes should be attributed to drift or other unknown confounding factors. We called QTL as regions where the minimum q -value across directions and parents was < 0.05 , combining consecutive sites with minimum q -values below the threshold into a single QTL. Ties (e.g., in cases of stretch of q -values = 0) were ordered by $\min_{r,t,p,s} P_{c,r,t,l,p,s}$ to prioritize loci.

We considered QTL to be shared between the previously described intercrossed F₁₂ WA × NA population (Parts *et al.* 2011) and the SGRP-4X when the distance between mapped intervals was < 10 kb.

QTL model selection

To compare goodness-of-fit of biallelic (1 vs. 3, 2 vs. 2) and more complicated (1 vs. 1 vs. 2, 1 vs. 1 vs. 1 vs. 1) linkage models, we used the Akaike information criterion, $2k - 2 \log \text{prob}(D|x)$. The log-likelihood was calculated from the assumed underlying normal distribution in the standard way, and k is the number of free parameters in the model. We calculated the Akaike information criterion (AIC) for all possible allele configurations and picked the one with the smallest score as the QTL model.

For similar comparison of the selection QTL models, we calculated the AIC as above, with a different likelihood model. We posited the existence of one, two, three, or four different driver alleles. For each model and replicate, we estimated a separate fitness parameter for each driver allele from its frequency changes between day 0 and day 8. We then calculated the expected frequencies of the other alleles under this model and calculated the probability of observing each allele frequency from a normal distribution $N(\text{expected change; observed change}, s_p^2)$, where s_p^2 is the variance in allele frequencies estimated from changes during the control experiment c separately for each allele, $s_p^2 = \text{var}_{l,r}(m_{c,r,1,l,p} - m_{c,r,0,l,p})$. We picked the model with the smallest average AIC across the two replicates, if it was substantially better supported (AIC difference at least 1.0) than the null model of no change.

Results and Discussion

Genetic background of founder strains

We selected four founder strains of distinct geographic and ecological origins (Figure 1A) that are segregating at 64% of the *S. cerevisiae* SNPs previously described (Liti *et al.* 2009a). We used the DBVPG6765 strain as representative of the WE lineage, YPS128 of the NA lineage, Y12 of the SA

lineage, and DBVPG6044 of the WA one. These strains have been extensively characterized at the genomic (Liti *et al.* 2009a) and phenotypic (Warringer *et al.* 2011) levels and successfully used for QTL mapping in two-parent crosses (Cubillos *et al.* 2011; Parts *et al.* 2011; Salinas *et al.* 2012). To further improve the reference genomes, we resequenced the founders at high coverage and generated high-quality assemblies, containing >95% of the sequence for each strain in one large scaffold per chromosome (A. Bergström, J. T. Simpson, F. Salinas, L. Parts, A. Zia, A. N. Nguyen Ba, A. M. Moses, E. J. Louis, V. Mustonen, J. Warringer, R. Durbin, and G. Liti, unpublished results). With the exception of the subtelomeric regions, we found no large structural variants that could prevent meiotic recombination and cause significant allelic distortions, making these strains highly suitable for genetic crosses.

We identified a total of 109,701 SNPs in the parents, most (86%) of which are private to a single lineage (Figure 1B). In comparison, a commonly used BY $\times$ RM cross has less than half the number ($\sim 47,000$) of segregating sites (Nagarajan *et al.* 2010). The majority, 68,727 SNPs, map within nondubious ORFs (average of 11.8 SNPs per gene) and nearly one-third, 21,467 SNPs, alter the protein sequence (average of 3.7 changes per protein). Of the 5793 nondubious ORFs in the genome, 5414 (93%) have at least one SNP and 4542 (78%) have at least one amino acid difference between the parents. Further, 16% of all nonsynonymous changes are predicted to be deleterious by assessing the conservation of the residue in homologous proteins with the SIFT score (Kumar *et al.* 2009; A. Bergström, J. T. Simpson, F. Salinas, L. Parts, A. Zia, A. N. Nguyen Ba, A. M. Moses, E. J. Louis, V. Mustonen, J. Warringer, R. Durbin, and G. Liti, unpublished results). Interestingly, derived alleles (as determined from comparisons with other *Saccharomyces* species) private to a single lineage are predicted to be deleterious more frequently than those present in multiple lineages (18% vs. 9%) (Figure 1B and A. Bergström, J. T. Simpson, F. Salinas, L. Parts, A. Zia, A. N. Nguyen Ba, A. M. Moses, E. J. Louis, V. Mustonen, J. Warringer, R. Durbin, and G. Liti, unpublished results).

Spatial statistics of genetic variation are important determinants of the mapping population quality. Nearly all previous yeast QTL mapping studies have utilized laboratory strains with mosaic genomes of variable local ancestry (Liti *et al.* 2009a). In contrast, genetic distances between our founder strains estimated in 10-kb windows are narrowly centered on the genome average, and thus the polymorphic sites are evenly distributed across the genome (Figure 1C and Figure S1). Therefore, we avoid problems arising from admixture that otherwise affect the nature and spatial distribution of identified QTL and interactions between coevolved alleles.

Genomic landscape of the mapping population

We generated the SGRP-4X mapping population using a funnel design similar to the mouse collaborative cross (Churchill *et al.* 2004) (Figure 1D and Figure S2). Initially,

we crossed haploid segregants from two F_1 crosses containing complementary auxotrophies and selected for diploids on minimal media (Materials and Methods, Figure S2). We then applied 11 rounds of random intercrosses as previously described (Parts *et al.* 2011). In addition to establishing the >10 million-segregant population for pooled studies, we isolated 192 haploid individuals, sequenced them to a median of $11\times$ coverage, and called genotypes (Materials and Methods). Thirteen segregants and 11 individual chromosomes were heterozygous for a large fraction of polymorphic sites, suggesting contamination or aneuploidy (Figure S3), and were filtered from further analyses.

The majority of the genome had all four parental backgrounds segregating in the pool, with median allele frequencies of 0.26 (WE), 0.21 (WA), 0.26 (NA), and 0.26 (SA) and parental allele frequencies between 0.1 and 0.5 for 89% of the sites (Figure 2A). The lower allele frequency of the WA genome is consistent with slightly slower growth of the WA parent on minimal medium and less efficient sporulation, resulting in negative selection against the loci harboring responsible WA alleles. There were 9 regions of at least 40 kb with more extreme allele frequencies (<10% or >40%, Table S1A) and 54 (<15% and >35%) after the intercrosses (Table S1B, Figure S4), some of which are analyzed further below. Allele frequencies estimated by deep sequencing total DNA from the entire pool were strongly correlated with estimates from segregants (median Pearson's r in 500 site blocks >0.75, Materials and Methods), indicating that the isolated individuals are representative of the population.

De novo SNP calling in the 11.2-Mb mappable genome of the 179 haploid individuals yielded 100 SNPs. This corresponds to a mutation rate of 3.4×10^{-10} per site per cell division (assuming 150 generations during the intercross process, Materials and Methods), very close to the commonly cited rate of 3.3×10^{-10} (Lynch *et al.* 2008). There were five *de novo* SNPs in the 591 kb of the genome subject to selection during the intercross (5.2 expected, Table S1A), and none of them were predicted to be deleterious by SIFT. Thus, we have no evidence supporting their contribution to formation of allele frequencies in SGRP-4X. A full description of these variants will be reported in a separate article (A. Bergström, J. T. Simpson, F. Salinas, L. Parts, A. Zia, A. N. Nguyen Ba, A. M. Moses, E. J. Louis, V. Mustonen, J. Warringer, R. Durbin, and G. Liti, unpublished results).

Whole-genome sequencing of the segregants allows accurate estimation of the genetic map. The outcrossing divided each genome into haplotype blocks identical by descent to one of the parental strains (Figure 2B), with an average segregant having 374 such blocks of median length 23 kb (Figure S5A). Using a recently developed inference method that leverages linkage disequilibrium between genotyped sites (Illingworth *et al.* 2013), we estimated an average genome-wide per-base per-generation recombination rate of 3.2×10^{-6} , giving a total number of recombination events approximately fourfold higher than that observed for F_1

Figure 1 Genetic diversity and breeding of the founder strains. (A) The four founder strains sample a large fraction of the genetic variation in the species. The phylogenetic tree includes all the strains sequenced in the SGRP project (Liti *et al.* 2009a), with major geographic clusters highlighted. (B) Distribution and summary statistics of single-nucleotide variants between the founders. The Venn diagram shows the sharing of derived alleles between parents, after rooting each variant using *S. paradoxus* as the outgroup. An additional 20,405 SNPs (19% of the total) are not included as they could not be rooted. Numbers in sectors give the count and fraction of nonsynonymous polymorphisms and fraction of nonsynonymous polymorphisms predicted to have a deleterious effect on protein function from SIFT analysis (Ng and Henikoff 2003). (C) Chromosome II sequence similarity between founders (logarithmic scale, full-genome plots given in Figure S1). The pairwise similarity between founder strain genome sequences was computed in windows of size 10 kb as $N/(d + 1)$, where N is the number of sites in the window with high-quality alignments and d is the number of single-nucleotide differences between the strains within these sites. Values corresponding to different levels of sequence divergence are indicated on the vertical axis. Strain colors are the same as in A. The bottom two similarity plots show the same statistic for S288c/RM11 and S288c/YJM789 crosses previously used in yeast QTL mapping, with an uneven distribution of sequence variation. (D) Cross design of SGRP-4X (see also Figure S2).

segregants (Mancera *et al.* 2008). Surprisingly, the recombination rate in SGRP-4X is 80% higher than estimated in the F_{12} WA $\times$ NA cross (1.7×10^{-6} , Figure S5B). We observed recombination hot and cold spots in concordance with previous reports, with centromeres exhibiting a lower than average recombination rate (Figure 2C), and a set of short haplotype blocks likely corresponding to noncrossovers (Figure S5A). A full description of the inference and analysis of recombination rates is given elsewhere (Illingworth *et al.* 2013).

Overall, parental contributions to the SGRP-4X population are close to the 25% expected under selectively neutral random outcrossing. This compares favorably with smaller multiparent populations in other model organisms (Kover *et al.* 2009; Philip *et al.* 2011; King *et al.* 2012), some of which have an unequal representation of founder strains due to selection and drift during breeding. While a handful of loci were selected in the SGRP-4X during intercross (Table S1), loss of genetic complexity by drift was negligible due to the extremely large population size (>10 million).

Figure 2 Genetic and phenotypic diversity of SGRP-4X. (A) Distribution of the fraction of the genome contributed by each founder to the sequenced segregants. (B) Fine mosaic structure of SGRP-4X individuals. Chromosome IIs from 10 individual segregants (y-axis) are painted according to the parental background at each locus, using the colors from Figure 1A. (C) Inferred scaled linkage disequilibrium measure D' for chromosome II, evaluated in 5-kb windows in the WA $\times$ NA cross, SGRP-4X, and the S288c $\times$ YJM789 cross from Mancera *et al.* (2008). Loci with minor allele private to one founder were used for the SGRP-4X inference. Blue crosses reflect the absence of measurement in regions that did not contain markers. (D) Distribution of normalized growth rate of individual isolated segregants under nonstress, heat, arsenite, and paraquat conditions. Average values for founder strains from multiple replicas are indicated.

maintained. The even contribution from founders coupled with the very high number of recombination events makes SGRP-4X well suited for QTL mapping.

Phenotypic diversity of SGRP-4X

Outcrossing a multifounder population over many generations breaks linkage disequilibrium and increases haplotype diversity. As a result, allele structures that have coevolved in

parental lineages over millions of years are disrupted or rearranged in new combinations, which can alter the distribution of phenotypes. To assess whether traits were affected by the genetic churning of the intercross, we quantified the mitotic growth properties (rate, lag, and efficiency) of the retained 179 haploid segregants in basal conditions (non-stress) and exposure to three stresses: a toxic natural variant of arsenic [arsenite, As(III)], an oxidative agent generating

superoxide radicals (paraquat), and high temperature (heat, 40°) (*Materials and Methods*, Table S2).

Overall, segregants were remarkably fit in nonstress conditions, with almost all of them (170/179) exceeding the growth rate of the notoriously slowly proliferating reference strain BY4741 (Figure 2D). Thus, the breakup of coevolved alleles had little overall effect on fitness, in contrast to severe growth defects detected in crosses between highly diverged lineages of the sister species *S. paradoxus* (Liti *et al.* 2009b). We observed a generally poor performance of the segregants in paraquat, to the extent that 35 F₁₂ recombinants were unable to proliferate in the concentration previously used for F₁ progeny (Cubillos *et al.* 2011). We hypothesize that this reflects the presence of epistatic interactions that have evolved in the parental populations for pathways underlying the oxidative stress response. These interactions are broken up in the multiparent cross due to the more rare cosegregation of compatible alleles and the reduction in physical linkage, which will target linked QTL.

Many crossing rounds can reduce phenotypic complexity in the population due to stabilizing and directional selection for particular trait values or outbreeding depression. The distributions of measured growth traits were unimodal and lacked skew in the nonstress condition, which would have been expected if substantial directional selection had been acting during the intercross (Figure 2D). To assess the evolution of trait complexity across generations, we compared the SGRP-4X phenotype distributions to those measured in the pairwise F₁ crosses between the four parents (Cubillos *et al.* 2011). The SGRP-4X segregants were 1.6 and 2.7 times more variable in the absence of stress and in high-temperature environments, whereas they were only half as variable in the presence of As(III) (Figure S6). These trends were also reflected in the degree of transgression, *i.e.*, the frequency of segregants with more extreme trait values than those of the parents (Figure S7), with more variable traits showing higher transgression. The markedly negative skew of paraquat growth phenotypes in SGRP-4X was evident in the extreme levels of negative transgression (47 transgressive segregants in paraquat vs. 3 and 1 in heat and arsenite, respectively), even under substantially milder paraquat exposure (Figure S7).

Taken together, the data from individually phenotyped segregants show that increasing genetic diversity by including multiple parents and performing many rounds of intercross did not substantially reduce fitness, had mostly limited impact on trait complexity, and imposed no strong directional selection during the process.

QTL mapping by linkage analysis

Availability of whole-genome genotype and phenotype data for the 179 segregants allowed us to assess the ability to detect QTL for the growth traits described above. While the segregants were sequenced primarily with the goal of assessing the recombination landscape (Illingworth *et al.* 2013) and we were powered only to detect large effects, it

is informative to assess the QTL mapping resolution and reproducibility properties when using individual segregants. We identified 6, 16, and 3 significant QTL for mitotic growth properties under exposure to heat, arsenite, and paraquat, respectively (FDR < 0.5, *Materials and Methods*, Table S3 and Figure S8A), with a median interval size of 27.8 kb. Five of the 7 previously detected QTL with LOD > 9 from the six pairwise F₁ crosses (Cubillos *et al.* 2011) were replicated in SGRP-4X at a FDR = 0.17 (*Materials and Methods*, Table S4A). This number increased to 16 of 24 QTL with a more lenient cutoff of LOD > 3 for F₁ crosses (Cubillos *et al.* 2011) and FDR = 0.5 for this study (Table S4B). The rest of the previously mapped QTL could be undetected due to the complexity of the genetic interactions, overestimates of the true effect size in a smaller sample (Beavis effect), or false positives in the previous screen.

In multiple cases, the QTL regions included known causal or strong candidate genes. One replicated QTL at chromosome IV (1505–1524 kb) included the polymorphic ARR gene cluster present in the SA background (Cubillos *et al.* 2011), which is known to be essential for arsenite tolerance (Bobrowicz *et al.* 1997). The arsenite growth rate QTL at chromosome XV (932–957 kb) contains the candidate gene *VMA4*, a subunit of the vacuolar H⁺-ATPase known to be involved in As(III) tolerance in the reference strain (Zhou *et al.* 2009), with the peak of strongest linkage only 340 bp away (Figure 3A).

We next asked whether some QTL were present in more than one condition. We defined a QTL as pleiotropic if the strongest linkage to growth in another environment within a 20-kb window was significant (*Materials and Methods*) and identified 22 trait-specific QTL and 3 pleiotropic QTL (Figure 3B, Table S5, and Figure S8B). For example, the arsenite QTL at chromosome XI 68–82 kb also had a strong linkage signal for heat growth efficiency at chromosome XI 48 kb. This region contains the gene *PEX1*, which is an AAA-peroxin whose deletion allele in the reference background has a fitness effect under both heat stress (Sinha *et al.* 2008) and arsenite (Pan *et al.* 2010).

A segregant population allows detection of antagonistic alleles that increase fitness in spite of coming from an unfit strain or vice versa (Rieseberg *et al.* 2003). For example, although the NA strain has superior resistance to heat, we found NA alleles that are detrimental for this phenotype (Table S3). Similarly, five of the QTL mapped in the arsenite-resistant SA and WA strains were antagonistic and contributed negatively to mitotic growth in the presence of arsenite. Such alleles may have arisen in yeast due to antagonistic pleiotropy (Orr 1998; Qian *et al.* 2012) or linkage to selected alleles (Liti and Louis 2012). Once released from their original genetic context and placed in an environment with a single stressor, they are free to exhibit their positive effect on growth.

The 12 rounds of intercrosses break linkage between nearby sites and improve QTL mapping resolution. Whereas we would expect a 3-LOD support interval of 64 kb in the

Figure 3 Linkage analysis in 179 SGRP-4X segregants. (A) Strength of linkage to arsenite resistance for a QTL region mapped on chromosome XV. For each parental allele, $-\log_{10}(P)$ values of linkage (y-axis) are given across a chromosome XV region (x-axis). Colors for alleles are as in the bottom panel and Figure 1. Bottom panel shows growth rate in arsenite (y-axis) for segregants with different genotypes at the site of strongest linkage at the QTL locus (x-axis, jitter added). Average values for each parent are given under the corresponding segregants. (B) Number and pleiotropy of linkage QTL. The number of genome-wide significant QTL found using linkage mapping for the three conditions is displayed on the x-axis, with colors indicating pleiotropic QTL significant in another condition at a lower threshold (*Materials and Methods*).

96-segregant F_1 populations (Cubillos *et al.* 2011) if recombination is uniform, it narrows down to 6 kb in the 179 F_{12} segregants from SGRP-4X. In practice, we observed longer support intervals (median 27.8 kb), as the heterogeneity in recombination rate results in the majority of the genome having stronger than average linkage, and some regions likely contain linked QTL (Liti and Louis 2012) that we merged into a single peak. While with 179 sequenced segregants we did not have enough power to map the abundant weaker-effect alleles that make up a large fraction of the narrow-sense heritability (Bloom *et al.* 2013), we did recover the strong QTL from our earlier work. Overall, we mapped 25 narrow QTL regions that replicated previous results and contained strong candidates for follow-up, demonstrating the utility of SGRP-4X for complex trait mapping.

Allele frequency dynamics in SGRP-4X under heat selection

QTL can be efficiently mapped by bulk genotyping selected populations (Brauer *et al.* 2006; Segre *et al.* 2006; Ehrenreich *et al.* 2010; Wenger *et al.* 2010), which is most powerful when selection operates for a prolonged period (Parts *et al.* 2011). To apply this approach in SGRP-4X, we used the 40° heat exposure environment, where two of our founder strains (NA and SA) are fit, while the other two are moderately unfit (WE) or strongly unfit (WA) (Figure 2D, *Materials and Methods*). In addition, growth under high temperature is a classical complex trait, for which many genes are known to be involved, and a large number of causal loci have been

characterized (Steinmetz *et al.* 2002; Sinha *et al.* 2006, 2008; Cubillos *et al.* 2011; Parts *et al.* 2011).

We identified 34 and 39 regions with significant allele frequency changes in haploid and diploid pools, respectively, and designated them as QTL (FDR 1%, *Materials and Methods*, Table S7, Figure 4A, and Figure S9). Nineteen of these overlap between ploidies, suggesting that heat-resistance QTL operate largely independently of ploidy, indicating dominant or additive effects and giving additional evidence for the reproducibility of our calls. The mapped regions have a median size of 4.8 kb, a resolution comparable to that expected under linkage mapping and finer than that of bulk segregant analyses in F_1 crosses (Ehrenreich *et al.* 2012).

The number of QTL found is a substantial increase compared to the 21 mapped using the same methodology in a F_{12} WA $\times$ NA cross (Parts *et al.* 2011). Seven and 10 of these previously identified loci were detected in the haploid and diploid outbred populations at 1% FDR, respectively (Figure 4B), which increased to 13 and 17 at a more lenient 5% FDR (Table S7). A further 3 were within 30 kb of a QTL, but outside its mapped region, suggesting either linked QTL or low resolution in one of the two studies. For three loci (*IRA1*, *IRA2*, and the subtelomeric region of chromosome XIII), we previously observed fixation of the beneficial allele upon selection (Parts *et al.* 2011; Illingworth *et al.* 2012). In contrast, while the allele frequencies in these regions changed by >15% in the current study, no allele was fixed or completely removed from the SGRP-4X pool even after long-term selection (Figure 4A).

Figure 4 QTL mapping from allele frequency changes upon heat selection. (A) Allele frequencies at *IRA2* region during heat selection. Top four panels represent allele frequencies (y-axis) of each parental allele in the pool at different time points [day 0 (T0), black; day 8 (T4) haploid, green; day 16 (T8) diploid, red; day 20 (T10) diploid, blue] around the *IRA2* locus (x-axis). The bottom panel represents genes around the interval and combined signal for allele frequency change (y-axis). Evidence for increase [$-\log_{10}(P)$] is given in the top part, while evidence for decrease is in the bottom (*Materials and Methods*). (B) Overlap of QTL mapped using selection. Venn diagram depicts QTL found previously in the NA $\times$ WA cross (purple) and in this study using haploid (gray) and diploid (orange) pools. (C) Decay of linkage in a region under selection for two cross designs. Thick blue lines show the LD' profile (y-axis) up- and downstream from the driver location of chromosome XV (*IRA2*, x-axis) for the WA $\times$ NA cross (left) and SGRP-4X (right). The fraction of observed WA alleles at each site is plotted for each segregating site with a black circle for pool after control experiment (WA $\times$ NA cross, replica 2, T4; SGRP-4X, replica 2, T4), and a red circle for pool after heat selection (WA $\times$ NA cross, replica 2, T6; SGRP-4X, replica 2, T10). Only the sites where WA is the derived allele are plotted for SGRP-4X. The LD' profiles are computed from the segregants' genotype data (Illingworth *et al.* 2013) with no knowledge of the selection data. The LD' curves demonstrate complete linkage at the driver locus and gradual decline as a function of the distance to the driver, which is faster in SGRP-4X and visually matches the difference in size and shape of the selective sweeps upon heat selection.

Both linkage and selection mapping approaches benefit from additional recombination events to give narrower QTL intervals. We found that the allele frequency changes in the *IRA2* region selected under heat stress were markedly different between the previous WA $\times$ NA cross and SGRP-4X (Figure 4C). The higher recombination rate in SGRP-4X resulted in a QTL centered more precisely on the validated *IRA2* gene. Also, a large upstream region was no longer selected for in SGRP-4X, which could be explained either by a linked upstream QTL or by differences in the local recombination rate between the two crosses. The large decrease in linkage disequilibrium (LD) in the region (Figure 4C) strongly supports the latter explanation, showing how

recombination reshapes the landscape of allele frequency changes under selection.

Surprisingly, all but one of the alleles increasing in frequency (7/8 and 6/6 in haploids and diploids) were from the only intermediate heat-resistant WE background, thus showing abundant advantageous variants in one parent. We found no positively selected alleles from the NA strain, which is most heat resistant. All the detrimental alleles were from heat-intermediate and heat-sensitive strains, with 17 and 25 from WE and 14 and 10 from WA in haploids and diploids, respectively (Table S7). The abundance of detrimental variants supports the recently found excess of loss-of-function alleles in the unfit WA strain (Warringer *et al.* 2011; Zorgo

et al. 2012), likely due to adaptation to a very specific niche that does not include a strong positive influence of high temperature. Overall, the large number of regions selected in the SGRP-4X population under high temperature deepens the characterization of the highly complex heat-resistance phenotype, with different wild strains harboring alleles that modulate growth and survival.

Complexity of genotype–phenotype relationship in the SGRP-4X

A key goal of quantitative genetic studies is to estimate the phenotypic contribution of different alleles at each QTL (King *et al.* 2012). In the SGRP-4X and other multiparent populations (Churchill *et al.* 2004; Kover *et al.* 2009; King *et al.* 2012), it is possible to distinguish between biallelic QTL, where there are only two distinct effect sizes that segregate either in 1:3 or in 2:2 ratio (Figure 5A), and multiallelic ones, where more than two alleles have an independent fitness effect (segregating 1:1:2 or 1:1:1:1).

Comparison of linkage models with varying numbers of free parameters (*Material and Methods*) revealed that most linkage QTL (5/6 for heat, 2/3 for paraquat, and 10/16 for arsenite) are best explained by the effect of a single allele (Table S8 and Figure 5A). The more complex 1:1:2 patterns, where backgrounds of two founders have an independent beneficial or deleterious effect, explained the remaining QTL, while the 2:2 segregation pattern, with the causal allele shared between two founders, was never observed. These results suggest that most strong genetic contributors to the phenotype in SGRP-4X are alleles private to a single founder. Abundant rare coding variants specific to individual human populations recently reported (Abecasis *et al.* 2012; Fu *et al.* 2012; Keinan and Clark 2012; Tennessen *et al.* 2012) could similarly underlie a substantial proportion of heritable trait variation.

Next, we used statistical model selection in QTL regions of the intercross (Table S1) and heat stress (Table S7) to infer which alleles have independent fitness effects. For example, we previously detected a chromosome V region strongly selected during consecutive rounds of crossing of the WA and NA strains (Parts *et al.* 2011; Illingworth *et al.* 2012), but we were unable to distinguish positive selection for one parent from negative selection against the other. Allele frequencies in SGRP-4X indicated at least three different fitness values for this locus, with strong selection for the NA parent, two neutral alleles (SA and WE), and negative selection against the WA background (Figure 5B). In general, we found almost all regions strongly selected during the crossing to have more than one driving allele (7/9 regions, Table S8B). For heat selection after the crossing, we could confidently identify 29 and 34 biallelic QTL and 5 and 5 multiallelic QTL in haploids and diploids, respectively (Table S8, C and D). For instance, we observed strong selection for the NA version of *IRA1* on chromosome II, weaker selection for the WE one, and negative selection against the SA and WA alleles.

Understanding the effects of multiple variants at a single locus has so far been limited by mapping precision and sensitivity, experimental variance in phenotyping, and low throughput of candidate validation (Mackay *et al.* 2009; Trontin *et al.* 2011). We have overcome some of these constraints in yeast by sensitively identifying allelic contributions in a pool under selection (Parts *et al.* 2011). Here, we observed a predominance of single alleles underlying QTL, consistent with a large fraction of heritable phenotypic variation being due to alleles private to individual founders. However, more complex allelic series for QTL in selection pools were also evident. Our results are consistent with previous findings of excess biallelic QTL compared to other allele distributions in F₁ crosses (Cubillos *et al.* 2011; Ehrenreich *et al.* 2012), although this could be affected by the choice of founder strains, crossing design, and mapping approach.

Narrow mapping intervals allow rapid causal gene identification

QTL mapping in SGRP-4X resulted in narrow regions ripe for following up with single-gene studies. First, we used reciprocal hemizygosity (Steinmetz *et al.* 2002) to test the effects of different *IRA1* and *IRA2* alleles on growth under heat stress. The *IRA* genes are key regulators of the RAS signaling pathway, whose defects result in hyperactive RAS. This in turn leads to high levels of cyclic AMP (cAMP) and high PKA activity that inhibits the heat-stress response. We have previously shown that WA alleles of *IRA1* and *IRA2* have a reduced fitness at high temperature (Parts *et al.* 2011) and mapped these loci again in SGRP-4X, where the frequency of *IRA2*^{WA} decreased from 0.47 to 0.32 under selection, while *IRA2*^{NA} and *IRA2*^{SA} frequencies increased by 6% and 27%, respectively.

Quantitative growth curves and plate spot dilutions of reciprocal hemizygous strains support the model where *IRA2*^{WA} is heat sensitive, while *IRA2*^{NA} and *IRA2*^{WE} have nearly identical and superior fitness (Figure 6A and Figure S10). Surprisingly, although *IRA2*^{SA} showed the greatest frequency increase, we did not find differences with any other allele in the reciprocal hemizygosity assay (Figure S10), even when compared to the weak *IRA2*^{WA}. In contrast, at the *IRA1* locus, allele frequency changes suggest that *IRA1*^{SA} and *IRA1*^{WA} are both deleterious (Table S8). We replicated this result by reciprocal hemizygosity in the WA × SA cross, where the *IRA1*^{SA} allele did not outperform the known deleterious *IRA1*^{WA} version (Figure 6B). These results suggest that in the SA background heat resistance might not be mediated through the cAMP/PKA pathway and instead be triggered by alternative mechanisms and complex genetic interactions.

Finally, we validated the pleiotropic effect of candidate gene *VPS53* in heat and arsenite in the WE × SA and WE × WA hybrids. While the *VPS53* region LOD score was below the genome-wide significance cutoff in QTL mapping, it had strong support in the early phenotyping data, which reduced to moderate linkage support in both conditions after observing

Figure 5 Allelic heterogeneity of QTL. (A) Number of QTL for different segregation models. The bi-allelic (1:3 or 2:2) and multi-allelic (1:1:2 or 1:1:1:1) models are defined based on phenotypic effects. Colored circles depict the four alleles distributed in all possible fitness-segregating scenarios, ">" and "<" indicate greater or lower fitness. Numbers refer to QTL fitting each model for linkage analysis, intercross, and heat selection in haploid and diploid populations. (B) Allele frequencies in three regions (chromosomes VIII, XII, and V) after intercross rounds, but before selection experiments. Chromosome VIII region was inferred to have a 1:3, chromosome XII region a 2:2, and chromosome V region a complex 1:2:1 allele segregation. Bottom panel of each plot highlights individual genes around the interval, with potential causal candidates in green.

all the replicates. Hybrids containing the WE allele had a substantial fitness advantage under both stresses, while no differences were observed for any of the other alleles (Figure 6C and Figure S11A). Interestingly, the effect manifested primarily on the lag time in arsenite, a compound mainly acting via a lag time increase, whereas it manifested principally as an efficiency effect when encountering heat stress (Figure S11B). Thus, the mapped QTL is strain dependent (Cubillos *et al.* 2011) with *VPS53*^{WE} being the fit-

test allele with a fully penetrant effect regardless of the cross combination.

The narrow mapping intervals in SGRP-4X allowed us to rapidly follow up promising candidate genes and validate the complexity of fitness contributions of a single locus. Full understanding of these complex architectures requires substantial follow-up work involving repetition of the selection experiments with fixation of different alleles to distinguish between interacting partners and independent fitness. The

Figure 6 Phenotypes of strains with reciprocally hemizygous genotypes for causal genes. (A) Lower fitness of the *IRA2^{WA}* allele. Shown is the average of each fitness component measurement ($n = 2$) for all combinations of reciprocal hemizygote pairs of *IRA2* alleles. Individual alleles were deleted from the heterozygous diploid strain and assessed for their individual contributions to high-temperature growth. (B) Fitness averages for the *IRA1* reciprocal hemizygotes in the WA $\times$ SA cross. (C) Mitotic growth curves for arsenite and high-temperature growth in reciprocal hemizygotes for the *VPS53* gene derived from the WE $\times$ SA and WA $\times$ SA crosses.

ability to quantify the effects of individual alleles and the narrow mapping intervals help in designing these studies and in constructing informative allele combinations.

Prospects of mapping complex traits in model organisms

The goal of using model organisms and controlled crosses for quantitative genetics research is to closely mimic either idealized population genetic models or natural populations in terms of genetic and phenotypic variation. The SGRP-4X population presented here has nearly equal contributions from all founders, uniform spatial distribution of segregating sites, little effect from drift on allele frequencies, and low linkage, making it a close model for an idealized multiparent advanced intercross. Any bias in the initial allele frequencies due to selection and associated hitchhiking of linked alleles during the intercross will not affect the usability of the population for mapping traits during pooled selection experiments so long as the initial frequencies remain substantially >0 . Having a control experiment and several time points enabled us to fully factor in the initial allele frequencies, and QTL are called only if during the selection phase there is a significant change against that baseline.

Even more genetic and trait diversity can be incorporated into the mapping population by crossing more distant parents. For example, several extremely divergent, but not reproductively isolated, strains have recently been recovered in China (Wang *et al.* 2012). To obtain model populations closer to that of humans, other natural isolates and mosaic

lab strains could be crossed in bulk to produce a population with variable allele frequencies. However, as the mapping approaches used here rely on at least moderate frequencies of causal alleles for sensitive detection, they have to be adapted to scenarios with rarer alleles.

In general, model populations remain fruitful for the study of isolated individual genetic mechanisms, be it rare variants, epistasis, or common small-effect alleles, and ideally, specialized controlled populations would be constructed for each. SGRP-4X is useful in a variety of settings, as we have shown for the study of weaker alleles and of multiple contributors at a locus and following up allelic series and interactions between loci. With rapid advances in highly multiplexed whole-genome sequencing, an even larger number of new individuals derived from the SGRP4-X could be genotyped in the future to approach nearly full power to map trait loci (Bloom *et al.* 2013; Ludlow *et al.* 2013; Wilkening *et al.* 2013).

We have harnessed the awesome power of baker's yeast for quantitative genetics studies. Full-genome sequences containing a large fraction of natural variation in the species, high-resolution phenotypes, an extremely large number of individuals, and low linkage disequilibrium have produced a very powerful mapping population. This resource can be further utilized for linkage or selection mapping of more phenotypes, creating the outbred diploid lines for extremely large-scale mapping, crossed to various deletion, overexpression, and tag collections for understanding phenotypes underlying the complex trait alleles and modified by

standard tools for validating and following up small-effect alleles.

Acknowledgments

We thank all the members of the Sanger Institute sequencing production teams for generating the sequence data. We thank Alex Mott and Agnes Llored for technical help. This work was funded by grants from Atip-Avenir, Association pour la Recherche Contre le Cancer (SFI20111203947) (to G.L.), and The Wellcome Trust (grant WT077192/Z/05/Z) (to R.D.). We further acknowledge the Wellcome Trust for support under grant 098051. This work was also supported in part by Region Midi Pyrénées (France) under grant 09005247 and was carried out in the frame of the European Cooperation in Science and Technology Action (FA0907-Bioflavour) under the European Union's Seventh Framework Programme for Research (FP7). F.A.C. is supported by Conicyt-Programa de Atracción e Inserción/Concurso Nacional de Apoyo al retorno de investigadores/as desde el extranjero grant 82130010. L.P. is supported by a fellowship from the Canadian Institute for Advanced Research and a Marie Curie International Outgoing Fellowship.

Literature Cited

- Abecasis, G. R., A. Auton, L. D. Brooks, M. A. DePristo, R. M. Durbin *et al.*, 2012 An integrated map of genetic variation from 1,092 human genomes. *Nature* 491: 56–65.
- Bloom, J. S., I. M. Ehrenreich, W. T. Loo, T. L. Lite, and L. Kruglyak, 2013 Finding the sources of missing heritability in a yeast cross. *Nature* 494: 234–237.
- Bobrowicz, P., R. Wysocki, G. Owsianik, A. Goffeau, and S. Ulaszewski, 1997 Isolation of three contiguous genes, ACR1, ACR2 and ACR3, involved in resistance to arsenic compounds in the yeast *Saccharomyces cerevisiae*. *Yeast* 13: 819–828.
- Brauer, M. J., C. M. Christianson, D. A. Pai, and M. J. Dunham, 2006 Mapping novel traits by array-assisted bulk segregant analysis in *Saccharomyces cerevisiae*. *Genetics* 173: 1813–1816.
- Brem, R. B., G. Yvert, R. Clinton, and L. Kruglyak, 2002 Genetic dissection of transcriptional regulation in budding yeast. *Science* 296: 752–755.
- Churchill, G. A., D. C. Airey, H. Allayee, J. M. Angel, A. D. Attie *et al.*, 2004 The Collaborative Cross, a community resource for the genetic analysis of complex traits. *Nat. Genet.* 36: 1133–1137.
- Cubillos, F. A., E. Billi, E. Zorgo, L. Parts, P. Fargier *et al.*, 2011 Assessing the complex architecture of polygenic traits in diverged yeast populations. *Mol. Ecol.* 20: 1401–1413.
- Donnelly, P., 2008 Progress and challenges in genome-wide association studies in humans. *Nature* 456: 728–731.
- Durrant, C., H. Tayem, B. Yalcin, J. Cleak, L. Goodstadt *et al.*, 2011 Collaborative Cross mice and their power to map host susceptibility to *Aspergillus fumigatus* infection. *Genome Res.* 21: 1239–1248.
- Ehrenreich, I. M., N. Torabi, Y. Jia, J. Kent, S. Martis *et al.*, 2010 Dissection of genetically complex traits with extremely large pools of yeast segregants. *Nature* 464: 1039–1042.
- Ehrenreich, I. M., J. Bloom, N. Torabi, X. Wang, Y. Jia *et al.*, 2012 Genetic architecture of highly complex chemical resistance traits across four yeast strains. *PLoS Genet.* 8: e1002570.
- Flint, J., and T. F. Mackay, 2009 Genetic architecture of quantitative traits in mice, flies, and humans. *Genome Res.* 19: 723–733.
- Fu, W., T. D. O'Connor, G. Jun, H. M. Kang, G. Abecasis *et al.*, 2013 Analysis of 6,515 exomes reveals the recent origin of most human protein-coding variants. *Nature* 493: 216–220.
- Gan, X., O. Stegle, J. Behr, J. G. Steffen, P. Drewe *et al.*, 2011 Multiple reference genomes and transcriptomes for *Arabidopsis thaliana*. *Nature* 477: 419–423.
- Huang, X., M. J. Paulo, M. Boer, S. Effgen, P. Keizer *et al.*, 2011 Analysis of natural allelic variation in *Arabidopsis* using a multiparent recombinant inbred line population. *Proc. Natl. Acad. Sci. USA* 108: 4488–4493.
- Huang, W., S. Richards, M. A. Carbone, D. Zhu, R. R. Anholt *et al.*, 2012 Epistasis dominates the genetic architecture of *Drosophila* quantitative traits. *Proc. Natl. Acad. Sci. USA* 109: 15553–15559.
- Huxley, C., E. D. Green, and I. Dunham, 1990 Rapid assessment of *S. cerevisiae* mating type by PCR. *Trends Genet.* 6: 236.
- Illingworth, C. J., L. Parts, S. Schiffels, G. Liti, and V. Mustonen, 2012 Quantifying selection acting on a complex trait using allele frequency time series data. *Mol. Biol. Evol.* 29: 1187–1197.
- Illingworth, C. J., L. Parts, A. Bergstrom, G. Liti, and V. Mustonen, 2013 Inferring genome-wide recombination landscapes from advanced intercross lines: application to yeast crosses. *PLoS ONE* 8: e62266.
- Keinan, A., and A. G. Clark, 2012 Recent explosive human population growth has resulted in an excess of rare genetic variants. *Science* 336: 740–743.
- King, E. G., C. M. Merkes, C. L. McNeil, S. R. Hoofer, S. Sen *et al.*, 2012 Genetic dissection of a model complex trait using the *Drosophila* Synthetic Population Resource. *Genome Res.* 22: 1558–1566.
- Kover, P. X., W. Valdar, J. Trakalo, N. Scarcelli, I. M. Ehrenreich *et al.*, 2009 A multiparent advanced generation inter-cross to fine-map quantitative traits in *Arabidopsis thaliana*. *PLoS Genet.* 5: e1000551.
- Kozarewa, I., Z. Ning, M. A. Quail, M. J. Sanders, M. Berriman *et al.*, 2009 Amplification-free Illumina sequencing-library preparation facilitates improved mapping and assembly of (G+C)-biased genomes. *Nat. Methods* 6: 291–295.
- Kumar, P., S. Henikoff, and P. C. Ng, 2009 Predicting the effects of coding non-synonymous variants on protein function using the SIFT algorithm. *Nat. Protoc.* 4: 1073–1081.
- Li, H., and R. Durbin, 2009 Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics* 25: 1754–1760.
- Liti, G., and E. J. Louis, 2012 Advances in quantitative trait analysis in yeast. *PLoS Genet.* 8: e1002912.
- Liti, G., D. M. Carter, A. M. Moses, J. Warringer, L. Parts *et al.*, 2009a Population genomics of domestic and wild yeasts. *Nature* 458: 337–341.
- Liti, G., S. Haricharan, F. A. Cubillos, A. L. Tierney, S. Sharp *et al.*, 2009b Segregating YKU80 and TLC1 alleles underlying natural variation in telomere properties in wild yeast. *PLoS Genet.* 5: e1000659.
- Ludlow, C. L., A. C. Scott, G. A. Cromie, E. W. Jeffery, A. Sirr *et al.*, 2013 High-throughput tetrad analysis. *Nat. Methods* 10: 671–675.
- Lynch, M., W. Sung, K. Morris, N. Coffey, C. R. Landry *et al.*, 2008 A genome-wide view of the spectrum of spontaneous mutations in yeast. *Proc. Natl. Acad. Sci. USA* 105: 9272–9277.
- Mackay, T. F., E. A. Stone, and J. F. Ayroles, 2009 The genetics of quantitative traits: challenges and prospects. *Nat. Rev. Genet.* 10: 565–577.
- Mancera, E., R. Bourgon, A. Brozzi, W. Huber, and L. M. Steinmetz, 2008 High-resolution mapping of meiotic crossovers and non-crossovers in yeast. *Nature* 454: 479–485.

- Manolio, T. A., F. S. Collins, N. J. Cox, D. B. Goldstein, L. A. Hindorf *et al.*, 2009 Finding the missing heritability of complex diseases. *Nature* 461: 747–753.
- Marullo, P., M. Bely, I. Masneuf-Pomarede, M. Pons, M. Aigle *et al.*, 2006 Breeding strategies for combining fermentative qualities and reducing off-flavor production in a wine yeast model. *FEMS Yeast Res.* 6: 268–279.
- Nagarajan, M., J. B. Veyrieras, M. de Dieuleveult, H. Bottin, S. Fehrmann *et al.*, 2010 Natural single-nucleosome epi-poly-morphisms in yeast. *PLoS Genet.* 6: e1000913.
- Naumov, G. I., T. A. Nikonenko, and V. I. Kondrat'eva, 1994 Taxonomic identification of *Saccharomyces* from yeast genetic stock centers of the University of California. *Genetika* 30: 45–48.
- Ng, P. C., and S. Henikoff, 2003 SIFT: predicting amino acid changes that affect protein function. *Nucleic Acids Res.* 31: 3812–3814.
- Nordborg, M., and D. Weigel, 2008 Next-generation genetics in plants. *Nature* 456: 720–723.
- Orr, H. A., 1998 Testing natural selection vs. genetic drift in phenotypic evolution using quantitative trait locus data. *Genetics* 149: 2099–2104.
- Pan, X., S. Reissman, N. R. Douglas, Z. Huang, D. S. Yuan *et al.*, 2010 Trivalent arsenic inhibits the functions of chaperonin complex. *Genetics* 186: 725–734.
- Parts, L., F. A. Cubillos, J. Warringer, K. Jain, F. Salinas *et al.*, 2011 Revealing the genetic structure of a trait by sequencing a population under selection. *Genome Res.* 21: 1131–1138.
- Perlstein, E. O., D. M. Ruderfer, D. C. Roberts, S. L. Schreiber, and L. Kruglyak, 2007 Genetic basis of individual differences in the response to small-molecule drugs in yeast. *Nat. Genet.* 39: 496–502.
- Philip, V. M., G. Sokoloff, C. L. Ackert-Bicknell, M. Striz, L. Branstetter *et al.*, 2011 Genetic analysis in the Collaborative Cross breeding population. *Genome Res.* 21: 1223–1238.
- Qian, W., D. Ma, C. Xiao, Z. Wang, and J. Zhang, 2012 The genomic landscape and evolutionary resolution of antagonistic pleiotropy in yeast. *Cell Rep.* 29: 1399–1410.
- Quail, M. A., T. D. Otto, Y. Gu, S. R. Harris, T. F. Skelly *et al.*, 2012 Optimal enzymes for amplifying sequencing libraries. *Nat. Methods* 9: 10–11.
- Rieseberg, L. H., A. Widmer, A. M. Arntz, and J. M. Burke, 2003 The genetic architecture necessary for transgressive segregation is common in both natural and domesticated populations. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* 358: 1141–1147.
- Salinas, F., F. A. Cubillos, D. Soto, V. Garcia, A. Bergstrom *et al.*, 2012 The genetic basis of natural variation in oenological traits in *Saccharomyces cerevisiae*. *PLoS ONE* 7: e49640.
- Segre, A. V., A. W. Murray, and J. Y. Leu, 2006 High-resolution mutation mapping reveals parallel experimental evolution in yeast. *PLoS Biol.* 4: e256.
- Simon, M., O. Loudet, S. Durand, A. Berard, D. Brunel *et al.*, 2008 Quantitative trait loci mapping in five new large recombinant inbred line populations of *Arabidopsis thaliana* genotyped with consensus single-nucleotide polymorphism markers. *Genetics* 178: 2253–2264.
- Sinha, H., B. P. Nicholson, L. M. Steinmetz, and J. H. McCusker, 2006 Complex genetic interactions in a quantitative trait locus. *PLoS Genet.* 2: e13.
- Sinha, H., L. David, R. C. Pascon, S. Clauser-Munster, S. Krishnakumar *et al.*, 2008 Sequential elimination of major-effect contributors identifies additional quantitative trait loci conditioning high-temperature growth in yeast. *Genetics* 180: 1661–1670.
- Steinmetz, L. M., H. Sinha, D. R. Richards, J. I. Spiegelman, P. J. Oefner *et al.*, 2002 Dissecting the architecture of a quantitative trait locus in yeast. *Nature* 416: 326–330.
- Storey, J. D., and R. Tibshirani, 2003 Statistical significance for genomewide studies. *Proc. Natl. Acad. Sci. USA* 100: 9440–9445.
- Tennessen, J. A., A. W. Bigham, T. D. O'Connor, W. Fu, E. E. Kenny *et al.*, 2012 Evolution and functional impact of rare coding variation from deep sequencing of human exomes. *Science* 337: 64–69.
- Trontin, C., S. Tisne, L. Bach, and O. Loudet, 2011 What does *Arabidopsis* natural variation teach us (and does not teach us) about adaptation in plants? *Curr. Opin. Plant Biol.* 14: 225–231.
- Wang, Q. M., W. Q. Liu, G. Liti, S. A. Wang, and F. Y. Bai, 2012 Surprisingly diverged populations of *Saccharomyces cerevisiae* in natural environments remote from human activity. *Mol. Ecol.* 21: 5404–5417.
- Warringer, J., and A. Blomberg, 2003 Automated screening in environmental arrays allows analysis of quantitative phenotypic profiles in *Saccharomyces cerevisiae*. *Yeast* 20: 53–67.
- Warringer, J., E. Zorgo, F. A. Cubillos, A. Zia, A. Gjuvsland *et al.*, 2011 Trait variation in yeast is defined by population history. *PLoS Genet.* 7: e1002111.
- Wenger, J. W., K. Schwartz, and G. Sherlock, 2010 Bulk segregant analysis by high-throughput sequencing reveals a novel xylose utilization gene from *Saccharomyces cerevisiae*. *PLoS Genet.* 6: e1000942.
- Wilkening, S., M. M. Tekkedil, G. Lin, E. S. Fritsch, W. Wei *et al.*, 2013 Genotyping 1000 yeast strains by next-generation sequencing. *BMC Genomics* 14: 90.
- Zhou, X., A. Arita, T. P. Ellen, X. Liu, J. Bai *et al.*, 2009 A genome-wide screen in *Saccharomyces cerevisiae* reveals pathways affected by arsenic toxicity. *Genomics* 94: 294–307.
- Zorgo, E., A. Gjuvsland, F. A. Cubillos, E. J. Louis, G. Liti *et al.*, 2012 Life history shapes trait heredity by accumulation of loss-of-function alleles in yeast. *Mol. Biol. Evol.* 29: 1781–1789.

Communicating editor: M. Johnston

GENETICS

Supporting Information

<http://www.genetics.org/lookup/suppl/doi:10.1534/genetics.113.155515/-/DC1>

High-Resolution Mapping of Complex Traits with a Four-Parent Advanced Intercross Yeast Population

Francisco A. Cubillos, Leopold Parts, Francisco Salinas, Anders Bergström,
Eugenio Scovacricchi, Amin Zia, Christopher J. R. Illingworth, Ville Mustonen, Sebastian Ibstedt,
Jonas Warringer, Edward J. Louis, Richard Durbin, and Gianni Liti

Figure S1 Sequence similarity along founder strain chromosomes (same as Figure 1D).

Haploid parental strains:
all strains are *hoΔ*, *ura3Δ*.
The *lys2Δ* were deleted
using URA3 (URA+)

Figure S2 Cross design of the SGRP-4X mapping population.

Figure S3 Chromosome I aneuploidy. (A) Major variant frequency along chromosomes I, II and III for one example segregant. The frequency of each base is counted at all polymorphic sites and the frequency of the most common one is plotted. (B) The depth of coverage of mapped reads for the same region and segregant. Coverage is averaged in non-overlapping windows of 10kb. An extra copy of a chromosome is manifested in a variant frequency pattern where parts of the chromosome have major variant frequencies close to 1 (homozygous stretches) while other parts have major variant frequencies fluctuating around 0.5 (heterozygous stretches), and a depth of coverage across the chromosome twice as high as that of the other chromosomes.

Figure S4 Genome-wide parental allele frequencies in the F12 SGRP-4X population. Allele frequencies for NA, WA, SA and WE parental strains are shown for each chromosome. Estimates were obtained for both replicas (black and light-blue lines) in haploid T4 mock samples. Dashed cut-off denotes regions with at least one parental allele frequency below 15% or above 35%.

Figure S5 Genome-wide recombination landscape (A) Recombination block size distribution across all segregants and entire genome (B) Linkage decay in the WA x NA cross, SGRP-4X, and the S288c x YJM789 cross from (MANCERA *et al.* 2008). SGRP-4X (blue) has less linkage disequilibrium left compared to the two parent cross (red) after 12 rounds of crossing. Both advanced intercrosses have less linkage disequilibrium left than the F1 cross of (MANCERA *et al.* 2008) shown in black. We computed the mean decay of scaled linkage disequilibrium D' across the genome for these crossing experiments. In order to also have the SGRP-4X curve starting from one (full linkage) only segregating site pairs where the minor allele at both loci stems from a single parental strain were considered for this cross.

Figure S6 (A) Phenotype distributions of SGRP-4X compared to two parent F1 crosses from the same founder strains reported in (CUBILLOS *et al.* 2011). (B) Phenotypic variance of SGRP-4X and two parent F1 crosses. Set 1 and set 2 indicate segregants with *ura3* and *lys2* genotypes, respectively.

Figure S7 Phenotypic landscape of SGRP-4X segregants for quantitative growth at different environments. X-axis depicts segregants ranked based on their relative growth value to the BY reference strain (y-axis). Arrows indicate the relative fitness of parental strains. Set 1 and set 2 indicate segregants with *ura3* and *lys2* genotypes, respectively.

Figure S8 Number of QTLs mapped by linkage analysis upon q-value cutoff. (A) Number of QTLs map for Heat, Arsenite and Paraquat resistance at different q-value cutoffs. (B) Number of pleiotropic QTLs depending on the q-value cutoff. The total number of mapped QTLs for each condition (0.5 cutoff from part A) is given with a dashed line.

Figure S9 Allele frequency change histogram for each parental strain on high temperature growth (heat) and non-selective media growth (control). The number of sites with the respective change from the x-axis is given in log scale on the y-axis; vast majority of the sites have changes smaller than 0.1.

A.

B.

Figure S10 Reciprocal hemizygosity and heat growth assay for different hybrid combinations. A. Mitotic growth curves. B. Serial dilutions.

Figure S11 Reciprocal hemizygosity for *VPS53* for high temperature growth and arsenite stress (3mM). A. Mitotic growth curves. B. Serial dilutions.

Table S1 Genomic regions selected during the intercross process. Allele frequencies were estimated by deep sequencing replica 1 (R1) and replica 2 (R2) pools. A. Regions with at least one parental allele frequency below 10% or above 40%. B. Regions with at least one parental allele frequency below 15% or above 35%.

A.

Genomic coordinates				Allele frequencies							
#Chr.	Start	End	Peak	R1-NA	R2-NA	R1-WA	R2-WA	R1-WE	R2-WE	R1-SA	R2-SA
IV	644963	736332	699005	0.43	0.39	0.03	0.04	0.31	0.23	0.24	0.32
V	69083	220165	172739	0.46	0.50	0.02	0.02	0.29	0.29	0.24	0.18
VI	47103	94543	70206	0.31	0.28	0.31	0.34	0.02	0.02	0.39	0.34
VIII	69085	128898	112561	0.14	0.16	0.15	0.14	0.04	0.04	0.69	0.71
XI	59875	126027	85720	0.09	0.08	0.06	0.04	0.70	0.73	0.19	0.11
XII	425940	490580	444427	0.04	0.04	0.09	0.09	0.42	0.44	0.45	0.43
XV	140619	193609	171462	0.28	0.28	0.53	0.53	0.02	0.02	0.19	0.17
XV	368035	434143	385888	0.37	0.40	0.16	0.15	0.04	0.04	0.40	0.37
XVI	170646	217074	205083	0.14	0.19	0.08	0.08	0.20	0.20	0.60	0.54

B.

Genomic coordinates				Allele frequencies							
#Chr.	Start	End	Peak	R1-NA	R2-NA	R1-WA	R2-WA	R1-WE	R2-WE	R1-SA	R2-SA
I	2323	71024	18367	0.08	0.26	0.31	0.29	0.2	0.07	0.28	0.22
I	71617	114885	60137	0.08	0.19	0.31	0.38	0.2	0.17	0.28	0.24
II	416370	587125	473193	0.21	0.24	0.18	0.18	0.5	0.49	0.12	0.11
III	51494	125974	120975	0.34	0.1	0.12	0.24	0.26	0.42	0.28	0.25
III	128486	216348	141113	0.25	0.1	0.16	0.24	0.21	0.42	0.37	0.25
III	262261	306853	306675	0.29	0.28	0.43	0.42	0.29	0.32	0.03	0.04
IV	195099	291961	220090	0.28	0.41	0.18	0.17	0.28	0.33	0.42	0.25
IV	292644	294427	265125	0.28	0.41	0.14	0.17	0.3	0.33	0.29	0.25
IV	352902	857195	836508	0.12	0.39	0.08	0.04	0.22	0.23	0.62	0.32
IV	859705	1521409	970414	0.12	0.09	0.08	0.1	0.22	0.26	0.62	0.65
V	7343	348693	278855	0.53	0.54	0.13	0.17	0.17	0.14	0.06	0.23
VI	27376	197390	49322	0.31	0.09	0.31	0.24	0.02	0.21	0.39	0.5
VII	60719	67418	66950	0.42	0.49	0.19	0.21	0.17	0.08	0.19	0.22
VII	83612	371940	145166	0.41	0.49	0.24	0.21	0.12	0.08	0.89	0.22
VII	372929	419737	289279	0.41	0.41	0.24	0.18	0.12	0.16	0.89	0.26
VII	431486	443950	382164	0.41	0.24	0.24	0.14	0.12	0.22	0.89	0.46
VII	444540	645680	583313	0.27	0.24	0.08	0.14	0.3	0.22	0.35	0.46
VII	646700	651438	625336	0.27	0.37	0.08	0.17	0.3	0.21	0.35	0.3
VII	653843	690509	670173	0.23	0.37	0.17	0.17	0.23	0.21	0.36	0.3
VII	690544	700282	712171	0.23	0.47	0.17	0.21	0.23	0.22	0.36	0.18
VII	746379	1001775	856542	0.22	0.47	0.18	0.21	0.19	0.22	0.45	0.18
VIII	23372	216237	112561	0.14	0.16	0.15	0.14	0.04	0.04	0.69	0.71
VIII	218122	383850	202868	0.07	0.16	0.18	0.14	0.46	0.04	0.24	0.71
IX	176252	220342	197884	0.21	0.2	0.16	0.19	0.22	0.17	0.42	0.42
IX	265507	334500	299349	0.41	0.45	0.16	0.13	0.28	0.28	0.12	0.12
IX	334500	353165	352596	0.46	0.45	0.08	0.13	0.3	0.28	0.07	0.12
IX	353527	431743	392101	0.46	0.33	0.08	0.09	0.3	0.27	0.07	0.21

X	384305	556857	425491	0.11	0.05	0.13	0.16	0.24	0.22	0.62	0.57
X	556857	649653	503092	0.12	0.05	0.2	0.16	0.31	0.22	0.38	0.57
XI	33220	264819	85720	0.09	0.08	0.06	0.04	0.7	0.73	0.19	0.11
XI	338910	418601	361437	0.42	0.41	0.27	0.29	0.19	0.15	0.32	0.32
XI	572381	663018	614768	0.21	0.25	0.35	0.42	0.08	0.1	0.27	0.28
XII	71640	275970	148591	0.49	0.54	0.16	0.14	0.17	0.21	0.19	0.11
XII	326930	522948	445103	0.04	0.05	0.08	0.1	0.41	0.44	0.49	0.46
XII	563210	624957	593223	0.38	0.35	0.19	0.23	0.34	0.33	0.06	0.03
XII	628515	636692	625505	0.39	0.35	0.15	0.23	0.24	0.33	0.2	0.03
XII	661866	672780	668651	0.39	0.3	0.15	0.14	0.24	0.33	0.2	0.22
XII	675396	705031	710591	0.28	0.3	0.25	0.14	0.38	0.33	0.11	0.22
XII	713182	772090	748082	0.28	0.27	0.25	0.26	0.38	0.31	0.11	0.13
XII	898244	1059052	948127	0.34	0.28	0.32	0.43	0.21	0.26	0.96	0.13
XIII	130360	202531	156687	0.28	0.26	0.3	0.31	0.15	0.12	0.29	0.24
XIII	202531	207796	237397	0.28	0.5	0.3	0.2	0.15	0.19	0.29	0.08
XIII	224729	388307	288300	0.47	0.5	0.26	0.2	0.21	0.19	0.13	0.08
XIII	492832	688003	538185	0.18	0.19	0.25	0.23	0.44	0.45	0.15	0.18
XIII	690563	812561	641567	0.38	0.19	0.2	0.23	0.36	0.45	0.09	0.18
XIII	871305	917558	904123	0.41	0.4	0.18	0.19	0.17	0.17	0.24	0.3
XIV	248283	342275	294659	0.17	0.11	0.18	0.17	0.21	0.22	0.45	0.49
XIV	569372	711472	624062	0.09	0.11	0.36	0.35	0.32	0.29	0.23	0.2
XV	44064	688426	262977	0.04	0.27	0.16	0.54	0.12	0.05	0.58	0.2
XV	692396	763132	448710	0.42	0.27	0.2	0.54	0.21	0.05	0.16	0.2
XV	955003	1075608	1069074	0.89	0.72	0.16	0.16	0.16	0.15	0.33	0.34
XVI	115934	290163	205083	0.14	0.19	0.08	0.08	0.2	0.2	0.6	0.54
XVI	332645	455472	295445	0.14	0.19	0.17	0.08	0.21	0.2	0.49	0.54
XVI	741530	822498	761835	0.39	0.3	0.14	0.12	0.26	0.24	0.41	0.4

Table S2 The mitotic phenotypes growth lag, growth rate and growth efficiency for non-stress, heat, arsenite 1.5 mM, Paraquat 100 µg/mL and Paraquat 400 µg/mL on every segregant, are given for both technical replicates. BY4741 strain was used to normalize growth values across experiments (see methods).

Table S2 is available for download at <http://www.genetics.org/lookup/suppl/doi:10.1534/genetics.113.155515/-/DC1>.

Table S3 QTL summary detected from linkage analysis. Genomic position (peak and confidence interval), growth parameter, q-value, number of segregants, average phenotype per genotype and LOD values are indicated for each condition. QTLs mapped in more than one growth parameter are indicated with (*).

Heat	Coordinates		NA					WA			WE			SA		
Chromosome	Peak	Confidence Interval	Growth Parameter	q-value	# segr.	Average phenotype	log10p	# segr.	Average phenotype	log10p	# segr.	Average phenotype	log10p	# segr.	Average phenotype	log10p
II	99632	89962-116942	rate	0.49	34	0.01	0.2	37	0.14	1.4	55	-0.20	4.5	43	0.13	1.4
IV	669598	629721-700566	rate	0.49	58	-0.21	4.7	17	0.12	0.7	40	0.06	0.4	55	0.13	2.1
XI	66234	57355-68340	lag	0.04	13	-0.68	7.9	23	-0.03	0.1	99	0.07	2.3	37	0.03	0.1
XIII	895642*	875433-913695	rate	0.27	54	0.13	2	32	-0.34	5.7	34	0.12	1	50	-0.01	0.1
XIII	895642	863207-913722	lag	0.35	54	0.06	0.8	33	-0.37	5.2	35	0.11	0.7	50	0.08	0.6
XIV	665508	658446-676654	rate	0.49	31	-0.30	4.5	43	0.12	1.4	66	0.03	0.3	28	0.08	0.4

Paraquat	Coordinates		NA					WA			WE			SA		
Chromosome	Peak	Confidence Interval	Growth Parameter	q-value	# segr.	Average phenotype	log10p	# segr.	Average phenotype	log10p	# segr.	Average phenotype	log10p	# segr.	Average phenotype	log10p
XII	612566	606828-644083	lag	0.05	55	0.06	1.7	21	0.08	1.7	58	-0.14	5.6	28	0.03	0.4
III	212260	209644-218031	lag	0.17	44	-0.05	0.5	39	0.16	4.8	40	-0.02	0	52	-0.09	2.4
XIV	356878	349229-424774	lag	0.05	32	0.03	0.4	39	0.08	1.9	58	0.03	0.7	42	-0.14	5.6

Arsenite	Coordinates		NA					WA			WE			SA		
Chromosome	Peak	Confidence Interval	Growth Parameter	q-value	# segr.	Average phenotype	log10p	# segr.	Average phenotype	log10p	# segr.	Average phenotype	log10p	# segr.	Average phenotype	log10p
II	752324	740638-769695	rate	0.3	40	0.13	0.5	30	-0.26	5.7	40	0.16	0.8	49	0.16	1.1
IV	1247236	1231010-1262173	lag	0.42	50	0.3	2.7	23	0.35	1.5	37	-0.4	4.3	62	-0.04	0.5
IV	1516800	1505710-1524277	lag	0.42	44	-0.04	0.2	24	-0.55	4.6	38	0.14	0.6	54	0.22	1.9
V	102405	100406-124766	rate	0.3	65	0.17	1.4	9	-0.5	4.8	47	0.09	0.1	45	0.07	0.2
VI	23330	20600-27541	rate	0.3	49	0.11	0.2	34	0.2	1.4	14	-0.41	5	55	0.08	0.1
VII	7936	7378-8879	rate	0.3	18	-0.28	4.5	27	0.05	0.3	58	0.17	1.2	31	0.2	1.2
XI	68139	68139-82467	rate	0.3	14	0.23	0.9	13	-0.41	4.8	101	0.09	0.2	25	0.18	0.8
XII	981727	955642-996974	rate	0.3	33	0.07	0.1	50	0.12	0.3	36	0.18	0.9	6	-0.63	4.6
XIII	346821	326055-355502	rate	0.3	51	0.16	0.9	40	0.24	2	53	-0.12	4.8	21	0.09	0.2
XIII	861124	852242-865392	rate	0.3	47	-0.14	4.5	45	0.15	0.6	41	0.21	1.5	32	0.14	0.6
XIV	560654	553404-560824	rate	0.37	27	0.15	0.4	38	0.15	0.4	4	-0.67	4.2	45	0.09	0.1
XIV	644582	637362-665955	lag	0.42	30	0.20	0.9	53	0.08	0.3	53	-0.32	4.3	37	0.27	1.7
XV	173494	167483-207895	lag	0.07	44	0.13	0.5	80	0.05	0.1	4	-1.76	6.1	46	0.07	0.2
XV	944695	932741-957388	rate	0.42	52	-0.10	4.1	35	0.18	0.9	27	0.11	0.2	53	0.19	1.6

XVI	342504	310689- 408109	lag	0.42	34	-0.42	4.6	37	0.33	2.2	49	0.18	1.1	53	-0.02	0.3
XVI	72835	48951- 109062	rate	0.32	40	0.19	1.1	34	0.16	0.7	54	-0.12	4.4	38	0.18	0.8

Table S4 Linkage analysis in SGRP-4X and two-parent crosses.

A. Genomic regions mapped by linkage analysis in the SGRP-4X that were previously detected in two parent crosses using the same founder strains (Cubillos et al, 2011).

Phenotype	Chromosome	Peak	Growth trait	-log10p	Nominal p-value	Strain	LOD (Cubillos et al, 2011)
Arsenite	III	302000	rate	3.5	0.12	SA	8.98
Arsenite	IV	1506000	rate	4	0.12	WA	14.21
Arsenite	VII	54000	rate	4.5	0.12	NA	3.66
Arsenite	XVI	899000	rate	3.3	0.12	NA	7.16
Arsenite	VI	196000	rate	2.9	0.26	WA	3.41
Arsenite	III	302000	efficiency	2.4	0.13	SA	9.26
Arsenite	IV	1506000	efficiency	3	0.069	WA	9.79
Arsenite	XVI	899000	efficiency	2.8	0.069	NA	4.81
Arsenite	VI	65000	efficiency	2.0	0.259	SA	3.15
Arsenite	III	302000	lag	3.3	0	WA	9.41
Arsenite	IV	1506000	lag	4.6	0	WA	11.82
Heat	XIII	875000	rate	5.7	0	WA	11.67
Heat	XIII	849000	efficiency	2.6	0.224	SA	3.31
Heat	VIII	336000	efficiency	3.8	0	WA	3.38
Paraquat	IV	625000	efficiency	3.1	0	WA	3.01
Paraquat	XII	606000	rate	3.0	0.18	WE	9.44

B. Number of QTLs mapped in both studies depending on LOD and FDR thresholds. Expected number of QTLs and FDR threshold values are indicated in () for each **P-value cutoff**.

F1 maximum LOD score	Total F1 QTLs with LOD	Replicated F1 QTLs at the below indicated P-value cutoff		
		0.01	0.12	0.3
0	26	5 (0.3, 0.05)	11 (3.1, 0.29)	16 (7.8, 0.49)
5	12	3 (0.1, 0.04)	7 (1.5, 0.21)	9 (3.6, 0.40)
9	7	3 (0.1, 0.02)	5 (0.8, 0.17)	7 (2.1, 0.30)

Table S5 Phenotype and genomic position for QTLs mapped in more than one stress condition (pleiotropic QTLs). (*)
denotes significant parameter.

Phenotype 1	Growth trait	Chromosome	Peak	Phenotype 2	Growth trait	Peak	-log10p value (Strongest linkage)	-log10p value (within 20 kb)	q-value
Paraquat	lag	XIV	356878	Heat	Rate	336972	1.6	3.7*	0.4
Heat	rate	VI	45521	Paraquat	Rate	25547	3.9	5*	0.14
Arsenite	rate	XI	68139	Heat	Lag	48148	2.4	7.9*	0.02
Heat	lag	XI	66234	Arsenite	Rate	46371	1.2	4.8*	0.14
Paraquat	rate	VI	44835	Heat	rate	44835	3.6*	4.1	0.18

Table S6 Average sequencing coverage of analysed pool samples.

Trait	Ploidy	Replica	Timepoint	Average sequencing coverage
Control	Diploid	R2	T0	724.5
Control	Diploid	R1	T4	512.1
Control	Diploid	R2	T4	426.4
Control	Diploid	R1	T8	134.3
Control	Diploid	R2	T8	157.4
Control	Haploid	R1	T0	251.6
Control	Haploid	R1	T4	151.3
Control	Haploid	R2	T4	136.2
Heat	Diploid	R1	T8	313.1
Heat	Diploid	R2	T8	485.4
Heat	Diploid	R1	T10	104.5
Heat	Diploid	R2	T10	123.1
Heat	Haploid	R1	T4	252.2
Heat	Haploid	R2	T4	384.6
Heat	Haploid	R2	T10	101.5

Table S7 Regions selected for upon heat selection in F12 pools. (*) denotes QTLs previously mapped in the NA x WA heat selection (Parts et al, 2011).

Haploids

Trait	Chromosome	Peak Location (bp)	Region start (bp)	Region end (bp)	q value	Allele decrease	Allele increase
Haploid_Heat	I	7730	3208	7730	0.0183	WA	-
Haploid_Heat	I	41803	41803	48116	0	WA	WE
Haploid_Heat	II	192041	191473	194229	0.0039	WE	-
Haploid_Heat	II	246577	246577	246581	0.0255	WE	-
Haploid_Heat	II	323106	322797	323214	0.0169	WE	-
Haploid_Heat	II	387448	387448	387930	0	WE	-
Haploid_Heat	II	425572	423048	426882	0.0407	WE	-
Haploid_Heat	II	469449	465411	469724	0.0251	WA	-
Haploid_Heat*	II	520956	520956	539099	0	WA	-
Haploid_Heat	II	586848	585904	589502	0.0194	WE	-
Haploid_Heat	II	691644	690896	692190	0.0308	WE	-
Haploid_Heat	III	103983	103937	103983	0.0053	-	WE
Haploid_Heat	III	204170	194461	204170	0.0316	WA	-
Haploid_Heat	IV	88326	88326	94650	0	WA	-
Haploid_Heat	IV	167026	167026	167026	0.0042	WE	-
Haploid_Heat	IV	193568	193568	193596	0.0058	WE	-
Haploid_Heat	IV	309560	308968	311212	0.0391	WE	-
Haploid_Heat	IV	341596	340208	345209	0.0024	WE	-
Haploid_Heat	IV	414378	411541	414543	0.0417	WA	-
Haploid_Heat*	IV	442270	440553	446871	0.0336	WA	-
Haploid_Heat	IV	482650	479691	482994	0.0045	WA	-
Haploid_Heat	IV	594972	594972	614480	0	WE	-
Haploid_Heat	IV	779615	778412	779615	0.0382	WE	-
Haploid_Heat	IV	1165241	1165241	1202931	0	WE	SA
Haploid_Heat*	IV	1299590	1299590	1300574	0.0086	WE	-
Haploid_Heat	IV	1320892	1318531	1323778	0.0238	WE	-
Haploid_Heat	IV	1424223	1424223	1443661	0	WA	-
Haploid_Heat	V	210929	207967	214084	0.0023	WE	-
Haploid_Heat	V	258507	258488	258551	0.0217	WE	-
Haploid_Heat	V	485594	485594	488977	0.0374	WA	-
Haploid_Heat*	V	527121	524479	528927	0.0334	WA	-
Haploid_Heat	V	551439	551439	551656	0.0081	WA	WE
Haploid_Heat	VI	255151	254056	256286	0.0442	WE	-
Haploid_Heat	VII	182663	179739	185385	0.029	WA	-
Haploid_Heat	VII	486824	486599	487208	0.0365	-	WE
Haploid_Heat	VII	548507	548507	548783	0.0065	-	WE
Haploid_Heat	VII	634484	633588	637654	0.0185	WA	-

Haploid_Heat	VII	670084	669746	672925	0.0292	WA	-
Haploid_Heat	VIII	245817	240977	251365	0.0289	WA	-
Haploid_Heat	VIII	293911	290684	293911	0.0042	WA	WE
Haploid_Heat	VIII	379356	377959	380388	0.0276	WE	-
Haploid_Heat	VIII	424239	424239	456475	0	WE	-
Haploid_Heat	IX	74155	73727	76912	0.0295	WA	-
Haploid_Heat	IX	110880	110880	111569	0.045	WA	-
Haploid_Heat	X	144887	144540	145037	0.0314	WA	-
Haploid_Heat	X	159751	158895	161248	0.0319	WA	-
Haploid_Heat*	X	236389	236389	237201	0	WA	-
Haploid_Heat*	X	412413	412413	431667	0	WE	-
Haploid_Heat	X	678785	675814	681146	0.0336	WA	-
Haploid_Heat	XI	40024	40024	45406	0	WE	-
Haploid_Heat	XI	495717	494304	495717	0.0375	WA	-
Haploid_Heat	XI	514756	512598	515249	0.0184	WA	-
Haploid_Heat	XII	168350	168350	169953	0.0214	WE	-
Haploid_Heat	XII	209043	201880	211227	0.0023	WE	-
Haploid_Heat	XII	508720	503088	509048	0.0292	WA	-
Haploid_Heat	XII	646369	645661	650453	0.045	WA	-
Haploid_Heat	XII	1044364	1044364	1045357	0.0265	-	WE
Haploid_Heat	XIII	49254	48927	49730	0.0265	WE	-
Haploid_Heat	XIII	79258	78303	82388	0.0336	WA	-
Haploid_Heat	XIII	137716	135459	139699	0.0395	WA	-
Haploid_Heat	XIII	393791	388784	395295	0.0152	WA	-
Haploid_Heat	XIII	632758	631614	633136	0.0328	WA	-
Haploid_Heat	XIII	691786	687027	693409	0.0184	WA	-
Haploid_Heat	XIII	721067	721067	721562	0	WA	-
Haploid_Heat*	XIII	751054	747251	752520	0.0184	WA	-
Haploid_Heat	XIII	811132	809125	814002	0.0264	WA	-
Haploid_Heat	XIII	826593	825733	829030	0.0216	WE	-
Haploid_Heat*	XIII	877042	877042	913941	0	WA	-
Haploid_Heat	XIV	115349	114050	116117	0.003	WE	-
Haploid_Heat	XIV	173831	173831	180003	0	WE	-
Haploid_Heat	XIV	220441	220211	223244	0.044	WA	-
Haploid_Heat	XIV	273491	269608	274329	0.0095	-	WE
Haploid_Heat	XIV	290463	289635	291362	0.0351	-	WE
Haploid_Heat	XIV	356012	352810	356728	0.037	WA	-
Haploid_Heat	XIV	393352	393352	394078	0.0183	WA	-
Haploid_Heat*	XIV	481655	480850	482296	0.0027	WE	-
Haploid_Heat	XIV	639071	637452	639071	0.0465	WA	-
Haploid_Heat*	XV	174678	174678	176616	0	WA	-
Haploid_Heat	XV	345603	345603	370360	0	WA	-
Haploid_Heat	XV	739750	738846	742605	0.0451	WA	-
Haploid_Heat	XV	791982	787944	793038	0.018	WA	-

Haploid_Heat	XV	837629	837629	862997	0	WA	-
Haploid_Heat	XV	910563	901122	915550	0.018	WA	-
Haploid_Heat	XV	954035	954010	954307	0.0075	-	WE
Haploid_Heat*	XV	1034398	1029988	1037749	0.0146	WA	-
Haploid_Heat	XVI	68965	66703	70600	0.0201	WA	-
Haploid_Heat	XVI	145598	143934	145716	0.0296	WE	-
Haploid_Heat	XVI	264571	264571	264725	0	WA/WE	-
Haploid_Heat	XVI	798875	797108	800153	0.0384	WA	-
Haploid_Heat	XVI	815371	812986	818489	0.0184	WA	-
Haploid_Heat	XVI	860139	857632	860652	0.0492	WA	-

Diploids

Trait	Chromosome	Peak Location (bp)	Region start (bp)	Region end (bp)	q value	Allele decrease	Allele increase
Diploid_Heat	I	8001	8001	57156	0	WA	WE
Diploid_Heat	I	156349	154768	160003	0.0202	WA	-
Diploid_Heat	II	192393	189166	194229	0.0337	WE	-
Diploid_Heat	II	241424	233408	252194	0.0033	WE	-
Diploid_Heat	II	287551	286762	289132	0.006	WE	-
Diploid_Heat	II	384879	384879	389907	0	WE	-
Diploid_Heat*	II	510231	510231	544469	0	WA	-
Diploid_Heat	II	670475	668591	671539	0.0268	WA	-
Diploid_Heat	III	167332	166887	171203	0.0169	WA	-
Diploid_Heat	IV	21677	19588	24628	0.0406	WE	-
Diploid_Heat	IV	78729	78729	101420	0	WA	-
Diploid_Heat	IV	161585	161585	162746	0.0044	WE	-
Diploid_Heat	IV	301375	301375	305375	0.0098	WE	-
Diploid_Heat	IV	342650	341565	342797	0.0222	WE	-
Diploid_Heat	IV	414378	412379	415405	0.0232	WA	-
Diploid_Heat	IV	508188	505545	510196	0.0252	WE	-
Diploid_Heat	IV	595513	594399	598488	0.0026	WE	-
Diploid_Heat	IV	659686	658417	661178	0.04	WE	-
Diploid_Heat*	IV	1043658	1042465	1046575	0.0061	-	WE
Diploid_Heat	IV	1075808	1075286	1075853	0.0254	WE	-
Diploid_Heat	IV	1136886	1136886	1139515	0	WE	-
Diploid_Heat*	IV	1293360	1292695	1299206	0.0026	WE	-
Diploid_Heat	IV	1321498	1317879	1321625	0.0319	WE	-
Diploid_Heat	IV	1433562	1433562	1443459	0	WA	-
Diploid_Heat	IV	1491973	1490677	1494771	0.0447	WA	-
Diploid_Heat	V	15966	15229	16323	0.0377	WA	-
Diploid_Heat	V	75659	75455	80303	0.0033	WE	-
Diploid_Heat	V	216732	216732	220165	0	WE	-

Diploid_Heat	V	485594	485594	486858	0.0333	WA	-
Diploid_Heat*	V	540523	537410	550561	0.0082	WA	WE
Diploid_Heat	VI	83366	80828	86372	0.0297	WA	-
Diploid_Heat	VI	194098	191621	195363	0.0424	WE	-
Diploid_Heat	VI	253031	249001	255005	0.0343	WE	-
Diploid_Heat	VII	12505	12505	12505	0.0057	WE	-
Diploid_Heat	VII	101057	95002	106423	0.0318	-	WE
Diploid_Heat*	VII	138161	122858	140637	0.0053	-	WE
Diploid_Heat	VII	281559	279321	284178	0.0439	WA	-
Diploid_Heat	VII	304250	303178	306344	0.0409	WA	-
Diploid_Heat	VII	349168	349047	352209	0.0064	-	WE
Diploid_Heat	VII	487208	483838	503759	0.0053	-	WE
Diploid_Heat	VII	766107	765420	766107	0.0227	WE	-
Diploid_Heat*	VII	860980	860539	862237	0.0182	-	WE
Diploid_Heat	VIII	286803	278463	287902	0.0377	WA	-
Diploid_Heat	VIII	421413	418995	428826	0.0307	WA	-
Diploid_Heat	VIII	481690	480039	482961	0.0248	WA	-
Diploid_Heat	IX	72405	61274	80199	0.0118	WA	-
Diploid_Heat	IX	132145	129781	133695	0.0032	WE	-
Diploid_Heat	IX	260231	258754	260413	0.0328	WE	-
Diploid_Heat	IX	359074	358447	361389	0.0269	WE	-
Diploid_Heat	X	30373	30116	30730	0.0258	-	WE
Diploid_Heat	X	49501	49357	51723	0.035	-	WE
Diploid_Heat	X	84308	83775	86415	0.0254	WA	-
Diploid_Heat*	X	216505	209216	265118	0.0026	WE	-
Diploid_Heat*	X	409187	409187	458557	0	WE	-
Diploid_Heat	X	606608	604873	608497	0.0281	WE	-
Diploid_Heat	XI	41491	39004	43452	0.0404	WE	-
Diploid_Heat	XI	72599	72599	96919	0	WE	-
Diploid_Heat	XI	148989	148762	148989	0.0469	WE	-
Diploid_Heat	XI	183853	183853	202219	0	WE	-
Diploid_Heat	XI	440857	439054	441454	0.0437	WA	-
Diploid_Heat	XI	512598	511724	518802	0.0133	WA	-
Diploid_Heat	XII	22222	21405	22491	0.0264	-	WE
Diploid_Heat	XII	204576	203683	204576	0.0083	WE	-
Diploid_Heat	XII	509221	505810	511267	0.0267	WA	-
Diploid_Heat	XII	650547	646161	650731	0.0114	WA	-
Diploid_Heat	XII	762443	761264	768100	0.0166	WE/WA	-
Diploid_Heat	XII	810266	808446	810423	0.005	WE	-
Diploid_Heat	XII	846811	845924	860243	0.0111	WA	-
Diploid_Heat	XIII	49730	46836	52335	0.0379	WA	-
Diploid_Heat	XIII	78509	78303	84178	0.0117	WA	-
Diploid_Heat	XIII	210922	210127	212014	0.0495	WA	-
Diploid_Heat	XIII	548998	545976	552304	0.0033	WE	-

Diploid_Heat	XIII	687330	685688	699414	0.011	WA	-
Diploid_Heat	XIII	718071	716977	721562	0.0384	WA	-
Diploid_Heat*	XIII	760318	757317	763742	0.0476	WA	-
Diploid_Heat	XIII	778455	775320	778751	0.0494	WA	-
Diploid_Heat*	XIII	885660	885660	885660	0.0143	WE	-
Diploid_Heat	XIV	70909	65700	74669	0.0148	WA	-
Diploid_Heat	XIV	114314	112908	116758	0.0026	WE	-
Diploid_Heat	XIV	159231	159231	178002	0	WE	-
Diploid_Heat	XIV	391906	387530	395589	0.0128	WA	WE
Diploid_Heat*	XIV	481662	480344	483244	0.0046	WE	-
Diploid_Heat	XIV	639071	633308	639553	0.0346	WA	-
Diploid_Heat*	XIV	681641	679997	684671	0.0256	WA	-
Diploid_Heat*	XV	162197	162197	187590	0	WA	-
Diploid_Heat	XV	369650	369650	369757	0	WA	-
Diploid_Heat	XV	518718	516666	518878	0.0409	WE	-
Diploid_Heat	XV	739495	735874	765691	0.0089	WA	-
Diploid_Heat	XV	837629	837629	862223	0	WA	-
Diploid_Heat*	XV	1032774	1032774	1035179	0	WA	-
Diploid_Heat	XVI	41379	38929	50394	0.0028	WE	-
Diploid_Heat	XVI	272770	272770	276004	0	WE	-
Diploid_Heat	XVI	374153	372739	375200	0.0261	-	WE
Diploid_Heat	XVI	554766	554347	557974	0.017	WA	-
Diploid_Heat	XVI	747555	747367	750031	0.0054	WE	-
Diploid_Heat	XVI	873626	871856	875074	0.0285	WA	-

Table S8 Segregation patterns for QTLs identified from: (A) linkage analysis; (B) selected regions during twelve intercross rounds; (C) haploid heat selection and (D) diploid heat selection. In the “Alleles” column a single allele indicate the allele with different fitness compared to the other three in the 1:3 segregation mode; a pair of alleles with symbol “&” in between, indicates alleles with equal fitness in the 2:2 segregation mode (also the other two have equal fitness); pair of alleles with symbol “-” indicates the alleles with distinct fitness effect in multi-allelic segregation mode (1:1:2 – 1:2:1 or 2:1:1).

A. Linkage Analysis

Phenotype	Chrom	Peak	Growth trait	AIC Weight														P-value	Alleles
				NA-vs- WA, WE, SA,	WA-vs- NA, WE, SA	WE-vs- NA, WA, SA	SA-vs- NA, WA, WE	NA, WA- vs- WE, SA	NA, WE- vs- WA, SA	NA,SA -vs- WA,W E	NA-vs- WA-vs- WE,SA	NA- vs- WE- vs- WA, SA	NA- vs- SA- vs- WA, WE	WA- vs- WE- vs- NA, SA	WA- vs- SA- vs- NA, WE	WE- vs- SA- vs- NA, WA	NA- vs- WA- vs- WE- vs- SA		
Arsenite	II	752324	rate	1.5	-9.3	1.1	0.4	-2	-0.1	-1.1	-8.4	0.9	-0.2	-8.3	-8.4	-1	-7.4	1.20E-05	WA
Arsenite	IV	1247236	lag	-2.7	-0.3	-6.2	1.5	-7.8	1.9	0.2	-6.8	-7.1	-1.9	-6.4	0.6	-9.9	-9	1.00E-05	WA-SA
Arsenite	IV	1516800	lag	1.9	-6.9	1.3	-1.1	-4	1.8	0	-7.3	2.3	-0.4	-5.9	-6.9	-3.1	-6.5	1.30E-04	NA-WA
Arsenite	V	102405	rate	-0.1	-7.2	2	1.9	2	-1.1	0.5	-7.2	-0.4	0.6	-6.3	-6.9	2.9	-6.3	1.00E-04	WA
Arsenite	VI	23330	rate	1.8	-0.1	-7.8	2	-0.6	0	1.6	-0.2	-6.8	2.6	-7.8	0.2	-7.1	-6.8	7.80E-05	WA-WE
Arsenite	VII	7936	rate	2	0.4	-8.6	1	0.5	-3.5	0.6	1.2	-7.8	1.6	-8	-2.7	-7.6	-7	1.60E-04	NA-WA
Arsenite	XI	68139	rate	0.9	-7.3	1.9	1	0.1	0.9	-0.4	-6.9	1.1	0.5	-7.5	-6.8	0.5	-6.5	1.10E-04	WA-WE
Arsenite	XII	981727	rate	1.5	-6.1	1.5	-6.7	-6	1.1	-6.8	-6.1	1.9	-6.7	-5.8	-6.6	-5.7	-5.8	1.30E-04	SA
Arsenite	XIII	346821	rate	0.9	-1.3	-7.2	1.9	-4.9	-1.4	0.6	-4.2	-6.3	1.5	-6.8	-0.9	-6.5	-5.9	9.70E-05	WE
Arsenite	XIII	861124	rate	-6.7	1.4	-0.2	1.4	-1.7	0.2	-2.1	-5.9	-6	-5.8	-1.4	1.2	-0.8	-5	1.60E-04	NA
Arsenite	XIV	560654	rate	0.8	1.8	-6.2	-0.6	0.4	-1.1	-3.1	1.1	-5.3	-2.1	-6.2	-0.8	-5.7	-5.2	3.30E-04	WE
Arsenite	XIV	644582	lag	1.6	2	-10.2	1.9	1.3	1.7	1.3	2.1	-9.4	2.3	-9.3	2.7	-9.2	-8.4	2.70E-04	WE
Arsenite	XV	173494	lag	1.4	1.7	-5.9	1.6	2	2	0.4	2.4	-5.2	1.4	-5.7	2.5	-5.1	-4.7	4.90E-06	WE
Arsenite	XV	944695	rate	-5.7	0.9	1.9	-0.5	-0.8	-3.4	0.8	-4.7	-5	-4.8	1.6	-2.4	-0.1	-4	4.60E-04	NA
Arsenite	XVI	342504	lag	-6.8	-1.7	0.5	1.8	1.5	0.6	-5.5	-7.4	-5.9	-8.2	-4.9	-0.8	1.5	-7.6	5.30E-05	NA-SA
Arsenite	XVI	72835	rate	0.4	1.2	-6.4	0.9	-1.3	-0.6	-1.8	-0.3	-5.4	-0.8	-5.4	0.4	-5.4	-4.4	2.30E-	WE

Heat	II	99632	rate	1.9	-0.2	-6.8	-0.1	-0.2	-4.3	-0.3	0.3	-6.4	0.3	-6	-3.3	-5.8	-5.4	04 6.00E-04	WE
Heat	IV	669598	rate	-7	1.2	1.6	-1.5	-3.4	-3.6	0.8	-6.1	-6.5	-6.2	1.6	-2.6	-2.7	-5.5	1.20E-04	NA
Heat	XI	66234	lag	-14.2	2	-2	2	-5.2	2	-2.3	-13.7	-	-	-1.8	3	-4.4	-13	9.00E-08	NA
Heat	XIII	895642	lag	-1.4	-9.4	0.6	2	1.3	-5.1	-0.7	-9	-4.2	-1	-8.6	-9.8	1.5	-8.8	3.60E-05	WA
Heat	XIII	895642	rate	1	-8.2	1.1	1.3	-0.4	-0.8	-1	-7.2	0.2	0	-7.3	-7.2	0.6	-6.3	1.10E-05	WA-SA
Heat	XIV	665508	rate	-6.6	-0.2	1.8	1.7	1.1	-1	-1.4	-6.1	-6	-5.6	-1	-0.2	2.1	-5.2	1.90E-04	NA
Paraquat	III	212260	lag	-0.7	-0.8	-9	1.7	-5.3	-0.9	-2.1	-4.7	-8	-1.3	-8.6	-0.6	-8.3	-7.7	1.70E-05	WA
Paraquat	XIV	356878	lag	1.5	-7.3	2	-2.2	-1.3	1.6	-4.6	-6.4	2.4	-4.1	-7.4	-7.7	-1.7	-6.9	9.10E-05	WA-SA
Paraquat	VI	45521	rate	1.7	-1.2	1.2	-9	-1.9	0.3	-3.7	-1.4	1.3	-8.2	-3.3	-8.9	-8.3	-7.9	1.60E-05	SA

B. Intercross

Chromosome	Peak	LOD	Alleles
IV	699005	12.9	NA-WA
V	172739	13.1	NA-WA
VI	70206	12.8	WE
VIII	112561	10.2	WE-SA
XI	85720	13.1	WE
XII	444427	14.4	NA&WA
XV	171462	12.3	WE-SA
XV	385888	13.1	WA&WE
XVI	205083	14.1	WA-SA

C. Heat Haploids

Chromosome	Peak	LOD	Alleles
I	41803	13.1	WA
II	192041	14.1	SA
II	387448	12.1	WA
II	520956	14	NA-WE
III	103983	14.1	NA&WA
IV	88326	12.9	WE&SA
IV	167026	14.8	WA
IV	193568	15.9	WE
IV	341596	15.2	SA
IV	482650	13.7	NA-WA
IV	594972	15.7	NA
IV	1165241	14.6	WA&WE
IV	1299590	14.7	NA&WA
IV	1424223	13.8	WE&SA
V	210929	15.1	SA
V	551439	15.3	NA&WA
VII	548507	13.9	WE
VIII	293911	14.5	WA
VIII	424239	13.2	WA-WE
X	236389	14.7	WA&WE
X	412413	13.9	NA-WE
XI	40024	15.3	WE
XII	209043	14.2	WA&SA
XIII	721067	14.7	WA
XIII	877042	13.2	WA
XIV	115349	13.6	NA
XIV	173831	15.5	WE
XIV	273491	15.1	SA
XIV	481655	14.5	NA&WE
XV	174678	11.8	NA-SA
XV	345603	14.9	NA
XV	837629	13.6	WA
XV	954035	14.6	WE
XVI	264571	15.5	SA

D. Heat Diploids

Chromosome	Peak	LOD	Alleles
I	8001	15.4	WA
II	241424	14.5	WE&SA
II	287551	13.8	WE
II	384879	13.7	NA-WA
II	510231	14.1	NA-WE
IV	78729	15.1	WA
IV	161585	15.3	WE
IV	301375	15.2	WA
IV	595513	14.6	WA&WE
IV	1043658	15.1	SA
IV	1136886	15.9	WE
IV	1293360	15	WE
IV	1433562	14.6	NA&WA
V	75659	15.4	WE
V	216732	14.2	WE
V	540523	15.3	WA
VII	12505	13.7	NA
VII	138161	13.7	SA
VII	349168	15.8	WE
VII	487208	14.3	SA
IX	132145	14.5	WE
X	216505	15.5	NA&SA
X	409187	13.9	WE-SA
XI	72599	16	WE
XI	183853	14.2	NA&WA
XII	204576	15.6	WE
XII	810266	15.7	NA
XIII	548998	15.8	WA&SA
XIV	114314	15.8	WE
XIV	159231	15.6	WE
XIV	481662	13.6	WA-SA
XV	162197	13.6	NA&SA
XV	369650	16	WA
XV	739495	13.5	SA
XV	837629	13.9	WA-WE
XV	1032774	14.1	WE
XVI	41379	14.3	SA
XVI	272770	14.7	SA
XVI	747555	15.5	WE&SA