Mycoflora of Soybeans Used for Meju Fermentation

Dae-Ho Kim^{1,2}, Seon-Hwa Kim¹, Soon-Wo Kwon¹, Jong-Kyu Lee² and Seung-Beom Hong^{1,*}

¹Korean Agricultural Culture Collection, Agricultural Microbiology Div. National Academy of Agricultural Science, RDA, Suwon 441-853, Korea

²Tree Pathology and Mycology Laboratory, Division of Forest Environment Protection, Kangwon National University, Chuncheon 200-701, Korea

Volume 41, No 2, p. 100-107, 2013.

Table 2 contained an error (bold in text).

Table 2. List of fungal species from soybeans and their isolation frequencies

Scientific name	Representative strains		Incidence on meju ^b	Isolation frequency from sodium hypochlorite untreated soybeans ^c				Isolation frequency from sodium hypochlorite untreated soybeans ^c			
	Strain number	Sequence number ^a		Number of factories ^d	DRBC	DG18	MEA	Number of factories ^d	DRBC	DG18	MEA

The corrected Table 2 follows below (bold in text).

Table 2. List of fungal species from soybeans and their isolation frequencies

Scientific name	Representative strains		Incidence on meju ^b	Isolation frequency from sodium hypochlorite untreated soybeans ^c				Isolation frequency from sodium hypochlorite-treated soybeans ^c			
	Strain number	Sequence number ^a		Number of factories ^d	DRBC	DG18	MEA	Number of factories ^d	DRBC	DG18	MEA