What Does the Cochrane Collaboration Say about Rehabilitation of the Arm after Stroke?

Mehrholtz J, Hädrich A, Platz T, et al. Electromechanical and robot-assisted arm training for improving generic activities of daily living, arm function, and arm muscle strength after stroke. *Cochrane Database Syst Rev.* 2012;6(6):CD006876.<http://dx.doi.org/10.1002/14651858.CD006876.pub3> Medline:22696362

Coupar F, Pollock A, van Wijck F, et al. Simultaneous bilateral training for improving arm function after stroke. *Cochrane Database Syst Rev.* 2010;(4):CD006432.<http://dx.doi.org/10.1002/14651858.CD006432.pub2> Medline:20393947

Barclay-Goddard RE, Stevenson TJ, Poluha W, et al. Mental practice for treating upper extremity deficits in individuals with hemiparesis after stroke. *Cochrane Database Syst Rev.* 2011;(5):CD005950.<http://dx.doi.org/10.1002/14651858.CD005950.pub4> Medline:21563146

Sirtori V, Corbetta D, Moja L, et al. Constraint-induced movement therapy for upper extremities in stroke patients. *Cochrane Database Syst Rev.* 2009;(4):CD004433.<http://dx.doi.org/10.1002/14651858.CD004433.pub2> Medline:19821326

Laver KE, George S, Thomas S, et al. Virtual reality for stroke rehabilitation. *Cochrane Database Syst Rev.* 2011;(9):CD008349.<http://dx.doi.org/10.1002/14651858.CD008349.pub2> Medline:21901720

Coupar F, Pollock A, Legg LA, et al. Home-based therapy programmes for upper limb functional recovery following stroke. *Cochrane Database Syst Rev.* 2012;5(5):CD006755.<http://dx.doi.org/10.1002/14651858.CD006755.pub2> Medline:22592715

Doyle S, Bennett S, Fasoli SE, et al. Interventions for sensory impairment in the upper limb after stroke. *Cochrane Database Syst Rev.* 2010;(6):CD006331.<http://dx.doi.org/10.1002/14651858.CD006331.pub2> Medline:20556766

Thieme H, Mehrholtz J, Pohl M, et al. Mirror therapy for improving motor function after stroke. *Cochrane Database Syst Rev.* 2012;3(3):CD008449.<http://dx.doi.org/10.1002/14651858.CD008449.pub2> Medline:22419334

French B, Thomas LH, Leathley MJ, et al. Repetitive task training for improving functional ability after stroke. *Cochrane Database Syst Rev.* 2007;(4):CD006073.<http://dx.doi.org/10.1002/14651858.CD006073.pub2> Medline:17943883

Winter J, Hunter S, Sim J, et al. Hands-on therapy interventions for upper limb motor dysfunction following stroke. *Cochrane Database Syst Rev.* 2011;(6):CD006609.<http://dx.doi.org/10.1002/14651858.CD006609.pub2> Medline:21678359

Price CIM, Pandyan AD. Electrical stimulation for preventing and treating post-stroke shoulder pain. *Cochrane Database Syst Rev.* 2000;(4):CD001698.<http://dx.doi.org/10.1002/14651858.CD001698> Medline:11034725

Ada L, Foongchomcheay A, Canning CG. Supportive devices for preventing and treating subluxation of the shoulder after stroke. *Cochrane Database Syst Rev.* 2005;(1):CD003863.<http://dx.doi.org/10.1002/14651858.CD003863.pub2> Medline:15674917

The Cochrane Collaboration is an international not-for-profit and independent organization dedicated to making up-to-date, accurate information about the effects of health care readily available worldwide. It produces and disseminates systematic reviews of health care interventions and promotes the search for evidence in the form of clinical trials and other studies of interventions. For more information, visit www.cochrane.org.

DOI:10.3138/ptc.65.4.cochrane