Published in final edited form as:

Front Biol (Beijing). 2013 June 1; 8(3): . doi:10.1007/s11515-012-1204-4.

ROS-mediated regulation of CXCR4 in cancer

Mahandranauth A. Chetram and Cimina V. Hinton*

Center for Cancer Research and Therapeutic Development, Clark Atlanta University, Atlanta, GA, 30314, USA

Abstract

Oxidative stress and the accumulation of reactive oxygen species (ROS) play a role in cancer cells developing an advanced, phenotypic signature that associates with metastasis and progression. Increased ROS concentrations are involved in promoting cancer development and metastasis by inducing expression of oncogenes, suppressing activity of anti-survival molecules and by activating various cell survival and proliferation signaling pathways. Oxidative stress is higher in the epithelium of cancer patients than patients without the disease, and antioxidant trials are currently being explored as a therapeutic option. However, studies have shown that ROS increases expression of CXCR4 in cancer and immune cells. CXCR4 expression in tumors strongly correlates to metastasis and poor prognosis. Herein, we discuss an emerging relationship between ROS and CXCR4 in cancer cells.

Keywords

Reactive oxygen species; CXCR4; HIF1 ; Metastasis; PI3K/AKT; ERK1/2

Introduction

The tumor microenvironment is a major player in cancer progression and metastasis. Some metastatic influences from the tumor microenvironment are due to various secreted molecules from immune cells, such as cytokines and chemokines, which results in decreased function of tumor suppressor genes, increased function of proto-oncogenes and cell survival signaling (Liu et al., 2011). The tumor microenvironment also secretes reactive oxygen species (ROS), which modulate anti-cancer and pro-cancer pathways to enhance tumorigenesis (Cook et al., 2004)[2]. ROS are a group of highly reactive oxygen containing molecules, where their accumulation typically functions as second messengers in intracellular signal transduction, and consequently, enhances cancer cell survival, proliferation and migration (Storz, 2005; Pan et al., 2009). Excessive accumulation of ROS, however, result in deleterious effects, such as genotoxicity and cell death (Simon et al., 2000).

Reactive oxygen species

Reactive oxygen species are oxygen-derived free radicals, such as superoxide anion ($O_2^{\cdot-}$), hydroxyl radicals ($OH^{\cdot}$), peroxy ($RO_2^{\cdot}$), and oxygen-derived nonradical species like hydrogen peroxide (H_2O_2) (Turrens, 2003). The major cellular source of ROS is through the mitochondrial complexes I and III, where electrons from NADH or $FADH_2$ react with oxygen to produce superoxide anions (Turrens, 2003). ROS is also produced in the cytosol

by intracellular enzymes such as oxidases, lipoxygenases and cyclooxygenases (Turrens, 2003). It is well established that ROS is implicated oxidizing biomolecules such as DNA, lipids and protein, which over time can lead to cell death (Davies, 1993; Lindahl, 1993; Wagner et al., 1994). Under physiologic conditions, to prevent cellular damage, cells have developed several antioxidant systems to defend against the deleterious effects of ROS. Some of the main antioxidant systems used to maintain ROS homeostasis, are the glutathione, thioredoxin, superoxide dismutase (SOD) and catalase systems (Circu and Aw, 2010).

Interestingly, accumulating studies show that ROS play a critical role in regulating normal cellular function through gene transcription, cell signaling and post-translational modification (Pan et al., 2009; Chetram et al., 2011). It has recognized that loss of ROS homeostasis results in pathogenic effects and fatal diseases, such as diabetes, cardiovascular disease and cancer (Galaris et al., 2008; Hambali et al., 2011; Rains and Jain, 2011).

Reactive oxygen species in cancer

Tumorigenesis is characterized by a profound reprogramming of the genome, which ultimately activates oncogenes and inhibits tumor suppressor genes. Part of this reprogramming is due to elevated levels of ROS, which occurs as a result of the close proximity of DNA to hydroxyl radical generation, consequently modifying purine, pyrimidine and deoxyribose residues (Fraga et al., 1990). In cancer cells, increased ROS accumulation is generally due to loss of antioxidant or increased expression of pro-oxidants. For instance, Nelson et al. (2004) has observed somatic silencing of glutathione S-transferase (GSTP1), a member of cellular antioxidant systems, in almost all prostate cancer cases that were investigated. In addition to prostate cancer, GSTP1 was also silenced in various other malignancies, such as hepatocellular carcinoma and breast cancer (Tchou et al., 2000; Cho et al., 2012). Conversely, cancer cells also induce ROS generation through the expression of various pro-oxidants molecules, such as NOX and PUMA, and through changes in mitochondrial functions. Additionally, current evidence suggest that the tumor microenvironment consisting of immune cells, fibroblast cells and stromal cells work together to enhance the production of ROS, thereby promoting cancer progression (Cook et al., 2004). Together, the loss of antioxidants mechanisms and the gain of pro-oxidant mechanisms regulate ROS to facilitate tumor development and progression (Liou and Storz, 2010).

It is well established that ROS functions as a secondary-messenger that regulates cell survival pathways and ultimately promoting malignant transformation and progression (Storz, 2005). Emerging evidence indicates that ROS also promotes post-translation modification of kinases, phosphatases, tumor suppressors and oncogenic molecules, such as kinase p21ras, PTEN, p53 and HIF1, respectively (Lee et al., 2002; Salmeen et al., 2003), and mediates cell survival and metastasis by regulating pathways such as PI3K/AKT and MAPK/ERK1/2 in the prostate, breast and leukemia (Kumar et al., 2008; Tomic et al., 2011). For instance, hydrogen peroxide activated the MAPK pathway to provide cellular protection and proliferation in response to oxidative stress, which correlated with sustained activation of Protein Kinase C (PKC) (Guyton et al., 1996; Wu et al., 2006). Ha and Yu (2010) described that ROS enhanced hepatocellular carcinogenesis via inhibition of PTEN in favor of AKT. Our laboratory, and others, also observed an inactivation of PTEN catalytic function by ROS, which correlated with increased expression of phosphorylated AKT (p-AKT) (Ha and Yu, 2010; Chetram et al., 2011). Moreover, ROS regulates transcription factors, directly or indirectly, resulting in cell cycle induction and apoptosis suppression (Gupta et al., 1999). Transcription factors such as Sp1, glucocorticoid receptor and Egr1 were described as redox-sensitive, and underwent modifications of their DNA binding

domains or posttranslational modification sites to promote or inhibit gene expression (Ammendola et al., 1994; Esposito et al., 1995; Landriscina et al., 2009).

Chemokine (CXC) receptor 4 (CXCR4)

The secreted chemokine protein family mediates cellular functions, such as: (i) regulation of hematopoiesis; (ii) leukocyte maturation; (iii) angiogenesis; (iv) cellular trafficking; (v) homing of immune cells; and (vi) development of lymphoid tissue (Hinton et al., 2010). Chemokine proteins were initially characterized by their ability to induce leukocyte migration recruitment (Loetscher et al., 2000). Since then, chemokines and their roles have been characterized in a variety of host tissues, and the roles that have been ascribed to chemokines make them favorable to cancer development, progression and metastasis: (i) immune cell induction and migration to a site of infection; and (ii) regulation of cell migration during normal tissue maintenance or development (Hinton et al., 2010).

Chemokine receptors are G protein-linked, transmembrane proteins found on the surfaces of endo- and epithelial cells of many tissues (Gupta et al., 1998). The family roster includes approximately 45 ligands and 19 receptors. As many as six receptors show affinity for one specific ligand. However, most ligands have an affinity for several receptors, and can bind to non-G protein coupled receptors (Lau et al., 2004). The subsequent activation of a chemokine receptor involves classical pathways of cell survival: phospholipase C (PLC- β), PI3K/AKT, and the MAPK cascade (Hinton et al., 2010). Stromal-cell derived factor 1 (SDF1 or CXCL12) is a conserved member of the β -chemokine subfamily and is considered the exclusive ligand for CXCR4 receptor (Balabanian et al., 2005). It has been suggested that SDF1 may also bind CXCR7; and ubiquitin was recently identified as an alternate ligand for CXCR4 (Cruz-Orengo et al., 2011; Saini et al., 2011).

Chemokine receptors are associated with a myriad of inflammatory illnesses (cardiovascular, allergic inflammatory, neuro-inflammatory and HIV-associated diseases) and cancer (Gerard and Rollins, 2001). The interaction between chemokines and their receptors have been shown to induce cell migration, cytoskeletal rearrangement, and adhesion of malignant cells onto endothelial cells for invasion, as activation of CXCR4-induced migration of immune cells, as well as embryonic development, growth regulation, angiogenesis, and hematopoiesis (Kucia et al., 2004; Wang et al., 2007). Studies have also implicated CXCR4 in malignant cancer development by its involvement in: (i) cell motility; (ii) adhesion; (iii) secretion of matrix metallo-proteinases (MMPs); (iv) angiogenesis and (v) activation of metastatic-associated signaling pathways (G-proteins, PI3K/AKT, JAK/STAT, Src kinase and HER2) (Busillo and Benovic, 2007; Wang et al., 2007), which has been well demonstrated in several human malignancies.

Cancer mortality is often a result of primary tumor cell metastasis to distal organs, where the expression of CXCR4 tightly correlates with a metastatic cell phenotype. Müller et al. (2001) observed that CXCR4 was higher in malignant breast tumors compared to their normal, healthy counterparts, suggesting that CXCR4 expression correlated with increased metastasis-associated mortality [43]. Tanabe et al. (1997) demonstrated that the expression of CXCR4 in surgically resected invasive ductal carcinomas significantly correlated with the degree of lymph node metastasis. Likewise, Taichman et al. (2002) observed that CXCR4 facilitated prostate cancer metastasis to the bone, the primary site of distal prostate cancer colonization.

ROS-mediated regulation of CXCR4

In cancer and other cell models, CXCR4 and SDF1 are upregulated in high ROS concentrations and/or hypoxic environments, where oxygen levels are reduced below normal

(Li et al., 2009). In these conditions, cells react by inducing expression of a hypoxia-inducible factor (HIF) transcription complex, which enhances the expression of targeted genes involved in adjusting to low oxygen concentrations (Fig. 1). A survey of malignant tissues demonstrated an increase in HIF1 α expression in bladder, breast, colon and prostate tumors, which correlated with CXCR4 and SDF1 α expression in breast tumors (Talks et al., 2000; Staller et al., 2003; Zhong et al., 2004). The heterodimeric HIF transcription complex consists of a regulatory HIF1 α subunit and the constitutively expressed HIF1 β subunit, both members of the basic helix–loop–helix (bHLH) and PER-ARNT-SIM (PAS) families of transcription factors (Wang et al., 1995). The HIF1 α subunit acts as a transcription factor for CXCR4 by binding to the HIF1 response element (HRE) found in the promoter region of CXCR4 (Staller et al., 2003). Therefore, a tumor microenvironment that is rich in ROS may critically influence CXCR4-mediated expression and functions, ultimately encouraging cancer progression.

ROS-mediated regulation of CXCR4 expression and function is fairly new in cancer research. Wu et al. (2006) described that ROS sustained PKC and ERK1/2 activation, which led to epithelial-mesenchymal transition (EMT)-like scattering, migration and reduced growth of hepatoma cells. Models of ROS regulation of CXCR4 are scarcely described, but have been shown in immune cells. Lee et al. (2007) demonstrated that depletion of ROS by a ROS scavenger, N-acetylcysteine (NAC), promoted CXCR4-mediated activation of AKT in B cells. Lin et al. (2011) described that CXCR4 induced ROS production in hematopoietic stem cells. Finally, loss of ROS reduced secretion of SDF1 α from human bone marrow osteoblasts and endothelial cells (Dar et al., 2011). We have shown in prostate cancer cells that ROS oxidized PTEN, rendering it inactive, which may have explained the concomitant increase in expression of p-AKT by ROS, and further suggest a regulatory role of ROS in cancer cells (Chetram et al., 2011). Likewise, we observed an increase in CXCR4 expression upon ROS treatment, which also correlated with increased expression of p-AKT (Chetram et al., 2011). ROS enhanced prostate cancer cell migration and invasion in a CXCR4-dependent manner, which was abrogated by CXCR4 antagonist AMD3100 (Chetram et al., 2011).

ROS is used in current therapy to induce apoptosis in tumors (Ozben, 2007). Drugs, such as anthracyclines and epipodophyllotoxins, are used to induce ROS accumulation in cancer cells, hoping to create oxidative stress and induce cell death (Ozben, 2007). Conversely, the accumulation of ROS has been shown to activate survival pathways in cancer cells (Liu et al., 2006). Our findings in prostate cancer, along with immune studies, demonstrate that ROS may act as an inducing molecule, resulting in increased CXCR4 expression and subsequent cell survival mechanisms. Therefore, cells may increase ROS generation and expression of survival genes, like CXCR4, to escape oxidative stress and avoid death (Pani et al., 2010). We and others observed an increase in p-AKT expression, which activates HIF1 α and may have led to the correlating increase in CXCR4 (Liu et al., 2006; Kumar et al., 2008; Chetram et al., 2011). We have also demonstrated that an absence of PTEN increased CXCR4 expression and function (Chetram et al., 2011). Interestingly, ROS inactivated PTEN through oxidation at cysteine residues, which further correlated with increased p-AKT expression and CXCR4 function, and elucidates a putative mechanism for cancer cell survival (Lee et al., 2002; Chetram et al., 2011). Moreover, p-AKT increased expression may be due to ROS-mediated inactivation of PTEN, and ultimately, CXCR4 expression. Therefore, it is possible that ROS may activate HIF1 α and subsequently induce the expression of CXCR4 for cell survival (Li et al., 2009). To our knowledge, we are first in describing a link between ROS and CXCR4, which may suggest a potential mechanism of how cells may acquire increased CXCR4 and a metastatic phenotype, and a tendency to move to a distal organ. Concentrations of ROS have been generally considered apoptotic in cancer cells. We believe that ROS concentrations should be considered during therapeutic

treatments, since survival mechanisms and metastasis can inadvertently be induced. Several key questions remain about ROS and CXCR4-mediated cancer progression, such as (i) whether survival mechanisms have already been activated in the primary tumor; or (ii) whether primary tumors induce survival mechanism after exposure to ROS, hypoxia and other deleterious microenvironments. ROS and oxidative stress have long been associated with tumor progression; however, the underlying mechanisms are well defined (Kumar et al., 2008). To improve the efficacy of oxidant cancer therapy, studies should consider neutralizing ROS molecules, instead of increasing concentrations, or take into account the stage and grade of tumors since oxidative stress can encourage a metastatic signature in cancer cells (Kumar et al., 2008).

Acknowledgments

Limitations of space preclude extensive citation of literature; we apologize to those whose work is not mentioned herein. Research in this laboratory is supported, in part, by National Institutes of Health grants RCM122622A (MAC), F31CA153908 (MAC), 2G12RR003062-22 (CVH) and 1P20MD002285 (CVH).

References

- Ammendola R, Mesuraca M, Russo T, Cimino F. The DNA-binding efficiency of Sp1 is affected by redox changes. *Eur J Biochem.* 1994; 225(1):483–489. [PubMed: 7925470]
- Balabanian K, Lagane B, Infantino S, Chow KY, Harriague J, Moepps B, Arenzana-Seisdedos F, Thelen M, Bachelier F. The chemokine SDF-1/CXCL12 binds to and signals through the orphan receptor RDC1 in T lymphocytes. *J Biol Chem.* 2005; 280(42):35760–35766. [PubMed: 16107333]
- Busillo JM, Benovic JL. Regulation of CXCR4 signaling. *Biochim Biophys Acta.* 2007; 1768(4):952–963. [PubMed: 17169327]
- Chetram MA, Odero-Marah V, Hinton CV. Loss of PTEN permits CXCR4-mediated tumorigenesis through ERK1/2 in prostate cancer cells. *Mol Cancer Res.* 2011; 9(1):90–102. [PubMed: 21076047]
- Cho YH, Shen J, Gammon MD, Zhang YJ, Wang Q, Gonzalez K, Xu X, Bradshaw PT, Teitelbaum SL, Garbowski G, Hibshoosh H, Neugut AI, Chen J, Santella RM. Prognostic significance of gene-specific promoter hypermethylation in breast cancer patients. *Breast Cancer Res Treat.* 2012; 131(1):197–205. [PubMed: 21837480]
- Circu ML, Aw TY. Reactive oxygen species, cellular redox systems, and apoptosis. *Free Radic Biol Med.* 2010; 48(6):749–762. [PubMed: 20045723]
- Cook JA, Gius D, Wink DA, Krishna MC, Russo A, Mitchell JB. Oxidative stress, redox, and the tumor microenvironment. *Semin Radiat Oncol.* 2004; 14(3):259–266. [PubMed: 15254869]
- Cruz-Orengo L, Holman DW, Dorsey D, Zhou L, Zhang P, Wright M, McCandless EE, Patel JR, Luker GD, Littman DR, Russell JH, Klein RS. CXCR7 influences leukocyte entry into the CNS parenchyma by controlling abluminal CXCL12 abundance during autoimmunity. *J Exp Med.* 2011; 208(2):327–339. [PubMed: 21300915]
- Dar A, Schajnovitz A, Lapid K, Kalinkovich A, Itkin T, Ludin A, Kao WM, Battista M, Tesio M, Kollet O. Rapid mobilization of hematopoietic progenitors by AMD3100 and catecholamines is mediated by CXCR4-dependent SDF-1 release from bone marrow stromal cells. *Leukemia.* 2011; 25(8):1286–1296. [PubMed: 21494253]
- Davies KJ. Protein modification by oxidants and the role of proteolytic enzymes. *Biochem Soc Trans.* 1993; 21(2):346–353. [PubMed: 8359494]
- Esposito F, Cuccovillo F, Morra F, Russo T, Cimino F. DNA binding activity of the glucocorticoid receptor is sensitive to redox changes in intact cells. *Biochim Biophys Acta.* 1995; 1260(3):308–314. [PubMed: 7873605]
- Fraga CG, Shigenaga MK, Park JW, Degan P, Ames BN. Oxidative damage to DNA during aging: 8-hydroxy-2'-deoxyguanosine in rat organ DNA and urine. *Proc Natl Acad Sci USA.* 1990; 87(12):4533–4537. [PubMed: 2352934]
- Galaris D, Skiada V, Barbouti A. Redox signaling and cancer: the role of “labile” iron. *Cancer Lett.* 2008; 266(1):21–29. [PubMed: 18374479]

- Gerard C, Rollins BJ. Chemokines and disease. *Nat Immunol.* 2001; 2(2):108–115. [PubMed: 11175802]
- Gupta A, Rosenberger SF, Bowden GT. Increased ROS levels contribute to elevated transcription factor and MAP kinase activities in malignantly progressed mouse keratinocyte cell lines. *Carcinogenesis.* 1999; 20(11):2063–2073. [PubMed: 10545407]
- Gupta SK, Lysko PG, Pillarisetti K, Ohlstein E, Stadel JM. Chemokine receptors in human endothelial cells. Functional expression of CXCR4 and its transcriptional regulation by inflammatory cytokines. *J Biol Chem.* 1998; 273(7):4282–4287. [PubMed: 9461627]
- Guyton KZ, Liu Y, Gorospe M, Xu Q, Holbrook NJ. Activation of mitogen-activated protein kinase by H₂O₂. Role in cell survival following oxidant injury. *J Biol Chem.* 1996; 271(8):4138–4142. [PubMed: 8626753]
- Ha HL, Yu DY. HBx-induced reactive oxygen species activates hepatocellular carcinogenesis via dysregulation of PTEN/Akt pathway. *World J Gastroenterol.* 2010; 16(39):4932–4937. [PubMed: 20954279]
- Hambali Z, Ahmad Z, Arab S, Khazaai H. Oxidative stress and its association with cardiovascular disease in chronic renal failure patients. *Indian J Nephrol.* 2011; 21(1):21–25. [PubMed: 21655165]
- Hinton CV, Avraham S, Avraham HK. Role of the CXCR4/CXCL12 signaling axis in breast cancer metastasis to the brain. *Clin Exp Metastasis.* 2010; 27(2):97–105. [PubMed: 18814042]
- Kucia M, Jankowski K, Reza R, Wysoczynski M, Bandura L, Allendorf DJ, Zhang J, Ratajczak J, Ratajczak MZ. CXCR4-SDF-1 signalling, locomotion, chemotaxis and adhesion. *J Mol Histol.* 2004; 35(3):233–245. [PubMed: 15339043]
- Kumar B, Koul S, Khandrika L, Meacham RB, Koul HK. Oxidative stress is inherent in prostate cancer cells and is required for aggressive phenotype. *Cancer Res.* 2008; 68(6):1777–1785. [PubMed: 18339858]
- Landriscina M, Maddalena F, Laudiero G, Esposito F. Adaptation to oxidative stress, chemoresistance, and cell survival. *Antioxid Redox Signal.* 2009; 11(11):2701–2716. [PubMed: 19778285]
- Lau EK, Allen S, Hsu AR, Handel TM. Chemokine-receptor interactions: GPCRs, glycosaminoglycans and viral chemokine binding proteins. *Adv Protein Chem.* 2004; 68:351–391. [PubMed: 15500866]
- Lee RL, Westendorf J, Gold MR. Differential role of reactive oxygen species in the activation of mitogen-activated protein kinases and Akt by key receptors on B-lymphocytes: CD40, the B cell antigen receptor, and CXCR4. *J Cell Commun Signal.* 2007; 1(1):33–43. [PubMed: 18481208]
- Lee SR, Yang KS, Kwon J, Lee C, Jeong W, Rhee SG. Reversible inactivation of the tumor suppressor PTEN by H₂O₂. *J Biol Chem.* 2002; 277(23):20336–20342. [PubMed: 11916965]
- Li S, Deng Y, Feng J, Ye W. Oxidative preconditioning promotes bone marrow mesenchymal stem cells migration and prevents apoptosis. *Cell Biol Int.* 2009; 33(3):411–418. [PubMed: 19356708]
- Lin W, Wu G, Li S, Weinberg EM, Kumthip K, Peng LF, Méndez-Navarro J, Chen WC, Jilg N, Zhao H, Goto K, Zhang L, Brockman MA, Schuppan D, Chung RT. HIV and HCV cooperatively promote hepatic fibrogenesis via induction of reactive oxygen species and NFκB. *J Biol Chem.* 2011; 286(4):2665–2674. [PubMed: 21098019]
- Lindahl T. Instability and decay of the primary structure of DNA. *Nature.* 1993; 362(6422):709–715. [PubMed: 8469282]
- Liou GY, Storz P. Reactive oxygen species in cancer. *Free Radic Res.* 2010; 44(5):479–496. [PubMed: 20370557]
- Liu LZ, Hu XW, Xia C, He J, Zhou Q, Shi X, Fang J, Jiang BH. Reactive oxygen species regulate epidermal growth factor-induced vascular endothelial growth factor and hypoxia-inducible factor-1α expression through activation of AKT and P70S6K1 in human ovarian cancer cells. *Free Radic Biol Med.* 2006; 41(10):1521–1533. [PubMed: 17045920]
- Liu J, Zhang Y, Zhao J, Yang Z, Li D, Katirai F, Huang B. Mast cell: insight into remodeling a tumor microenvironment. *Cancer Metastasis Rev.* 2011; 30(2):177–184. [PubMed: 21267769]
- Loetscher P, Moser B, Baggiolini M. Chemokines and their receptors in lymphocyte traffic and HIV infection. *Adv Immunol.* 2000; 74:127–180. [PubMed: 10605606]

- Müller A, Homey B, Soto H, Ge N, Catron D, Buchanan ME, McClanahan T, Murphy E, Yuan W, Wagner SN, Barrera JL, Mohar A, Verástegui E, Zlotnik A. Involvement of chemokine receptors in breast cancer metastasis. *Nature*. 2001; 410(6824):50–56. [PubMed: 11242036]
- Nelson WG, De Marzo AM, DeWeese TL, Isaacs WB. The role of inflammation in the pathogenesis of prostate cancer. *J Urol*. 2004; 172(5):6–12. [PubMed: 15201726]
- Ozben T. Oxidative stress and apoptosis: impact on cancer therapy. *J Pharm Sci*. 2007; 96(9):2181–2196. [PubMed: 17593552]
- Pan JS, Hong MZ, Ren JL. Reactive oxygen species: a double-edged sword in oncogenesis. *World J Gastroenterol*. 2009; 15(14):1702–1707. [PubMed: 19360913]
- Pani G, Galeotti T, Chiarugi P. Metastasis: cancer cell's escape from oxidative stress. *Cancer Metastasis Rev*. 2010; 29(2):351–378. [PubMed: 20386957]
- Rains JL, Jain SK. Oxidative stress, insulin signaling, and diabetes. *Free Radic Biol Med*. 2011; 50(5):567–575. [PubMed: 21163346]
- Saini V, Staren DM, Ziarek JJ, Nashaat ZN, Campbell EM, Volkman BF, Marchese A, Majetschak M. The CXC chemokine receptor 4 ligands ubiquitin and stromal-cell derived factor-1 {alpha} function through distinct receptor interactions. *J Biol Chem*. 2011; 286(38):33466–33477. [PubMed: 21757744]
- Salmeen A, Andersen JN, Myers MP, Meng TC, Hinks JA, Tonks NK, Barford D. Redox regulation of protein tyrosine phosphatase 1B involves a sulphenyl-amide intermediate. *Nature*. 2003; 423(6941):769–773. [PubMed: 12802338]
- Simon HU, Haj-Yehia A, Levi-Schaffer F. Role of reactive oxygen species (ROS) in apoptosis induction. *Apoptosis*. 2000; 5(5):415–418. [PubMed: 11256882]
- Staller P, Sulitkova J, Lisztwan J, Moch H, Oakeley EJ, Krek W. Chemokine receptor CXCR4 downregulated by von Hippel-Lindau tumour suppressor pVHL. *Nature*. 2003; 425(6955):307–311. [PubMed: 13679920]
- Storz P. Reactive oxygen species in tumor progression. *Front Biosci*. 2005; 10(1–3):1881–1896. [PubMed: 15769673]
- Taichman RS, Cooper C, Keller ET, Pienta KJ, Taichman NS, McCauley LK. Use of the stromal cell-derived factor-1/CXCR4 pathway in prostate cancer metastasis to bone. *Cancer Res*. 2002; 62(6):1832–1837. [PubMed: 11912162]
- Talbot KL, Turley H, Gatter KC, Maxwell PH, Pugh CW, Ratcliffe PJ, Harris AL. The expression and distribution of the hypoxia-inducible factors HIF-1alpha and HIF-2alpha in normal human tissues, cancers, and tumor-associated macrophages. *Am J Pathol*. 2000; 157(2):411–421. [PubMed: 10934146]
- Tanabe S, Heesen M, Yoshizawa I, Berman MA, Luo Y, Bleul CC, Springer TA, Okuda K, Gerard N, Dorf ME. Functional expression of the CXC-chemokine receptor-4/fusin on mouse microglial cells and astrocytes. *J Immunol*. 1997; 159(2):905–911. [PubMed: 9218610]
- Tchou JC, Lin X, Freije D, Isaacs WB, Brooks JD, Rashid A, De Marzo AM, Kanai Y, Hirohashi S, Nelson WG. GSTP1 CpG island DNA hypermethylation in hepatocellular carcinomas. *Int J Oncol*. 2000; 16(4):663–676. [PubMed: 10717233]
- Tomic J, Lichty B, Spaner DE. Aberrant interferon-signaling is associated with aggressive chronic lymphocytic leukemia. *Blood*. 2011; 117(9):2668–2680. [PubMed: 21205928]
- Turrens JF. Mitochondrial formation of reactive oxygen species. *J Physiol*. 2003; 552(2):335–344. [PubMed: 14561818]
- Wagner BA, Buettner GR, Burns CP. Free radical-mediated lipid peroxidation in cells: oxidizability is a function of cell lipid bisallylic hydrogen content. *Biochemistry*. 1994; 33(15):4449–4453. [PubMed: 8161499]
- Wang GL, Jiang BH, Rue EA, Semenza GL. Hypoxia-inducible factor 1 is a basic-helix-loop-helix-PAS heterodimer regulated by cellular O2 tension. *Proc Natl Acad Sci USA*. 1995; 92(12):5510–5514. [PubMed: 7539918]
- Wang J, Wang J, Dai J, Jung Y, Wei CL, Wang Y, Havens AM, Hogg PJ, Keller ET, Pienta KJ, Nor JE, Wang CY, Taichman RS. A glycolytic mechanism regulating an angiogenic switch in prostate cancer. *Cancer Res*. 2007; 67(1):149–159. [PubMed: 17210694]

- Wu WS, Tsai RK, Chang CH, Wang S, Wu JR, Chang YX. Reactive oxygen species mediated sustained activation of protein kinase C alpha and extracellular signal-regulated kinase for migration of human hepatoma cell Hepg2. *Mol Cancer Res.* 2006; 4(10):747–758. [PubMed: 17050668]
- Zhong H, Semenza GL, Simons JW, De Marzo AM. Up-regulation of hypoxia-inducible factor 1alpha is an early event in prostate carcinogenesis. *Cancer Detect Prev.* 2004; 28(2):88–93. [PubMed: 15068831]

Figure 1.

A schematic of ROS-mediated regulation of CXCR4 expression in cancer cells as described by (1) Chetram et al. (2011), (2) Lui et al. (2006) and (3) Staller et al. (2003).