Published in final edited form as:

Angew Chem Int Ed Engl. 2013 July 22; 52(30): . doi:10.1002/anie.201302749.

Anthracimycin, a Potent Anthrax Antibiotic from a Marine-Derived Actinomycete**

Dr. Kyoung Hwa Jang[†], Dr. Sang-Jip Nam[†], Dr. Jeffrey B. Locke[‡], Mr. Christopher A. Kauffman[†], Ms. Deanna S. Beatty[†], Ms. Lauren A. Paul[†], and Prof. Dr. William Fenical^{†,*}

[†]Center for Marine Biotechnology and Biomedicine, Scripps Institution of Oceanography, University of California at San Diego, 9500 Gilman Drive, La Jolla, CA 92093-0204 (USA)

[‡]Trius Therapeutics, 6310 Nancy Ridge Dr., Ste. 105, San Diego, CA 92121 (USA)

Keywords

Anthrax; antibiotics; *Bacillus anthracis*; polyketides; marine-derived *Streptomyces* sp

The human infectious disease Anthrax is caused by the spore-forming, Gram-positive bacterium *Bacillus anthracis*. The disease, which is most common in those that handle infected farm animals, has also been used as a bioterrorism weapon. In 2001, just a week after the World Trade Center attacks, Anthrax was deliberately spread through the US postal system by sending letters with powdered *B. anthracis* spores. This caused 22 cases of Anthrax infection and ultimately claimed five lives.^[1] Depending upon the method of exposure (inhalation of spores, direct bacterial contact, etc.) *B. anthracis* infections can require prolonged treatment often for 6 months with a variety of antibiotics. The pulmonary form of Anthrax is considered a medical emergency that may require continuous intravenous therapy with potent antibiotics. In a potential bioterrorism attack, individuals exposed to *B. anthracis* will be given antibiotics prior to onset of the illness. A vaccine has been developed but is not yet available to the general public.^[2] Given the severity of this disease, and the fact that it can be spread by aerosol dispersal, the development of an effective arsenal of new antibiotics remains a high public priority.

As a new, developing resource for microbial antibiotics, we have focused our efforts on marine microorganisms, particularly those that are in the deep oceans and specific to the marine environment. Examination of a *Streptomyces* species (our strain CNH365), isolated from near-shore marine sediments found near Santa Barbara, CA, showed that culture extracts possessed significant activity against *B. anthracis* and methicillin-resistant *Staphylococcus aureus* in broth dilution assays. Subsequent fractionation of the extract following antibacterial activities yielded a pure antibiotic, anthracimycin (**1**). Anthracimycin was isolated as a white solid that analyzed for the molecular formula C₂₅H₃₂O₄ by HR-ESI-MS analysis (−0.2 mmu). The ¹³C NMR spectrum of **1** (Table 1) displayed what appeared to be two ketone functionalities, one ester or lactone group (δ 194.1, 190.9, and 168.9), nine methine carbon signals in the aromatic/olefinic region (δ 140 ~ 103), and an oxymethine

**This work was supported by the NIH (NIGMS) under grant 1RO1 GM084350, and in part by the Transformational Medical Technologies program contract HDTRA1-10-C-004 from the Department of Defense Chemical and Biological Defense Program through the Defense Threat Reduction Agency (DTRA). The authors thank Drs. Arnold Rheingold and Curtiss Moore (UCSD) for performing X-ray diffraction analysis.

*Fax: (+1) 858-534-1318, wfenical@ucsd.edu.

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.xxxxxxx>.

group (δ_C 70.0). Analysis of gCOSY NMR data for the olefinic protons at δ_H 6.5~5.4 revealed the presence of a conjugated diene, an isolated *Z*-olefin and two trisubstituted double bonds. These spectroscopic data, taking into consideration the ten degrees of unsaturation from the molecular formula, revealed that anthracimycin was a tricarboxylic metabolite.

Clean separation of the signals in the 1H NMR spectrum allowed identification of the main structural fragments by analysis of strong vicinal and weaker long-range correlations in the gCOSY NMR experiment. All gCOSY derived connectivities within the structural fragments were also supported by gHMBC data (Table 1). The interpretation of gCOSY, gHSQC, and gHMBC NMR data allowed a 14-membered macrolide ring containing an enolized α -diketone and a lactone to be identified and confirmed. The methyl proton signal at δ_H 1.39 (CH₃-23) exhibited long-range correlations with two carbonyl carbon signals δ_C 168.9 (C-1) and 190.9 (C-3), and a methine carbon signal δ_C 49.2 (CH-2). Additionally, H-21 (δ_H 5.57) correlated to the lactone carbonyl (C-1, δ_C 168.9), confirming the lactone macrolide linkage.

Anthracimycin (**1**) possesses asymmetric carbon centers at C-2, C-6, C-7, C-12, C-15, C-16, and C-21, the relative configurations of which were assigned on the basis of ROESY experiments and proton coupling constant analysis. Interpretation of 2D ROESY NMR cross peaks for H-12/H-8 and H-12/H-6, allowed these axial protons to be positioned on the bottom face of the bicyclic system, whereas correlations of H-25/H-7, H-25/H-8, and H-25/H-11 oriented these protons on the top face. These results coincided well with the coupling constants between key protons ($J_{6,7} = 11.8$ Hz, $J_{6,15} = 6.7$ Hz, $J_{7,8} = 10.5$ Hz, $J_{7,12} = 14.5$ Hz, $J_{11,12} = 10.3$ Hz), and allowed the assignment of a *cis* orientation for the A/B ring juncture and a *trans* orientation for the B/C ring juncture. Analysis of 1H NMR data, allowed a *cis* double bond to be placed at the C-17 - C-18 position, and a *trans* double bond at C-19 - C-20. These assignments were made based upon analysis of the vicinal coupling constants of 10.5 and 15.2 Hz, respectively. These assignments were also supported by an observed NOE correlation between H-16 and H-19. NMR data, however, could not be utilized to define the relative configurations of the adjacent C-2 and C-21 methyl groups.

The full structure and absolute configurations at all asymmetric carbon centers of **1** were ultimately assigned based upon X-ray crystallographic data. X-ray quality crystals were obtained from slow recrystallization of anthracimycin from CH₂Cl₂/MeOH/Et₂O solutions. The X-ray experiment defined the absolute configurations of the stereocenters in **1** as 2*R*, 6*R*, 7*S*, 12*S*, 15*R*, 16*R*, and 21*R* (Figure 1 and supporting information). In addition, the X-ray experiment indicated significant disorder in the location of the enol proton of the enolized α -diketone. Thus, while NMR data have not been observed to confirm keto-enol tautomerization, X-ray data showed the proton to be virtually equidistant between each oxygen atom, thus suggesting rapid keto-enol tautomerization.

The structure of anthracimycin is composed of a rare combination of a 14, 6 and 6-membered rings of an apparent polyketide origin. Searching the scientific literature does not result in the discovery of journal publications on this structure. A patent in 2011, however was discovered that shows the planar structure of a compound, similar or possibly identical to **1**, devoid of stereochemical details and not illustrating the keto-enol tautomerization.^[3] Because the full stereostructure of this compound was not defined and not peer reviewed, we cannot confidently compare these molecules.

While this class of compounds is extremely rare, a tricyclic metabolite with a similar carbon skeleton is found in chlorotonil (**2**), a rare metabolite produced by the unrelated myxobacterium *Sorangium cellulosum*.^[4] Chlorotonil differs from **1** by the presence of an additional methyl group at C-8, it possesses a gem-dichloro functionality at C-4 and is enantiomeric at most relevant stereocenters. The synthesis of this unique metabolite was also reported^[5] but no mention of bioactivity has appeared.

Anthracimycin is a potent antibiotic against *Bacillus anthracis* (strain UM23C1-1) showing MIC = 0.031 $\mu\text{g/mL}$ in a broth microdilution assay.^[6] The antimicrobial spectrum of anthracimycin also includes Gram-positive genera such as staphylococci, enterococci, and streptococci, but it either lacked activity or was weakly active against the Gram-negative species tested. (Table 2). While the mechanism of action remains to be fully defined, it has been observed that the compound inhibits DNA/RNA synthesis in metabolic labeling experiments.^[7] Follow up gel-based studies demonstrated that the nucleic acid inhibition does not likely occur through DNA intercalation.^[7] Despite a large MIC shift in the presence of mouse serum, early *in vivo* results from MRSA infected CD1 mice showed that **1** provided significant protection (90% survival) at 10 mpK.^[7] A more thorough examination of the efficacy of **1** in whole animal models will clearly be important.^[8]

To predict the bioactivity impact of chlorination as found in **2**, we prepared the analogous dichloro-anthracimycin derivative **3** by treatment of **1** with *N*-chlorosuccinimide in dichloromethane (see supporting information). Comparison bioassays conducted with **1** and **3** showed that the dichloro derivative was still significantly active, but had lost about half of its potency against *B. anthracis*. Interestingly, the potency of **3** was identical to **1** against *Streptococcus pneumoniae*, and was more active against the Gramnegative pathogens *Haemophilus influenzae*, *Burkholderia thailandensis*, *E. coli*, and *Pseudomonas aeruginosa*, perhaps indicating a greater ability to penetrate the gram-negative cell wall.

Anthracimycin is a structurally-unique compound that does not compare with any reported antibacterial natural products. Although somewhat similar in structure to chlorotonil (**2**), anthracimycin is a potent antibacterial metabolite with potential in the treatment of Gram-positive pathogens such as *Bacillus anthracis* and methicillin-resistant *Staphylococcus aureus* (MRSA). That the compound shows promising preliminary activity *in vivo* (MRSA)^[7] and that the dichloro derivative **3** has surprising activity against problematic Gram-negative pathogens, suggests that there could be significant potential utility found in this new antibacterial structure class.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

References

- [1]. As defined by the US Center for Disease Control, see: <http://www.bt.cdc.gov/agent/anthrax/needtoknow.asp>
- [2]. See Medicine Net website: <http://www.medicinenet.com/anthrax>
- [3]. Igarashi Y, Iida T, Miyanochi K, Sudo Y. PCT/JP 2011/010586.
- [4]. Gerth K, Steinmetz H, Höfle G, Jansen R. Angew. Chem. Int. Ed. 2008; 47:600–602.
- [5]. Rahn N, Kalesse M. Angew. Chem. Int. Ed. 2008; 47:597–599.
- [6]. Clinical and Laboratory Standards Institute (CLSI). Methods for dilution antimicrobial susceptibility tests for bacteria that grow aerobically; approved standard – ninth edition, M07-A9. CLSI; Wayne, PA: 2012.
- [7]. Hensler, M.; Soubih, E.; Jang, KH.; Thienphrapa, W.; Locke, J.; Cunningham, M.; Kwan, B.; Nam, S-J.; Kauffman, C.; Beatty, D.; Shaw, KJ.; Nizet, V.; Fenical, W. Poster presentation # F-1514 at the Annual ICAAC Meeting in San Francisco; September, 2012;
- [8]. Comprehensive *in vivo* experiments, currently in progress, will be published elsewhere.

Figure 1.
X-ray drawing of anthracimycin illustrating its absolute configuration.

Table 1NMR Assignments for Anthracimycin (**1**) in CDCl₃.^[a]

No.	^1H (J in Hz)	c	HMBC
1		168.9	
2	3.53, q, (7.0)	49.2	C-3,23
3		190.9	
4	5.96, s	103.0	C-2, 3, 5, 6
5		194.1	
6	2.58, dd (11.8, 6.7)	52.6	C-4, 5, 12, 15, 16
7	1.98, ddd (11.8, 12.5, 14.5)	37.4	C-9
8	1.52, ddd (4.5, 10.5, 16.0)	31.4	C-10
8	2.39, ddd (16.0, 4.5, 4.5)		
9	5.36, br d (4.5)	121.0	C-7, 11, 24
10		134.0	
11	2.02, dd (16.5, 4.0)	37.5	C-9, 24
11	1.82, dd (16.5, 10.3)		
12	2.64 ^[b]	33.0	C-6, 10, 14
13	5.71, d (10.0)	133.0	C-15
14	5.53, dd (10.0, 5.0)	124.9	C-12, 15
15	2.60 ^[b]	46.0	C-6, 13, 14, 16, 17
16	1.93, m	32.8	C-6, 15, 17, 18, 25
17	5.40, dd (10.5, 9.8)	139.1	C-15, 16, 19
18	5.87, dd (10.5, 11.0)	126.1	C-16, 19, 20
19	6.45, dd (15.2, 11.0)	123.7	C-17, 18, 20, 21
20	5.56, dd (15.2, 2.4)	131.7	C-18, 19, 21, 22
21	5.57, dq (2.4, 6.5)	70.0	C-1, 19, 20, 22
22	1.33, d (6.5)	21.0	C-20, 21
23	1.39, d (7.0)	11.8	C-1, 2, 3
24	1.67, s	23.5	C-9, 10, 11
25	0.94, d (7.0)	16.4	C-16, 17

^[a]500 MHz for ¹H NMR and 125 MHz for ¹³C NMR.^[b]The coupling constant could not be measured because of overlapping signals.

Table 2

In vitro antibacterial activities of **1** and **3** against a diversity of gram-positive and gram-negative pathogenic bacteria.

Pathogen	Strain	1, MIC (µg/ml)	3, MIC (µg/ml)
<i>B. anthracis</i>	UM23C1-1	0.03125	0.0625
<i>S. aureus</i>	ATCC 13709	0.0625	0.125
<i>E. faecalis</i>	ATCC 29212	0.125	0.5
<i>S. pneumoniae</i>	ATCC 51916	0.25	0.25
<i>E. coli</i>	MCR106 <i>imp</i>	>128	16
<i>E. coli</i>	MG1655 <i>tolC</i>	>128	>128
<i>H. influenzae</i>	ATCC 31517	>256	32
<i>H. influenzae</i>	ATCC 31517 KO	4	4
<i>Burkholderia thailandensis</i>	E264 KO	>256	32
<i>Pseudomonas aeruginosa</i>	PAO1 KO	>256	32