CASE REPORT

Huge undifferentiated fibroblastic sarcoma of the foot and lower leg: impact of diagnostic delay and discussion of an entity

Cornelia S L Müller,¹ Claudia Pföhler,¹ Dieter Kohn,¹ Thomas Vogt¹

¹Department of Dermatology, Saarland University, Homburg/Saar, Germany

²Department of Orthopaedics and Orthopaedic Surgery, Saarland University, Homburg, Germany

Correspondence to

Dr Cornelia S L Müller, c_mueller1977@yahoo.de, cornelia.mueller@uks.eu

SUMMARY

We report a case of a 40-year-old female patient with a huge fibroblastic differentiated sarcoma of the left foot. After sampling all clinical, pathological and imaging information, the case was presented to the interdisciplinary tumour conference. Further treatment and medical care took place in the department of orthopaedics and orthopaedic surgery of our university hospital. Treatment consisted of amputation of the left foot and lower leg as an extremity-preserving surgical procedure was not possible because clear surgical margins could not be achieved with preservation of a functional rest of ankle and foot. Till date, no metastases have occurred.

BACKGROUND

The presented case should sensitise clinicians to possible exceptional settings of clinical symptoms in non-typical age categories. It is mandatory to have a high degree of suspicion in cases of unilateral, non-healing, tumorous tissue proliferations and perform early biopsies to avoid fatal diagnostic delay with severe outcomes for the patient. Herein, we discussed the role of diagnostic delay, clinical and histological differentials and the entity of undifferentiated fibroblastic sarcomas.

This case is worth conferring for the following three reasons:

1. A prolonged delay in diagnosis and need for ablation of the foot and lower leg due to previous false diagnoses and a neglect of histological examination;
2. The clinical differential diagnoses of the swollen leg must be elucidated;
3. With the advent of modern immunopathological, molecular diagnostics and molecular-based therapy, the entity of undifferentiated fibroblastic soft tissue tumours must be critically discussed.

CASE PRESENTATION

A 40-year-old female patient presented with a significant swelling of the left foot that began over the left ankle and extended to the interphalangeal spaces of nearly all the toes with extension to the plantar surface of the foot. The skin above the swelling was tense and had a yellowish ulceration for several weeks (figure 1). The tumour itself was compact and displayed a doughy consistency. The swelling appeared approximately 3 years ago and grew slowly, but steadily for years. A few weeks prior to a dermatological consultation, the left foot

and lower leg caused severe pain, especially when at rest. The mobility of the ankle joint was limited to dorsal extension-plantar flexion of 10°. There were no sensomotoric deficits. A fracture of the left ankle with Volkmann's triangle 16 years ago healed without complications. The patient did not have any related diseases. Of note, the patient has been under the care of a family physician continuously, as well as specialists in internal medicine, immunology and orthopaedics through all the time. The clinical differential diagnoses and previous working diagnoses included chronic lymphatic oedema and erysipelas lacking typical clinical and laboratory signs of inflammation (dolor, calor and redness). Systemic and topical glucosteroids, antiphlogistics, lymphatic drainage or physiotherapy did not lead to a substantial improvement in symptoms.

In 2011, the patient presented to the outpatient service of our dermatology department. We obtained MRI and performed multiple skin biopsies of the tumour. A coronal (figure 2A) and axial (figure 2B)

Figure 1 Clinical appearance after first dermatological consultation displaying the immense tumorous proliferation involving the entire foot, inclusive of the plantar surface and lower leg.

To cite: Müller CSL, Pföhler C, Kohn D, et al. *BMJ Case Rep* Published online: [please include Day Month Year] doi:10.1136/bcr-2013-200192

Figure 2 Coronal (A) and axial (B) T1-weighted (T1W) MRI demonstrate a hypointense tumour, diffusely infiltrating the soft tissue at the metacarpal level, with an extension from the dorsal surface down to the plantar region (B). Postcontrast medium injection (C) the T1W fat suppressed coronal image shows inhomogeneous contrast medium uptake of the tumour without evidence of larger necrotic areas.

T1-weighted (T1W) MRI demonstrated a hypointense tumour, diffusely infiltrating the soft tissue at the metacarpal level, with an extension from the dorsal surface down to the plantar region (figure 2B). Postcontrast medium injection (figure 2C) the T1W fat suppressed coronal image shows inhomogeneous contrast medium uptake of the tumour without evidence of larger necrotic areas.

Several specimens obtained from the back of the foot displayed a malignant tumour composed of spindle cells in a myxoid stroma (figure 3A–C). A three-phase bone scintigraphy was compatible with a soft tissue tumour of the left foot without any evidence for osseous metastases. Finally, a low grade, diffuse, proliferating, fibroblastic differentiated sarcoma was diagnosed.

TREATMENT

After sampling all clinical, pathological and imaging information, the case was presented to the interdisciplinary tumour conference in our university clinic. Further treatment and medical

care took place in the department of orthopaedics and orthopaedic surgery of our university hospital. Treatment consisted of amputation of the left foot and lower leg as an extremity-preserving surgical procedure was not possible because clear surgical margins could not be achieved with preservation of a functional rest of ankle and foot (figure 4).

OUTCOME AND FOLLOW-UP

Till date, no metastases have occurred.

DISCUSSION

Diagnostic delay

Typically, such ‘giant’ tumours are noted in people that are very old and suffer from psychiatric and psychosocial comorbidities, such as dementia. In cases of giant melanoma, especially patients with depression, the prognosis is worse because of a delayed diagnosis.¹ Thick melanomas are relatively common, but giant melanomas covering entire parts of the body are rarely observed.

Figure 3 (A) and (B) Histopathology reveals a fibroblastic-differentiated spindle cells (H&E stain). (C) Mucin stain highlighting the myxoid areas.

These patients are at high risk for metastatic disease.² In cases of breast cancer, several studies have been conducted to examine the delay in diagnosis of breast cancer, especially in young females. Andrews and Bates presented these findings in 2000 for breast cancer, with special emphasis on medicolegal implications. Although patient delay is generally associated with more advanced lesions at the time of presentation, the effect of delay on survival or the length of delay which is clinically significant is uncertain.^{3 4} In addition, publications on delay in diagnosis of oral or lung cancer exist.^{5 6} Soft tissue sarcoma patients primarily consult physicians in advanced stages of the disease due to missing clinical symptoms or an inability to recognise these tumours that are often located within the body cavities. Nakamura *et al*⁷ reported that 18% of the sarcoma patients investigated in their study had distant metastases at the time of

diagnosis. They showed that tumour size, tumour site and the interval between the onset of the initial symptoms and the first medical consultation to the hospital were all found to be significant predictors of distant metastases at diagnosis. Finally, Nakamura *et al*⁷ concluded that “a shorter delay may have a beneficial effect on treatment options and outcome, improving survival in some sarcoma patients.”

The follow-up period for the patient presented herein has been too short and the outcome has to be monitored in short intervals. Patients, especially those at a younger age, are often in ongoing multidisciplinary medical care by various disciplines due to a sensitive subjective health perception. Physicians of all disciplines should therefore be aware of the possibility of the occurrence of even rare tumours particularly in the age group.

Figure 4 Macroscopy of the amputate for pathological examination.

In the case presented herein, the possibility of a malignant soft tissue tumour was neglected by all parties involved due to the young age of the patient, thus leading to a diagnostic delay of several years. Provided that tumour size and the interval between the onset of the initial symptom and the first medical consultation serve as significant predictors of survival (as proposed by Nakamura *et al*⁷), this almost certainly would have had an immense impact for the patient. Apart from these thoughts, an earlier diagnosis would have probably protected the patient from extensive ablative surgery. As a rule, we would suggest that any swelling of soft parts being present far more than 4 weeks and being not otherwise explained do need histological clarification with no exception.

Clinical differential diagnoses

The major clinical differential diagnoses of a unilateral swollen leg can be clinically differentiated by using three leading symptoms (acute and chronically swollen leg and a swollen leg without associated oedema) or differentiated by affected skin colour (see tables 1 and 2).

Swollen legs are frequently encountered in daily routine. Manifold possible causes of swollen legs are known, and in many cases, the clinician is faced with a challenging scenario. Important points are the history of the symptom (swollen leg), distribution, course, associated skin lesions. In addition, previous trauma, surgery, radiation therapies, infections and thrombosis have to be excluded.⁸⁻⁹ Primarily inspection of both legs is mandatory. Depending on associated symptoms, such as fever or pain, laboratory investigations and duplex sonography should be added. In all cases of lymphatic stasis, repression of the lymphatic vessels by malignant tumors must be considered.¹⁰ In all cases that cannot be clearly clarified, biopsies in the early stages of the disease are mandatory.

Histomorphology revisited

On the basis of the histopathological evaluation, the fibroblastic sarcoma was a diagnostic and nosological challenge. In the broadest sense, this diagnosis was characterised by the exclusion of distinct sarcomatous entities. Conventional histomorphology is characterised by a dermal and subcutaneous cell-rich, spindle-shaped, mesenchymal tumour that is focally composed of fascicles of mesenchymal cells with pale eosinophilic cytoplasm. Mitotic figures were observed consistently. Focal sclerosis and hyalinisation was observed due to the long standing, aged lesion. The immunohistochemistry features were unremarkable. Not surprisingly, the

Table 2 Differential diagnoses of swollen legs differentiated by affected skin colour

Swelling with changes in skin colour	Swelling without changes in skin colour
▶ Arthritis ▶ Acute dermatitis ▶ Infections (bacterial or fungal) ▶ Eosinophilic fasciitis ▶ Necrotising fasciitis ▶ Arthropod bites ▶ Oedema in paralysed extremities ▶ Post-thrombotic syndrome ▶ Malignancies	▶ Cardiac insufficiency ▶ Primary or secondary lymphatic oedema (Nonne-Milroy's type); can be associated with a marked papillomatosis cutis lymphostatica ▶ Sudeck's syndrome ▶ Podagra ▶ Malignancies

expression of vimentin and smooth muscle actin was observed, while all other differentiating markers of melanocytic, epithelial or vascular differentiation stained negatively. Of interest are the myxoid areas. An inflammatory infiltrate was absent.

Molecular studies failed to demonstrate fused in sarcoma (FUS) translocation which would have confirmed the low-grade fibromyxoid sarcoma.

According to these immunohistological and molecular features, we diagnosed a low-grade undifferentiated fibroblastic sarcoma. Less data exist due to the rarity of the tumour and inconsistent nomenclature of these soft tissue sarcomas.

In our view, the tumour was most likely comparable to the low-grade fibromyxoid sarcoma and adult-type fibrosarcoma, displaying a morphological spectrum of sarcoma entities, with undifferentiated subentities at the cutting edges. On a molecular basis, these sarcomas belong to the group of soft tissue tumours with simple genomic aberrations and specific translocations.¹¹ For the low-grade fibromyxoid sarcoma specific translocation of t(7;16)(q33;p11.2) with formation of the fusion protein FUS-CREB3L2 and for superficial examples of the adult-type fibrosarcoma. COL1A1-PDGFB fusion transcripts such as those occurring in dermatofibrosarcoma protuberans are known.¹¹

None of these molecular changes existed in the case presented here. Hence, the diagnosis depended on classic morphology and exclusion of other tumours. Histopathological differential diagnoses include low-grade myofibrosarcomas, synovial sarcomas, malignant peripheral nerve sheath tumours, myxoinflammatory fibroblastic sarcomas, myxoid leiomyosarcomas and dedifferentiated/pleomorphic liposarcomas.¹²⁻¹³

Table 1 Differential diagnoses of swollen legs according to leading symptoms (acute and chronically swollen leg and a swollen leg without associated oedema)

Leading symptom	Examples
Acute, unilateral swollen leg	▶ Arterial closure ▶ Erysipelas ▶ Phlebothrombosis ▶ Traumata
Chronic, unilateral swollen leg	▶ Peripheral artery occlusive disease ▶ Chronic venous insufficiency ▶ Post-thrombotic syndrome ▶ Arthrogenic stasis
Swollen leg without associated oedema	▶ Angiodysplasia ▶ Lipedema ▶ Lipomatosis (dolorosa)

Learning points

- ▶ Any swellings of soft parts being present far more than 4 weeks and being not otherwise explained do need histological clarification with no exception.
- ▶ The major clinical differential diagnoses of a unilateral swollen leg can be clinically differentiated by using the three leading symptoms: acute and chronically swollen leg and a swollen leg without associated oedema or differentiated by affected skin colour.
- ▶ In conclusion, the presented case should sensitise clinicians for possible exceptional settings of clinical symptoms in non-typical age categories.

Acknowledgements The authors would like to thank Anne Kerber for excellent techniques in histological and immunohistological staining. They also thank the extraordinary collaboration with the Praxis für Dermatopathologie in Friedrichshafen, Germany for performing molecular investigations and reference pathology.

Contributors CSLM wrote major parts of the manuscript and prepared the pathology report. CP and TV cared for the patient in the outpatient unit and contributed to the manuscript. DK cared for the patient in orthopaedic department and performed amputation of the leg and also contributed to the manuscript.

Competing interests None.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- del Boz J, García JM, Martínez S, *et al.* Giant melanoma and depression. *Am J Clin Dermatol* 2009;10:419–20.
- Grisham AD. Giant melanoma: novel problem, same approach. *South Med J* 2010;103:1161–2.
- Tartter PJ, Pace D, Frost M, *et al.* Delay in diagnosis of breast cancer. *Ann Surg* 1999;229:91–6.
- Andrews BT, Bates T. Delay in the diagnosis of breast cancer: medico-legal implications. *Breast* 2000;9:223–37.
- Gomez I, Warnakulasuriya S, Varela-Centelles PI, *et al.* Is early diagnosis of oral cancer a feasible objective? Who is to blame for diagnostic delay? *Oral Dis* 2010;16:333–42.
- Valdes S, Garcia E, Perez H, *et al.* Length of diagnostic delay in patients with non-small-cell lung cancer. *MEDICC Rev* 2010;12:29–32.
- Nakamura T, Matsumine A, Matsubara T, *et al.* The symptom-to-diagnosis delay in soft tissue sarcoma influence the overall survival and the development of distant metastasis. *J Surg Oncol* 2011;104:771–5.
- Khandker SR. Images in emergency medicine. A case of a painful swollen leg. Phlegmasia cerulea dolens. *Ann Emerg Med* 2012;59:176, 208.
- Tomlow B, Rellum M. [A woman with painful, swollen legs]. *Ned Tijdschr Geneesk* 2011;155:A1336.
- Altmeyer P. *Dermatologische Differenzialdiagnose. Der Weg zur klinischen Diagnose.* Heidelberg: Springer, 2007.
- Taylor BS, Barretina J, Maki RG, *et al.* Advances in sarcoma genomics and new therapeutic targets. *Nat Rev Cancer* 2011;11:541–57.
- Mentzel T, Calonje E, Wadden C, *et al.* Myxofibrosarcoma. Clinicopathologic analysis of 75 cases with emphasis on the low-grade variant. *Am J Surg Pathol* 1996;20:391–405.
- Mansoor A, White CR Jr. Myxofibrosarcoma presenting in the skin: clinicopathological features and differential diagnosis with cutaneous myxoid neoplasms. *Am J Dermatopathol* 2003;25:281–6.

Copyright 2013 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow