CORRESPONDENCE

Fluid Replacement With Hydroxyethyl Starch in Critical Care—a Reassessment

by PD Dr. med. Christiane S. Hartog, Prof. Dr. med. Tobias Welte,
Prof. Dr. med. MSc Peter Schlattmann, Prof. Dr. med. Konrad Reinhart
in volume 26/2013

Methodological Issues of the Studies Cited

The doubts raised concerning the use of hydroxyethyl starch (HES) for fluid replacement in critical care (1) rely almost exclusively on data from the 6S (2) and CHEST studies (3). In both studies, central venous pressure, central venous saturation, urine output, and serum lactate concentrations were all within the ranges recommended by the Surviving Sepsis Campaign for the initial therapy of sepsis (4). This indicates that at the time of randomization these patients were already hemodynamically stabilized and normovolemic. This would not be surprising because before randomization, patients in the 6S study (2) had received on average 3.0–3.5 L of fluids, and 42% of them had received colloids. Consequently, in both studies fluid replacement with HES was contraindicated. In neither of the two studies was fluid administration standardized or goal-directed, nor were criteria for renal replacement therapy defined *a priori*, even though this was a secondary outcome parameter. Furthermore, the CHEST study data (3) do not provide evidence of a statistically significant increase in renal replacement therapy (relative risk [RR], 1.20; 95% confidence interval [CI] 1.00 to 1.44; $p = 0.05$), and based on some of the RIFLE criteria, HES administration was even associated with less renal injury.

It is entirely incomprehensible why in the CHEST study (3) two hyperchloremic solutions were used for fluid replacement in critical care, although chloride-induced renal vasoconstriction has been known for a long time. Even in healthy individuals, the infusion of saline may result in a 40% decrease in renal perfusion (5). For these (and other unmentioned) considerable methodological issues, no conclusions can be drawn from the results of the two studies regarding the efficacy of balanced HES solutions, especially when it comes to hypovolemic, hemodynamically unstable patients in whom HES solutions are indicated.

DOI: 10.3238/arztebl.2013.0735a

REFERENCES

1. Hartog CS, Welte T, Schlattmann P, Reinhart K: Fluid replacement with hydroxyethyl starch in critical care—a reassessment. *Dtsch Arztebl Int* 2013; 110(26): 443–50.
2. Perner A, Haase N, Guttormsen AB, et al.: Hydroxyethyl starch 130/0.42 versus Ringer's acetate in severe sepsis. *N Engl J Med* 2012; 367: 124–34.
3. Myburgh JA, Finfer S, Bellomo R, et al.: Hydroxyethyl starch or saline for fluid resuscitation in intensive care. *N Engl J Med* 2012; 367: 1901–11.

4. Chowdhury AH, Cox EF, Francis ST, Lobo DN: A randomized, controlled, double-blind crossover study on the effects of 2-L infusions of 0.9% saline and Plasma-Lyte(R) 148 on renal blood flow velocity and renal cortical tissue perfusion in healthy volunteers. *Ann Surg* 2012; 256: 18–24.

Prof. em. Dr. med. Hans-Joachim Priebe
Merzhausen
hans-joachim.priebe@uniklinik-freiburg.de

Studies Are Carried out in Hospitals

Apart from 6% HES in balanced electrolyte solutions, “Hyperhes” is available, a preparation containing 6% HES in 7.2% hypertonic saline.

This infusion solution has certainly saved the lives of many patients, when given as a concentrate for fluid resuscitation.

Freshly polytraumatized patients are at a high risk of bleeding to death at the accident site and the first blood unit only becomes available after arrival in the emergency trauma room. Until that point in time, only Hyperhes, HES, gelatin or crystalloids are at hand, and the best fluid replacement effect is achieved with HES solutions.

Should HES be really taken off the market, we will definitely miss it at the actual accident site, but these studies are undertaken in hospitals, and rarely in out-of-hospital situations.

DOI: 10.3238/arztebl.2013.0735b

REFERENCES

1. Hartog CS, Welte T, Schlattmann P, Reinhart K: Fluid replacement with hydroxyethyl starch in critical care—a reassessment. *Dtsch Arztebl Int* 2013; 110(26): 443–50.

Mark Schorr
Anästhesist, Honorararzt
Pirmasens
docteurmark@me.com

In Reply:

Here our reply to our esteemed colleague Prof. Priebe: The serious concerns expressed by the regulatory authorities were not only triggered by 2 large studies, but also by the meta-analysis of Zarychanski et al. (1), revealing a significantly increased mortality and a significantly increased need for renal replacement therapy in HES-treated patients compared with patients treated with crystalloid solutions, based on 31 studies with altogether 10 290 patients. A recent Cochrane review about renal failure, which we could not take into account in our paper, includes 42 studies with 11 399 patients. The authors arrived at the conclusion that “it is likely that these risks outweigh any benefits [of HES]” and that thus “alternate volume replacement therapies should be used in place of HES products” (2).

In patients with severe sepsis, volume deficits are not limited to the initial phase, but often persist for two to three days because of fluid losses into the tissue (“capillary leak”). It is not unusual to find a positive fluid balance of 10–20 and more liters in these patients during the acute phase of sepsis, regardless of the type of fluid used. Apart from the statistically significant negative effects of HES in the CHEST study, it should also be considered that adverse events observed in the HES group occurred with a cumulative dose of 5 mL/kg, i.e. at a dosage far less than half of the presently approved daily dose. The use of hyperchloremic solutions in both arms of a blinded, randomized study cannot explain the adverse events in one of these groups.

For Mr. Schorr, we would like to highlight two important studies conducted in out-of-hospital environment: In an out-of-hospital setting, 1282 patients with traumatic brain injury not accompanied by hypovolemic shock (3) and 853 severely injured trauma patients with hypovolemic shock (4) received either 0.9 %, 7.5% saline or 7.5% saline/6% dextran-70. Having met the futility criteria, the first study was terminated early, because the 6-month neurologic outcome and 28-day mortality (75.1%, 75.7 % and 74.3 %, respectively) were almost identical (3). Likewise, the second study was discontinued because of comparable 28-day mortality; in addition, safety concerns were raised, since among patients who received hypertonic fluids and no blood transfusions a fatal outcome was observed about twice as often as in patients who received

isotonic saline (4). Therefore, the current German poly-trauma guidelines recommend crystalloids as first-line treatment. Having a similar risk profile as HES, gelatin solutions should no longer be used either. They prolong the bleeding time and are therefore no longer approved for sale in the USA.

DOI: 10.3238/arztebl.2013.0735c

REFERENCES

1. Zarychanski R, Abou-Setta AM, Turgeon AF, et al.: Association of hydroxyethyl starch administration with mortality and acute kidney injury in critically ill patients requiring volume resuscitation: a systematic review and meta-analysis. *JAMA* 2013; 309: 678–88.
2. Mutter TC, Ruth CA, Dart AB: Hydroxyethyl starch (HES) versus other fluid therapies: effects on kidney function. *Cochrane Database Syst Rev* 2013; 7(CD007594).
3. Bulger EM, May S, Brasel KJ, et al.: Out-of-hospital hypertonic resuscitation following severe traumatic brain injury: a randomized controlled trial. *JAMA* 2010; 304: 1455–64.
4. Bulger EM, May S, Kerby JD, et al.: Out-of-hospital hypertonic resuscitation after traumatic hypovolemic shock: a randomized, placebo controlled trial. *Annals of Surgery* 2010; 253: 431–41.
5. Hartog CS, Welte T, Schlattmann P, Reinhart K: Fluid replacement with hydroxyethyl starch in critical care—a reassessment. *Dtsch Arztebl Int* 2013; 110(26): 443–50.

Prof. Dr. med. Konrad Reinhart

Klinik für Anästhesiologie und Intensivmedizin
Universitätsklinikum Jena
Konrad.Reinhart@med.uni-jena.de

Conflict of interest statement

The authors of all contributions declare that no conflict of interest exists.