Published in final edited form as:

Biochem Biophys Res Commun. 2012 October 12; 427(1): . doi:10.1016/j.bbrc.2012.09.021.

Recombinant expression of TLR5 proteins by ligand supplementation and a leucine-rich repeat hybrid technique

Minsun Hong^{#a}, Sung-il Yoon^{#a,b,*}, and Ian A. Wilson^{a,c,*}

^aDepartment of Molecular Biology, The Scripps Research Institute, La Jolla, CA 92037, USA

^bDepartment of Systems Immunology and Institute of Antibody Research, College of Biomedical Science, Kangwon National University, Chuncheon 200-701, Republic of Korea

^cThe Skaggs Institute for Chemical Biology, The Scripps Research Institute, La Jolla, CA 92037, USA

These authors contributed equally to this work.

Abstract

Vertebrate TLR5 directly binds bacterial flagellin proteins and activates innate immune responses against pathogenic flagellated bacteria. Structural and biochemical studies on the TLR5/flagellin interaction have been challenging due to the technical difficulty in obtaining active recombinant proteins of TLR5 ectodomain (TLR5-ECD). We recently succeeded in production of the N-terminal leucine rich repeats (LRRs) of *Danio rerio* (*dr*) TLR5-ECD in a hybrid with another LRR protein, hagfish variable lymphocyte receptor (VLR), and determined the crystal structure of its complex with flagellin D1–D2–D3 domains. Although the structure provides valuable information about the interaction, it remains to be revealed how the C-terminal region of TLR5-ECD contributes to the interaction. Here, we present two methods to obtain recombinant TLR5 proteins that contain the C-terminal region in a baculovirus expression system. First, production of biologically active full-length *dr*TLR5-ECD was substantially enhanced by supplementation of expression culture with purified flagellin proteins. Second, we designed TLR5-VLR hybrids using an LRR hybrid technology by single and double LRR fusions and were able to express diverse regions of *dr*TLR5-ECD, allowing us to detect a previously unidentified TLR5/flagellin interaction. The *dr*TLR5-VLR hybrid technique was also successfully applied to human TLR5-ECD whose expression has been highly problematic. These alternative TLR5 expression strategies provide an opportunity to obtain a complete view of the TLR5/flagellin interaction and can be applied to other LRR proteins.

Keywords

Toll-like receptor 5; Flagellin; Leucine-rich repeat; LRR hybrid; Variable lymphocyte receptor; Innate immunity

© 2012 Elsevier Inc. All rights reserved.

*Corresponding authors. Address: Department of Systems Immunology and Institute of Antibody Research, College of Biomedical Science, 1 Kangwondaehakgil, Biomedical Science Building A-204, Chuncheon 200-701, Republic of Korea. Fax: +82 33 250 8380 (S.-i. Yoon). Address; Department of Molecular Biology, The Scripps Research Institute, 10550 N. Torrey Pines Road, BCC-206, La Jolla, CA 92037, USA. Fax: +1 858 784 2980 (I.A. Wilson). sungil@kangwon.ac.kr (S.-i. Yoon), wilson@scripps.edu (I.A. Wilson).

Appendix A. Supplementary data Supplementary data associated with this article can be found, in the online version, at <http://dx.doi.org/10.1016/j.bbrc.2012.09.021>.

1. Introduction

Toll-like receptors (TLRs) play a key role in the first line of defense against pathogens by recognizing common molecular structures of pathogenic microbes and subsequently activating innate immune responses [1]. Ten TLRs (TLR1–TLR10) have been identified in humans and are classified into 5 subgroups (TLR1/2/6/10, TLR3, TLR4, TLR5, and TLR7/8/9) depending on the nature of their ligands [2].

TLR5 is found in vertebrates from fish to mammals and directly binds a bacterial protein, flagellin, which polymerizes into a flagellar filament in a spiral manner [3,4]. Flagellin-induced TLR5 signaling activates NF- κ B, a proinflammatory transcription factor, and promotes production of inflammatory cytokines for the activation of mucosal immunity against pathogenic flagellated bacteria in epithelial cells of the intestine and lung. TLR5 also exhibits anti-apoptotic radioprotective activity and protects the hematopoietic system and gastrointestinal tissues from radiation damages [5]. Thus, biochemical and structural studies on the TLR5/flagellin interaction is critical for the development and improvement of vaccines and vaccine adjuvants as well as therapeutics that would alleviate side effects of anticancer radiotherapy [5,6].

TLR5 is a type I receptor that consists of three domains including an N-terminal flagellin-binding ectodomain (ECD), a single-pass transmembrane domain, and an intracellular signaling domain. Biochemical and structural studies on the TLR5-ECD/flagellin interactions have been limited due to technical difficulty in obtaining TLR5-ECD proteins by conventional protein expression approaches. To overcome the expression challenges, we screened expression of diverse TLR5-ECD orthologs and found that only zebrafish (*Danio rerio*) TLR5-ECD (*dr*TLR5-ECD) proteins were produced in sufficient quantities in a baculovirus expression system [4]. However, the *dr*TLR5-ECD aggregated as a nonfunctional protein. To improve the TLR5-ECD protein behavior in solution, we utilized a leucine-rich repeat (LRR) hybrid technique that fuses two different LRR proteins, the TLR5 protein of interest with LRR repeats from a variable lymphocyte receptor (VLR), to produce TLR5-VLR hybrids [7–10]. As a result, several *dr*TLR5-VLR hybrid proteins were purified as functionally active monomers in substantially improved yields and used for structure determination of a complex between the N-terminal 14 LRR modules of *dr*TLR5 and the D1–D2–D3 domains of *Salmonella* flagellin, FliC [4]. The primary binding interaction is mainly mediated by a protruding loop of TLR5 LRR9 and the FliC D1 domain, contributing to formation of a high-affinity 1:1 TLR5/FliC heterodimer. Two copies of the heterodimers further assemble into a 2:2 complex using TLR5 LRR12–LRR13.

The structure provided important information about the TLR5/flagellin interaction at atomic resolution. Nevertheless, the complex lacks the C-terminal region of *dr*TLR5-ECD and the FliC D0 domain. Thus, it still remains to be revealed how the C-terminal region of TLR5-ECD contributes to TLR5/flagellin binding and TLR5 signaling. To provide a complete view on the interaction, we further screened and improved expression of *dr*TLR5-ECD and its fragments by ligand supplementation and an innovative protein engineering technique. First, we produced functional *dr*TLR5-ECD proteins in sufficient quantity for structural studies by adding exogenous purified FliC proteins into the *dr*TLR5-ECD expression culture. Second, we fused *dr*TLR5-ECD and hagfish VLR by single or double fusions and were able to express the C-terminal region of *dr*TLR5-ECD. Analogously, the *dr*TLR5-VLR hybridization strategy was successfully applied to expression and purification of human (*Homo sapiens*) TLR5-ECD (*hs*TLR5-ECD) fragments. These versatile approaches to recombinant TLR5 protein expression have already contributed to understanding of the structural and molecular mechanism for the TLR5/flagellin interaction and should enable

advances in protein preparation of other challenging LRR receptors for biochemical or biophysical studies.

2. Materials and methods

2.1. Expression and purification of FliC and CBLB502

FliC (residues 1–504) of *Salmonella enterica* subspecies *enterica* serovar Dublin and its variant CBLB502 (FliC residues 1–175, a 16-residue linker, and FliC residues 401–504) were expressed and purified as previously described [4,5]. FliC and CBLB502 were produced in an *Escherichia coli* expression system using a modified expression vector, pET49b, which contains an N-terminal His₆ tag and an enterokinase or thrombin cleavage site. FliC proteins were obtained from soluble fractions of cell lysates, whereas CBLB502 was refolded from inclusion bodies with 2 M urea. FliC and CBLB502 proteins were purified by Ni-NTA and size-exclusion chromatography.

2.2. Construction of TLR5-ECD and TLR5-VLR hybrid expression vectors

TLR5-ECD and TLR5-VLR hybrids were expressed in a baculovirus expression system. A baculovirus transfer vector, pAcGP67A, that allows recombinant proteins to be extracellularly expressed under the control of the GP67 signal peptide, was modified to contain the thrombin cleavage site, Strep-Tactin II tag, and His₆ tag at the C-terminus and was used throughout [11]. To construct full-length TLR5-ECD expression vector, DNAs that encode *dt*TLR5b residues 22–652 [UniProt ID F8W3J5] or *hs*TLR5 residues 21–639 (UniProt ID Q32MI2) (Fig. S1) were amplified by PCR and ligated into the modified pAcGP67A vector between the N-terminal GP67 signal peptide and the C-terminal thrombin cleavage site.

To generate TLR5-VLR hybrid protein expression constructs, DNAs that encode the TLR5-VLR fusion proteins were inserted into the modified pAcGP67A vector. We prepared numerous inserts that can be grouped into three different types including an N-terminal single fusion, a C-terminal single fusion, and a double fusion at both N- and C-terminal regions (Fig. 1). The first type of insert is a single-fusion hybrid T^N-x^V that contains the N-terminal fragment of TLR5-ECD (the N-terminal residue 22 to LRR module number x) and the C-terminal region of hagfish VLR B.61 (LRR4 to the residue 200). The C-terminal region of VLR was first amplified by PCR and the N-terminal part of TLR5 was then extended in the second PCR using the first PCR products as a reverse primer. The second type of insert is the C-terminal single fusion product V^T_x-C between the N-terminal region of VLR (the N-terminal residue 22 to LRR2) and the C-terminal fragment of TLR5-ECD (LRR module number x to the C-terminus). The inserts were created by two consecutive PCRs as described above for T^N-x^V . The third type of TLR5-VLR hybrids is a double fusion product $V^T_a-b^V$ where the central LRR fragments of TLR5-ECD from LRR 'a' to LRR 'b' is flanked by the N-terminal (the N-terminus to LRR2) and C-terminal (LRR4 to residue 200) regions of VLR. $V^T_a-b^V$ inserts were also made by two consecutive PCRs but with DNA templates of T^N-x^V or V^T_x-C for the second PCR.

2.3. Expression and purification of TLR5 proteins

TLR5-ECD and TLR5-VLR hybrids were expressed in a baculovirus expression system and purified by consecutive chromatography using Ni-NTA, Strep-Tactin II, and size-exclusion chromatography columns as previously described [4]. In some TLR5 constructs including TLR5-ECD and its VLR hybrid V^T_1-C , purified FliC proteins were added into expression culture at ~24 h postinfection at which proteins begin to be synthesized under a polyhedrin promoter of pAcGP67 expression vector and TLR5/FliC complexes were then purified in the same manner as TLR5-VLR hybrids.

2.4. Native PAGE

Binding of TLR5-VLR hybrids to FliC or CBLB502 was analyzed by native PAGE [11,12]. TLR-VLR hybrids, FliC (or CBLB502), and their complexes were loaded on 4–20% polyacrylamide gels and native PAGE was carried out without SDS in Tris–glycine buffer at pH 8.8 for 2–3 h at 100 V.

3. Results

3.1. Improvement in production of active drTLR5-ECD proteins by FliC supplementation

The major roadblock in *drTLR5*-ECD expression is that a majority of the recombinant *drTLR5*-ECD proteins are not properly folded, thereby forming aggregates by themselves or nonspecifically with cellular proteins resulting in non-functional proteins during expression even in a baculovirus expression system that uses eukaryotic insect cells. Only one third of the expressed proteins formed biologically active monomers by size-exclusion chromatography, even although the proteins had been previously purified by Ni-NTA and Strep-Tactin affinity chromatography (Fig. 2A). One litre expression culture provided a small quantity of purified *drTLR5*-ECD proteins ($<50 \mu\text{g}$ from 1 L expression culture) that was not sufficient for structural studies in stark contrasts to a TLR5-VLR hybrid, $^{\text{T}}\text{N-14}^{\text{V}}$, that was purified mainly as monomers (Fig. 2B) with a purification yield of $\sim 1.4 \text{ mg}$ per 1 L culture (Table S1).

Given that *drTLR5*-ECD was expressible, we searched for a method that would improve TLR5-ECD protein stability to enhance the final yield of active *drTLR5*-ECD proteins. For this purpose, we supplemented expression culture with purified FliC proteins at $\sim 24 \text{ h}$ postinfection. As a result, a peak that corresponds to *drTLR5*-ECD complexed with FliC became notably higher and larger in a FliC dose-dependent manner (Fig. 3). We obtained $\sim 0.4 \text{ mg}$ of *drTLR5*-ECD/FliC complex after three consecutive purification steps (Ni-NTA, Strep-Tactin, and size-exclusion chromatography) from 1 L culture when 0.6 mg of FliC was added as a supplement. We conclude that the purification yield increased by at least four-fold by FliC supplementation. This substantial improvement in production of active *drTLR5*-ECD proteins by FliC supplementation during culture enabled us to perform X-ray crystallographic studies on the TLR5/FliC complex, which required a large quantity of protein.

3.2. Expression of the drTLR5-ECD N-terminal fragments by a single LRR fusion

Both *drTLR5*-ECD and VLR adopt a typical LRR domain structure consisting of 22 and 5 LRR modules, respectively, and are protected by N- and C-terminal capping motifs (LRRNT and LRRCT, respectively). The LRRNT and LRRCT motifs are indispensable for stability of the LRR domains and are one of determining factors for levels of protein expression. We hypothesized that the capping motifs of TLR5 and their connecting regions are responsible for aggregation and low protein yields. To improve solubility and yields of *drTLR5*-ECD, we designed and screened *drTLR5*-VLR hybrids, in which LRRNT or LRRCT of TLR5 was swapped with that of a high expresser LRR protein, VLR, using one of the 2nd (leucine), 7th (leucine), and 10th (asparagine) residues of the LRR consensus sequence ($\text{xL}_2\text{xxL}_5\text{xL}_7\text{xxN}_{10}$) as a fusion point (Fig. 1). Such technique has been already utilized in other TLRs, including TLR4 and TLR1/2/6 subfamilies, and enhanced protein expression levels and crystallizability for structural studies [7–9,13].

First, we designed nine $^{\text{T}}\text{N-x}^{\text{V}}$ hybrids ($^{\text{T}}\text{N-4}^{\text{V}}$, $^{\text{T}}\text{N-6}^{\text{V}}$, $^{\text{T}}\text{N-8}^{\text{V}}$, $^{\text{T}}\text{N-10}^{\text{V}}$, $^{\text{T}}\text{N-12}^{\text{V}}$, $^{\text{T}}\text{N-14}^{\text{V}}$, $^{\text{T}}\text{N-17}^{\text{V}}$, $^{\text{T}}\text{N-19}^{\text{V}}$, and $^{\text{T}}\text{N-21}^{\text{V}}$) that contain residues from the *drTLR5* N-terminal residue 22 through one of the 2nd and 10th LRR consensus residues of LRR module 4–21 fused with VLR B.61 LRR4-LRRCT (Fig.

1A and B and Table S1) [14]. Six (T_N-4^V , T_N-6^V , T_N-10^V , T_N-12^V , T_N-14^V , and T_N-17^V) of the nine tested T_N-x^V LRR hybrids were successfully expressed in a baculovirus expression system. Except for T_N-10^V that is very unstable with low purification yield (less than 0.1 mg from 1 L culture), T_N-4^V , T_N-6^V , T_N-12^V , T_N-14^V , and T_N-17^V were stable in solution and were purified as monomers (Fig. 2B). T_N-4^V , T_N-6^V , and T_N-14^V exhibited the highest expression levels with a purification yield of at least 1 mg from 1 L culture. T_N-12^V and T_N-16^V yielded 0.2–0.5 mg. The T_N-x^V hybridization showed a high success rate of at least 60%.

We analyzed binding of purified T_N-x^V hybrid proteins to FliC by native PAGE and size-exclusion chromatography (Table S1). Consistent with the critical role of a protruding loop of LRR9 in FliC binding by creating a binding groove [4], the T_N-12^V , T_N-14^V , and T_N-17^V fusion proteins with the LRR9 loop formed a stable complex with CBLB502 that exhibited the same full TLR5-signaling activity as the full-length FliC (Fig. 4). In contrast, T_N-4^V and T_N-6^V that lack the LRR9 loop did not interact with CBLB502. T_N-10^V did not bind FliC although T_N-10^V contains the LRR9 loop, but this result is consistent with our prediction that the LRR9 loop in T_N-10^V is not expected to adopt an optimal FliC-binding conformation for TLR5/FliC interaction due to significant steric clashes of LRR9 and its proximal region with VLR residues (Fig. S2).

3.3. Expression of the drTLR5-ECD C-terminal fragments by a single LRR fusion

To address the contribution of the TLR5 C-terminal region to flagellin binding, we designed LRR hybrid constructs that encode a fusion protein of the *drTLR5*-ECD C-terminal region and the VLR N-terminal region, and tested their expression (Fig. 1C). We designed eight V_Tx-C hybrids (V_T1-C , V_T3-C , V_T5-C , V_T7-C , V_T9-C , V_T11-C , V_T13-C , and V_T18-C) that extend either from the 2nd, 7th, or 10th LRR consensus residue in TLR5 LRR module number x through the C-terminal residue 652 of TLR5-ECD in fusion with VLR B.61 LRRNT-LRR2 (Fig. 1D and Table S1).

Only two V_Tx-C hybrids, V_T1-C and V_T11-C , were expressed, exhibiting a much lower success rate compared to V_N-x^T hybridization. Although V_T1-C was expressible, it formed aggregates as observed for *drTLR5*-ECD. Similar to *drTLR5*-ECD, aggregation formation was prevented by FliC supplementation during protein expression. The purification yield was enhanced to ~0.3 mg per 1 L culture in the presence of FliC. V_T11-C was purified with a higher proportion of a monomeric form with much less aggregation compared to *drTLR5*-ECD (Fig. 2A and C). The final purification yield of V_T11-C was ~0.6 mg from 1 L culture. Since V_T11-C does not have the LRR9 loop, it did not form a tight complex with FliC as efficiently as other *drTLR5*-VLR hybrids that contain the LRR9 loop, such as T_N-12^V , T_N-14^V , and T_N-17^V (Fig. 4). Instead, only some of the V_T11-C material formed complexes with CBLB502 even at molar excess of CBLB502, suggesting a weak interaction (Fig. 4). This previously unidentified TLR5/flagellin interaction is therefore localized on the C-terminal region of TLR5-ECD after LRR17, given that a TLR5-VLR hybrid, V_T11-17^V , that contains TLR5 LRR11–LRR17 did not bind FliC (see below).

3.4. Expression of the central region of drTLR5-ECD by double hybridization with VLR

T_N-x^V and V_Tx-C hybridization screening provides valuable information about fusion points that are able to express stable monomeric TLR5 proteins. By combination of two sets of hybrids (the first set, V_T1-C and V_T11-C ; the second set, $N-12^{TV}$, $N-14^{TV}$, and $N-17^{TV}$), we designed five double hybrids (V_T1-12^V , V_T1-14^V , V_T1-17^V , V_T11-14^V , and V_T11-17^V) that contain the central LRR region of TLR5-ECD flanked by N- and C-terminal regions of VLR (Table S1). For example, V_T11-17^V contains TLR5 LRR11 through LRR17 for which the N- and C-terminal ends are fused with VLR LRRNT-LRR2 and LRR4-LRRCT, respectively.

As expected, all the double hybrids were expressible with purification yields of 0.2–3.0 mg from 1 L culture. They exhibited expected FliC binding behavior depending on the presence of the LRR9 loop. $^{VT}1-12^V$, $^{VT}1-14^V$, and $^{VT}1-17^V$ formed complexes with FliC, whereas $^{VT}11-14^V$ and $^{VT}11-17^V$ did not.

3.5. Application of the drTLR5-VLR hybrid technique to hsTLR5

We also applied ligand supplementation and the LRR hybrid technique to *hsTLR5*-ECD to improve protein expression and purification. It is well known that *hsTLR5*-ECD is not expressible in a functional form even in a mammalian expression system. FliC supplementation did not significantly enhance *hsTLR5*-ECD expression, whereas VLR fusion allowed *hsTLR5*-ECD fragments to be expressed (Table S2). Nine *hsTLR5*-VLR hybrids ($^{TN}4^V$, $^{TN}6^V$, $^{TN}12^V$, $^{TN}14^V$, $^{TN}17^V$, $^{VT}1-C$, $^{VT}11-C$, $^{VT}11-14^V$, $^{VT}11-17^V$) were generated using identical designs that provided high expression levels of *drTLR5*, and their expression levels were analyzed (Table S2). $^{TN}4^V$, $^{TN}6^V$, and $^{VT}11-14^V$ hybrids that contain short fragments of human *TLR5*-ECD were expressed at high levels and purified as monomers with purification yields of ~0.3–0.8 mg from 1 L culture.

4. Discussion

Here, we present the use of alternative, powerful expression techniques to overcome the technical difficulty in production of functionally active recombinant proteins of an LRR innate immunity receptor. Since *TLR5* was first identified as an innate immune receptor for bacterial flagellin in 2001 [3], biophysical characterizations, including structural studies, of *TLR5* itself and its interaction with flagellin have been challenging due to the limited quantity of active *TLR5*-ECD proteins that could be obtained by conventional expression methods. Instead, mutational studies combined with cell-based singling assays were used to identify *TLR5* residues that may be involved in flagellin interaction [15]. However, such results must be carefully interpreted since any decrease or a lack of cellular activity in mutants could not only be interpreted as disruption of *TLR5*/flagellin interaction but also as a result of mutation-induced protein misfolding [4]. Thus, it is essential that such mutation results should be confirmed by direct interaction studies that employ purified functional proteins, emphasizing the importance of studies to improve particularly challenging protein expression.

It has been reported that the yield of biologically relevant forms can be enhanced by external supplementation of culture medium with a small chemical ligand or by co-expression with protein ligands mainly in the *E. coli* expression system [16,17]. Binding of a ligand to the active site of an enzyme or a receptor results, in most cases, in stabilization of unstable region in a protein and improves the yield of active proteins. Consistent with that notion, we obtained active *TLR5*-ECD proteins in large quantity by supplementation of culture with a protein ligand, flagellin, in a baculovirus expression system. Given that flagellin buries extensive *TLR5* surfaces (at least 1200 Å²) upon complex formation, flagellin is likely to mask and stabilize some flexible regions, i.e. the protruding loops of LRR7 or LRR9, of *drTLR5* [4], providing a potential explanation for how flagellin supplementation improves the yield of biologically active *drTLR5*-ECD proteins.

LRR domains are formed by successive assembly of multiple LRR modules and provide a scaffold that exerts versatile functions by modulating its surface residues. Each LRR is a structure motif that stacks consecutively with other LRRs into a solenoid-like shape with parallel β -strands on the concave side and an array of various secondary structures on the convex side. This LRR structure is stabilized at the N- and C-terminal ends by LRRNT and LRRCT capping motifs, respectively, that mask the hydrophobic ends of the solenoid. Protein engineering of LRR domains was first introduced in ribonuclease inhibitor [18]. The

number of the central LRR modules was manipulated between the N-terminal and the C-terminal capping motifs. Later, more extensive LRR engineering was tested in TLR4 and TLR1/2/6 using LRR hybridization between two different LRR proteins to fuse an unstable LRR protein of interest and a more stable LRR protein, hoping to improve the stability and yield of the protein of interest [7–9]. By a single fusion that replaces either the N-terminal or C-terminal region of TLR-ECD with the corresponding region of a VLR, TLR-VLR hybrids exhibited substantial improvement in protein expression and crystallizability without adversely affecting the intrinsic biological activity of the TLR. In our study, we extended this methodology to a double fusion by inserting TLR5 LRR modules between the N-terminal and the C-terminal regions of a VLR. The double fusion hybrids were expressible if each of their two fusions yielded high expression levels. Using this double fusion, we were able to expand the crystallization search space and crystallize ^{VT}11-17^V.

We further extended the LRR hybrid technique to *hs*TLR5-ECD. *hs*TLR5-ECD is extremely difficult to express in an active form even in a mammalian expression system. Because *dr*TLR5 and *hs*TLR5 contain homologous sequences (but with relatively low sequence identity, 37%), we tested expression of *hs*TLR5-VLR hybrids using an identical fusion strategy as with *dr*TLR5-ECD. We were able to express and purify fusion proteins in around a third of designed *hs*TLR5-VLR hybrid constructs. *hs*TLR5 LRRNT-LRR6 or LRR11–LRR14 regions were expressible using ^TN-4^V, ^TN-6^V, and ^{VT}11-14^V *hs*TLR5-VLR hybrids, suggesting that other regions, such as LRR6–LRR11 and LRR14-LRRCT, are the main cause of poor expression of *hs*TLR5-ECD.

The improvement of TLR5 expression levels and purification yields by LRR hybridization with a VLR has been essential for structural and binding studies. TLR5-VLR hybrids provide an important tool to biochemically and biophysically study the TLR5/flagellin interaction. In particular, the LRR hybrid technique was critical for determination of a crystal structure of the ^TN-14^V/flagellin complex [4]. Furthermore, this LRR hybrid technique provides highly versatile applications for expression of LRR-containing immune receptors whose heterologous expression has been elusive. For examples, TLR7/8/9-ECD can be fused with relatively stable VLR or TLR3 proteins. NOD-like receptor (NLR) hybrids can be designed in fusion with its structurally homologous LRR protein, ribonuclease inhibitor (RI), for expression screening. Along with the LRR hybrid technique, the increase in production of active TLR5-ECD proteins by flagellin supplementation would benefit ongoing structural studies that require a large amount of protein sample.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank Andrei L. Osterman and Oleg Kurnasov (Sanford-Burnham Medical Research Institute) for CBLB502 proteins. The work was funded by an NIH grant (AI042266 to I.A.W.), the Skaggs Institute for Chemical Biology (I.A.W.), and Basic Science Research Program through the National Research Foundation of Korea (2012R1A1A1003701 to S.I.Y.). This is manuscript no. 21896 from The Scripps Research Institute.

References

- [1]. Takeda K, Kaisho T, Akira S. Toll-like receptors. *Annu. Rev. Immunol.* 2003; 21:335–376. [PubMed: 12524386]
- [2]. Botos I, Segal DM, Davies DR. The structural biology of Toll-like receptors. *Structure.* 2011; 19:447–459. [PubMed: 21481769]

- [3]. Hayashi F, Smith KD, Ozinsky A, Hawn TR, Yi EC, Goodlett DR, Eng JK, Akira S, Underhill DM, Aderem A. The innate immune response to bacterial flagellin is mediated by Toll-like receptor 5. *Nature*. 2001; 410:1099–1103. [PubMed: 11323673]
- [4]. Yoon SI, Kurnasov O, Natarajan V, Hong M, Gudkov AV, Osterman AL, Wilson IA. Structural basis of TLR5-flagellin recognition and signaling. *Science*. 2012; 335:859–864. [PubMed: 22344444]
- [5]. Burdelya LG, Krivokrysenko VI, Tallant TC, Strom E, Gleiberman AS, Gupta D, Kurnasov OV, Fort FL, Osterman AL, Didonato JA, Feinstein E, Gudkov AV. An agonist of Toll-like receptor 5 has radioprotective activity in mouse and primate models. *Science*. 2008; 320:226–230. [PubMed: 18403709]
- [6]. Turley CB, Rupp RE, Johnson C, Taylor DN, Wolfson J, Tussey L, Kavita U, Stanberry L, Shaw A. Safety and immunogenicity of a recombinant M2e-flagellin influenza vaccine (STF2.4xM2e) in healthy adults. *Vaccine*. 2011; 29:5145–5152. [PubMed: 21624416]
- [7]. Jin MS, Kim SE, Heo JY, Lee ME, Kim HM, Paik SG, Lee H, Lee JO. Crystal structure of the TLR1–TLR2 heterodimer induced by binding of a tri-acylated lipopeptide. *Cell*. 2007; 130:1071–1082. [PubMed: 17889651]
- [8]. Kang JY, Nan X, Jin MS, Youn SJ, Ryu YH, Mah S, Han SH, Lee H, Paik SG, Lee JO. Recognition of lipopeptide patterns by Toll-like receptor 2-Toll-like receptor 6 heterodimer. *Immunity*. 2009; 31:873–884. [PubMed: 19931471]
- [9]. Kim HM, Park BS, Kim JI, Kim SE, Lee J, Oh SC, Enkhbayar P, Matsushima N, Lee H, Yoo OJ, Lee JO. Crystal structure of the TLR4-MD-2 complex with bound endotoxin antagonist Eritoran. *Cell*. 2007; 130:906–917. [PubMed: 17803912]
- [10]. Yoon SI, Hong M, Wilson IA. An unusual dimeric structure and assembly for TLR4 regulator RP105-MD-1. *Nat. Struct. Mol. Biol.* 2011; 18:1028–1035. [PubMed: 21857663]
- [11]. Yoon SI, Hong M, Han GW, Wilson IA. Crystal structure of soluble MD-1 and its interaction with lipid IVA. *Proc. Natl. Acad. Sci. USA*. 2010; 107:10990–10995. [PubMed: 20534476]
- [12]. Ainge GD, Martin WJ, Compton BJ, Hayman CM, Larsen DS, Yoon SI, Wilson IA, Harper JL, Painter GF. Synthesis and Toll-like receptor 4 (TLR4) activity of phosphatidylinositol dimannoside analogues. *J. Med. Chem.* 2011; 54:7268–7279. [PubMed: 21936536]
- [13]. Jin MS, Lee JO. Application of hybrid LRR technique to protein crystallization. *BMB Rep.* 2008; 41:353–357. [PubMed: 18510864]
- [14]. Kim HM, Oh SC, Lim KJ, Kasamatsu J, Heo JY, Park BS, Lee H, Yoo OJ, Kasahara M, Lee JO. Structural diversity of the hagfish variable lymphocyte receptors. *J. Biol. Chem.* 2007; 282:6726–6732. [PubMed: 17192264]
- [15]. Andersen-Nissen E, Smith KD, Bonneau R, Strong RK, Aderem A. A conserved surface on Toll-like receptor 5 recognizes bacterial flagellin. *J. Exp. Med.* 2007; 204:393–403. [PubMed: 17283206]
- [16]. Hozjan V, Guo K, Wu X, Oppermann U. Ligand supplementation as a method to increase soluble heterologous protein production. *Expert Rev. Proteomics*. 2008; 5:137–143. [PubMed: 18282129]
- [17]. Li C, Schwabe JW, Banayo E, Evans RM. Coexpression of nuclear receptor partners increases their solubility and biological activities. *Proc. Natl. Acad. Sci. USA*. 1997; 94:2278–2283. [PubMed: 9122185]
- [18]. Stumpp MT, Forrer P, Binz HK, Pluckthun A. Designing repeat proteins: modular leucine-rich repeat protein libraries based on the mammalian ribonuclease inhibitor family. *J. Mol. Biol.* 2003; 332:471–487. [PubMed: 12948496]

Fig. 1. T^N-x^V (A, B) and V^T-x-C (C, D) hybrid designs that allow linkage of LRR fragments of TLR5 and VLR. (A) Schematic representation of T^N-x^V hybrids. In T^N-17^V as an example of T^N-x^V hybrids, the TLR5 N-terminal region from the N-terminus to LRR17 (yellow in continuous lines) was fused with the VLR C-terminal region from LRR4 to the C-terminus (cyan in continuous lines) at the 10th consensus residue (asparagine in red). (B) Amino acid sequences of T^N-x^V hybrids at a fusing LRR module. TLR5 and VLR sequences are highlighted in yellow and cyan, respectively. A consensus residue where fusion occurs is shown in red. (C) A schematic representation of V^T-x-C hybrids. In V^T-1-C as an example of V^T-x-C hybrids, the TLR5 C-terminal region from the LRR1 to the C-terminus (yellow in continuous lines) was fused with the VLR N-terminal region from the N-terminus to LRR2 (cyan in continuous lines) at the 10th consensus residue (asparagine in red). (D) Amino acid sequences of V^T-x-C hybrids in the fusing LRR module.

Fig. 2. Size-exclusion chromatography profiles of *dr*TLR5-ECD (A), ^TN-14^V (B), and ^{VT}11-C (C). Expressed proteins were first purified by Ni-NTA and Strep-Tactin affinity chromatography. The resulting proteins were digested by thrombin and injected into a Superdex 200 16/600 size-exclusion chromatography column.

Fig. 3.

Improved expression yield of functional *drTLR5-ECD* proteins by FliC supplementation. Different amounts of purified FliC proteins (70 µg, 200 µg, or 600 µg) were added into 1 L of *drTLR5-ECD* expression culture at ~24 h postinfection. At 48 h postinfection, culture medium that contains *drTLR5-ECD* and FliC was harvested and subjected to purification by Ni-NTA affinity chromatography. The resulting material was analyzed using a Superdex 200 10/300 column to estimate the yield of *drTLR5-ECD/FliC* complex (A). Peak fractions that correspond to the *drTLR5-ECD/FliC* complex were analyzed by SDS-PAGE, indicating the presence of both *drTLR5-ECD* and FliC in the peak (B). Peak height and width increased in a FliC dose-dependent manner, demonstrating that FliC supplementation is a crucial step to improve *drTLR5-ECD* protein yield.

Fig. 4.

Native PAGE analysis of interaction between *d*TLR5-VLR hybrids and CBLB502. CBLB502 was added in molar excess compared to *d*TLR5-VLR hybrids in each reaction. CBLB502 completely shifted ^TN-12^V, ^TN-14^V, and ^TN-17^V proteins by forming complexes. Noticeably, only a fraction of the ^VT11-C protein was complexed with CBLB502, suggesting a weak interaction.