Published in final edited form as:

Org Lett. 2013 September 20; 15(18): . doi:10.1021/ol4023084.

Copper-Catalyzed Electrophilic Amination of Organolithiums Mediated by Recoverable Siloxane Transfer Agents

Minh H. Nguyen and Amos B. Smith III

Department of Chemistry, University of Pennsylvania, Philadelphia, Pennsylvania 19104, United States

Amos B. Smith: smithab@sas.upenn.edu

Abstract

The development and validation of copper-catalyzed, electrophilic amination of aryl and heteroaryl organolithiums with N,N -dialkyl- O -benzoylhydroxylamines have been achieved exploiting recoverable siloxane transfer agents. Given the ready availability of organolithium compounds, the mild reaction conditions, the ease of product purification and the ready recovery of the siloxane transfer agents, this transformation comprises a useful tactic to access diverse aryl and heteroaryl amines.

Transition metal-mediated C-N bond construction is one of the most important transformations in organic synthesis for the elaboration of diverse aryl and heteroaryl amines, given the prevalence of this structural motif in natural products, pharmaceutical agents, and functional materials.¹ Among them, the Buchwald-Hartwig palladium-catalyzed cross-coupling of aryl halides with amines, the Ullmann-type copper-catalyzed cross-coupling of aryl halides with amines and the Chan-Lam copper-mediated oxidative coupling of aryl metals with amines have proven to be powerful methods for C-N bond formation.² An umpolung tactic, involving transition metal-catalyzed amination of organometallic reagents with electrophilic aminating reagents (e.g., chloroamines and hydroxylamines), provides a valuable complementary approach.³ Indeed, various organometallic reagents are known to undergo electrophilic amination, including organomagnesiums,⁴ organozincs,⁵ organotinns,⁶ organoborons,^{6,7} organosilicons,⁸ and organotitaniums.⁹ These reagents however are frequently prepared from the corresponding organolithium reagents. The direct use of organolithiums would eliminate the need for additional synthetic manipulations and/or the isolation of suitable nucleophilic partners. Although organolithium reagents are commercially available and/or readily accessible via simple metal-halogen exchange or metallation, the direct application of these agents in electrophilic amination remains a

Correspondence to: Amos B. Smith, III, smithab@sas.upenn.edu.

Supporting Information Available. Experimental procedures and characterization data for all new compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

challenge, often hampered by low yields, limited substrate scope, and in some cases require excess reagents.^{3a, 3d}

Recently we reported a highly atom-efficient tactic for palladium-catalyzed cross-coupling of aryl and alkenyl organolithium reagents with aryl and alkenyl iodides (Scheme 1), exploiting readily recoverable and/or polymeric siloxane transfer agents.^{10,11} As part of the ongoing program to expand the siloxane transfer agent tactic, we report here the copper-catalyzed electrophilic amination of aryl and heteroaryl organolithiums with electrophilic aminating reagents, mediated by recoverable siloxanes.

Based on the similarities between the reactive intermediate involved in siloxane-based cross-coupling reactions (alkoxide **1a**, Scheme 1) and arylsilanes **2**, recently employed in electrophilic aminations reported by Miura and co-workers (Scheme 2),⁸ we reasoned that we could intersect this reaction manifold via a siloxane transfer agent. Success would provide a valuable and practical method for the direct amination of organolithium compounds with electrophilic aminating reagents, such as *N,N*-dialkyl-*O*-benzoyl hydroxylamines. To the best of our knowledge, there are no reports of an effective, general method to employ *directly* organolithium reagents for the preparation of *N,N*-dialkyl aryl and heteroaryl amines, an important structural motif in many biologically active compounds.¹²

To explore this thesis, we began by subjecting alkoxide intermediate **1a**, derived from the reaction of PhLi and **1** to conditions similar to those reported by Miura and coworkers, employing *N,N*-dibenzyl-*O*-benzoyl hydroxylamine as the electrophile (Scheme 3). Pleasingly, the desired product, dibenzyl aniline, was obtained in 63% yield. Although the majority of siloxane **1** could be recovered from the reaction mixture, the siloxane-derived benzoate **5** and triphenyl methanol were also isolated. We envision that under the conditions the reaction proceeds via two pathways (Scheme 4). The desired reaction involves transmetalation from silicon to copper, with subsequent electrophilic amination of the aryl copper intermediate to generate **4a**. Alternatively, the lithium alkoxide (**1a**) can *O*-acylate **3a** to generate silane **5** or transfer the phenyl group to **3a** to generate triphenyl methanol **6**.

With these preliminary, albeit intriguing results in hand, we turned to optimize the reaction, including both product yield and siloxane recovery (Table 1). Changing the solvent from 1,4-dioxane to THF increased the yield of **4a** (entry 1). Importantly the reaction proceeded effectively at room temperature within a 2 h time period. An increase in product yield and siloxane recovery was also realized when the catalyst loading was increased (entries 2 and 3); a further increase in the copper catalyst (40 mol %) however proved ineffective (entry 4). Particularly pleasing, a significant increase in the efficiency of the process was observed when *N*-[2-(diphenylphosphino)benzylidene]cyclohexylamine (dpca), the optimal ligand employed in siloxane mediated cross-coupling reactions,^{10,11} was employed. The use of 10 mol % CuI and 10 mol % dpca proved optimal, leading to complete conversion of **3a** within 2 h at room temperature to furnish **4a** in 98% isolated yield, with full recovery of the siloxane transfer agent (entry 5). Attempts to reduce the amount of PhLi or the catalyst loading, while maintaining the efficiency of the process, provided no enhancement (entries 6 and 7). Importantly, as a control experiment only 7% of **4a** was obtained in the absence of the siloxane transfer agent (entry 8). Also as expected, reaction of PhLi with the electrophilic aminating reagent **3a** provided triphenyl methanol as the major product; no amination product could be detected (entry 9).

We next turned to explore the scope of the Cu-catalyzed electrophilic amination of organolithium agents, employing siloxane transfer agent **1** under the optimized reaction conditions (Scheme 5). Various acyclic as well as cyclic amines underwent effective

coupling with PhLi to form the corresponding anilines in good to excellent yield (**4a–i**). The resultant *N*-allyl and *N*-benzyl moieties (**4a–c**) can serve as useful synthetic handles upon removal of the benzyl or allyl groups to access either unsubstituted or monosubstituted anilines (e.g., **4c**).¹⁴ Notably, the reaction of PhLi and the sterically hindered electrophile *N,N*-diisopropyl-*O*-benzoyl hydroxylamine proceeds to form the corresponding aniline in good yield (**4e**). Electrophilic amination was also effective with a variety of organolithium agents. In particular, the electron-rich *p*-methoxyphenyl lithium readily underwent electrophilic amination to provide the corresponding amination products in good yield (**4j** and **4k**). On the other hand, electron-deficient organolithiums proceeded with somewhat lower efficiency (**4l–n**). Pleasingly however, the aryl-Cl moiety remained intact in the amination of *p*-chlorophenyl lithium with piperidine-1-yl benzoate, providing an opportunity for additional transformations, such as palladium-catalyzed cross-coupling reactions (**4n**). Sterically hindered organolithiums such as *o*-methoxyphenyl lithium and 1-naphthyl lithium also proceeded albeit in moderate yield (**4o** and **4q**). Importantly, heteroaryl lithiums such as 3-pyridyl lithium and 2-benzofuryl lithium were also tolerated in the amination (**4p** and **4r**), the latter accessed via a direct ortho lithiation/amination sequence of benzofuran. Use of 3-thienyl lithium and the sterically hindered mesityl lithium however proved unsuccessful (**4s** and **4t**). Gratifyingly, the amination protocol can be carried out on a gram-scale to provide an excellent yield of the desired product (**4f**, 94%), indicating both scalability and reliability of the method. Equally important, in this case, the siloxane transfer agent was recovered in 92% yield via Kugelrohr distillation, with the amination product **4f** conveniently isolated via acid-base extraction.

We next explored the use of other siloxane transfer agents, focusing on the ability to recycle the transfer agent via a simple and easy protocol. Success here should facilitate adoption of the siloxane tactic by the chemical community.

As shown in Scheme 6, Brønsted base-incorporated siloxane transfer agents **7** and **8** were found to be effective in the electrophilic amination between PhLi and *N,N*-dibenzyl-*O*-benzoyl hydroxylamine, providing the amination product in 64% and 82% yield, respectively with near quantitative recovery of the transfer agents via acid-base extraction,^{10b} exploiting the differential basicity of the siloxanes and the amine products.

Finally we examined the utility of our recently introduced polymer-supported siloxane transfer agent PSTA-I₂₀₀,¹¹ now for electrophilic amination between PhLi and piperidine-1-yl benzoate (Table 2). Pleasingly, the polymer can be used for at least 3 cycles, providing excellent to good yields of the desired amination product **4g**, albeit with a small decrease in efficiency after each cycle. In all cases, the amination product was purified via acid-base extraction and the polymer recovered in near quantitative yield via precipitation in CH₃CN, thus featuring a chromatography-free purification strategy.

In summary, we have developed and validated the direct copper-catalyzed, electrophilic amination of aryl and heteroaryl organolithiums with *N,N*-dialkyl-*O*-benzoyl hydroxylamines. Importantly, the use of siloxane transfer agents offers, for the first time, a viable solution for the direct application of organolithiums in electrophilic amination, and in turn eliminates the need for multiple synthetic manipulation and/or isolation of the required organometallic reagents (e.g., organoborons and organosilicons), as well as the requirement for stoichiometric amount of toxic, heavy metals (e.g., Zn, Sn, and Ti). The mild, room temperature reaction conditions, in conjunction with the ease of product purification, make this approach a valuable addition to methods for C–N bond construction. Studies to expand the scope of this reaction, as well as the application of siloxane transfer agents in other transformations continue in our Laboratory.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Financial support was provided by the NIH through Grant CA-19033. We thank Dr. Rakesh Kohli at the University of Pennsylvania for assistance in obtaining HRMS.

References

1. (a) Hili R, Yudin AK. *Nat Chem Biol.* 2006; 2:284. [PubMed: 16710330] (b) Ricci, A., editor. *Amino Group Chemistry. From Synthesis to the Life Sciences.* Wiley-VCH; Weinheim: 2007.
2. For reviews, see: Hartwig JF. *Acc Chem Res.* 2008; 41:1534. [PubMed: 18681463] Surry DS, Buchwald SL. *Chem Sci.* 2011; 2:27. [PubMed: 22432049] Ley SV, Thomas AW. *Angew Chem, Int Ed.* 2003; 42:5400. Monnier F, Taillefer M. *Angew Chem, Int Ed.* 2009; 48:6954. Qiao JX, Lam PYS. *Synthesis.* 2011:829.
3. For reviews, see: Erdik E, Ay M. *Chem Rev.* 1989; 89:1947. Dembech P, Seconi G, Ricci A. *Chem Eur J.* 2000; 6:1281. [PubMed: 10840948] Barker TJ, Jarbo ER. *Synthesis.* 2011:3954. Daskapan T. *Arkivoc.* 2011:230.
4. (a) Tsutsui H, Hayashi Y, Narasaka K. *Chem Lett.* 1997; 26:317. (b) Campbell MJ, Johnson JS. *Org Lett.* 2007; 9:1521. [PubMed: 17362022] (c) Hatakeyama T, Yoshimoto Y, Ghorai SK, Nakamura M. *Org Lett.* 2010; 12:1516. [PubMed: 20222741]
5. (a) Berman AM, Johnson JS. *J Am Chem Soc.* 2004; 126:5680. [PubMed: 15125656] (b) Berman AM, Johnson AS. *Synlett.* 2005:1799. (c) Berman AM, Johnson JS. *J Org Chem.* 2006; 71:219. [PubMed: 16388639] (d) Barker TJ, Jarvo ER. *J Am Chem Soc.* 2009; 131:15598. [PubMed: 19824677]
6. (a) Liu S, Yu Y, Liebeskind LS. *Org Lett.* 2007; 9:1947. [PubMed: 17444649] (b) Zhang Z, Yu Y, Liebeskind LS. *Org Lett.* 2008; 10:3005. [PubMed: 18557628]
7. (a) Liu S, Liebeskind LS. *J Am Chem Soc.* 2008; 130:6918. [PubMed: 18465855] (b) He C, Chen C, Cheng J, Liu C, Liu W, Li Q, Lei A. *Angew Chem, Int Ed.* 2008; 47:6414. (c) Rucker RP, Whittaker AM, Dang H, Lalic G. *Angew Chem, Int Ed.* 2012; 51:3953. (d) Xiao Q, Tian L, Tan R, Xia Y, Qiu D, Zhang Y, Wang J. *Org Lett.* 2012; 14:4230. [PubMed: 22870926] (e) Mlynarski SN, Karns AS, Morken JP. *J Am Chem Soc.* 2012; 134:16449. [PubMed: 23002712] (f) Zhu C, Li G, Ess DH, Falck JR, Kurti L. *J Am Chem Soc.* 2012; 134:18253. [PubMed: 23082853]
8. Miki Y, Hirano K, Satoh T, Miura M. *Org Lett.* 2013; 15:172. [PubMed: 23256842]
9. Barker TJ, Jarvo ER. *Angew Chem, Int Ed.* 2011; 50:8325.
10. Smith AB III, Hoyer AT, Martinez-Solorio D, Kim W-S, Tong R. *J Am Chem Soc.* 2012; 134:4533. [PubMed: 22352347] Martinez-Solorio D, Hoyer AT, Nguyen MH, Smith AB III. *Org Lett.* 2013; 15:2454. For the closely related Hiyama cross-coupling reaction, see: Nakao Y, Hiyama T. *Chem Soc Rev.* 2011; 40:4893. and references cited therein. [PubMed: 21755089]
11. Nguyen MH, Smith AB III. *Org Lett.* 2013; 15:4258. [PubMed: 23901881]
12. (a) Trump RP, Blanc JE, Stewart EL, Brown PJ, Caivano M, Gray DW, Hoekstra WJ, Willson TM, Han B, Turnbull P. *J Comb Chem.* 2007; 9:107. [PubMed: 17206838] (b) Dolle RE, Bourdonnec BL, Goodman AJ, Morales GA, Thomas CJ, Zhang W. *J Comb Chem.* 2008; 10:753. [PubMed: 18991466]
13. Tamao-Fleming oxidation: Tamao K, Ishida N, Tanaka T, Kumada M. *Organometallics.* 1983; 2:1694. Fleming I, Henning R, Plaut H. *J Chem Soc, Chem Commun.* 1984:29.
14. (a) Jaime-Figueroa S, Liu Y, Muchowski JM, Putman DG. *Tetrahedron Lett.* 1998; 39:1313. (b) Cadierno V, Garcia-Garrido SE, Gimeno J, Nebra N. *Chem Commun.* 2005:4086. (c) Wang J-Y, Wang D-X, Zheng Q-Y, Huang Z-T, Wang M-X. *J Org Chem.* 2007; 72:2040. [PubMed: 17286438]

Scheme 1.
Siloxane based cross-coupling reactions

Scheme 2.
Miura's electrophilic amination of aryl silanes

Scheme 3.

Initial attempt at the electrophilic amination of phenyl lithium employing siloxane transfer agent

Scheme 4.
Proposed reaction pathways

Scheme 5.

Substrate-Scope Study of the electrophilic amination of organolithiums employing siloxane transfer agent ^a

^aAll reactions were performed on 0.3 mmol scale. The organolithiums were generated via lithium-halogen exchange between *t*-BuLi and the corresponding organohalides. The siloxane was recovered via flash chromatography (average recovery = 85%). ^bSiloxane coelutes with product and was oxidized¹³ with H₂O₂ in NaOH to facilitate product purification. ^cReaction was run on 3 mmol scale. ^d20% CuI was used. ^eThe organolithium was generated by treating 2,3-benzofuran with *n*-BuLi. ^fEstimated via H-NMR analysis of the crude reaction mixture.

Scheme 6.
Performance of Brønsted base containing siloxanes

Table 1

Optimization of the electrophilic amination of organolithiums employing siloxane transfer agent^a

entry	equiv PhLi	equiv 1	mol % CuI	ligand (mol%)	yield of 4a	product ratio ^b 1:5
1	1.5	1.8	5%	JohnPhos (10%)	67%	5:1
2	1.5	1.8	10%	JohnPhos (10%)	81%	10:1
3	1.5	1.8	20%	JohnPhos (10%)	91%	19:1
4	1.5	1.8	40%	JohnPhos (10%)	84%	>20:1
5	1.5	1.8	10%	dpca (10%)	98%	>20:1
6	1.2	1.5	10%	dpca (10%)	88%	>20:1
7	1.5	1.8	5%	dpca (5%)	73%	11:1
8 ^c	1.5	---	10%	dpca (10%)	7%	---
9 ^c	1.5	---	---	---	nd	---

^a All reactions were performed on 0.3 mmol scale.

^b Determined by ¹H-NMR analysis of the reaction mixture following an aqueous workup and extraction with Et₂O.

^c Ph₃COH as the major product. nd = not detected.

Table 2

Recyclability of Polymer-supported Siloxane Transfer Agent

cycle	isolated yield of 4g	polymer recovered
1 st	92%	95%
2 nd	80%	92%
3 rd	71%	90%