Published in final edited form as:

Curr Opin Cell Biol. 2013 October ; 25(5): . doi:10.1016/j.ceb.2013.06.003.

Guiding cell migration by tugging

Sergey V. Plotnikov and Clare M. Waterman*

Cell Biology and Physiology Center, National Heart Lung and Blood Institute, National Institutes of Health, Bethesda, MD

Abstract

The ability of cells to move directionally towards areas of stiffer extracellular matrix (ECM) via a process known as “durotaxis” is thought to be critical for development and wound healing, but durotaxis can also drive cancer metastasis. Migration is driven by integrin-mediated focal adhesions (FAs), protein assemblies that couple contractile actomyosin bundles to the plasma membrane, transmit force generated by the cytoskeleton to the ECM, and convert the mechanical properties of microenvironment into biochemical signals. To probe the stiffness of the ECM, motile fibroblasts modulate FA mechanics on the nanoscale and exert forces that are reminiscent of repeated tugging on the ECM. Within a single cell, all FAs tug autonomously and thus act as local rigidity sensors, allowing discernment of differences in the extracellular matrix rigidity at high spatial resolution. In this article, we review current advances that may shed light on the mechanism of traction force fluctuations within FAs. We also examine plausible downstream effectors of tugging forces which may regulate cytoskeletal and FA dynamics to guide cell migration up ECM stiffness gradients.

Introduction

Directional cell movement is critical to embryonic development, immune system function, angiogenesis, and wound healing, as well as cancer metastasis. Cell migration is induced by a variety of signaling mechanisms that receive and process information from the cell's environment and provide specific control of cytoskeletal and adhesion machineries within the cell [1]. Historically, attention has been focused on understanding how diffusible or ECM-associated biochemical cues are transduced into activity of intracellular signaling networks that regulate cytoskeletal and adhesion dynamics. However, recent studies have highlighted the importance of physical cues such as ECM topology or rigidity in guiding cell migration. In particular, the propensity of cells to migrate towards areas of higher ECM rigidity via a process known as “durotaxis” has garnered interest [2]. Durotaxis is thought to contribute to physiological processes including stem cell differentiation [3,4], epithelial-to-mesenchymal transition [5,6], development of the nervous system [7,8], innate immunity [9], as well as promoting breast cancer or glioblastoma metastases [10,11].

The ability of cells to durotax up rigidity gradients requires mechanisms for constant surveillance of the variability in the stiffness landscape of the ECM in the cellular microenvironment. Several cellular structures have been proposed as force or rigidity sensors, including the plasma membrane [12], actin filaments [13,14], the cortical

*Correspondence to: watermancm@nhlbi.nih.gov, National Heart, Lung and Blood Institute, National Institutes of Health, Building 50 South Drive, Room 4537 MSC 8019, Bethesda Maryland 20892-8019, T: (301)-435-2949, F: (301)-480-6012.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

cytoskeleton [15,16], the nucleus [17], and cadherin-based adherens junctions [18]. However, there is extensive evidence that actomyosin-based contractility and integrin-based FAs are essential for ECM rigidity sensing [19,20]. Durotaxis is known to require myosin contractility [21], and the activity of FA proteins including FAK [22], paxillin, and vinculin [23], indicating that integrin-based FAs serve as the rigidity sensors that specifically guide durotaxis. In this review, we focus on recent observations of the spatial and temporal dynamics of forces exerted by FAs during ECM rigidity sensing. We discuss possible molecular mechanisms that could mediate force dynamics in FAs and how force dynamics could be translated into polarized regulation of cytoskeletal and FA dynamics that drive directed cell migration.

Traction force fluctuations guide durotaxis

We recently used high-resolution traction force microscopy to characterize the nanoscale dynamics of cell-generated forces on the ECM [23]. Our studies revealed that mature FAs which appear static by other methods of microscopy may actually possess internal fluctuations in mechanics. FAs within a single cell were found to adopt one of two states: a stable state where traction was spatially and temporally static, and a dynamic state in which the pattern of traction fluctuation was reminiscent of repeated, centripetal tugging on the ECM. The choice between tugging and stable FA states could be predictably controlled by modulating ECM rigidity, myosin contractility, or a FAK/phosphopaxillin/vinculin pathway. Tugging traction in FAs was found to be dispensable for FA maturation, chemotaxis and haptotaxis, but critical for directed cell migration towards stiff ECM, i.e. durotaxis (Figure 1). Repeated FA tugging on the ECM suggests a means of repeatedly sampling the local ECM rigidity landscape over time. ECM rigidity sensing by individual tugging FAs could allow tight control of directional migration to guide cells along highly localized or dynamically changing ECM rigidity gradients during development or in tumors. FA-mediated sensing of local stiffness cues may also be utilized in addition to biochemical gradient sensing of diffusible and immobilized cues to fine-tune cell path-finding during development, morphogenesis, and pathological processes such as metastasis.

Mechanistic Basis of Force Fluctuations

There are three basic components contributing to force on the ECM at an FA: 1) Myosin II, which produces force on 2) Actin filaments, which act as a conduit of the force to 3) FA proteins and integrins, which comprise the linkage between actin and the ECM through the plasma membrane. Dynamic changes in assembly/disassembly, activity, or protein-protein interactions within any of these three components could be responsible for mediating the fluctuations in force transmission seen in FAs (Figure 2).

Fluctuations in myosin contractility

Temporal variations in myosin II contractility represents the most obvious mechanism for applying fluctuating pulling forces to the ECM. At the whole cell level, oscillating cell contractions driven by cyclical accumulation of myosin II on the actin network have recently been observed in non-excitable cells [24–26]. However, global changes in actomyosin contractility are likely not responsible for fluctuations in traction force in FA, since force fluctuations in neighboring FAs, which are mechanically coupled to each other via the actin cytoskeleton [27], are not temporally correlated [23]. This suggests that dynamics of traction force are regulated locally, within single FA. Myosin II is closely associated with FAs [28] and interacts with FA components such as Rac guanine exchange factor α -PIX [29]. Yet, myosin II is generally not considered a *bona fide* FA protein, because it tends to localize behind FAs within stress fibers and it is thought to transmit force from a distance through the stress fiber to the FA [30].

Mathematical modeling predicts that the activity of local contractile units could oscillate spontaneously due to the collective activity of motors acting on an elastic material [31]. Remarkably, such oscillations may be mechanosensitive, as for any given myosin activity the predicted oscillation frequency is determined by the elasticity of the FA and the ECM [32]. A direct demonstration that myosin accumulation at a FA affects the dynamics of its traction force is required to validate the notion that local fluctuations in contractility produce tugging forces at FA.

Fluctuations in actin mechanics

Since actin serves as the mechanical conduit between force generated by myosin and ECM-bound integrins, it is possible that cyclic variation in the mechanics of the actin cytoskeleton underlie traction fluctuations. Thus, cyclic variations in the assembly/disassembly of actin to modulate actin mechanics are a plausible mechanism for the origin of traction fluctuations at FA. There is mounting evidence that lamellipodial actin assembly occurs in cycles [33–35]. However, force fluctuations observed recently in fibroblasts occurred in mature FAs that are not associated with lamellipodial actin, but which are instead linked to actin stress fibers.

Actin filaments within stress fibers terminate with their assembly-competent barbed ends linked to the FA [36]. These filaments could undergo cycles of assembly that would allow cyclic slack to develop in the stress fiber, and thus cyclic reduction in myosin-generated force transmitted to the ECM. Members of the Ena/VASP and formin families have been implicated as the main molecular players in the assembly of stress fibers at FAs [37–39] and could mediate cyclic assembly of F-actin at FAs. Ena/VASP proteins localize to FAs [40], and the related protein Mena binds directly to integrins [41]. Formins have been identified in FA proteomic screens [42–44], and inhibiting the activity of formin proteins disrupts FA traction and maturation [45,46]. It was also recently demonstrated that VASP proteins are implicated in force-dependent stress fiber remodeling via interactions with zyxin [38,47], and that formins allow accelerated barbed end elongation when under tension [48]. This suggests the intriguing possibility that fluctuations in contraction and actin assembly in stress fibers could cooperate via a positive feedback loop to mediate traction fluctuations at FAs [49,50].

Fluctuations in the FA “molecular clutch”

If contraction and actin polymerization are constant, tugging traction at FAs could also result from temporal variations in the strength or number of linkages between actin and the ECM via changes in protein-protein interactions within FAs. The putative chain of protein-protein interactions making up this linkage has been termed the “molecular clutch” based on its role in transmitting force from actomyosin to the ECM, analogous to the clutch of a car that transmits force generated by the engine to the drive shaft. There is extensive evidence for transient, regulatable interactions between actin and the ECM through FA proteins. Studies conducted by using FRAP (fluorescence recovery after photobleaching) imaging and fluorescent speckle microscopy have demonstrated that FA proteins bind and dissociate from FAs on a much faster time-scale than FA turnover and exhibit partial coupling to the actin cytoskeleton [51–53]. In particular, talin and vinculin are extensively implicated as load-bearing components mediating the regulatable tunable link between actin and ECM-bound integrins [54–56]. Talin and vinculin interactions with actin, integrins, and other FA proteins are highly regulated through phosphorylation, lipid binding, and small GTPase signaling pathways [57,58]. Thus, it is possible that collective regulation of the integrin- or actin-binding affinity of talin and/or vinculin through oscillations in signal transduction could produce tugging force dynamics in FAs.

An alternative mechanism for generating tugging traction force from a simple ‘motor—clutch’ system that does not require positive or negative feedback or signaling pathways for regulation was recently predicted by a stochastic mathematical model [59]. The model consisted of actin undergoing retrograde flow, “clutch molecules” whose engagement to actin and the ECM was dictated by association rate constants and whose disengagement was dictated by both rate constants and force (i.e. breaking strength), and ECM stiffness. The model predicted two distinct regimes of force transmission to the ECM within an individual FA: static and oscillating. The static force regime was promoted by stochastic association and disassociation of many clutches across a FA resulting in even, constant traction force centered within the FA. This state occurred when engaged clutch lifetime was short and ECM stiffness was high. The oscillating regime, which was promoted by long engagement lifetime and low ECM stiffness, was the result of cooperative engagement and simultaneous failure of local clusters of clutches within a FA, generating oscillations in traction magnitude and peak traction force position within the FA. Thus, the simple motor-clutch model generally accounts for stiffness-dependent changes in traction dynamics that have been observed in neurons and fibroblasts [23,59]. However, order-of-magnitude differences between model predictions and experimental measurements in the rate of traction force increase call in to question whether this simple mechanism operates *in vivo*. Furthermore, this model does not account for the polymerization of actin that is known to occur at FAs.

Decoding traction dynamics by downstream effectors

To migrate directionally along gradients of ECM stiffness, cells require a mechanism to continuously measure variability in the stiffness landscape of the ECM and control cytoskeletal and adhesion dynamics. Cells sample stiffness by exerting actomyosin-generated pulling forces on the surrounding ECM through FAs [19,20]. Fluctuations of traction stress within FA may be a means by which cells repeatedly tug at the ECM to detect spatial and temporal changes in rigidity. But why is fluctuating FA traction required for durotaxis on a spatially and temporally *stable* ECM rigidity gradient but not for chemotaxis or haptotaxis [23]? Tension on FA proteins is thought to drive conformational changes including stretching or unfolding, which alter protein-protein interactions, induce recruitment of cytosolic proteins, and activate signaling pathways [60]. This suggests that fluctuating or oscillating signals may be specifically required for durotaxis. Although we still do not know which signaling pathways transduce dynamics of tugging forces into cellular behavior, only those which are: (1) mechanically activated; (2) originate from integrin FAs; and (3) exhibit fluctuating activity could be regulated by the dynamics of traction forces. Among known signaling pathways involved in the regulation of cell migration, integrins, their effectors and stretch-activated ion channels satisfy these criteria and will be discussed below.

Cyclic activation of integrins

One obvious candidate for an effector of tugging traction dynamics that may mediate directed cell migration is integrins themselves. Indeed, it is well established that integrin affinity for ECM is enhanced by tension [61–63]. However, a recent elegant study showed that while a single tug on an $\alpha 5 \beta 1$ integrin increased its affinity for ligand, the cyclic application of force reinforced the integrin-ligand bond, prolonging bond lifetime by two orders of magnitude to maintain single bonds for minutes [64]. This suggests that repeated application of tension on integrins by FA tugging could be a mechanism to prolong adhesion lifetime. Thus, in a cell on an ECM stiffness gradient, tugging on the FAs coupled to a stiffer ECM region may overcome some threshold for cyclic mechanical reinforcement, while this threshold may not be reached in FAs coupled to the softer matrix area. This would result in preferential reinforcement of FAs in the direction of greater stiffness to promote directed migration.

Several signaling cascades downstream of integrin activation have been shown or predicted to exhibit oscillating activity and are known to regulate cytoskeletal and adhesion dynamics. Thus, in addition to reinforcing integrin adhesion, repeated tugging on integrins could induce cyclic activation of integrin signaling. FAK and Src tyrosine kinase activities are induced by integrin activation and force [65] and are known to play an important role in cell mechanosensing, migration, and invasion [22,66]. Although oscillations of Src/FAK activation have been predicted by mathematical modeling [67,68], they have not been observed experimentally. The Rho-family GTPases RhoA, Rac1, and Cdc42 have been implicated as master regulators of cell motility by controlling cytoskeleton and FA dynamics and are known to be activated by integrins and mechanical stimuli [69]. Live-cell imaging of biosensors has recently revealed coordinated cyclic activity of these three GTPases at the leading edge of migrating fibroblasts [70]. Although these oscillations could be induced by force fluctuation on integrins, the presence of both positive regulation by GEFs and negative regulation by GAPs and GDIs suggest that a robust biological oscillator in GTPase activity could arise in the absence of external triggers [71,72]. Indeed, RhoA activity oscillations that are critical to leading edge protrusion have been shown to be dependent on cycles of PKA-mediated RhoA phosphorylation, which induces a RhoA-RhoGDI interaction to inhibit Rho activity [73]. Since PKA activation is integrin-dependent, this suggests that cyclic force on integrins could be the upstream activator of these signal oscillations. However, whether signal oscillations are required for motility and how they are decoded to produce directional migration in response to a rigidity gradient is not known.

Cyclic activation of stretch-activated Ca^{2+} channels

Regulation of calcium signaling by tugging forces is also a plausible mechanism for how traction dynamics direct cell migration towards stiff ECM. Ca^{2+} is a well-known master regulator of cell migration, and front-to-rear gradients of intracellular Ca^{2+} underlie polarization and migration during chemotaxis [74,75]. Cytoplasmic Ca^{2+} controls actomyosin by activating myosin II [76], as well as the gelsolin/villin family of F-actin severing proteins [77]. Ca^{2+} also promotes FA turnover by activating calpain family proteases [78], which cleave several FA proteins including talin [79]. Thus, it is possible that a gradient of ECM stiffness could give rise to a gradient of local calcium influx, which in turn could promote local contraction as well as actin filament and FA turnover to cause cells to move towards stiffer ECM.

It is tempting to speculate that intracellular calcium gradients could be produced by stretch-activated calcium channels (SACs), whose activity is induced by repeated tugging on the ECM [80]. SACs are activated by mechanical stimulation, which triggers transient opening and ion flux [81]. Blocking SAC activity suppresses several mechanically-induced cellular responses, including cell migration, FA maturation and traction force development [82,83], suggesting that SACs could both cause traction fluctuations as well as mediate their downstream effects. Importantly, sustained mechanical stimulus does not maintain SAC activity due to channel adaptation [81]; thus, maintenance of a localized Ca^{2+} gradient in cells via SACs would require repeated local mechanical stimulus. Recently, striking transient calcium “flickers” mediated by the TRPM7 SAC were visualized at the front of migrating fibroblasts [84]. Indeed, TRPM7 has been implicated in FA function, as it localizes to FAs and promotes calpain protease activity and FA turnover when overexpressed, while TRPM7 silencing increases FA strength [85]. Thus, fluctuations in traction force at FAs could promote repeated activation of SACs, possibly TRPM7, to maintain locally increased Ca^{2+} at the leading edge, thereby affecting FA and cytoskeletal dynamics to direct cell migration.

Open Questions and Future Perspectives

Although transient changes in FA mechanics have a role in ECM-rigidity sensing and durotaxis, whether this mechanism contributes to other mechanosensitive processes, such as cell differentiation or epithelial-to-mesenchymal transition, remains to be investigated. Demonstrating that tugging forces regulate a specific cellular function, such as durotaxis, without affecting other mechanosensitive responses would be an important advance for future development of therapeutics aimed at modulating ECM-guided cell migration. Another open question is whether soluble factors can modulate migratory or invasive properties of cells by regulating the dynamics of traction forces at FAs. Indeed, although traction fluctuations are required for durotaxis, they strongly decrease the velocity of random cell migration [23]. Thus, growth factors may impact on cell motility through suppression of tugging traction, either by activating actomyosin contractility or by decreasing the strength of FAs [86,87]. In agreement with this hypothesis, decreased FA strength correlates with the onset of cancer cell motility, suggesting a functional link between cell invasiveness and the dynamics of traction forces at FAs [87]. It remains to be determined if traction fluctuations are required for metastasis, in which case they could become a therapeutic target.

Acknowledgments

The authors are grateful to Dr. Jian Liu for helpful discussion on the molecular basis of force fluctuations. We thank Dr. Gregory Alushin for careful reading and commenting on the manuscript. This research was supported by the Division of Intramural Research, National Heart Lung and Blood Institute.

References and recommended reading

Papers of particular interest have been highlighted as:

- of special interest
  - of outstanding interest
1. Petrie RJ, Doyle AD, Yamada KM. Random versus directionally persistent cell migration. *Nat Rev Mol Cell Biol.* 2009; 10:538–549. [PubMed: 19603038]
  - 2••. Lo CM, Wang HB, Dembo M, Wang YL. Cell movement is guided by the rigidity of the substrate. *Biophys J.* 2000; 79:144–152. In this seminal paper that established significance of mechanosensing for cell migration, live cell imaging was used to follow the directional movement of fibroblasts plated on an extracellular matrix with a steep gradient of rigidity. The authors found that migrating cells translocate towards areas of stiffer extracellular matrix and their movement is guided by purely physical interactions with the ECM substrate. [PubMed: 10866943]
  3. Tse JR, Engler AJ. Stiffness gradients mimicking in vivo tissue variation regulate mesenchymal stem cell fate. *PLoS ONE.* 2011; 6:e15978. [PubMed: 21246050]
  4. Choi YS, Vincent LG, Lee AR, Kretschmer KC, Chirasatitsin S, Dobke MK, Engler AJ. The alignment and fusion assembly of adipose-derived stem cells on mechanically patterned matrices. *Biomaterials.* 2012; 33:6943–6951. [PubMed: 22800539]
  5. Guo WH, Frey MT, Burnham NA, Wang YL. Substrate rigidity regulates the formation and maintenance of tissues. *Biophys J.* 2006; 90:2213–2220. [PubMed: 16387786]
  6. de Rooij J, Kerstens A, Danuser G, Schwartz MA, Waterman-Storer CM. Integrin-dependent actomyosin contraction regulates epithelial cell scattering. *J Cell Biol.* 2005; 171:153–164. [PubMed: 16216928]
  7. Flanagan LA, Ju YE, Marg B, Osterfield M, Janmey PA. Neurite branching on deformable substrates. *Neuroreport.* 2002; 13:2411–2415. [PubMed: 12499839]

8. Koch D, Rosoff WJ, Jiang J, Geller HM, Urbach JS. Strength in the periphery: growth cone biomechanics and substrate rigidity response in peripheral and central nervous system neurons. *Biophys J*. 2012; 102:452–460. [PubMed: 22325267]
9. Mandeville JT, Lawson MA, Maxfield FR. Dynamic imaging of neutrophil migration in three dimensions: mechanical interactions between cells and matrix. *J Leukoc Biol*. 1997; 61:188–200. [PubMed: 9021925]
10. Paszek MJ, Zahir N, Johnson KR, Lakins JN, Rozenberg GI, Gefen A, Reinhart-King CA, Margulies SS, Dembo M, Boettiger D, et al. Tensional homeostasis and the malignant phenotype. *Cancer Cell*. 2005; 8:241–254. [PubMed: 16169468]
11. Ulrich TA, de Juan Pardo EM, Kumar S. The mechanical rigidity of the extracellular matrix regulates the structure, motility, and proliferation of glioma cells. *Cancer Res*. 2009; 69:4167–4174. [PubMed: 19435897]
12. Houk AR, Jilkine A, Mejean CO, Boltyanskiy R, Dufresne ER, Angenent SB, Altschuler SJ, Wu LF, Weiner OD. Membrane tension maintains cell polarity by confining signals to the leading edge during neutrophil migration. *Cell*. 2012; 148:175–188. [PubMed: 22265410]
- 13•. Galkin VE, Orlova A, Egelman EH. Actin filaments as tension sensors. *Curr Biol*. 2012; 22:R96–R101. A comprehensive review of recent advances in analysis of filamentous actin conformational change under mechanical tension. [PubMed: 22321312]
- 14•. Hayakawa K, Tatsumi H, Sokabe M. Actin filaments function as a tension sensor by tension-dependent binding of cofilin to the filament. *J Cell Biol*. 2011; 195:721–727. By using an elegant set of biophysical approaches, these authors showed that mechanical forces affect the structure and dynamics of the actin cytoskeleton. Tension applied to an actin filament was shown to decrease its susceptibility to the actin-severing protein cofilin, suggesting that filamentous actin can act as a tension sensor. [PubMed: 22123860]
15. Byfield FJ, Wen Q, Levental I, Nordstrom K, Arratia PE, Miller RT, Janmey PA. Absence of filamin A prevents cells from responding to stiffness gradients on gels coated with collagen but not fibronectin. *Biophys J*. 2009; 96:5095–5102. [PubMed: 19527669]
16. Ehrlicher AJ, Nakamura F, Hartwig JH, Weitz DA, Stossel TP. Mechanical strain in actin networks regulates FilGAP and integrin binding to filamin A. *Nature*. 2011; 478:260–263. [PubMed: 21926999]
17. Kim DH, Khatau SB, Feng Y, Walcott S, Sun SX, Longmore GD, Wirtz D. Actin cap associated focal adhesions and their distinct role in cellular mechanosensing. *Sci Rep*. 2012; 2:555. [PubMed: 22870384]
18. Liu Z, Tan JL, Cohen DM, Yang MT, Sniadecki NJ, Ruiz SA, Nelson CM, Chen CS. Mechanical tugging force regulates the size of cell–cell junctions. *Proc Natl Acad Sci USA*. 2010; 107:9944–9949. [PubMed: 20463286]
19. Hoffman BD, Grashoff C, Schwartz MA. Dynamic molecular processes mediate cellular mechanotransduction. *Nature*. 2011; 475:316–323. [PubMed: 21776077]
20. Moore SW, Roca-Cusachs P, Sheetz MP. Stretchy proteins on stretchy substrates: the important elements of integrin-mediated rigidity sensing. *Dev Cell*. 2010; 19:194–206. [PubMed: 20708583]
21. Raab M, Swift J, Dingal PC, Shah P, Shin JW, Discher DE. Crawling from soft to stiff matrix polarizes the cytoskeleton and phosphoregulates myosin-II heavy chain. *J Cell Biol*. 2012; 199:669–683. [PubMed: 23128239]
22. Wang HB, Dembo M, Hanks SK, Wang YL. Focal adhesion kinase is involved in mechanosensing during fibroblast migration. *Proc Natl Acad Sci USA*. 2001; 98:11295–11300. [PubMed: 11572981]
23. Plotnikov SV, Pasapera AM, Sabass B, Waterman CM. Force fluctuations within focal adhesions mediate ECM-rigidity sensing to guide directed cell migration. *Cell*. 2012; 151:1513–1527. [PubMed: 23260139]
24. He L, Wang X, Tang HL, Montell DJ. Tissue elongation requires oscillating contractions of a basal actomyosin network. *Nat Cell Biol*. 2010; 12:1133–1142. [PubMed: 21102441]
- 25•. Martin AC, Kaschube M, Wieschaus EF. Pulsed contractions of an actin-myosin network drive apical constriction. *Nature*. 2009; 457:495–499. By combining live-cell imaging with quantitative image analysis, this elegant study unraveled a mechanism driving apical constriction

and epithelial cell sheet invagination during *Drosophila* gastrulation. These authors demonstrated repeated pulses of epithelial cell constriction powered by oscillating actomyosin contractions and controlled by the coordinated activation of two transcription factors, Trio and Snail. The authors proposed that the oscillating contractions of the cortical cytoskeleton act as a developmentally regulated cellular ratchet, which drives and controls incremental reduction of the apical cell area to drive tissue morphogenesis. [PubMed: 19029882]

26. Rauzi M, Lenne PF, Lecuit T. Planar polarized actomyosin contractile flows control epithelial junction remodelling. *Nature*. 2010; 468:1101–1114.
27. Schwingel M, Bastmeyer M. Force mapping during the formation and maturation of cell adhesion sites with multiple optical tweezers. *PLoS ONE*. 2013; 8:e54850. [PubMed: 23372781]
28. Tojkander S, Gateva G, Schevzov G, Hotulainen P, Naumanen P, Martin C, Gunning PW, Lappalainen P. A molecular pathway for myosin II recruitment to stress fibers. *Curr Biol*. 2011; 21:539–550. [PubMed: 21458264]
29. Lee CS, Choi CK, Shin EY, Schwartz MA, Kim EG. Myosin II directly binds and inhibits Dbl family guanine nucleotide exchange factors: a possible link to Rho family GTPases. *J Cell Biol*. 2010; 190:663–674. [PubMed: 20713598]
30. Vicente-Manzanares M, Ma X, Adelstein RS, Horwitz AR. Non-muscle myosin II takes centre stage in cell adhesion and migration. *Nat Rev Mol Cell Biol*. 2009; 10:778–790. [PubMed: 19851336]
31. Jülicher F, Prost J. Cooperative molecular motors. *Phys Rev Lett*. 1995; 75:2618–2621. [PubMed: 10059357]
32. Jülicher F, Prost J. Spontaneous oscillations of collective molecular motors. *Phys Rev Lett*. 1997; 78:4510–4513.
33. Giannone G, Dubin-Thaler BJ, Döbereiner H-G, Kieffer N, Bresnick AR, Sheetz MP. Periodic lamellipodial contractions correlate with rearward actin waves. *Cell*. 2004; 116:431–443. [PubMed: 15016377]
34. Ponti A, Machacek M, Guppton SL, Waterman-Storer CM, Danuser G. Two distinct actin networks drive the protrusion of migrating cells. *Science*. 2004; 305:1782–1786. [PubMed: 15375270]
35. Westendorf C, Negrete J, Bae AJ, Sandmann R, Bodenschatz E, Beta C. Actin cytoskeleton of chemotactic amoebae operates close to the onset of oscillations. *Proc Natl Acad Sci USA*. 2013; 110:3853–3858. [PubMed: 23431176]
36. Guppton SL, Eisenmann K, Alberts AS, Waterman-Storer CM. mDia2 regulates actin and focal adhesion dynamics and organization in the lamella for efficient epithelial cell migration. *J Cell Sci*. 2007; 120:3475–3487. [PubMed: 17855386]
37. Price CJ, Brindle NP. Vasodilator-stimulated phosphoprotein is involved in stress-fiber and membrane ruffle formation in endothelial cells. *Arterioscler Thromb Vasc Biol*. 2000; 20:2051–2056. [PubMed: 10978248]
38. Hoffman LM, Jensen CC, Kloeker S, Wang CL, Yoshigi M, Beckerle MC. Genetic ablation of zyxin causes Mena/VASP mislocalization, increased motility, and deficits in actin remodeling. *J Cell Biol*. 2006; 172:771–782. [PubMed: 16505170]
39. Tojkander S, Gateva G, Lappalainen P. Actin stress fibers - assembly, dynamics and biological roles. *J Cell Sci*. 2012; 125:1855–1864. [PubMed: 22544950]
40. Reinhard M, Halbrügge M, Scheer U, Wiegand C, Jockusch BM, Walter U. The 46/50kDa phosphoprotein VASP purified from human platelets is a novel protein associated with actin filaments and focal contacts. *EMBO J*. 1992; 11:2063–2070. [PubMed: 1318192]
41. Guppton SL, Riquelme D, Hughes-Alford SK, Tadros J, Rudina SS, Hynes RO, Lauffenburger D, Gertler FB. Mena binds $\beta 5$ integrin directly and modulates $\beta 5$ function. *J Cell Biol*. 2012; 198:657–676. [PubMed: 22908313]
42. Kuo JC, Han X, Hsiao CT, Yates JR, Waterman CM. Analysis of the myosin-II-responsive focal adhesion proteome reveals a role for β -Pix in negative regulation of focal adhesion maturation. *Nat Cell Biol*. 2011; 13:383–393. [PubMed: 21423176]
43. Schiller HB, Friedel CC, Boulegue C, Fässler R. Quantitative proteomics of the integrin adhesome show a myosin II-dependent recruitment of LIM domain proteins. *EMBO Rep*. 2011; 12:259–266. [PubMed: 21311561]

44. Humphries JD, Byron A, Bass MD, Craig SE, Pinney JW, Knight D, Humphries MJ. Proteomic analysis of integrin-associated complexes identifies RCC2 as a dual regulator of Rac1 and Arp6. *Sci Signal.* 2009; 2(ra51)
- 45•. Oakes PW, Beckham Y, Stricker J, Gardel ML. Tension is required but not sufficient for focal adhesion maturation without a stress fiber template. *J Cell Biol.* 2012; 196:363–374. This study sheds the light on the role of mechanical tension and lamellar actin architecture in focal adhesion maturation. It demonstrated that while inhibiting formin or α -actinin activity suppresses stress fiber assembly, it does not perturb cellular contractility, yet it impedes maturation of focal adhesions. The authors proposed that focal adhesion maturation is not driven by contractility-dependent mechanotransduction pathways alone, but requires the assembly of a stress fiber at the adhesion sites. [PubMed: 22291038]
46. Riveline D, Zamir E, Balaban NQ, Schwarz US, Ishizaki T, Narumiya S, Kam Z, Geiger B, Bershadsky AD. Focal contacts as mechanosensors: externally applied local mechanical force induces growth of focal contacts by an mDia1-dependent and ROCK-independent mechanism. *J Cell Biol.* 2001; 153:1175–1186. [PubMed: 11402062]
- 47•. Smith MA, Blankman E, Gardel ML, Luetjohann L, Waterman CM, Beckerle MC. A zyxin-mediated mechanism for actin stress fiber maintenance and repair. *Dev Cell.* 2010; 19:365–376. This study identified a novel molecular mechanism for rapid repair and maintenance of the structural integrity of the actin cytoskeleton. The authors showed that local damage of stress fibers, induced by a mechanical stress, can be stabilized and repaired by recruiting the LIM protein zyxin, the actin polymerizing factor VASP, and the actin cross-linking protein α -actinin into compromised stress fiber zones. Suppression of stress fiber repair was shown to decrease traction forces on the ECM, suggesting that this mechanism plays an important role in tensional homeostasis. [PubMed: 20833360]
48. Jégou A, Carlier MF, Romet-Lemonne G. Formin mDia1 senses and generates mechanical forces on actin filaments. *Nat Commun.* 2013; 4:1883. [PubMed: 23695677]
- 49•. Shemesh T, Otomo T, Rosen MK, Bershadsky AD, Kozlov MM. A novel mechanism of actin filament processive capping by formin: solution of the rotation paradox. *J Cell Biol.* 2005; 170:889–893. In this paper, Shemesh and colleagues developed a theoretical framework for describing formin-mediated actin filament assembly controlled by elastic stresses. Based on structural organization of an actin-FH2-domain complex, the authors suggested that torsion elastic stresses determine the regime of actin polymerization upon processive capping by a formin dimer. The optimal regime of actin polymerization was shown to require cyclic switch between two distinct modes of capping by formin, which causes oscillations in torsion elastic stresses. [PubMed: 16157699]
50. Shemesh T, Geiger B, Bershadsky AD, Kozlov MM. Focal adhesions as mechanosensors: a physical mechanism. *Proc Natl Acad Sci USA.* 2005; 102:12383–12388. [PubMed: 16113084]
51. Guo WH, Wang YL. Retrograde fluxes of focal adhesion proteins in response to cell migration and mechanical signals. *Mol Biol Cell.* 2007; 18:4519–4527. [PubMed: 17804814]
52. Hu K, Ji L, Applegate KT, Danuser G, Waterman-Storer CM. Differential transmission of actin motion within focal adhesions. *Science.* 2007; 315:111–115. [PubMed: 17204653]
53. Wolfenson H, Bershadsky AD, Henis YI, Geiger B. Actomyosin-generated tension controls the molecular kinetics of focal adhesions. *J Cell Sci.* 2011; 124:1425–1432. [PubMed: 21486952]
54. Grashoff C, Hoffman BD, Brenner MD, Zhou R, Parsons M, Yang MT, McLean MA, Sligar SG, Chen CS, Ha T, et al. Measuring mechanical tension across vinculin reveals regulation of focal adhesion dynamics. *Nature.* 2010; 466:263–266. [PubMed: 20613844]
55. Humphries JD, Wang P, Streuli C, Geiger B, Humphries MJ, Ballestrem C. Vinculin controls focal adhesion formation by direct interactions with talin and actin. *J Cell Biol.* 2007; 179:1043–1057. [PubMed: 18056416]
56. Jiang G, Giannone G, Critchley DR, Fukumoto E, Sheetz MP. Two-piconewton slip bond between fibronectin and the cytoskeleton depends on talin. *Nature.* 2003; 424:334–337. [PubMed: 12867986]
57. Peng X, Nelson ES, Maiers JL, DeMali KA. New insights into vinculin function and regulation. *Int Rev Cell Mol Biol.* 2011; 287:191–231. [PubMed: 21414589]

58. Moser M, Legate KR, Zent R, Fässler R. The tail of integrins, talin, and kindlins. *Science*. 2009; 324:895–899. [PubMed: 19443776]
- 59•. Chan CE, Odde DJ. Traction dynamics of filopodia on compliant substrates. *Science*. 2008; 322:1687–1691. This study combined live-cell imaging and mathematical modelling to reveal the dynamics of traction forces exerted by growth cone filopodia on a compliant substrate. The authors predicted that focal adhesions can exhibit either stable or oscillating force transmission to the extracellular matrix through an adhesion site, and the choice between these states is controlled by ECM stiffness. They demonstrated the predictions by using traction force microscopy in living neuronal growth cones. [PubMed: 19074349]
60. Vogel V, Sheetz MP. Cell fate regulation by coupling mechanical cycles to biochemical signaling pathways. *Curr Opin Cell Biol*. 2009; 21:38–46. [PubMed: 19217273]
61. Kong F, García AJ, Mould AP, Humphries MJ, Zhu C. Demonstration of catch bonds between an integrin and its ligand. *J Cell Biol*. 2009; 185:1275–1284. [PubMed: 19564406]
62. Chen X, Xie C, Nishida N, Li Z, Walz T, Springer TA. Requirement of open headpiece conformation for activation of leukocyte integrin α_2 . *Proc Natl Acad Sci USA*. 2010; 107:14727–14732. [PubMed: 20679211]
63. Friedland JC, Lee MH, Boettiger D. Mechanically activated integrin switch controls $\alpha_5\beta_1$ function. *Science*. 2009; 323:642–644. [PubMed: 19179533]
- 64•. Kong F, Li Z, Parks WM, Dumbauld DW, García AJ, Mould AP, Humphries MJ, Zhu C. Cyclic mechanical reinforcement of integrin-ligand interactions. *Mol Cell*. 2013; 49:1060–1068. This paper describes a phenomenon of ‘cyclic mechanical reinforcement’ of single integrin-ligand bond. The authors used single-molecule AFM to demonstrate that application of cyclic force strengthened the integrin–fibronectin bond and prolonged bond lifetime by two orders of magnitude, maintaining single bonds for minutes. Bond reinforcement was shown to accumulate over repetitive force applications, suggesting that the ability to integrate cyclic forces may allow the cell to ‘count’ the force cycles and therefore sense mechanical properties of the microenvironment. [PubMed: 23416109]
65. Giannone G, Sheetz MP. Substrate rigidity and force define form through tyrosine phosphatase and kinase pathways. *Trends Cell Biol*. 2006; 16:213–223. [PubMed: 16529933]
66. Chan KT, Cortesio CL, Huttenlocher A. FAK alters invadopodia and focal adhesion composition and dynamics to regulate breast cancer invasion. *J Cell Biol*. 2009; 185:357–370. [PubMed: 19364917]
67. Caron-Lormier G, Berry H. Amplification and oscillations in the FAK/Src kinase system during integrin signaling. *J Theor Biol*. 2005; 232:235–248. [PubMed: 15530493]
68. Kaimachnikov NP, Kholodenko BN. Toggle switches, pulses and oscillations are intrinsic properties of the Src activation/deactivation cycle. *FEBS J*. 2009; 276:4102–4118. [PubMed: 19627364]
69. Lessey EC, Guilluy C, Burridge K. From mechanical force to RhoA activation. *Biochemistry*. 2012; 51:7420–7432. [PubMed: 22931484]
70. Machacek M, Hodgson L, Welch C, Elliott H, Pertz O, Nalbant P, Abell A, Johnson GL, Hahn KM, Danuser G. Coordination of Rho GTPase activities during cell protrusion. *Nature*. 2009; 461:99–103. [PubMed: 19693013]
71. Guilluy C, Garcia-Mata R, Burridge K. Rho protein crosstalk: another social network? *Trends Cell Biol*. 2011; 21:718–726. [PubMed: 21924908]
72. Tsai TY, Choi YS, Ma W, Pomerening JR, Tang C, Ferrell JE. Robust, tunable biological oscillations from interlinked positive and negative feedback loops. *Science*. 2008; 321:126–129. [PubMed: 18599789]
- 73•. Tkachenko E, Sabouri-Ghomi M, Pertz O, Kim C, Gutierrez E, Machacek M, Groisman A, Danuser G, Ginsberg MH. Protein kinase A governs a RhoA-RhoGDI protrusion-retraction pacemaker in migrating cells. *Nat Cell Biol*. 2011; 13:660–667. In this study, by using a combination of live-cell imaging of protein activity biosensors and sophisticated image analysis, the authors revealed a role of cyclic activation of cAMP-activated protein kinase A (PKA) in regulation of leading edge protrusion dynamics. The authors demonstrated that PKA governs the timing of cell protrusions by phosphorylating RhoA on Ser¹⁸⁸ residue, which downregulates RhoA activity via RhoGDI-mediated mechanism. Thus, PKA and RhoGDI form a negative

feedback loop that controls the cycling of RhoA activity at the leading edge of migrating cell and acts as a pacemaker of the protrusion-retraction cycle. [PubMed: 21572420]

74. Brundage RA, Fogarty KE, Tuft RA, Fay FS. Calcium gradients underlying polarization and chemotaxis of eosinophils. *Science*. 1991; 254:703–706. [PubMed: 1948048]
75. Hahn K, DeBiasio R, Taylor DL. Patterns of elevated free calcium and calmodulin activation in living cells. *Nature*. 1992; 359:736–738. [PubMed: 1436037]
76. Somlyo AP, Somlyo AV. Signal transduction by G-proteins, Rho-kinase and protein phosphatase to smooth muscle and non-muscle myosin II. *J Physiol*. 2000; 522:177–185. [PubMed: 10639096]
77. Ono S. Mechanism of depolymerization and severing of actin filaments and its significance in cytoskeletal dynamics. *Int Rev Cytol*. 2007; 258:1–82. [PubMed: 17338919]
78. Giannone G, Rondé P, Gaire M, Haiech J, Takeda K. Calcium oscillations trigger focal adhesion disassembly in human U87 astrocytoma cells. *J Biol Chem*. 2002; 277:26364–26371. [PubMed: 12011063]
79. Franco SJ, Huttenlocher A. Regulating cell migration: calpains make the cut. *J Cell Sci*. 2005; 118:3829–3838. [PubMed: 16129881]
80. Kobayashi T, Sokabe M. Sensing substrate rigidity by mechanosensitive ion channels with stress fibers and focal adhesions. *Curr Opin Cell Biol*. 2010; 22:669–676. [PubMed: 20850289]
81. Hamill OP, Martinac B. Molecular basis of mechanotransduction in living cells. *Physiol Rev*. 2001; 81:685–740. [PubMed: 11274342]
82. Lee J, Ishihara A, Oxford G, Johnson B, Jacobson K. Regulation of cell movement is mediated by stretch-activated calcium channels. *Nature*. 1999; 400:382–386. [PubMed: 10432119]
83. Munevar S, Wang Y-L, Dembo M. Regulation of mechanical interactions between fibroblasts and the substratum by stretch-activated Ca^{2+} entry. *J Cell Sci*. 2004; 117:85–92. [PubMed: 14627625]
- 84••. Wei C, Wang X, Chen M, Ouyang K, Song LS, Cheng H. Calcium flickers steer cell migration. *Nature*. 2009; 457:901–905. This study describes the dynamics of calcium influx within a migrating fibroblast. The authors visualized transient calcium microdomains, calcium flickers, in the leading lamella and showed that local activation of the mechano-gated ion channel, TRPM7 at the front of a cell correlates with the direction of cell migration in response to biochemical or physical cues. [PubMed: 19118385]
85. Su LT, Agapito MA, Li M, Simonson WT, Huttenlocher A, Habas R, Yue L, Runnels LW. TRPM7 regulates cell adhesion by controlling the calcium-dependent protease calpain. *J Biol Chem*. 2006; 281:11260–11270. [PubMed: 16436382]
86. Schneider IC, Hays CK, Waterman CM. Epidermal growth factor–induced contraction regulates paxillin phosphorylation to temporally separate traction generation from de-adhesion. *Mol Biol Cell*. 2009; 20:3155–3167. [PubMed: 19403690]
87. Lynch L, Vodyanik PI, Boettiger D, Guvakova MA. Insulin-like growth factor I controls adhesion strength mediated by $\alpha_5\beta_1$ integrins in motile carcinoma cells. *Mol Biol Cell*. 2005; 16:51–63. [PubMed: 15509657]

Figure 1. Nanoscale fluctuations of traction forces mediate ECM rigidity sensing and guide directed cell migration. Dynamics of traction forces within individual FAs are essential to direct cells towards stiff ECM. Zoomed insert depicts repetitive movement of force peak along individual FA (shown in green).

Figure 2.

Possible molecular mechanisms mediating dynamics of traction forces within FAs. **(a)** Fluctuations of traction forces due to oscillations in local actomyosin contractility. The model shows minifilaments of non-muscle myosin II closely associated with FAs. Mathematical modeling predicts that activity of these local contractile units could oscillate spontaneously in a stiffness-dependent manner. **(b)** Fluctuations of traction forces due to cyclic variation in actin assembly at stress fibers. The model shows actin filaments within a stress fiber undergoing cycles of elongation mediated by a putative actin elongation factor, possibly Ena/VASP or formin family proteins. **(c)** Fluctuations of traction forces due to temporal variation in engagement between the actin cytoskeleton and the ECM. The model shows actin filaments being variably engaged to integrins via talin and vinculin.