Published in final edited form as:

World J Pediatr Congenit Heart Surg. 2013 January ; 4(1): . doi:10.1177/2150135112461924.

REPORT FROM THE STS NATIONAL DATABASE WORK FORCE:

Clarifying the Definition of Operative Mortality

David M. Overman, MD¹, Jeffrey P. Jacobs, MD², Richard L. Prager, MD³, Cameron D. Wright, MD⁴, David R. Clarke, MD⁵, Sara Pasquali, MD, MHS⁶, Sean M. O'Brien, PhD⁷, Rachel S. Dokholyan, MPH⁸, Paul Meehan⁸, Donna E. McDonald, MPH, RN⁹, Marshall L. Jacobs, MD¹⁰, Constantine Mavroudis, MD¹¹, and David M. Shahian, MD⁴

¹The Children's Heart Clinic, Children's Hospitals and Clinics of Minnesota, Minneapolis, Minnesota

²The Congenital Heart Institute of Florida (CHIF), All Children's Hospital, University of South Florida College of Medicine, Saint Petersburg and Tampa, Florida

³University of Michigan, Ann Arbor, Michigan

⁴Massachusetts General Hospital, Harvard Medical School, Boston, Massachusetts

⁵The Children's Hospital, University of Colorado Denver School of Medicine, Aurora, Colorado

⁶Departments of Pediatrics, Duke University School of Medicine, and Duke Clinical Research Institute, Durham, North Carolina

⁷Departments of Biostatistics, Duke University School of Medicine, and Duke Clinical Research Institute, Durham, North Carolina

⁸Duke University School of Medicine, and Duke Clinical Research Institute, Durham, North Carolina

⁹The Society of Thoracic Surgeons, Chicago, Illinois

¹⁰Center for Pediatric and Congenital Heart Diseases, Children's Hospital, Cleveland Clinic, Cleveland, Ohio

¹¹The Congenital Heart Institute of Florida (CHIF), Walt Disney Pavilion Florida Hospital for Children, Orlando, Florida

Abstract

Several distinct definitions of postoperative death have been used in various quality reporting programs. Some have defined a postoperative mortality as a patient who expires while still in the hospital, while others have considered all deaths occurring within a predetermined, standardized time interval after surgery. While if continues to collect mortality data using both these individual definitions, the Society of Thoracic Surgeons (STS) believes that either alone may be inadequate. Accordingly, the STS prefers a more encompassing metric, Operative Mortality, which is defined as (1) all deaths occurring during the hospitalization in which the operation was performed, even if after 30 days; and (2) all deaths occurring after discharge from the hospital, but before the end of the thirtieth postoperative day. This manuscript provides clarification for some uncommon but important scenarios where the correct application of this definition may be problematic.

Keywords

cardiac disease; thoracic disease; congenital heart disease; outcomes analysis; quality improvement; database; mortality

Mortality is the most commonly assessed outcome in healthcare. Definitions of mortality have differed according to the specific disease or condition, the reporting entity, and data availability. The two most common definitions are in-hospital death, which is easily and accurately collected, but which may be impacted by local transfer practices and the availability of post-discharge extended care; and mortality within a standardized time frame (e.g., 30 days) following admission or procedure. The latter is advocated by the American Heart Association's Standards for Statistical Models Used for Public Reporting [1]. This approach is less impacted by differences in local post-acute care facilities and it also mitigates the potential for gaming through postoperative transfer of critically ill patients to another facility. The major disadvantages of 30-day mortality reporting are the loss of those deaths that occur after 30 days but within the index hospitalization; the difficulty and expense associated with accurate collection of such data; and the need for very precise definitions of critical time points, the major focus of this brief report.

Recognizing the advantages and disadvantages of these individual definitions, the Society of Thoracic Surgeons (STS) believes that each, used in isolation, is inadequate. As the primary mortality metric for its provider feedback and voluntary public reporting initiatives, STS employs a more inclusive definition that encompasses all the patients who would be defined by *either* in-hospital or 30-day mortality. This metric is Operative Mortality, and it is currently the most frequently reported outcome measure used by STS. The purpose of this manuscript is to clarify the definition of Operative Mortality used by STS.

Current Definitions

The STS Congenital Heart Surgery Database tracks three mortality metrics, utilizing the following definitions [2, 3, 4]:

1. Mortality Status At Hospital Discharge

“Indicate whether the patient was Alive or Dead at date and time of “Date of Hospital Discharge” for this operation.”

2. Mortality - 30-Day Status

“Indicate whether the patient was alive or dead on the 30th day post surgical procedure whether in hospital or not.”

3. Mortality - Operative Death

“Operative Mortality: Includes both (1) all deaths occurring during the hospitalization in which the operation was performed, even if after 30 days; and (2) those deaths occurring after discharge from the hospital, but within 30 days of the procedure.”

The STS Adult Cardiac Surgery Database also separately collects both in-hospital and 30-day mortality status and defines Operative Mortality as:

“(1) all deaths occurring during the acute episode of care in which the operation was performed (this includes patients transferred to other acute care facilities), even if after 30 days; and (2) those deaths occurring after discharge from the hospital, but within 30 days of the procedure unless the cause of death is clearly unrelated to the operation.”

The STS General Thoracic Database also reports both discharge mortality and 30 day mortality. In the STS General Thoracic Database Report, operative mortality is defined for the pulmonary resection model as “in hospital death or death within 30 days of surgery.”

Defining the problem

Occasionally, 30-day mortality rates present definitional issues. For example, a patient who was alive at the start of the 30th postoperative day, but died later during the same day, can be considered to have been both alive and dead on the 30th day post procedure, a potentially serious inconsistency in interpretation.

In addition, the unit of measure of postoperative days has not been adequately defined and might be:

1. 24 hour increments of time after the operation, or
2. calendar days, beginning at midnight on the day of the operation or conceivably even the first day after the operation.

The Solution

STS recommends that the unit of measure of postoperative days should be calendar days beginning the first midnight after the patient enters the operating room. Using this strategy, the first postoperative day would be the day after the operation, which is how most surgeons currently count these days. Consider a patient who enters the operating room at 8 am, comes out of the operating room at 2 PM, and dies at 8 PM of the same day; this death occurs on postoperative day (POD) zero. Meanwhile, consider a patient who enters the operating room at 8 PM, leaves the operating room at 11:45 PM, and dies 30 minutes later at 12:15 AM; this death this occurs on POD 1.

To address issues related to these uncommon but important timing scenarios, STS has adopted the following clarifications of the aforementioned definitions:

1. *Mortality - Operative Death*

Operative Mortality is defined as (1) all deaths occurring during the hospitalization in which the operation was performed, even if after 30 days; and (2) all deaths occurring after discharge from the hospital, but *before the end of the thirtieth postoperative day*.

2. *Mortality - 30-Day Status*

Indicate whether the patient was alive or dead *at the end of the 30th postoperative day* whether in hospital or not.

3. “*Date of Surgery*” is defined as the date of the data field “Time Patient Entered the OR”, and

4. *Calendar days* are the unit of measure for postoperative days. Calendar days will begin and end at midnight in the time zone of the hospital where the patient had surgery. The first postoperative day will commence at the first midnight after the patient enters the operating room (the first calendar date after the date of the field of data “Time Patient Entered the OR”).

This methodology may result in a patient emerging from the operating room on POD 1 (e.g. if their operation began at 11 PM and concluded at 2 AM). Situations such as this have generated considerable discussion in the past, as there was no clear Database documentation

on this issue. With the current clarification, such cases can now be accurately and consistently coded.

These new definitions also have implications for the measurement of postoperative length of stay (PLOS). Based on the rationale described above, PLOS is defined as the number of days from date of operation to date of discharge. By the new definitions, PLOS may be zero if the patient dies on the date of surgery, either in the operating room or immediately thereafter.

Accurate and consistent measurement of outcomes is a central pursuit of STS, and Operative Mortality is the single most important metric. The STS Database is increasingly used to measure quality [5, 6, 7] and document variation in outcomes [8, 9]; therefore, it is critical that definitions are precisely specified. With this clarification, one potential ambiguity in coding practices has been addressed.

References

1. Krumholz HM, Brindis RG, Brush JE, et al. Standards for statistical models used for public reporting of health outcomes: an American Heart Association Scientific Statement from the Quality of Care and Outcomes Research Interdisciplinary Writing Group: cosponsored by the Council on Epidemiology and Prevention and the Stroke Council. Endorsed by the American College of Cardiology Foundation. *Circulation*. 2006; 113(3):456–462. [PubMed: 16365198]
2. [Accessed January 25, 2011] STS Congenital Heart Surgery Database Data Specifications Version 3.0. [http://www.sts.org/sites/default/files/documents/pdf/CongenitalDataSpecificationsV3_0_20090904.pdf]
3. Jacobs JP, Mavroudis C, Jacobs ML, Maruszewski B, Tchervakov CI, Lacour-Gayet FG, Clarke DR, Yeh T, Walters HL 3rd, Kurosawa H, Stellin G, Ebels T, Elliott MJ. What is Operative Mortality? Defining Death in a Surgical Registry Database: A Report from the STS Congenital Database Task Force and the Joint EACTS-STS Congenital Database Committee. *The Annals of Thoracic Surgery*. May; 2006 81(5):1937–41. [PubMed: 16631716]
4. Jacobs JP, Jacobs ML, Mavroudis C, Maruszewski B, Tchervakov CI, Lacour-Gayet FG, Clarke DR, Yeh T, Walters HL 3rd, Kurosawa H, Stellin G, Ebels T, Elliott MJ, Vener DF, Barach P, Benavidez OJ, Bacha EA. What is Operative Morbidity? Defining Complications in a Surgical Registry Database: A Report from the STS Congenital Database Task Force and the Joint EACTS-STS Congenital Database Committee. *The Annals of Thoracic Surgery*. Oct.2007 84:1416–1421.
5. Jacobs JP, Jacobs ML, Austin EH, Mavroudis M, Pasquali SK, Lacour-Gayet FG, Tchervakov CI, Walters HW III, Bacha EA, del Nido PJ, Fraser CD, Gaynor JW, Hirsch JC, Morales DLS, Pourmoghadam KK, Tweddell JT, Prager RL, Mayer JE. Quality Measures for Congenital and Pediatric Cardiac Surgery. *The World Journal for Pediatric and Congenital Heart Surgery* 2012. Jan; 2012 3(1):32–47.
6. O'Brien SM, Clarke DR, Jacobs JP, Jacobs ML, Lacour-Gayet FG, Pizarro C, Welke KF, Maruszewski B, Tobota Z, Miller WJ, Hamilton L, Peterson ED, Mavroudis C, Edwards FH. An empirically based tool for analyzing mortality associated with congenital heart surgery. *The Journal of Thoracic and Cardiovascular Surgery*. 2009 Nov; 138(5):1139–53. November 2009. [PubMed: 19837218]
7. Jacobs ML, O'Brien SM, Jacobs JP, Mavroudis C, Lacour-Gayet FG, Pasquali SK, Welke KF, Pizarro C, Tsai F, Clarke DR. An empirically based tool for analyzing morbidity associated with operations for congenital heart disease. *The Journal of Thoracic and Cardiovascular Surgery*. In Review.
8. Jacobs JP, O'Brien SM, Pasquali SK, Jacobs ML, Lacour-Gayet FG, Tchervakov CI, Austin EH 3rd, Pizarro C, Pourmoghadam KK, Scholl FG, Welke KF, Mavroudis C, Richard E. Clark Paper: Variation in Outcomes for Benchmark Operations: An analysis of the Society of Thoracic Surgeons Congenital Heart Surgery Database. Richard Clark Award recipient for best use of the STS Congenital Heart Surgery Database. *The Annals of Thoracic Surgery*. 2011 Dec; 92(6):2184–92. December 2011. [PubMed: 22115229]

9. Jacobs JP, O'Brien SM, Pasquali SK, Jacobs ML, Lacour-Gayet FG, Tchervenkov CI, Austin EH, Pizarro C, Pourmoghadam KK, Scholl FG, Welke KF, Gaynor JW, Clarke DR, Mayer JE, Mavroudis C. Variation in Outcomes for Complexity-stratified Pediatric Cardiac Surgical Operations: An analysis of the STS Congenital Heart Surgery Database. *The Annals of Thoracic Surgery*. Accepted for publication, In press.