Case Report

Video-Assisted Thoracoscopic Surgery for Localized Neurofibroma of the Esophagus: Case Report and Review of the Literature

Katsunori Nishikawa¹, Noburo Omura¹, Masami Yuda¹, Yujiro Tanaka¹, Akira Matsumoto¹, Yuichiro Tanishima¹, Yoshiro Ishibashi¹, Katsuhiko Yanaga¹, Masahiro Ikegami²

¹Department of Surgery and ²Department of Pathology, Jikei University, Tokyo, Japan

Esophageal submucosal tumors are less common than other gastrointestinal tract tumors. Leiomyoma is the most common benign esophageal SMT, accounting for more than 70% of these tumors. We report on a case of a 56-year-old woman with a 3-cm diameter midthoracic esophageal submucosal tumor. Magnetic resonance imaging suggested leiomyoma or neurofibroma. Video-assisted thoracoscopic surgery was performed to enucleate the tumor from the esophageal wall by splitting the muscle layers. The postoperative course was uneventful, and the patient was discharged on postoperative day 8. Immunohistochemical staining confirmed the diagnosis of esophageal neurofibroma. Gastrointestinal tract involvement of neurofibromatous lesions is rare and occurs most frequently as a systemic manifestation of von Recklinghausen disease. Cases of localized esophageal neurofibroma with prior or subsequent evidence of generalized neurofibromatosis have rarely been documented. This is a rare case of isolated esophageal neurofibroma without classic systemic manifestations of generalized neurofibromatosis, and it is the first reported case treated by video-assisted thoracoscopic surgery.

Key words: Neurofibroma – Esophagus – Thoracoscopic surgery

Benign esophageal tumors are relatively rare, and leiomyomas account for most of these tumors. Rarely, gastrointestinal stromal tumors, papillomas, hemangioma, granular cell tumors, neurofibromas, and myxofibromas occur. Neurofibromas are generally associated with hereditary diseases and are usually a manifestation of von Recklinghausen disease (VRD).¹ However, neurofi-

Reprint requests: Katsunori Nishikawa, Department of Surgery, Jikei University School of Medicine, 3-25-8, Nishi-Shinbashi Minato-ku, Tokyo 1058471, Japan.

Tel.: 03 3433 1111 (ext. 3401); Fax: 03 5472 4140; E-mail: ka2nissy@gmail.com

Fig. 1 Endoscopic examination revealed an esophageal tumor at the midthoracic esophagus, which was covered with normal esophageal epithelium.

bromas rarely occur in the gastrointestinal tract (GI) as isolated tumors outside the classic clinical picture of the VRD.²⁻⁸

We report on a rare case of isolated esophageal neurofibroma in a patient without systemic manifestations of generalized neurofibromatosis, which is the first reported case treated by video-assisted thoracoscopic surgery (VATS).

Case Report

A 56-year-old woman underwent GI endoscopy for epigastric discomfort at an outpatient clinic in her neighborhood. Endoscopic examination revealed a 2-cm-long submucosal tumor (SMT) in the mid-thoracic esophagus. Since then, the tumor size was examined annually by GI endoscopy. GI endoscopy in the fourth year showed tumor enlargement, and the patient was eventually referred to our institution. Endoscopic examination was repeated for precise tumor review. The esophageal SMT measuring 3 cm in diameter was located in the midthoracic esophageal region 25 cm from the incisor teeth (Fig. 1). The tumor appeared to have grown approximately 1 cm relative to the size observed the previous year. Endoscopic ultrasonography demonstrated a hypoechoic and homogeneous mass in the submucosal layer. Following endoscopic ultraso-

Fig. 2 Chest computed tomography shows a well-circumscribed tumor (arrows) measuring $25 \times 33 \times 24$ mm located in the posterior mediastinum at the level of the tracheal bifurcation (A). A heterogeneous inner structure was revealed by magnetic resonance imaging (B).

nography, a fine-needle aspiration biopsy of the tumor was performed. However, the biopsy specimen size was too small for diagnosis. Computed tomographic and magnetic resonance imaging revealed a well-circumscribed tumor measuring $25 \times 33 \times 24$ mm located in the posterior mediastinum at the level of tracheal bifurcation, and with a heterogeneous inner structure (Fig. 2).

Radiologic and endoscopic findings suggested esophageal leiomyoma or neurogenic tumor. Although the patient did not show any clinical symptoms, surgical treatment was chosen because of the tumor enlargement.

The patient underwent enucleation of the tumor by VATS. A small 2.5-cm diameter thoracotomy was first created at the right fifth intercostal space on the anterior axillary line, with the patient in the left lateral decubitus position. Four additional thoracic ports (11.5 mm) were created on the posterior axillary line at the fifth intercostal spaces, medial

Fig. 3 Macroscopic appearance of the tumor. The resected tumor was soft in elasticity and measured $34 \times 28 \times 22$ mm.

axillary line at the fourth and seventh intercostal spaces, and proximal axillary line at the third intercostal line for the endoscope, lung retraction, and surgical assistance, respectively. After the esophagus was freed from the posterior mediastinal space, the tumor was found to bulge dorsally from the upper thoracic esophagus. A Sengstaken-Blakemore tube was inserted orally, and esophageal balloon inflation was used to push the tumor outward from the esophagus. The tumor was easily enucleated from the esophageal mucosa and muscular layer. The esophageal muscular layer was closed roughly with 3 single stitches using extracorporeal knotting technique. The resected tumor was soft in elasticity, it measured $34 \times 28 \times 22$ mm, and the cut surface was whitish (Fig. 3). Histologic examination revealed that the tumor was ill defined with indistinct borders and comprised mixed fibrillary collagen sheets and cords of spindle cells with nodular growth. No signs of atypia or significant mitotic activity were observed. Immunohistochemical staining results were positive for S-100 but negative for c-KIT, CD34, α SMA, and desmin (Fig. 4). The above morphologic and immunohistochemical characteristics were consistent with the diagnosis of neurofibroma.

Postoperative nasogastric tube or esophageal contrast study was considered unnecessary because

Fig. 4 Histologic appearance of the tumor. A tumor comprised a mixture of fibrillary collagen sheets and cords of spindle cells with nodular growth (A). Immunohistochemical staining results were positive for S-100 (B).

the tumor was enucleated without mucosal damage and additional muscular coverage was obtained. Therefore, the patient was allowed to consume water on the next day after surgery and a solid meal on the second day after surgery. The postoperative course was uneventful, and the patient was discharged on the eighth postoperative day. She remains tumor free without any postoperative complications 2 years after surgery.

Discussion

Esophageal SMTs are less common than other GI tract tumors and are classified as benign or malignant disease. Gastrointestinal stromal tumors are the most common malignant SMTs. Leiomyoma is the most common benign esophageal SMT, accounting for more than 70% of these tumors.⁹ Neurofibromas associated with esophageal SMTs have been reported by Plachta¹⁰ to be 0.9%.

Neurofibromas are generally associated with hereditary diseases and are usually manifestations of VRD.¹ They are associated with benign tumors comprising neural and connective tissue components, such as Schwann and perineural cells and myofibroblasts. Gastrointestinal involvement is reported in 25% patients with VRD.¹¹ There are 3 types of neurofibromas: localized, diffuse, and plexiform. Localized neurofibromas of GI tracts, unlike VRD, are benign nerve sheath tumor in the peripheral nervous system originating from the Auerbach plexus and Meissner submucosal plexus.^{12,13} Our patient had a localized neurofibroma,

and a type that is far more common in GI tracts. Cases of localized neurofibromas of the esophagus with prior or subsequent evidence of generalized neurofibromatosis, including our case, have rarely been documented.²⁻⁸ According to our Medline search, 1 case of plexiform neurofibroma of the esophagus was reported in association with neurofibroma and VRD.¹⁴ Most neurofibromas are benign tumors; however, nodular and particularly plexiform types may demonstrate a malignant transformation in 2% to 16% of affected individuals.¹⁵

We found 16 reported cases of localized esophageal neurofibromas, and Fujita et al² and other researchers³⁻⁸ summarized 11 cases in the literature. The mean age of the affected patients was 51.8 years (range, 26–75 years), and there was no gender preference (male to female, 7:9). Symptoms of dysphagia were observed in 32% of patients. Tumors were located mainly in the upper to middle sections of the thoracic esophagus (83%), and the mean tumor size was 5.7 cm (range, 0.5–22.5 cm), whereas leiomyomas were located in the middle to distal regions of the esophagus.¹⁶

Histopathologic diagnosis of neurofibroma is necessary in most cases because neurofibroma or other submucosal tumors, such as leiomyoma, are often difficult to diagnose using diagnostic imaging. The histopathologic characteristic appearance of neurofibroma mainly comprises spindle-shaped cells associated with collagen fibrils. In addition, immunohistochemical staining helps distinguish neurogenic from myogenic tumors. In our case, the tumor comprised a mixture of fibrillary collagen sheets and cords of spindle cells with modularly growth; it was positive for S-100 but negative for c-KIT, CD34, α SMA, and desmin.

The treatment for localized neurofibromas in GI tracts is primarily surgery and depends on the location and size of the lesions. Solitary well-circumscribed lesions may be only incidental findings at the time of screening endoscopy and require no further therapy, but larger solitary lesions may be noticed clinically because of intestinal obstruction or impingement on adjacent structures, and they may require resection. In fact, 14 of 16 reported cases were surgically treated. In the case of esophageal neurofibromatosis, especially associated with VRD, surgical resection should be performed promptly even if there are no clinical symptoms because malignant degeneration may occur prior to surgery. In our case, we selected surgical treatment because the tumor had enlarged over a relatively short

period and may have had the potential for malignant transformation.

To date, all reported surgical treatments have been performed through thoracotomy or laparotomy. Our case appears to be the first where VATS was used for successfully treating a patient with esophageal neurofibroma. Surgical procedures can be performed in a manner similar to those used for leiomyomas and other benign tumors.¹⁷

We conclude that enucleation of esophageal neurofibromas is feasible and is an ideal indication for a thoracoscopic approach. However, the procedure should be performed with extra care, especially if malignancy is suspected.

References

1. Hochberg FH, Dasilva AB, Galdabini J, Richardson EP Jr. Gastrointestinal involvement in von Recklinghausen's neurofibromatosis. *Neurology* 1974; **24**(12):1144–1151
2. Fujita H, Kako H, Misumi A, Harada K, Ishimaru Y, Ogawa M. A resected case of esophageal neurofibroma, and review of the Japanese literature. *Nippon Shokakibyo Gakkai Zasshi* 1993; **90**(4): 795–799
3. Saitoh K, Nasu M, Kamiyama R, Hatakeyama S, Maruyama M, Tsuruta K. Solitary neurofibroma of the esophagus. *Acta Pathol Jpn* 1985; **35**(2):527–531
4. Madrid G, Pardo J, Perez C, Pereda G, Galbe R, Ros LH. The neurofibroma of the oesophagus. *Eur J Radiol* 1986; **6**(1):67–69
5. Hishikawa Y, Miura T, Kakudo K, Matsushita H. Neurofibroma of the esophagus. *Radiat Med* 1984; **2**(4):224–225
6. Ishii T, Nitta M, Masaki T, Nakayama T. A case of multiple neurofibroma of the larynx and cervical esophagus. *Acta Otolaryngol Suppl* 2002; **547**:54–56
7. Lee R, Williamson WA. Neurofibroma of the esophagus. *Ann Thorac Surg* 1997; **64**(4):1173–1174
8. Ramírez Rodríguez JM, Deus Fombellida J, Lozano Mantecón R, Galle Vicente P. Solitary neurofibroma of the esophagus. *Rev Esp Enferm Dig* 1992; **82**(1):47–49
9. Seremetis MG, Lyons WS, de Guzman VC, Peabody JW Jr. Leiomyomata of the esophagus: an analysis of 838 cases. *Cancer* 1976; **38**(5):2166–2177
10. Plachta A. Benign tumors of the esophagus: review of literature and report of 99 cases. *Am J Gastroenterol* 1962; **38**: 639–652
11. Gogus S, Sarikayalar F, Akcoren Z, Yalnizoglu D, Hicsonmez A. Intestinal involvement and vasculopathy in von Recklinghausen's neurofibromatosis. *Turk J Pediatr* 1997; **39**:117–122
12. Butler D, Hanna E. Neurogenic tumor of the rectum. *Dis Colon Rectum* 1959; **2**(3):291–294
13. Grodsky L. Neurofibroma of the rectum in a patient with von Recklinghausen's disease. *Am J Surg* 1958; **5**(3):474–476

14. Ganeshan A, Hon LQ, Soonawalla Z, De'Costa H. Plexiform neurofibroma of the oesophagus: a mimicker of malignancy. *Br J Radiol* 2005;**78**(936):1095–1097
15. Reynolds RM, Browning GG, Nawroz I, Campbell IW. Von Recklinghausen's neurofibromatosis: neurofibromatosis type 1. *Lancet* 2003;**361**(9368):1552–1554
16. Punpale A, Rangole A, Bhambhani N, Karimundackal G, Desai N, de Souza A *et al.* *Leiomyoma of esophagus.* *Ann Thorac Cardiovasc Surg* 2007;**13**(2):78–81
17. Schmid RA, Schob OM, Klotz HP, Vogt P, Weder W. VATS resection of an oesophageal leiomyoma in a patient with neurofibromatosis Recklinghausen. *Eur J Cardiothorac Surg* 1997;**12**(4):659–662