CORRESPONDENCE

Fear of Dental Treatment—an Underrecognized Symptom in People With Impaired Mental Health

by Maria Lenk, PD Dr. rer. medic. Hendrik Berth,
Prof. Dr. med. Peter Joraschky, PD Dr. phil. Katja Petrowski,
Prof. Dr. med. Kerstin Weidner, Prof. Dr. med. dent. Christian Hannig
in volume 31–32/2013

Not a Disease Entity

Patients with underlying psychological disorders are highly significantly more likely to experience fear of dental treatment than healthy persons (1). For the affected patients, anesthesiological support with general anesthesia for dental treatment is often very important, as otherwise, avoidance behaviors may be expected to result in severe dental disorders with possible deterioration of the psychological disorder.

In contrast to phobia, “fear of dental treatment” is not in itself a disease. No indication exists for uncritically prescribing general anesthesia (2), especially in view of a multitude of other, effective therapeutic measures. Health insurers and associations of health insurance companies understood this as early as in 2007 and will reimburse anesthetic services for dental procedures in patients with fear of dental treatment only in cases where a specialist has certified the presence of a psychological or psychiatric disorder (Chapter 5, EBM).

Giving general anesthesia for dental treatment is often trivialized as being more comfortable for the patient, but from the perspective of anesthesiology a critical view is indicated because general anesthesia is not suitable for helping to reduce fear of dental treatment and additionally entail the risk of serious complications. Dental treatment can be more difficult in inactive patients (3), and additional medical equipment and space would be required in the dental surgery.

Regardless of all this, the indication for general anesthesia is uncontroversial, among others, for dental treatment in children younger than 12 years, in patients with disabilities, in cases where local anesthesia is contraindicated, or in cases of major surgery.

The indication “fear” should, however, be reserved exclusively for patients with psychiatric disorders who are receiving treatment. No anxiety disorder should be diagnosed and certified in healthy patients for billing purposes.

It is imperative, however, that patients with a phobia are referred for early detection and treatment. This is the only way in which such patients can be provided with adequate dental treatment and oral health.

DOI: 10.3238/arztebl.2013.0751a

REFERENCES

1. Lenk M, Berth H, Joraschky P, Petrowski K, Weidner K, Hannig C: Fear of dental treatment—an underrecognized symptom in people with impaired mental health. *Dtsch Arztebl Int* 2013; 110(31–32): 517–22.
2. Gordon D: A critical review of approaches to treatment of dental anxiety in adults. *J Anxiety Disord* 2013; 27: 365–78
3. Lee HH: Trends in death associated with pediatric dental sedation and general anesthesia. *Pediatr Anaesth* 2013; (23): 741–746.

On behalf of the board of the interdisciplinary working group Zahnärztliche Anästhesie (IAZA, dental anesthesia) of the specialist medical societies DGZMK and DGA

Prof. Dr. med. Grietje Beck

Klinik für Anästhesiologie, Intensivmedizin und Schmerztherapie
Dr. Horst Schmidt Kliniken GmbH
Klinikum der Landeshauptstadt Wiesbaden
grietje.beck@hsk-wiesbaden.de

In Reply:

The focus of our original article (1) was on the high rates of fear of dental treatment in patients with mental disorders. We pointed out possible options for improving the early detection of such dental anxieties. A critical evaluation of therapeutic options and indications was not the subject of our study.

Professor Beck raises an important issue when she discusses the indication for general anesthesia in patients with dental phobia, which doctors and dentists have to handle in a responsible fashion. We mostly agree with her sentiments. In view of the risks the indication should be defined critically and other therapeutic options considered.

Anesthetics cannot be used to treat the causes of any fear or anxiety, whereas psychotherapeutic interventions represent promising therapeutic options (2). If such therapeutic services are used early then highly acute, and complication-prone, dental situations can be prevented, which often make general anesthesia inevitable.

Fear of dental treatment is easily detected in routine clinical practice (1). Such fears should be considered especially in patients with posttraumatic stress disorder, anxiety disorders, or depressive disorders, as they are often associated with just such disorders (1).

Doctors from all medical specialties who are in contact with such patients can help with early detection. We were particularly pleased to receive questions from general practitioners and anesthesiologists who emphatically expressed their interest in psychotherapeutic interventions.

Identifying fear of dental treatment early is an important task, and appropriate measures should be taken, relative to the severity of the fear.

Not all fear patients require long term therapy, but less severe cases should also be taken seriously.

Everybody should benefit equally from the advances of modern dentistry. This entails enabling and supporting patients to make use of preventive measures, such as their annual checkup with their dentist, especially if they have existing fears. DOI: 10.3238/arztebl.2013.0751b

REFERENCES

1. Lenk M, Berth H, Joraschky P, Petrowski K, Weidner K, Hannig C: Fear of dental treatment—an underrecognized symptom in people with impaired mental health. *Dtsch Arztebl Int* 2013; 110(31–32): 517–22.
2. Kvale G, Berggren U, Milgrom P: Dental fear in adults: a meta-analysis of behavioral interventions. *Community Dent Oral Epidemiol* 2004; 32: 250–64.

Maria Lenk

Prof. Dr. med. Joraschky

Prof. Dr. med. Weidner

Klinik und Poliklinik für Psychotherapie und Psychosomatik
Universitätsklinikum Carl Gustav Carus Dresden
maria.lenk@mailbox.tu-dresden.de

Prof. Dr. med. dent. Hannig

Poliklinik für Zahnerhaltung
Universitätsklinikum Carl Gustav Carus Dresden

Conflict of interest statement

The authors of both contributions declare that no conflict of interest exists.