Published in final edited form as:

Bone. 2013 November ; 57(1): . doi:10.1016/j.bone.2013.07.030.

Dynamic Hydraulic Fluid Stimulation Regulated Intramedullary Pressure

Minyi Hu, Ph.D., Frederick Serra-Hsu, Ph.D., Neville Bethel, B.E./B.S., Liangjun Lin, M.S., Suzanne Ferreri, Ph.D., Jiqi Cheng, Ph.D., and Yi-Xian Qin, Ph.D.*

Department of Biomedical Engineering Stony Brook University Stony Brook, NY 11794-5281

Abstract

Physical signals within bone, i.e. generated from mechanical loading, have the potential to initiate skeletal adaptation. Strong evidence has pointed to bone fluid flow (BFF) as a media between an external load and the bone cells, in which altered velocity and pressure can ultimately initiate the mechanotransduction and the remodeling process within bone. Load-induced BFF can be altered by factors such as intramedullary pressure (ImP) and/or bone matrix strain, mediating bone adaptation. Previous studies have shown that BFF induced by ImP alone, with minimum bone strain, can initiate bone remodeling. However, identifying induced ImP dynamics and bone strain factor *in vivo* using a non-invasive method still remains challenging. To apply ImP as a means for alteration of BFF, it was hypothesized that non-invasive dynamic hydraulic stimulation (DHS) can induce local ImP with minimal bone strain to potentially elicit osteogenic adaptive responses via bone-muscle coupling. The goal of this study was to evaluate the immediate effects on local and distant ImP and strain in response to a range of loading frequencies using DHS. Simultaneous femoral and tibial ImP and bone strain values were measured in three 15-month-old female Sprague Dawley rats during DHS loading on the tibia with frequencies of 1Hz to 10Hz. DHS showed noticeable effects on ImP induction in the stimulated tibia in a nonlinear fashion in response to DHS over the range of loading frequencies, where peaked at 2Hz. DHS at various loading frequencies generated minimal bone strain in the tibiae. Maximal bone strain measured at all loading frequencies was less than 8μ . No detectable induction of ImP or bone strain was observed in the femur. This study suggested that oscillatory DHS may regulate the local fluid dynamics with minimal mechanical strain in bone, which serves critically in bone adaptation. These results clearly implied DHS's potential as an effective, non-invasive intervention for osteopenia and osteoporosis treatments.

Keywords

bone fluid flow; hydraulic fluid stimulation; intramedullary pressure; bone remodeling; mechanical loading; loading rate

© 2013 The Authors. Published by Elsevier Inc. All rights reserved.

*Corresponding Author: Dept. of Biomedical Engineering Stony Brook University Bioengineering Bldg., Rm 215 Stony Brook, NY 11794-5281 Phone: 631-632-1481 Fax: 631-632-8577 Yi-Xian.Qin@stonybrook.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Disclosure: All authors have no conflicts of interest.

Introduction

Aging or functional disuse of bone can subsequently create a number of physiological or pathophysiological changes in the skeleton of the affected subjects (e.g. elderly, long-term bed-rest patients, and astronauts who participate in long-duration spaceflight missions), leading to conditions such as osteopenia [1]. Studies of mechanobiology and novel modalities of mechanical loading have demonstrated their abilities in regulating bone strength [2-10]. However, the underlying mechanotransductive mechanisms, namely, how mechanical signals are delivered to bone cells and how bone cells respond to such signals, remain unclear.

As a potent regulator in bone adaptation, bone fluid flow (BFF) with altered velocity or pressure acts as a communication media between an external load and the bone cells, which then regulate bone remodeling [11-16]. In converse, discontinuous BFF can initiate bone turnover and result in osteopenia [17-20]. Physical signals such as intramedullary fluid pressure (ImP) have been suggested to initiate BFF and to influence the osteogenic signals within bone [18]. A few studies using surgical methods on *in vivo* animal models have shown that BFF can be altered by ImP without bone deformation, and that ImP alone is sufficient to induce potent adaptive responses in bone. Qin et al applied a direct alteration of ImP to an isolated turkey ulna without deforming the bone tissue and found increased bone formation in response to the applied pressure [18]. Similarly, a more recent study introduced a novel microfluidic system for generating dynamic ImP and BFF within the femurs of alert mice to induce osteogenic responses [21].

Therefore, ImP-induced BFF provides a great potential in developing novel mechanical stimuli as countermeasures for disuse bone loss. Previous *in vivo* study using oscillatory electrical muscle stimulation (MS) in a hindlimb suspension (HLS) functional disuse rat model has demonstrated that oscillatory MS-induced muscle contraction can generate ImP and bone strain to mitigate disuse osteopenia [9, 22, 23]. A potential mechanism of the interrelationship between vasculature adaptation and applied ImP alteration has been suggested in a later study [24]. Induced ImP possibly triggers the transformation of the bone nutrient vasculature, leading to the ultimate alteration in blood supply to bone. However, to non-invasively isolate the ImP factor from the bone strain factor in an *in vivo* setting still remains challenging. Furthermore, in order to establish the translational potential of ImP, a novel and non-invasive method that directly couples an external load and internal BFF would be an attractive intervention. To reach this goal, our group has recently developed a novel, non-invasive dynamic hydraulic stimulation (DHS). Its promising effects on mitigating disuse bone loss have been shown via a 4-week rat HLS study followed by μ CT and histomorphometry analyses [25]. The results indicated a great potential of DHS to become a novel countermeasure for clinical osteoporosis/osteopenia treatments.

Identifying stimulation parameters within an optimal loading regimen, e.g. frequency, is important to maximize the effectiveness of the stimulation and to generate beneficial adaptive responses in bone. As an important determinant of bone adaptation to mechanical loading, loading frequency has been shown with a positive correlation to the improved bone quality in various studies [26-28]. Interestingly, effective loading frequencies seem to differ among various loading modalities [8, 26, 27, 29]. Whole-body vibrations are known to be more effective at high frequencies (>30Hz). On the other hand, an ulna axial loading study in mice reported a higher effect at lower loading frequencies (5Hz and 10Hz) as opposed to higher frequencies (20Hz or 30Hz) [29]. Related to ImP and BFF loading theory, relatively lower frequencies (5Hz and 10Hz) were reported to be more effective in bone adaptation in the femur [21] and tibia [8], respectively.

To test the hypothesis that non-invasive DHS can induce local ImP with minimal strain to potentially elicit osteogenic adaptive responses, the objective of this study was to evaluate the immediate effects on femoral and tibial ImP and bone strain induced by DHS over the tibia within a broad range of loading frequencies.

Materials and Methods

Animals

All surgical and experimental procedures were approved by the Institutional Animal Care and Use Committee (IACUC) at Stony Brook University. Surgical experiments were performed on three 15-month-old female Sprague Dawley virgin rats (Charles River, MA; body mass 426 ± 36 g) to measure the ImP and bone strain simultaneously during DHS loading.

ImP Measurements

Each animal was anesthetized through standard isoflurane inhalation. An approximately 1cm incision was made at the anterior knee region of the animal's right leg to expose the distal femur and proximal tibia. From the distal end of the femur and the proximal end of the tibia, a 1mm hole was carefully drilled into each of the right femoral and tibial marrow cavities, respectively. Guided by a 16-gauge catheter, a micro-cardiovascular pressure transducer (Millar Instruments, SPR-524, Houston, TX) was inserted into each of the femoral and tibial marrow cavities (Figure 1). The catheter and the pressure transducer apparatus were sealed tightly within the drilled holes.

Bone Strain Measurements

Similar to the above surgical procedure, a 2cm incision was made at the anterior side of the right tibia and the lateral side of the right femur. A linear single element strain gauge (120 Ω , factor 2.06, Kenkyojo Co., Tokyo) was firmly attached to the flat surface of each of the same tibia and femur within the mid-diaphyseal regions (Figure 1). The exposed muscles underwent minimal disruption, and the open skin was sutured before applying DHS.

DHS Loading

DHS was achieved through a costume-made inflatable cuff placed around the right tibia, with the similar setup in our recently published study [25]. Briefly, the inflation and deflation of the cuff was driven by a syringe pump with the loading magnitudes and frequencies delivered by a programmable waveform/signal generator (HP33120A, Hewlett-Packard, Palo Alto, CA). A pressure sensor was included to monitor the stimulation amplitude throughout the entire stimulation process. The pressure stimulation was achieved by 40mmHg static pressure + the peak-to-peak dynamic pressures given by the function generator that was set at 1.5V constant voltage. To start the experiment, DHS was applied to the operated tibia by placing the stimulation cuff around the mid-tibia region and loaded at frequencies of 0.5Hz, 1Hz, 1.5Hz, 2Hz, 2.5Hz, 3Hz, 3.5Hz, 4Hz, 5Hz, 6Hz, 7Hz, 8Hz, 9Hz, and 10Hz. Measurements of ImP and bone strain of the stimulated right tibia and unstimulated right femur were recorded simultaneously using a strain gauge amplifier (SCXI-1000, National Instruments, Austin, TX). For each animal, the entire frequency spectrum was repeated for at least three times.

Fast Fourier Transform (FFT) Analysis

Each run of data recording was put through a Fast Fourier Transform using MatLab. The random noise was removed by zeroing any values in each power spectrum that were below a threshold. This threshold was defined as the middle point between the lowest peak of the

signal and the average of the random noise. Once the noise was removed, each frequency step was analyzed individually. Each step was divided into twenty intervals. The difference between the maximum and minimum values was calculated and averaged over all intervals for that particular frequency step. This was taken as the peak to peak value for the according loading frequency (Figure 2).

Statistical Analysis

The values of ImP measurements were reported as mean $\pm$ SD. The effects of treatments were evaluated using a Kruskal-Wallis one-way ANOVA on ranks and Dunn's pairwise comparison post-hoc test using GraphPad Prism 3.0 Software (GraphPad Software Inc., La Jolla, CA). Power analysis was performed using Statistical Power Calculator of Researcher's Toolkit provided by DSS Research (Fort worth, TX and Arlington, VA) to calculate the statistical power of each ImP measurement value at each frequency point compared to the baseline value with 95% confidence interval.

Results

Tibial and Femoral ImP Induced by DHS

Approximately 1mmHg tibial ImP and 5mmHg femoral ImP were generated by normal heart beat within the marrow cavities. Oscillatory DHS over the tibial region of each rat's right hindlimb, loaded at various frequencies, generated additional fluid pressure in the tibial marrow cavity but not in the femoral marrow cavity. Tibial ImP generated from DHS at each frequency was plotted and shown in Figure 3. The observed responding trend of the ImP (peak-to-peak) values against frequency was induced in a nonlinear fashion during DHS loading. The induced ImP values (peak-to-peak) were in the order of 1.98 $\pm$ 1.57mmHg at 0.5Hz ($p>0.05$), 9.20 $\pm$ 4.58mmHg at 1Hz ($p>0.05$), 13.98 $\pm$ 3.23mmHg at 1.5Hz ($p<0.01$), 14.48 $\pm$ 3.10mmHg at 2Hz ($p<0.01$), 13.99 $\pm$ 2.58mmHg at 2.5Hz ($p<0.01$), 13.18 $\pm$ 2.07mmHg at 3Hz ($p<0.01$), 12.52 $\pm$ 2.08mmHg at 3.5Hz ($p<0.01$), 12.09 $\pm$ 2.18mmHg at 4Hz ($p<0.05$), 11.52 $\pm$ 2.48mmHg at 5Hz ($p<0.05$), 10.56 $\pm$ 2.67mmHg at 6Hz ($p>0.05$), 9.59 $\pm$ 3.21mmHg at 7Hz ($p>0.05$), 8.55 $\pm$ 2.97mmHg at 8Hz ($p>0.05$), 7.76 $\pm$ 3.70mmHg at 9Hz ($p>0.05$), and 7.28 $\pm$ 4.85mmHg at 10Hz ($p>0.05$). The ImP reached the peak at 2Hz.

Statistical Power Analysis

The power analysis showed sufficient statistical power of each ImP measurement value compared to the baseline value with 95% confidence interval in the following frequencies: 1.0, 1.5, 2.0, 2.5, 3.0, 3.5, 4.0, 5.0, 6.0, 7.0, 8.0, 9.0, and 10.0Hz. The strong statistical power indicates that our current sample size is sufficient to reject the null hypothesis when false. The effect of DHS on ImP induction can well be determined over the frequency range of 1Hz to 10Hz.

Effect of DHS on Tibial and Femoral Bone Strain

Oscillatory DHS over the tibial region of the rats' hindlimbs at various loading frequencies generated minimal bone strain within the tibia. Maximal bone strain measured at all loading frequencies was smaller than 8 μ . Similar to the femoral ImP measurements, no detectable induction of bone strain was observed in the un-stimulated femur.

Discussion

For the first time, this study demonstrated a non-invasive method that can be applied to an *in vivo* rodent model to isolate ImP and bone deformation, which are the two key determinants for BFF. The promising results from this study indicated that DHS can generate local fluid pressure in bone with simultaneously minimal bone strain. DHS over the rat tibia induced a

peak of tibial ImP at 2Hz. While the observed tibial bone strain over the range of loading frequencies was smaller than 8μ , relatively high ImP values were observed as a function of loading frequency. Tibial ImP measurements between the loaded 1.5Hz and 5Hz showed significant increases compared to the baseline level. These results suggested that DHS can potentially produce high fluid pressure gradients within the local marrow cavity. As a key determinant of BFF, loading generated fluid pressure in bone may have strong potentials in attenuating disuse bone loss, if loaded at proper frequencies.

The results from this study coincided with one of our recently published *in vivo* experiment, in which DHS loading at frequency of 2Hz was shown to have mitigation effect on disuse trabecular bone loss in a rat HLS functional disuse model [25]. For example, trabecular bone volume fraction in the stimulated tibia was increased by 83% when a DHS protocol at 2Hz was applied for four weeks in conjunction with HLS. The high surface area of the trabecular network allows it to expose to the rapid change in fluid pressure, leading to the pronounced effects of DHS observed in the trabecular bone. The DHS stimulation cuff was designed as dynamic circular compressions around the diaphysis region of the loaded bone, which does not provide direct physical contact over the metaphyseal region. However, the anabolic/anti-catabolic effects of DHS on disuse trabecular bone strongly support the ImP-induced BFF mechanism. According to the beneficial role of mechanotransduction in triggering bone remodeling [30, 31], strong evidence has suggested that interstitial fluid flow in bone can be altered by external muscular activities through various mechanisms [32, 33]. Based on the muscle pump theory, it is thought that muscle contraction compresses the blood vessels in muscle, which generates an arteriovenous pressure gradient that further increases the hydraulic pressure in skeletal nutrient vessels and amplifies the capillary filtration in bone [19, 20, 34]. Increased vessel pressure can directly increase the ImP that further drives BFF [35]. Direct circular compressions provided by DHS over the surrounding muscles have demonstrated its ability on ImP induction and potential in attenuating disused bone loss [25].

The observed ImP induction have indicated a nonlinear response in the DHS loading spectrum between 0.5Hz and 10Hz, in which peaked at 2Hz. The nonlinear relationship between DHS loading frequencies and the according ImP inductions is presented in Figure 3. The frequencies and ImP values are positively correlated, however, it is highly frequency dependent and they do not correlate into a linear relationship. This response to the loading frequency range is different from our previous observation using oscillatory muscle stimulation (MS), in which oscillatory MS induced maximal ImP at 20Hz. Oscillatory MS was achieved via two disposable needle-sized electrodes inserted into the quadriceps of the stimulated rats. The electrodes were then connected to the waveform generator with 2V supplies to induce muscle contraction [9, 22]. DHS provides an external stimulus that creates dynamic compressions on the surrounding muscles over the tibia, as opposed to direct muscle stimulation provided by oscillatory Muscle Stimulation. Based on the characteristics of the tissue material, e.g. the viscoelastic nature of the surrounding muscle tissue, the loads at high frequencies could be quickly damped. Due to the different physical orientations of how oscillatory MS and DHS contact the loaded tissue, as well as the different material densities and viscosities within hard and soft tissues, maximal DHS-induced ImP may result at relatively lower frequency compared to MS. This also suggests that direct hydraulic coupling may influence bone adaptation in a more physiological frequency range, where normal heart rate is 360 times per minute. Furthermore, the mechanotransductive sequence through different connective tissues during DHS may attenuate the high frequency response in bone, e.g. via the connective pathway from the contacted muscle vs. tendon to bone, resulting in peak ImP at a different frequency. To further reveal this complex mechanism, future investigation may need to focus on muscle kinetics and BFF generation.

Previous studies from our group have shown that BFF can be altered by ImP with minimal bone strain, and that ImP alone is sufficient to induce bone adaptation. A loss of cortical bone by 5.7% was resulted in a disuse avian ulna [18]. An increase in cortical bone mass by 18% was shown with a direct fluid loading into the ulna at 20Hz for 4 weeks. Moreover, the study also observed a strong correlation between the transcortical fluid pressure gradient and total bone formation. Similarly, generation of dynamic ImP and BFF within the femurs of HLS mice using a novel microfluidic system significantly reduced the disuse bone mass loss in both trabecular bone and cortical bone [21]. Our present study demonstrated DHS's ability in ImP induction, which strongly suggests its potential in attenuating disuse bone loss. This hypothesis is further supported by the mitigation effect of DHS at 2Hz, indicated in our recent publication [25]. The response of bone cells to dynamic fluid flow mechanical stimulation was further demonstrated by a longitudinal *in vivo* study which indicated that the MSC population was positively influenced by the DHS-derived mechanical signals. Changes of MSC number in response to DHS bias their differentiation towards osteoblastogenesis, leading to bone formation even under disuse conditions [36]. Further study on the underlying mechanism in response to mechanical signals, e.g. mechanobiological modulation of cytoskeleton and calcium influx into osteoblastic cells [37], can provide insights for the future developments of novel and effective osteoporosis treatments. Interesting, DHS on the tibia only affected the tibial ImP but not the femoral ImP, indicating the local effect of DHS. This is suggesting that the function of DHS may be site specific. More detailed *in vivo* experiments are needed to validate this observation. Moreover, while increased bone fluid pressure and bone fluid flow regulation are strongly correlated, our current results of DHS-driven ImP induction suggest a possible mechanism that the induced ImP may subsequently enhance bone fluid flow. Future studies may extend our current experimental focus and incorporate further experiments of tracer transport determinations to verify this hypothesized mechanism [31, 38].

In summary, oscillatory DHS can generate local ImP as a function of stimulation frequency with minimal strain, where the induced dynamic ImP may subsequently enhance BFF. DHS, if applied at an optimal frequency, has strong potential in preventing and attenuating bone loss under disuse osteopenia condition. Results from this study provide evidence that may facilitate the development of a biomechanical based intervention for osteoporosis prevention and treatment, which provide great insights for future clinical application.

Acknowledgments

This research is kindly supported by the National Institute of Health (R01 AR52379 and AR61821, YXQ), the US Army Medical Research and Materiel Command, and the National Space Biomedical Research Institute through NASA Cooperative Agreement NCC 9-58 (YXQ). The authors are grateful to Drs. Clinton Rubin and Stefan Judex for the stimulating discussions.

References

- [1]. Lau RY, Guo X. A review on current osteoporosis research: with special focus on disuse bone loss. *J Osteoporos*. 2011; 2011:293808. [PubMed: 21876833]
- [2]. Pedersen EA, Akhter MP, Cullen DM, Kimmel DB, Recker RR. Bone response to *in vivo* mechanical loading in C3H/HeJ mice. *Calcif Tissue Int*. 1999; 65:41–6. [PubMed: 10369732]
- [3]. Kameyama Y, Hagino H, Okano T, Enokida M, Fukata S, Teshima R. Bone response to mechanical loading in adult rats with collagen-induced arthritis. *Bone*. 2004; 35:948–56. [PubMed: 15454102]
- [4]. Hsieh YF, Robling AG, Ambrosius WT, Burr DB, Turner CH. Mechanical loading of diaphyseal bone *in vivo*: the strain threshold for an osteogenic response varies with location. *J Bone Miner Res*. 2001; 16:2291–7. [PubMed: 11760844]

- [5]. Xie L, Rubin C, Judex S. Enhancement of the adolescent murine musculoskeletal system using low-level mechanical vibrations. *J Appl Physiol*. 2008; 104:1056–62. [PubMed: 18258802]
- [6]. Tanaka SM, Sun HB, Yokota H. Bone formation induced by a novel form of mechanical loading on joint tissue. *Biol Sci Space*. 2004; 18:41–4. [PubMed: 15308820]
- [7]. Yokota H, Tanaka SM. Osteogenic potentials with joint-loading modality. *J Bone Miner Metab*. 2005; 23:302–8. [PubMed: 15981026]
- [8]. Zhang P, Tanaka SM, Sun Q, Turner CH, Yokota H. Frequency-dependent enhancement of bone formation in murine tibiae and femora with knee loading. *J Bone Miner Metab*. 2007; 25:383–91. [PubMed: 17968490]
- [9]. Lam H, Qin YX. The effects of frequency-dependent dynamic muscle stimulation on inhibition of trabecular bone loss in a disuse model. *Bone*. 2008; 43:1093–100. [PubMed: 18757047]
- [10]. Ozcivici E, Luu YK, Adler B, Qin YX, Rubin J, Judex S, Rubin CT. Mechanical signals as anabolic agents in bone. *Nat Rev Rheumatol*. 2010; 6:50–9. [PubMed: 20046206]
- [11]. Pollack SR, Korostoff E, Sternberg ME, Koh J. Stress-generated potentials in bone: effects of collagen modifications. *J Biomed Mater Res*. 1977; 11:677–70. [PubMed: 561074]
- [12]. Montgomery RJ, Sutker BD, Bronk JT, Smith SR, Kelly PJ. Interstitial fluid flow in cortical bone. *Microvasc Res*. 1988; 35:295–307. [PubMed: 3393091]
- [13]. Reich KM, Gay CV, Frangos JA. Fluid shear stress as a mediator of osteoblast cyclic adenosine monophosphate production. *J Cell Physiol*. 1990; 143:100–4. [PubMed: 2156870]
- [14]. Rubin J, Biskobing D, Fan X, Rubin C, McLeod K, Taylor WR. Pressure regulates osteoclast formation and MCSF expression in marrow culture. *J Cell Physiol*. 1997; 170:81–7. [PubMed: 9012787]
- [15]. Jacobs CR, Yellowley CE, Davis BR, Zhou Z, Cimbala JM, Donahue HJ. Differential effect of steady versus oscillating flow on bone cells. *J Biomech*. 1998; 31:969–76. [PubMed: 9880053]
- [16]. Burger EH, Klein-Nulend J. Mechanotransduction in bone--role of the lacuno-canalicular network. *FASEB J*. 1999; 13(Suppl):S101–12. [PubMed: 10352151]
- [17]. Rubin CT, Sommerfeldt DW, Judex S, Qin Y. Inhibition of osteopenia by low magnitude, high-frequency mechanical stimuli. *Drug Discov Today*. 2001; 6:848–858. [PubMed: 11495758]
- [18]. Qin YX, Kaplan T, Saldanha A, Rubin C. Fluid pressure gradients, arising from oscillations in intramedullary pressure, is correlated with the formation of bone and inhibition of intracortical porosity. *J Biomech*. 2003; 36:1427–37. [PubMed: 14499292]
- [19]. Winet H. A bone fluid flow hypothesis for muscle pump-driven capillary filtration: II. Proposed role for exercise in erodible scaffold implant incorporation. *Eur Cell Mater*. 2003; 6:1–10. discussion 10-1. [PubMed: 14562269]
- [20]. Otter MW, Qin YX, Rubin CT, McLeod KJ. Does bone perfusion/reperfusion initiate bone remodeling and the stress fracture syndrome? *Med Hypotheses*. 1999; 53:363–8. [PubMed: 10616033]
- [21]. Kwon RY, Meays DR, Tang WJ, Frangos JA. Microfluidic enhancement of intramedullary pressure increases interstitial fluid flow and inhibits bone loss in hindlimb suspended mice. *J Bone Miner Res*. 2010; 25:1798–807. [PubMed: 20200992]
- [22]. Qin YX, Lam H. Intramedullary pressure and matrix strain induced by oscillatory skeletal muscle stimulation and its potential in adaptation. *J Biomech*. 2009; 42:140–5. [PubMed: 19081096]
- [23]. Lam H, Hu M, Qin YX. Alteration of contraction-to-rest ratio to optimize trabecular bone adaptation induced by dynamic muscle stimulation. *Bone*. 2011; 48:399–405. [PubMed: 20850577]
- [24]. Lam H, Brink P, Qin YX. Skeletal nutrient vascular adaptation induced by external oscillatory intramedullary fluid pressure intervention. *J Orthop Surg Res*. 2010; 5:18. [PubMed: 20222973]
- [25]. Hu M, Cheng J, Qin YX. Dynamic hydraulic flow stimulation on mitigation of trabecular bone loss in a rat functional disuse model. *Bone*. 2012; 51:819–25. [PubMed: 22820398]
- [26]. Turner CH, Owan I, Takano Y. Mechanotransduction in bone: role of strain rate. *Am J Physiol*. 1995; 269:E438–42. [PubMed: 7573420]
- [27]. Hsieh YF, Turner CH. Effects of loading frequency on mechanically induced bone formation. *J Bone Miner Res*. 2001; 16:918–24. [PubMed: 11341337]

- [28]. Rubin CT, McLeod KJ. Promotion of bony ingrowth by frequency-specific, low-amplitude mechanical strain. *Clin Orthop Relat Res*. 1994;165–74. [PubMed: 8118971]
- [29]. Warden SJ, Turner CH. Mechanotransduction in the cortical bone is most efficient at loading frequencies of 5-10 Hz. *Bone*. 2004; 34:261–70. [PubMed: 14962804]
- [30]. Wang L, Fritton SP, Weinbaum S, Cowin SC. On bone adaptation due to venous stasis. *J Biomech*. 2003; 36:1439–51. [PubMed: 14499293]
- [31]. Wang L, Ciani C, Doty SB, Fritton SP. Delineating bone's interstitial fluid pathway in vivo. *Bone*. 2004; 34:499–509. [PubMed: 15003797]
- [32]. Stevens HY, Meays DR, Frangos JA. Pressure gradients and transport in the murine femur upon hindlimb suspension. *Bone*. 2006; 39:565–72. [PubMed: 16677866]
- [33]. Valic Z, Buckwalter JB, Clifford PS. Muscle blood flow response to contraction: influence of venous pressure. *J Appl Physiol*. 2005; 98:72–6. [PubMed: 15377645]
- [34]. Laughlin MH. The muscle pump: what question do we want to answer? *J Appl Physiol*. 2005; 99:774. [PubMed: 16020444]
- [35]. Qin YX, Lam H, Ferreri S, Rubin C. Dynamic skeletal muscle stimulation and its potential in bone adaptation. *J Musculoskelet Neuronal Interact*. 2010; 10:12–24. [PubMed: 20190376]
- [36]. Minyi Hu RY, Michelle Lien, Morgan Teeratananon, Kunal Agarwal, Yi-Xian Qin. Dynamic Fluid Flow Mechanical Stimulation Modulates Bone Marrow Mesenchymal Stem Cells. *Bone Research*. 2013:98–104.
- [37]. Zhang S, Cheng J, Qin YX. Mechanobiological modulation of cytoskeleton and calcium influx in osteoblastic cells by short-term focused acoustic radiation force. *PLoS One*. 2012; 7:e38343. [PubMed: 22701628]
- [38]. Knothe Tate ML, Steck R, Forwood MR, Niederer P. In vivo demonstration of load-induced fluid flow in the rat tibia and its potential implications for processes associated with functional adaptation. *J Exp Biol*. 2000; 203:2737–45. [PubMed: 10952874]

Highlights

- Dynamic hydraulic stimulation (DHS) induced local intramedullary pressure (ImP) with minimal bone strain.
- Intramedullary pressure can be raised externally through skeletal muscle coupling.
- ImP induction in response to DHS was in a non-linear fashion over the loading frequencies, where peaked at 2Hz.
- The results from this study strongly suggest DHS's potential as an effective, non-invasive intervention for osteopenia and osteoporosis treatments.

Figure 1.

Surgical experiment setup. For the ImP measurement, a micro-cardiovascular pressure transducer was inserted into the marrow cavity of both tibia and femur. For bone strain measurement, a single element strain gauge was firmly attached to the flat surface of the same tibia and femur at the mid-diaphyseal region. Simultaneous ImP and bone strain were then measured in both tibia and femur under DHS on the tibia.

Figure 2.
Representative traces of ImP measurements from DHS-loading at various frequencies over time.

Figure 3.

Graph shows mean $\pm$ SD values of the ImP measurements. ImP in the tibia increased significantly with DHS loading frequency at 1.5Hz, 2Hz, 2.5Hz, 3Hz, 3.5Hz, 4Hz, and 5Hz. In the loading frequency spectrum from 0.5Hz to 10Hz, a maximum ImP of 14.48 ± 3.10 mmHg was observed at 2Hz, which was around 7 folds higher than 0.5Hz (1.98 ± 1.57 mmHg). ^a $p<0.01$ vs. baseline ImP; ^b $p<0.05$ vs. baseline ImP.