Published in final edited form as:

Cell Mol Life Sci. 2011 March ; 68(5): . doi:10.1007/s00018-010-0500-x.

Cigarette Smoking Reprograms Apical Junctional Complex Molecular Architecture in the Human Airway Epithelium *In Vivo*

Renat Shaykhiev¹, Fouad Otaki², Prince Bonsu¹, David T. Dang¹, Matthew Teater¹, Yael Strulovici-Barel¹, Jacqueline Salit¹, Ben-Gary Harvey^{1,2}, and Ronald G. Crystal^{1,2}

¹Department of Genetic Medicine, Department of Medicine, Weill Cornell Medical College, New York, New York

²Division of Pulmonary and Critical Care Medicine, Department of Medicine, Weill Cornell Medical College, New York, New York

Summary

The apical junctional complex (AJC), composed of tight and adherens junctions, maintains epithelial barrier function. Since cigarette smoking and chronic obstructive pulmonary disease (COPD), the major smoking-induced disease, are associated with increased lung epithelial permeability, we hypothesized that smoking alters the transcriptional program regulating airway epithelial AJC integrity. Transcriptome analysis revealed global down-regulation of physiological AJC gene expression in the airway epithelium of healthy smokers (n=59) compared to nonsmokers (n=53) in association with changes in canonical epithelial differentiation pathways such as PTEN signaling accompanied by induction of cancer-related AJC components. The overall expression of AJC-related genes was further decreased in COPD smokers (n=23). Exposure of airway epithelial cells to cigarette smoke extract *in vitro* resulted in down-regulation of several AJC genes paralleled by decreased transepithelial resistance. Thus, cigarette smoking induces transcriptional reprogramming of airway epithelial AJC architecture from its physiological pattern necessary for barrier function toward disease-associated molecular phenotype.

Keywords

tight junctions; adherens junctions; airway epithelium; epithelial polarity; cigarette smoking; transcriptional regulation; chronic obstructive pulmonary disease

Introduction

The apical junctional complex (AJC), a distinct multicomponent structure localized to the apex of the lateral membrane compartment of polarized epithelial cells, includes 2 kinds of intercellular junctions, tight junctions (TJ) and adherens junctions (AJ), a unique array of transmembrane proteins connected to an extensive network of cytoplasmic scaffold and signaling proteins [1,2]. The TJ, the most apical AJC structure, constitute a semipermeable barrier for solutes and ions and physically separate the apical and basolateral membranes [3,4]. Just below the TJ are the AJ which mediate cell-cell adhesion [5]. Detailed studies have defined the molecular composition of AJC [2], the posttranslational events governing AJC assembly in polarized epithelia [6–8], and the transcriptional changes associated with the AJC maturation during epithelial differentiation *in vitro* [9]. However, little is known about the global gene expression program encoding AJC components and AJC-associated

molecular pathways and transcriptional networks (the “AJC gene expression architecture”) in human epithelial tissues *in vivo* in health and under conditions characterized by an altered epithelial barrier function.

The airway epithelium constitutes an essential tissue barrier protecting the lung from inhaled environmental challenges [10,11]. Of these, cigarette smoke is a major risk factor for chronic obstructive pulmonary disease (COPD) and lung cancer [12–14]. Smoking can substantially compromise lung epithelial barrier function leading to increased epithelial permeability [15,16]. These observations have been commonly interpreted as a result of a smoking-induced damage to the junctional structure [17,18] due to direct cytotoxic effect of cigarette smoke on lung epithelial cells [19,20]. In the present study, we hypothesized that cigarette smoking might also have a more targeted effect on the airway epithelial barrier by modifying the AJC gene expression architecture in the small airway epithelium (SAE), the primary site of cigarette smoking-associated changes in the lung [21], and that this altered AJC-related gene expression contributes to a COPD-relevant molecular phenotype.

Results

Physiological AJC Gene Expression Architecture in the Human SAE

Microarray analysis revealed a distinct pattern of AJC gene expression in the SAE of healthy nonsmokers: 31 of 63 (49%) genes encoding known AJC components were expressed in all individuals, with considerable variability in expression levels of individual AJC genes (Figure 1). Among the constitutively expressed genes encoding transmembrane TJ proteins, CXADR (coxsackie virus and adenovirus receptor) was most abundant followed by the claudin genes CLDN -7, -3, -8, -1, and F11R (JAM-1; Figure 1A). Of the 12 AJC genes encoding cytoplasmic TJ components, 11 (92%) including TJP1, TJP2, and TJP3 [zona occludens (ZO) -1, -2, and -3, respectively], CSDA (ZONAB) and others were constitutively expressed (Figure 1B). Although expression levels of the epithelial polarity complexes genes were relatively low, genes encoding the major epithelial polarity complex PAR3/PAR6/aPKC and the core elements of the Crumbs (CRB3/INADL/MPP5) polarity complex were detected in the SAE of all healthy nonsmokers, whereas the Scribble complex (SCRIB/LLGL1/DLG) was detected in only a few subjects (Figure 1C). Among the AJ genes, CDH1 (E-cadherin), displayed the highest relative expression levels followed by MLLT4 (afadin; AF-6), CTNNB1 and other catenin family genes (Figure 1D). Consistent with the normal epithelial phenotype, CDH2 (N-cadherin), abundantly expressed in mesenchymal and neuronal tissues [5], was barely detected in the SAE of healthy nonsmokers (Figure 1D). Four genes implicated in regulation of apical epithelial polarity, e.g. NUMB, PTEN, FOXA1 and FOXA2 [22–25] were constitutively expressed at low levels (Figure 1E). Genes coding for AJC elements such as claudins 2, -5, -6, -11, -14, -17, -20, and nectin-1 (PVRL1) as well as putative AJC-regulating genes AMOT, HNF4A were not detected (Figure 1).

Smoking-dependent Derangement of the AJC Gene Expression Architecture in the Healthy SAE

Genome-wide analysis revealed that, whereas 19% (6,038 of 32,436) of all genes detected in the SAE were affected significantly by smoking (Figure 2A, left panel), the impact of smoking on the AJC transcriptional program was more profound, with 71% (39 of 55) of the detected AJC genes differentially expressed in healthy smokers *vs* nonsmokers (Figure 2A, right panel). Remarkably whereas at the genome-wide level, smoking was associated with up-regulation of a considerable number of genes (Figure 2A, left panel) that were mostly related to oxidative stress and xenobiotic metabolism (Figure 2B), there was almost a uniform suppression of the physiological AJC gene expression program, with 37 of 39

(95%) smoking-responsive AJC genes significantly down-regulated in healthy smokers compared to nonsmokers (Figure 2A, right panel; Figure 2B).

All AJC categories were broadly affected in healthy smokers, including 8 of 25 (32%) genes encoding transmembrane TJ proteins, 11 of 12 (92%) genes encoding cytoplasmic TJ, 6 of 12 (50%) polarity complex genes including all 3 members of the PAR3/PAR6/aPKC polarity complex, and 9 of 14 (64%) AJ genes, suggesting a net effect of smoking on the physiological SAE AJC program (Figure 2C; Table I). From the functional perspective, smoking was associated with down-regulation of the entire molecular AJC machinery, including the proteins that physically constitute cell-cell junctions, receptors for extracellular ligands, adaptor proteins, kinases, phosphatases and transcription factors (Figure 2C). In contrast to the physiological AJC genes, expression of the TJ gene CLDN10 and the AJ gene CDH2, both associated with epithelial carcinogenesis [26–28], was significantly up-regulated in the SAE of healthy smokers (Table I).

Differential expression of several AJC genes in healthy smokers *vs* nonsmokers was confirmed by TaqMan real-time PCR (Supplemental Figure 1). The expression of CLDN1 and CDH1, the two major components of TJ and AJ, respectively, was markedly decreased in the smokers' SAE cells at the protein level (Figure 2D; Supplemental Figure 2). Together, these data suggest that the physiological AJC molecular architecture in the human SAE is broadly deranged by cigarette smoking *in vivo*.

Dysregulation of the SAE AJC Transcriptional Program in COPD Smokers

Microarray analysis revealed that 18 of 37 (49%) AJC-encoding genes suppressed in the SAE of healthy smokers were further down-regulated in smokers with COPD (Table I). Also the putative AJC regulating genes PTEN, FOXA1 and FOXA2 demonstrated progressive smoking-dependent down-regulation in COPD smokers (Table I). By contrast, CLDN10 and CDH2, the 2 disease-associated AJC genes induced in the SAE of healthy smokers, were further up-regulated in COPD smokers (Table I) along with CLDN7, a TJ gene, implicated in epithelial carcinogenesis [29,30]. (See Supplemental Figure 4 for examples of AJC genes dysregulated in the SAE of COPD smokers.)

Based on the expression of all AJC genes detected in the SAE, unsupervised clustering analysis effectively segregated nonsmokers from both healthy and COPD smokers, with the COPD smokers forming a group at the pole opposite to that of nonsmokers in the clustering map (Figure 3A). Healthy smokers were distributed throughout the map with subsets having either nonsmoker-like or COPD-like molecular patterns (Figure 3A). Principal component analysis (PCA) revealed that the groups could be separated from each other more effectively based on the SAE expression of AJC genes (Figure 3B, right panel) compared to separation based on the expression of all genes in the SAE transcriptome (Figure 3B, left panel). Remarkably, the first PC increased from 12.2% in the transcriptome-wide analysis to 28.5% in the AJC gene-restricted analysis (Figure 3B), indicating that variability in AJC gene expression had a substantial contribution to genome-wide differences between the study groups. Similar to the cluster analysis, PCA revealed a progressive shift in AJC gene expression from healthy nonsmokers to smokers with COPD, whereas such trend was not prominent at the genome-wide scale (Figure 3B). This trend was not due to differences in gene number between the genome-wide and the 54-gene AJC-restricted datasets since PCA on randomly selected datasets with the same number of genes as the AJC-restricted dataset did not reveal group-specific variability between the subjects except for the dataset composed of 54 randomly selected oxidative stress-related genes (Supplemental Figure 3).

SAE AJC Index as a Measure of Smoking-Dependent Dysregulation of the SAE AJC Transcriptional Program

The SAE AJC gene expression index (I_{AJC}) was developed as a global measure of the coordinate smoking-dependent changes in the physiological SAE AJC gene expression by integrating information on expression levels of all smoking-responsive genes that encode AJC components. Consistent with the trends revealed by the clustering analysis and PCA, the average I_{AJC} progressively increased in a smoking-dependent manner from healthy smokers to COPD smokers ($p < 0.04$, Figure 3C). This analysis also revealed the contribution of individual AJC genes to the progressive smoking-dependent abnormality in the global SAE AJC gene expression architecture (Figure 3D). Notably, the I_{AJC} -based analysis revealed a higher magnitude of intergroup variability, suggesting that it can capture more information related to differential expression of AJC genes compared to traditional fold-change-based analysis.

Molecular Pathways Associated with Smoking-induced Changes in the SAE AJC Gene Expression Architecture

To determine whether down-regulation of the physiological AJC transcriptional program in the SAE of smokers is associated with smoking-induced changes in the canonical molecular pathways, we first performed a genome-wide screening for smoking-responsive genes negatively correlating ($r < -0.5$) with the I_{AJC} , i.e., having the expression pattern similar to AJC genes in the SAE of healthy smokers. The derived gene dataset (“AJC-associated genes”) included 992 probe sets corresponding to 825 unique annotated genes (Supplemental Table 3), indicating that suppression of the AJC transcriptional program has transcriptional coherence with at least 15% of other smoking-responsive genes. Among the top AJC-associated genes was peroxiredoxin 5 ($r = -0.66$; $Rho = -0.53$, $p < 0.0001$), an antioxidant gene proposed to protect the airway epithelium from smoking-induced injury [19]. Ingenuity pathway analysis revealed that the AJC-associated gene dataset (796 mapped genes) is enriched in elements of canonical molecular pathways implicated in regulation of epithelial differentiation, with the PTEN pathway showing the strongest association ($p < 10^{-4}$; Figure 4A). Other enriched pathways included protein kinase A signaling, insulin receptor signaling, phosphoinositide 3-kinase (PI3K)/AKT signaling (all $p < 10^{-3}$), Rac- and CDC42 signaling pathways (both $p < 0.005$, Figure 4A). Expression of the founding member of the PTEN pathway, the tumor suppressor PTEN, significantly correlated with overall AJC gene expression (Figure 4B). To construct a model of smoking-sensitive molecular interactions between individual AJC-encoding genes and associated molecular pathways, the set of AJC-encoding genes, PTEN pathway member genes and putative AJC-regulating genes correlating with the I_{AJC} ($r = -0.5$; $Rho = -0.5$; $p < 0.0001$) were subjected to STRING network analysis. STRING is a web-based analytic tool that accumulates information on functional protein–protein associations based on the knowledge of physical interactions between individual proteins including their interaction in curated biological pathways complemented by interactions that are predicted computationally using a set of prediction algorithms [31]. This analysis revealed that, based on known and predicted protein-protein interactions, a number of smoking-sensitive AJC-encoding genes encode proteins that interact with each other and are directly or indirectly linked to PTEN (Figure 4C), implying that AJC-associated protein-protein interactions are reprogrammed by smoking at the transcriptional level.

SAE Smoking-sensitive AJC Transcriptional Network

A uniform down-regulation of the physiological AJC gene expression pattern in the SAE of smokers suggests that transcriptional mechanisms determining the expression of individual AJC genes in the SAE are suppressed by smoking in a coordinate manner. This assumption was supported by the GO analysis, which indicated that biological functions “Nucleic acid

metabolism” (38% of dataset genes; $p < 10^{-7}$) and “Transcription” (27% of dataset genes; $p < 10^{-5}$) dominated among the functional GO categories significantly enriched in the AJC-associated gene dataset (Figure 5A). Based on the concept that high gene expression correlation may provide unbiased information on possible gene regulatory networks independent of prior knowledge about the physical interaction between the gene products [32,33], and since I_{AJC} provides a cumulative measure of AJC gene expression, we performed a genome-wide co-expression analysis that identified a subset of 21 smoking-responsive genes (“core AJC network genes”; Figure 5B), whose expression paralleled changes in the global AJC transcriptional program. This was evident by a strong negative correlation of their expression with the I_{AJC} ($Rho = -0.7$; $p < 0.0001$) and a strong positive correlation with at least 2 AJC-encoding genes ($Rho > 0.7$; $p < 0.0001$) in the SAE of healthy smokers (Supplemental Table 4). Of note, 15 of 21 (71%) core AJC network genes encoded nuclear proteins, consistent with a significant enrichment of categories “RNA binding” ($p < 10^{-5}$), “Nucleus” ($p < 10^{-5}$), and “Transcription” ($p < 0.001$) in the 21-gene core AJC network dataset (Supplemental Table 5). The core AJC network included genes encoding well-characterized transcriptional regulators such as CRSP3, MATR3, NUCKS, and PSIP1, as well as genes related to cytokine signaling and tissue morphogenesis, such as IL6ST and PKD2, respectively (Figure 5B; Supplemental Table 4).

Overall AJC network gene expression in the SAE of healthy smokers strongly correlated with expression of PTEN ($Rho = 0.77$, $p < 0.0001$; Figure 5C), indicating a link between the AJC transcriptional network and AJC-associated signaling pathways sensitive to smoking. The overall expression of the core AJC transcriptional network genes was further decreased in smokers with COPD ($p < 0.0003$ vs healthy smokers; Figure 5D), in agreement with progressive smoking-dependent suppression of the SAE AJC gene expression program in COPD (Figure 3C).

A subset of AJC-encoding genes, e.g., TJP1, MPP5, CDH1, MPP7, DLG1, and MPDZ, exhibited a particularly high degree of co-expression with the core AJC network elements (Figure 5B). Among these “intrinsic” AJC network genes, TJP1 and MPP5 exhibited the highest degree of interaction intensity, being strongly co-expressed ($Rho > 0.7$; $p < 0.0001$) with 18 (86%) and 16 (76%) of core AJC network genes, respectively (Figure 5B; Supplemental Table 4). Based on clustering analysis, individual AJC-encoding genes within the network displayed preferential interaction with specific core network elements (Figure 5B), suggesting the heterogeneity of regulatory pathways within the network.

Cigarette Smoke Modulates Airway Epithelial AJC *In Vitro*

To assess whether smoking can induce changes in the AJC transcriptional program and physiological properties independent of inflammatory cells or direct cytotoxicity effect on the airway epithelial barrier, we exposed the apical surface of differentiated 16HBE14o! human airway epithelial cells to cigarette smoke extract (CSE) daily at non-toxic concentrations (Figure 6A). After 6 days of exposure, expression of genes encoding AJC components CLDN1, CLDN8, CGN, and CDH1 was significantly down-regulated (Figure 6B). All these genes were also among the AJC genes significantly down-regulated in the SAE of smokers *in vivo* (Table I; Supplemental Figure 4). Down-regulation of these AJC genes was accompanied by CSE-induced suppression of genes encoding PTEN and transcription factor FOXO3A (Figure 6B), a downstream member of the PTEN pathway [34,35]. By contrast, expression of 2 smoking-inducible oxidative stress-related genes, CYP1B1 and NQO1 [36], was up-regulated (Figure 6B), indicating that the observed effect of CSE on AJC genes was not due to the general CSE-induced transcriptional suppression. This down-regulation of physiological AJC gene expression by non-cytotoxic CSE exposure *in vitro* was accompanied by a significant decrease of transepithelial electrical resistance (Figure 6C), a measure of the epithelial junctional barrier integrity.

Discussion

Apical-basal polarity, a hallmark of differentiated epithelia, is manifested by asymmetric distribution of specialized protein domains along the plasma membrane [7]. Localized to the apical membrane compartments of neighboring epithelial cells, the AJC is essential for epithelial barrier integrity and function, with the TJ governing the transport of solutes and ions across epithelia, and the AJ mediating cell-cell adhesion [1–4]. AJC protein complexes also function as signaling platforms that regulate gene expression, cell proliferation and differentiation [37,38] and provide a tumor-suppressing mechanism [39,40]. All of these functions are altered in the airway epithelium of smokers, which is leakier, hyperproliferative, abnormally differentiated and more susceptible to malignant transformation [16,21,41,42] compared to nonsmokers. The present study suggests that one of the mechanisms central to these airway epithelial barrier abnormalities in smokers is the dysregulation of the physiological AJC transcriptional program in the airway epithelium, an alteration that progresses during the development of smoking-induced lung disease.

Physiological SAE AJC Gene Expression Architecture

AJC barrier integrity is determined by its unique molecular composition, which varies from one anatomic location to another [4]. Under physiological conditions, the SAE appears to be leakier than the epithelium of proximal airways, indicative of a distinct SAE AJC molecular program that determines a balance between physiological permeability necessary for constitutive metabolic processes, and function as a barrier limiting diffusion of solutes, ions, and preventing penetration of inhaled microbes [1,43,44]. Although expression of several AJC proteins in the airway epithelium has been analyzed in detail previously [44], the entire AJC molecular composition of SAE *in vivo* under physiologic conditions has not been previously assessed. Microarray analysis of the present study revealed that 49% of genes encoding known AJC components are expressed in the SAE of healthy nonsmokers, with considerable variability in expression levels of individual AJC genes.

Transcriptional Suppression of Physiological AJC Program by Smoking

Cigarette smoke increases airway epithelial permeability *in vivo* and *in vitro* [15–19,45,46]. Since smoking is associated with dramatic gene expression changes in the airway epithelium [36], we hypothesized that chronic exposure to cigarette smoke modulates the transcriptional program necessary for the maintenance of AJC in the SAE. The data revealed that 67% of the AJC-encoding genes physiologically expressed in the SAE are significantly down-regulated in healthy smokers. All categories of AJC-encoding genes were broadly affected, with the genes encoding cytoplasmic TJ components being almost uniformly (92%) down-regulated. This was accompanied by down-regulation of putative regulators of epithelial polarity (e.g., PTEN, FOXA1, and FOXA2), suggesting some of them might represent smoking-sensitive regulators of AJC genes in the airway epithelium. Impressively, smoking-induced differences in AJC gene expression had a higher impact on the variability between smokers and nonsmokers than the global SAE transcriptome changes induced by smoking.

The data of the present study suggest a model in which the modulation of airway epithelial barrier by chronic cigarette smoking represents a relatively specific molecular process involving the transcriptional reprogramming of the AJC molecular architecture. Supporting this concept, we observed that 6-day exposure of differentiated human airway epithelium to non-cytotoxic concentrations of cigarette smoke extract *in vitro* resulted in down-regulation of several AJC genes, and was accompanied by the reduction of transepithelial resistance. These *in vitro* data indicate that the effect of cigarette smoke on AJC molecular integrity in the SAE does not require the presence of inflammatory cells and their mediators previously reported to down-regulate expression of AJC genes and increase airway epithelial

permeability [21,47,48]. Down-regulation of physiological AJC genes by smoking *in vivo* and *in vitro* was paralleled by decreased expression of PTEN and FOXO3A, a transcription factor in the PTEN pathway [34,35], as well as an up-regulation of oxidative stress-related genes, with a strong association between the I_{AJC} and PRDX5, an antioxidant gene proposed to limit smoking-increased airway epithelial permeability [19]. Although oxidative stress is known to induce disassembly of AJC at the protein level [49], our data suggests the possibility of a reciprocal relationships between transcriptional programs regulating AJC integrity and oxidative stress responses to smoking.

Cancer-like AJC Gene Expression Pattern in Smokers' SAE

The broad suppression of the physiological SAE AJC transcriptional program in healthy smokers was accompanied by an up-regulation of N-cadherin and CLDN10, AJC genes linked to cancer-related pathological changes in epithelial tissues. Epithelial expression of N-cadherin, a marker of neuronal and mesenchymal cells, is a common indicator of epithelial-mesenchymal transition (EMT) [5,50]. Other molecular changes typical for EMT, such as down-regulation of E-cadherin, occludin, TJP1 and CXADR [51], were also detected in the smokers' SAE, suggesting that dysregulation of SAE AJC gene expression in smokers might result from common transcriptional derangement associated with smoking-induced EMT. Consistent with this concept, nicotine, the major cigarette smoke constituent, induces EMT-like molecular and morphological changes in human lung cancer cell lines [52]. Up-regulation of N-cadherin and down-regulation of E-cadherin are the central EMT features in human lung cancer [28]. Our data provide evidence that smoking induces EMT-typical molecular changes in the healthy airway epithelium *in vivo* prior to clinical manifestations of lung cancer.

Another smoking-induced AJC gene was CLDN10, whose expression has been associated with increased epithelial permeability [53]. Recently, CLDN10 has been described as a marker of Clara cells in the mouse airways [54]. Given that the numbers of Clara cells are considerably reduced in the SAE of smokers [55], it is possible that smoking induces *de novo* expression of CLDN10 in other cell populations similar to various epithelial cancers, in which CLDN10 is overexpressed [26,27].

Smoking-sensitive SAE AJC-associated Molecular Pathways and Networks

Broad suppression of biologically related groups of genes suggests that central mechanisms controlling their expression, e.g. signaling pathways or transcriptional regulators, are dysregulated in a coordinated manner [32,33,56]. No central transcriptional mechanism regulating AJC biogenesis in the adult epithelia has been defined. The present study reveals that suppression of the physiological AJC gene expression program in the SAE of smokers is paralleled by down-regulation of 15% of other smoking-responsive genes and that this AJC-associated dataset is significantly enriched with the elements of canonical molecular pathways critical for epithelial differentiation. The top enriched pathways (PTEN, PI3K/AKT, Rac- and CDC42) have been implicated in posttranslational regulation of apical epithelial polarity and AJC assembly [6,8], but not yet linked to AJC transcriptional control. Among these pathways, the PTEN signaling pathway had the highest degree of association with smoking-induced changes in the AJC transcriptional program. The tumor suppressor PTEN, is an important mediator of epithelial apical membrane identity [24]. Our data indicate an interesting possibility that cigarette smoking reprograms AJC-associated protein-protein interactions necessary for the generation of airway epithelial polarity.

Based on the concept that co-expressed genes might share common mechanisms of transcriptional regulation [33], we identified a “core AJC transcriptional network” composed of 21 smoking-responsive genes that are strongly correlating with the I_{AJC} in the SAE of

healthy smokers. One of the critical hallmarks of biologically meaningful transcriptional networks is the enrichment of certain gene ontology categories [32,56]. Consistent with this requirement, the core AJC transcriptional network was significantly enriched in genes encoding nuclear proteins with nucleic acid-binding activity, including the transcriptional regulators MATR3, PSIP1, NUCKS [57–59], tumor suppressor CRSP3 [60], and PKD2, a gene mediating mechanosensation in primary cilia [61]. Remarkably, MPP5, a component of the Crumbs3 polarity complex localized in the primary cilium [62], and TJP1, a TJ component capable of modulating gene expression [37], were the intrinsic AJC network components most intensively interacting with the core AJC network genes. None of identified network genes has been previously implicated in regulation of the AJC, and the network structure was different from that of the epithelial junctional complex network determined based on protein-protein interactions [63]. The overall AJC network gene expression in the healthy smokers SAE strongly correlated with PTEN expression, indicating a molecular link between the AJC transcriptional network and AJC-associated signaling pathways.

It has been previously shown that cigarette smoke, in combination with the inflammatory cytokine interleukin-1 (IL-1), induces adherens junctions disassembly in endothelial cells by suppressing PTEN activity [64], indicating the possibility that cigarette smoke can utilize different molecular mechanisms to regulate AJC integrity depending on the targeted cell type and/or cellular microenvironment. Cigarette smoke is commonly associated with inflammatory responses in various tissues, including airway epithelium [65] and inflammatory cytokines, including IL-1 can induce significant derangement in the AJC structural integrity [48]. Our study extends these observations by providing evidence that smoking can induce a broad disarray in the airway epithelial AJC architecture by reprogramming transcriptional elements encoding essential components of AJC in a non-inflammatory and non-cytotoxic manner. It is remarkable that changes in the PTEN pathway are associated with both transcription-dependent and -independent mechanisms of AJC regulation by smoking.

In vivo gene co-expression data in our study helped identify a previously unknown relationship between the AJC complex genes, AJC transcriptional network and the PTEN pathway that exist in the human airway epithelium. Although additional mechanistic studies are necessary to determine whether PTEN is a direct regulator of AJC transcriptional program (or vice versa), our data indicate that mechanisms that regulate AJC transcriptional program and PTEN signaling pathway – the two cardinal features of the epithelial apical membrane identity – are coordinate and integrated into the complex transcriptional response of the human airway epithelium to the stress of cigarette smoking *in vivo*.

AJC Reprogramming in COPD

Another observation of the present study is that smoking-induced reprogramming of the physiological AJC architecture in the SAE progresses with the development of COPD, the major smoking-induced lung disease [13,21,65]. Not only the physiological AJC-encoding genes, but also the core AJC transcriptional network genes identified for healthy smokers, were progressively suppressed in the SAE of smokers with COPD, suggesting that down-regulation of AJC gene expression in COPD might be a consequence of progressive dysregulation of transcriptional regulatory mechanisms that starts in smokers prior to clinical evidence of disease. Reciprocally to the suppressed physiological AJC genes, the expression of smoking-induced cancer-associated AJC components N-cadherin and CLDN10 further increased in the SAE of COPD smokers, as was CLDN7, another AJC gene implicated in the promotion of epithelial carcinogenesis [29,30]. Together, these observations suggest that the overall AJC molecular architecture in the airway epithelium in COPD is reprogrammed in a smoking-dependent manner from its normal gene expression

pattern essential for maintaining an appropriate barrier function toward a cancer-like molecular phenotype.

Materials & Methods

Study Population

A total of 135 subjects were included in this study including 53 healthy nonsmokers, 59 healthy smokers, and 23 COPD smokers (see Supplemental Table 1, and the inclusion and exclusion criteria).

SAE Sampling, cDNA Preparation and Microarray Processing

SAE cells (10th to 12th order bronchi) were collected by fiberoptic bronchoscopy by brushing and processed for microarray analysis as described in the Supplemental Methods. Total RNA was extracted using a modification of the TRIzol method (Invitrogen, Carlsbad, CA) and processed to generate cDNA. Genome-wide gene expression analysis was performed using HG-U133 Plus 2.0 array Affymetrix, Santa Clara, CA) according to Affymetrix protocols, hardware and software. Overall microarray quality was verified by the criteria: (1) 3/5 ratio for GAPDH 3; and (2) scaling factor 10.0 [66]. The captured image data from the HG-U133 Plus 2.0 arrays was processed using MAS5 algorithm. The data was normalized using GeneSpring version 7.3.1 (Agilent technologies, Palo Alto, CA). See Supplemental Methods for further details. The raw data are available at the Gene Expression Omnibus (GEO) site (<http://www.ncbi.nlm.nih.gov/geo/>), accession number for this dataset is GSE20257.

Characterization of the Apical Junctional Complex Gene Expression

A total of 69 AJC-related genes were selected based on the literature (Table I). Using GeneSpring 7.3.1 software, genes were considered “expressed” if detected (P call of “Present”) in 20% of subjects in each study group. Differentially expressed genes between 2 groups were identified by the criteria: (1) P call 20% of samples in any of the groups; and (2) $p < 0.05$ between the groups with a Benjamini-Hochberg correction. A gene was defined as “smoking-responsive” if its expression changed significantly in healthy smokers compared to healthy nonsmokers. Unsupervised hierarchical clustering (with standard correlation as similarity measure and the complete linkage clustering algorithm) and PCA were carried out using GeneSpring software.

AJC Gene Expression Index

The SAE “AJC gene expression index” (AJC index, I_{AJC}) was determined as a global measure of the coordinate smoking-induced changes in the SAE AJC gene expression by integrating information on expression levels of all smoking-responsive genes encoding physiological AJC components. For genes represented by more than one probe set, the probe set with the lowest p value was used. Expression values were $\log_2$ transformed. For each gene, a mean and standard deviation were calculated from the values in healthy nonsmokers, and the “normal range” was defined as within 2 standard deviations (SD) of the mean, in the direction of the smoking-induced change. I_{AJC} for each subject was defined as a percentage of AJC genes with expression levels outside the normal range, using the formula:

$$I_{AJC}(\%) = \sum_{n=1}^n cEn$$

where $E1$ has a value of 1 if the expression level for gene 1 is >2 SD below that of healthy nonsmokers or a value of 0 if the expression level is ≤ 2 SD below that of healthy nonsmokers; $E2$ is the index for gene 2, etc., and the constant ($c=100$ divided by the number of smoking-responsive AJC genes, n) normalizes the index to the percent of AJC genes that are outside of the range of healthy nonsmokers. Statistical significance of differences in I_{AJC} between the groups was determined using the Wilcoxon signed-rank test.

Functional Analysis of Gene Expression Data

Genome-wide correlation analysis was performed to identify smoking-responsive gene probe sets having gene expression pattern similar to AJC genes down-regulated by smoking ("AJC-associated genes"), i.e., genes negatively correlated with the SAE I_{ajc} in healthy smokers using the Pearson correlation coefficient (r) ≤ -0.5 as a screening cut-off criteria. To determine canonical molecular pathways enriched within the AJC-associated gene dataset, the derived gene list was subjected to Ingenuity Pathway Analysis (www.ingenuity.com). Criteria for pathway enrichment were: (1) significance of association between the canonical pathway and the dataset ($p < 0.05$, Fischer's exact test); and (2) degree of enrichment ($>1\%$ unique genes from the input dataset belonging to the pathway). Functional categories enriched in the gene datasets were determined by Gene Ontology (GO) analysis using GeneSpring software and ProfCom analytic tool [67]. For selected gene sets, Spearman's rank correlation coefficient (Rho) was determined using StatView (SAS Institute Inc, Cary, NC). A smoking-sensitive network of AJC-associated protein-protein interactions was constructed using STRING (Search Tool for the Retrieval of Interacting Genes/Proteins; <http://string-db.org>) database [68], applying a gene list containing smoking-responsive AJC genes and genes in the top AJC-associated molecular pathways as a combined input dataset using the Markov clustering algorithm or, for recently discovered interactions, based on the literature data.

AJC Transcriptional Network

The core AJC network genes were identified based on: (1) P call $\geq 20\%$; (2) significantly down-regulated in the SAE of healthy smokers vs healthy nonsmokers ($p < 0.05$, Benjamini-Hochberg correction); (3) correlation with I_{AJC} ($r \leq -0.6$; Rho ≤ -0.7 , $p < 0.0001$); and (4) correlation with ≥ 2 known AJC genes (Rho ≤ -0.7 , $p < 0.05$). The intrinsic AJC network genes (nodes) were identified based on the criteria: (1) well-characterized gene encoding a defined AJC component (Table I); (2) P call $\geq 20\%$; (3) significantly down-regulated in the SAE of healthy smokers vs healthy nonsmokers ($p < 0.05$ with Benjamini-Hochberg correction); (4) correlation with I_{AJC} ($r \leq -0.5$; Rho ≤ -0.6 , $p < 0.0001$); (5) correlation with ≥ 3 core AJC network genes (Rho ≤ -0.7 , $p < 0.05$); and (6) correlation with ≥ 2 known AJC genes (Rho ≤ -0.6 , $p < 0.05$). AJC network genes were clustered into subgroups based on unsupervised hierarchical clustering.

Human Airway Epithelial Cell Cultures and *In Vitro* Studies

The human airway epithelial cell line 16HBE14o⁻ was chosen because these cells can differentiate in a 3D-culture system acquiring essential characteristics of the normal airway epithelium, including functional TJ and AJ [69]. The cells were grown in Eagle's minimum essential medium supplemented with 2 mM L-glutamine (Gibco, Carlsbad, CA), 10% fetal calf serum, 100 μ g/ml streptomycin, and 100 U/ml penicillin G, at 37°C in a 5% CO₂ incubator. After reaching $\sim 90\%$ confluence, they were passaged and seeded (10^5 cells/cm²) onto Transwell permeable filter inserts (12 mm diameter, pore size 0.4 μ m; Costar, Cambridge, MA) and grown at 37°C, 5% CO₂ under the liquid-covered conditions. The volumes of the apical (400 μ l) and basolateral (1200 μ l) chambers were set to avoid significant hydrostatic pressure gradient across cell layers [69]. Epithelial junctional

integrity was determined by measuring transepithelial electrical resistance (Millicell ERS-2 epithelial volt-ohm-meter; Millipore, Bedford, MA).

Aqueous cigarette cell extract (CSE) was generated from the combustion of 1 cigarette (Marlboro Red) bubbled through 12.5 ml of culture medium [70]. This medium, defined as “100% CSE”, was adjusted to pH 7.4 and filtered through a 0.22 μ m filter. Experiments were started once transepithelial resistance reached $700 \Omega \times \text{cm}^2$, a level corresponding to the differentiation of 16HBE14o⁻ cells expressing functional AJC components [69]. To mimic *in vivo* cigarette smoking, differentiated 16HBE14o⁻ cells were exposed to CSE from the apical surface by adding 1, 2 or 4% CSE, concentrations nontoxic for these cells [46]. Control cells were treated with culture medium only. To model chronic cigarette smoke exposure, the apical medium containing CSE remained in the upper chamber until the next day, when the resistance was measured and the medium was replaced with a fresh CSE-containing medium. The experiments were carried out for 6 days. After CSE exposure was completed, cell viability was assessed by trypan blue exclusion and RNA extracted and processed for TaqMan RT-PCR analysis. Experiments were repeated 3 times.

TaqMan Real-time PCR

TaqMan real-time RT-PCR was performed to confirm differential expression of a subset of genes found to be differentially expressed by microarray, using the same RNA samples that were used for the microarray analysis, and to determine the effect of *in vitro* CSE exposure of cultured airway epithelial cells on expression of selected AJC genes and known smoking-inducible oxidative stress-related genes. See Supplemental Methods for details.

Immunofluorescence Analysis

Cytospin preparations of SAE obtained from randomly selected healthy smokers (n=3) and healthy nonsmokers (n=3) were processed for immunofluorescence analysis. Samples were blocked with donkey serum for 30 min (Jackson ImmunoResearch, West Grove, PA) and incubated with primary antibodies mouse monoclonal antibodies against CLDN1 (10 μ g/ml; Abnova, Taipei, Taiwan) and CDH1 (10 μ g/ml; Invitrogen, Carlsbad, CA) at 4°C overnight in a humid chamber, with mouse IgG2a and mouse IgG1 (both from R&D; Minneapolis, MN) isotype controls, respectively. Nuclei were counterstained with 4',6-diamidino-2-phenylindole (DAPI; Invitrogen, Carlsbad, CA). Images were captured using an Olympus IX70 fluorescence microscope and analyzed using MetaMorph software (Universal Imaging Corporation, Downingtown, PA). Pseudocolor images were formed with red channel Cy5 fluorescence.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank RJ Kaner, AE Tilley, MW Butler, M O'Mahony, B. Witover and B. Ferris for help with this study; and N. Mohamed for help in preparing this manuscript. These studies were supported, in part, by NIH R01 HL074326; P50 HL084936 and UL1-RR024996.

References

1. Schneeberger EE, Lynch RD. The tight junction: a multifunctional complex. *Am J Physiol Cell Physiol.* 2004; 286:C1213–C1228. [PubMed: 15151915]
2. Hartsock A, Nelson WJ. Adherens and tight junctions: structure, function and connections to the actin cytoskeleton. *Biochim Biophys Acta.* 2008; 1778:660–669. [PubMed: 17854762]

3. Shin K, Fogg VC, Margolis B. Tight junctions and cell polarity. *Annu Rev Cell Dev Biol.* 2006; 22:207–235. [PubMed: 16771626]
4. Anderson JM, Van Itallie CM. Physiology and function of the tight junction. *Cold Spring Harbor Perspect Biol.* 2009; 1:a002584.
5. Gumbiner BM. Regulation of cadherin adhesive activity. *J Cell Biol.* 2000; 148:399–404. [PubMed: 10662767]
6. Mellman I, Nelson WJ. Coordinated protein sorting, targeting and distribution in polarized cells. *Nat Rev Mol Cell Biol.* 2008; 9:833–845. [PubMed: 18946473]
7. Nelson WJ. Remodeling epithelial cell organization: transitions between front-rear and apical-basal polarity. *Cold Spring Harbor Perspect Biol.* 2009; 1:a000513.
8. Samarín S, Nusrat A. Regulation of epithelial apical junctional complex by Rho family GTPases. *Front Biosci.* 2009; 14:1129–1142.
9. Halbleib JM, Saaf AM, Brown PO, Nelson WJ. Transcriptional modulation of genes encoding structural characteristics of differentiating enterocytes during development of a polarized epithelium in vitro. *Mol Biol Cell.* 2007; 18:4261–4278. [PubMed: 17699590]
10. Knight DA, Holgate ST. The airway epithelium: structural and functional properties in health and disease. *Respirology.* 2003; 8:432–446. [PubMed: 14708552]
11. Crystal RG, Randell SH, Engelhardt JF, Voynow J, Sunday ME. Airway epithelial cells: current concepts and challenges. *Proc Am Thorac Soc.* 2008; 5:772–777. [PubMed: 18757316]
12. Hecht SS. Tobacco carcinogens, their biomarkers and tobacco-induced cancer. *Nat Rev Cancer.* 2003; 3:733–744. [PubMed: 14570033]
13. Rabe KF, Hurd S, Anzueto A, Barnes PJ, Buist SA, Calverley P, Fukuchi Y, Jenkins C, Rodriguez-Roisin R, van WC, Zielinski J. Global strategy for the diagnosis, management, and prevention of chronic obstructive pulmonary disease: GOLD executive summary. *Am J Respir Crit Care Med.* 2007; 176:532–555. [PubMed: 17507545]
14. Lee G, Walser TC, Dubinett SM. Chronic inflammation, chronic obstructive pulmonary disease, and lung cancer. *Curr Opin Pulm Med.* 2009; 15:303–307. [PubMed: 19417670]
15. Jones JG, Minty BD, Lawler P, Hulands G, Crawley JC, Veall N. Increased alveolar epithelial permeability in cigarette smokers. *Lancet.* 1980; 1:66–68. [PubMed: 6101416]
16. Knowles MR, Buntin WH, Bromberg PA, Gatzky JT, Boucher RC. Measurements of transepithelial electric potential differences in the trachea and bronchi of human subjects in vivo. *Am Rev Respir Dis.* 1982; 126:108–112. [PubMed: 7091895]
17. Boucher RC, Johnson J, Inoue S, Hulbert W, Hogg JC. The effect of cigarette smoke on the permeability of guinea pig airways. *Lab Invest.* 1980; 43:94–100. [PubMed: 7392577]
18. Petecchia L, Sabatini F, Varesio L, Camoirano A, Usai C, Pezzolo A, Rossi GA. Bronchial airway epithelial cell damage following exposure to cigarette smoke includes disassembly of tight junction components mediated by the extracellular signal-regulated kinase 1/2 pathway. *Chest.* 2009; 135:1502–1512. [PubMed: 19447922]
19. Serikov VB, Leutenegger C, Krutilina R, Kropotov A, Pleskach N, Suh JH, Tomilin NV. Cigarette smoke extract inhibits expression of peroxiredoxin V and increases airway epithelial permeability. *Inhal Toxicol.* 2006; 18:79–92. [PubMed: 16326404]
20. Sun W, Wu R, Last JA. Effects of exposure to environmental tobacco smoke on a human tracheobronchial epithelial cell line. *Toxicology.* 1995; 100:163–174. [PubMed: 7624873]
21. Hogg JC, Timens W. The pathology of chronic obstructive pulmonary disease. *Annu Rev Pathol.* 2009; 4:435–459. [PubMed: 18954287]
22. Gao N, Ishii K, Mirosevich J, Kuwajima S, Oppenheimer SR, Roberts RL, Jiang M, Yu X, Shappell SB, Caprioli RM, Stoffel M, Hayward SW, Matusik RJ. Forkhead box A1 regulates prostate ductal morphogenesis and promotes epithelial cell maturation. *Development.* 2005; 132:3431–3443. [PubMed: 15987773]
23. Wells CD, Fawcett JP, Traweger A, Yamanaka Y, Goudreau M, Elder K, Kulkarni S, Gish G, Virag C, Lim C, Colwill K, Starostine A, Metlakov P, Pawson T. A Rich1/Amot complex regulates the Cdc42 GTPase and apical-polarity proteins in epithelial cells. *Cell.* 2006; 125:535–548. [PubMed: 16678097]

24. Martin-Belmonte F, Gassama A, Datta A, Yu W, Rescher U, Gerke V, Mostov K. PTEN-mediated apical segregation of phosphoinositides controls epithelial morphogenesis through Cdc42. *Cell*. 2007; 128:383–397. [PubMed: 17254974]
25. Bartscher I, Lickert H. Foxa2 regulates polarity and epithelialization in the endoderm germ layer of the mouse embryo. *Development*. 2009; 136:1029–1038. [PubMed: 19234065]
26. Cheung ST, Leung KL, Ip YC, Chen X, Fong DY, Ng IO, Fan ST, So S. Claudin-10 expression level is associated with recurrence of primary hepatocellular carcinoma. *Clin Cancer Res*. 2005; 11:551–556. [PubMed: 15701840]
27. Sugita M, Geraci M, Gao B, Powell RL, Hirsch FR, Johnson G, Lapadat R, Gabrielson E, Bremnes R, Bunn PA, Franklin WA. Combined use of oligonucleotide and tissue microarrays identifies cancer/testis antigens as biomarkers in lung carcinoma. *Cancer Res*. 2002; 62:3971–3979. [PubMed: 12124329]
28. Prudkin L, Liu DD, Ozburn NC, Sun M, Behrens C, Tang X, Brown KC, Bekele BN, Moran C, Wistuba II. Epithelial-to-mesenchymal transition in the development and progression of adenocarcinoma and squamous cell carcinoma of the lung. *Mod Pathol*. 2009; 22:668–678. [PubMed: 19270647]
29. Darido C, Buchert M, Pannequin J, Bastide P, Zalali H, Mantamadiotis T, Bourgaux JF, Garambois V, Jay P, Blache P, Joubert D, Hollande F. Defective claudin-7 regulation by Tcf-4 and Sox-9 disrupts the polarity and increases the tumorigenicity of colorectal cancer cells. *Cancer Res*. 2008; 68:4258–4268. [PubMed: 18519685]
30. Nubel T, Preobraschenski J, Tuncay H, Weiss T, Kuhn S, Ladwein M, Langbein L, Zoller M. Claudin-7 regulates EpCAM-mediated functions in tumor progression. *Mol Cancer Res*. 2009; 7:285–299. [PubMed: 19276185]
31. Jensen LJ, Kuhn M, Stark M, Chaffron S, Creevey C, Muller J, Doerks T, Julien P, Roth A, Simonovic M, Bork P, von MC. STRING 8—a global view on proteins and their functional interactions in 630 organisms. *Nucleic Acids Res*. 2009; 37:D412–D416. [PubMed: 18940858]
32. Jansen R, Greenbaum D, Gerstein M. Relating whole-genome expression data with protein-protein interactions. *Genome Res*. 2002; 12:37–46. [PubMed: 11779829]
33. Xulvi-Brunet R, Li H. Co-expression networks: graph properties and topological comparisons. *Bioinformatics*. 2010; 26:205–214. [PubMed: 19910304]
34. Emerling BM, Weinberg F, Liu JL, Mak TW, Chandel NS. PTEN regulates p300-dependent hypoxia-inducible factor 1 transcriptional activity through Forkhead transcription factor 3a (FOXO3a). *Proc Natl Acad Sci U S A*. 2008; 105:2622–2627. [PubMed: 18268343]
35. Nho RS, Kahm J. beta1-Integrin-collagen interaction suppresses FoxO3a by the coordination of Akt and PP2A. *J Biol Chem*. 2010; 285:14195–14209. [PubMed: 20223831]
36. Harvey BG, Heguy A, Leopold PL, Carolan BJ, Ferris B, Crystal RG. Modification of gene expression of the small airway epithelium in response to cigarette smoking. *J Mol Med*. 2007; 85:39–53. [PubMed: 17115125]
37. Balda MS, Matter K. Tight junctions and the regulation of gene expression. *Biochim Biophys Acta*. 2009; 1788:761–767. [PubMed: 19121284]
38. Koch S, Nusrat A. Dynamic regulation of epithelial cell fate and barrier function by intercellular junctions. *Ann N Y Acad Sci*. 2009; 1165:220–227. [PubMed: 19538310]
39. Itoh M, Bissell MJ. The organization of tight junctions in epithelia: implications for mammary gland biology and breast tumorigenesis. *J Mammary Gland Biol Neoplasia*. 2003; 8:449–462. [PubMed: 14985640]
40. Lee M, Vasioukhin V. Cell polarity and cancer—cell and tissue polarity as a non-canonical tumor suppressor. *J Cell Sci*. 2008; 121:1141–1150. [PubMed: 18388309]
41. Auerbach O, Stout AP, Hammond EC, Garfinkel L. Changes in bronchial epithelium in relation to cigarette smoking and in relation to lung cancer. *N Engl J Med*. 1961; 265:253–267. [PubMed: 13685078]
42. Wistuba II, Mao L, Gazdar AF. Smoking molecular damage in bronchial epithelium. *Oncogene*. 2002; 21:7298–7306. [PubMed: 12379874]
43. Ballard ST, Taylor AE. Bioelectric properties of proximal bronchiolar epithelium. *Am J Physiol*. 1994; 267:L79–L84. [PubMed: 8048545]

44. Coyne CB, Gambling TM, Boucher RC, Carson JL, Johnson LG. Role of claudin interactions in airway tight junctional permeability. *Am J Physiol Lung Cell Mol Physiol.* 2003; 285:L1166–L1178. [PubMed: 12909588]
45. Olivera DS, Boggs SE, Beenhouwer C, Aden J, Knall C. Cellular mechanisms of mainstream cigarette smoke-induced lung epithelial tight junction permeability changes in vitro. *Inhal Toxicol.* 2007; 19:13–22. [PubMed: 17127639]
46. Gangl K, Reininger R, Bernhard D, Campana R, Pree I, Reisinger J, Kneidinger M, Kundi M, Dolznig H, Thurnher D, Valent P, Chen KW, Vrtala S, Spitzauer S, Valenta R, Niederberger V. Cigarette smoke facilitates allergen penetration across respiratory epithelium. *Allergy.* 2009; 64:398–405. [PubMed: 19120070]
47. Mankertz J, Tavalali S, Schmitz H, Mankertz A, Riecken EO, Fromm M, Schulzke JD. Expression from the human occludin promoter is affected by tumor necrosis factor alpha and interferon gamma. *J Cell Sci.* 2000; 113(Pt 11):2085–2090. [PubMed: 10806119]
48. Coyne CB, Vanhook MK, Gambling TM, Carson JL, Boucher RC, Johnson LG. Regulation of airway tight junctions by proinflammatory cytokines. *Mol Biol Cell.* 2002; 13:3218–3234. [PubMed: 12221127]
49. Rao R. Oxidative stress-induced disruption of epithelial and endothelial tight junctions. *Front Biosci.* 2008; 13:7210–7226. [PubMed: 18508729]
50. Zeisberg M, Neilson EG. Biomarkers for epithelial-mesenchymal transitions. *J Clin Invest.* 2009; 119:1429–1437. [PubMed: 19487819]
51. Vincent T, Neve EP, Johnson JR, Kukalev A, Rojo F, Albanell J, Pietras K, Virtanen I, Philipson L, Leopold PL, Crystal RG, de Herreros AG, Moustakas A, Pettersson RF, Fuxe J. A SNAIL1-SMAD3/4 transcriptional repressor complex promotes TGF-beta mediated epithelial-mesenchymal transition. *Nat Cell Biol.* 2009; 11:943–950. [PubMed: 19597490]
52. Dasgupta P, Rizwani W, Pillai S, Kinkade R, Kovacs M, Rastogi S, Banerjee S, Carless M, Kim E, Coppola D, Haura E, Chellappan S. Nicotine induces cell proliferation invasion and epithelial-mesenchymal transition in a variety of human cancer cell lines. *Int J Cancer.* 2009; 124:36–45. [PubMed: 18844224]
53. Gunzel D, Stuijver M, Kausalya PJ, Haisch L, Krug SM, Rosenthal R, Meij IC, Hunziker W, Fromm M, Muller D. Claudin-10 exists in six alternatively spliced isoforms that exhibit distinct localization and function. *J Cell Sci.* 2009; 122:1507–1517. [PubMed: 19383724]
54. Zemke AC, Snyder JC, Brockway BL, Drake JA, Reynolds SD, Kaminski N, Stripp BR. Molecular staging of epithelial maturation using secretory cell-specific genes as markers. *Am J Respir Cell Mol Biol.* 2009; 40:340–348. [PubMed: 18757308]
55. Lumsden AB, McLean A, Lamb D. Goblet and Clara cells of human distal airways: evidence for smoking induced changes in their numbers. *Thorax.* 1984; 39:844–849. [PubMed: 6505991]
56. Eisen MB, Spellman PT, Brown PO, Botstein D. Cluster analysis and display of genome-wide expression patterns. *Proc Natl Acad Sci U S A.* 1998; 95:14863–14868. [PubMed: 9843981]
57. Ge H, Si Y, Roeder RG. Isolation of cDNAs encoding novel transcription coactivators p52 and p75 reveals an alternate regulatory mechanism of transcriptional activation. *EMBO J.* 1998; 17:6723–6729. [PubMed: 9822615]
58. Grundt K, Haga IV, Huitfeldt HS, Ostvold AC. Identification and characterization of two putative nuclear localization signals (NLS) in the DNA-binding protein NUCKS. *Biochim Biophys Acta.* 2007; 1773:1398–1406. [PubMed: 17604136]
59. Zeitz MJ, Malyavantham KS, Seifert B, Berezney R. Matrin 3: chromosomal distribution and protein interactions. *J Cell Biochem.* 2009; 108:125–133. [PubMed: 19562669]
60. Goldberg SF, Miele ME, Hatta N, Takata M, Paquette-Straub C, Freedman LP, Welch DR. Melanoma metastasis suppression by chromosome 6: evidence for a pathway regulated by CRSP3 and TXNIP. *Cancer Res.* 2003; 63:432–440. [PubMed: 12543799]
61. Nauli SM, Alenghat FJ, Luo Y, Williams E, Vassilev P, Li X, Elia AE, Lu W, Brown EM, Quinn SJ, Ingber DE, Zhou J. Polycystins 1 and 2 mediate mechanosensation in the primary cilium of kidney cells. *Nat Genet.* 2003; 33:129–137. [PubMed: 12514735]

62. Fan S, Hurd TW, Liu CJ, Straight SW, Weimbs T, Hurd EA, Domino SE, Margolis B. Polarity proteins control ciliogenesis via kinesin motor interactions. *Curr Biol*. 2004; 14:1451–1461. [PubMed: 15324661]
63. Paris L, Bazzoni G. The protein interaction network of the epithelial junctional complex: a system-level analysis. *Mol Biol Cell*. 2008; 19:5409–5421. [PubMed: 18923145]
64. Barbieri SS, Ruggiero L, Tremoli E, Weksler BB. Suppressing PTEN activity by tobacco smoke plus interleukin-1beta modulates dissociation of VE-cadherin/beta-catenin complexes in endothelium. *Arterioscler Thromb Vasc Biol*. 2008; 28:732–738. [PubMed: 18202321]
65. Yoshida T, Tudor RM. Pathobiology of cigarette smoke-induced chronic obstructive pulmonary disease. *Physiol Rev*. 2007; 87:1047–1082. [PubMed: 17615396]
66. Raman T, O Connor TP, Hackett NR, Wang W, Harvey BG, Attiyeh MA, Dang DT, Teater M, Crystal RG. Quality control in microarray assessment of gene expression in human airway epithelium. *BMC Genomics*. 2009; 10:493. [PubMed: 19852842]
67. Antonov AV, Schmidt T, Wang Y, Mewes HW. ProfCom: a web tool for profiling the complex functionality of gene groups identified from high-throughput data. *Nucleic Acids Res*. 2008; 36:W347–W351. [PubMed: 18460543]
68. Jensen LJ, Kuhn M, Stark M, Chaffron S, Creevey C, Muller J, Doerks T, Julien P, Roth A, Simonovic M, Bork P, von MC. STRING 8—a global view on proteins and their functional interactions in 630 organisms. *Nucleic Acids Res*. 2009; 37:D412–D416. [PubMed: 18940858]
69. Ehrhardt C, Kneuer C, Fiegel J, Hanes J, Schaefer UF, Kim KJ, Lehr CM. Influence of apical fluid volume on the development of functional intercellular junctions in the human epithelial cell line 16HBE14o-: implications for the use of this cell line as an in vitro model for bronchial drug absorption studies. *Cell Tissue Res*. 2002; 308:391–400. [PubMed: 12107432]
70. Gelbman BD, Heguy A, O'Connor TP, Zabner J, Crystal RG. Upregulation of pirin expression by chronic cigarette smoking is associated with bronchial epithelial cell apoptosis. *Respir Res*. 2007; 8:10. [PubMed: 17288615]
71. Vogelmann R, Nelson WJ. Fractionation of the epithelial apical junctional complex: reassessment of protein distributions in different substructures. *Mol Biol Cell*. 2005; 16:701–716. [PubMed: 15548593]

Non-standard abbreviations

AJC	apical junctional complex
AJ	adherens junctions
CDH	cadherin
CGN	cingulin
CLDN	claudin
COPD	chronic obstructive pulmonary disease
I^{AJC}	small airway epithelium apical junctional complex gene expression index
PCA	principal component analysis
PTEN	phosphatase and tensin homolog
SAE	small airway epithelium
TJ	tight junctions
TJp	tight junction protein

Figure 1.

Expression of AJC-related genes in the SAE of healthy nonsmokers. Ordinate represents P calls (% subjects in each group expressing a given gene; yellow bars) and normalized log-transformed expression levels (mean $\pm$ SD) of each gene [red - genes with relatively high expression (>20); blue - genes with relatively low expression (<20)].

Figure 2.

Smoking alters SAE AJC gene expression. **A.** Volcano plot comparing the normalized expression of all gene probe sets (left panel) or AJC-encoding genes (right panel) in the SAE of healthy smokers vs healthy nonsmokers; y-axis - negative log of p value; x-axis - log₂-transformed fold-change; red dots - significant differentially expressed probe sets; grey dots - not significant probe sets. **B.** Differentially expressed physiological AJC-encoding genes (blue bars) and other genes belonging to various functional categories (black bars) in the SAE of healthy smokers vs healthy nonsmokers; y-axis - log₂-transformed fold-change. **C.** Diagram of the AJC architecture; significantly up- and down-regulated in the SAE of healthy smokers are depicted as red- and blue-colored boxes, respectively; yellow boxes indicate genes with no significant change; **D.** Immunofluorescence analysis of differential expression of CLDN1 and CDH1 proteins (red color) in the cytospin preparations of SAE of healthy smokers vs healthy nonsmokers. Nuclei stained with DAPI (blue color); scale bar -10 μ m. More examples are shown in Supplemental Figure 2.

Figure 3. Smoking-dependent changes in the SAE AJC gene expression in COPD. **A.** Unsupervised hierarchical clustering of healthy nonsmokers (green lines), healthy smokers (orange lines) and COPD smokers (blue lines) based on the SAE expression of AJC-encoding genes. Genes expressed above the average are represented in red, below average in blue, and average in white. **B.** Principal component analysis of healthy nonsmokers (green circles), healthy smokers (orange circles) and COPD smokers (blue circles) on all expressed gene probe sets (left panel) and on AJC-encoding gene probe sets (right panel). The %-contributions of the first 3 principal components are indicated. **C.** Distribution (% of subjects/group; ordinate) of healthy nonsmokers (green), healthy smokers (orange) and COPD smokers (blue) based on the SAE AJC index (I_{AJC} ; % of abnormally expressed AJC-encoding genes; abscissa). Inserted box-plot graph shows the I_{AJC} distribution in each group: the bottom and top of each box represent the 25th and 75th percentile (the lower and upper quartiles, respectively), and the band near the middle of the box is the 50th percentile (the median); the ends of the whiskers represent the 10th percentile (lower) and the 90th percentile (upper); data not included between the whiskers are plotted as an outlier with a dot; p values determined based on the Wilcoxon signed-rank test. **D.** Abnormal expression frequency (ordinate; % of subjects/group having expression of a given AJC-encoding gene below the normal range).

Figure 4.

Smoking-sensitive AJC-associated molecular pathways. **A.** IPA-based canonical pathways enriched among AJC-associated genes (smoking-sensitive genes negatively correlating with the I_{AJC} with $r < -0.5$). Ordinate represents enrichment (% of AJC-associated genes belonging to the pathway; orange boxes) and -log p value of enrichment. Axonal guidance pathway (enrichment - 3.3%; $p < 0.01$) was excluded as being specific to the neuronal signaling. **B.** Spearman correlation (Rho) analysis of the PTEN gene normalized expression in the SAE of healthy smokers with the I_{AJC} . **C.** STRING8.2-based model of relationships between smoking-responsive physiological AJC-encoding genes and selected AJC-related genes correlating with the I_{AJC} with $r < -0.5$. Each circle corresponds to individual gene with the size proportional to correlation (r) with the I_{AJC} , and color corresponding to the clusters the genes were grouped based on known protein-protein interactions (Markov clustering); structure of lines reflects the confidence of association [from thick (most confident) to dashed (least confident)].

Figure 5.

Smoking-sensitive AJC-associated transcriptional network in the SAE. **A.** Biological processes-related GO categories significantly enriched in the dataset of AJC-associated genes (smoking-responsive; correlating with the I_{AJC} with $r < -0.5$); x-axis - enrichments (% of AJC-associated genes in the particular GO category); y-axis shows -log p value of enrichment. Grey lines determine the arbitrary defined area of significant enrichment (enrichment > 10%; p < 0.05). **B.** The core AJC network genes and their co-expression-based “interactions” with the intrinsic AJC network genes were identified based on their correlation with the I_{AJC} and each other as described in Materials & Methods. Each circle represents individual gene with the size differences proportional to inter-network interaction intensity, and color corresponding to the clusters based on the unsupervised hierarchical clustering; structure of lines reflects the strength of correlation between 2 genes (see Supplemental Table 4). **C.** Spearman rank correlation (Rho) analysis of the PTEN gene normalized expression and the mean AJC network gene expression in the SAE of healthy smokers. **D.** Mean normalized expression levels of AJC network genes in the study groups; indicated are mean $\pm$ SD indicated for each group and p values indicate statistical differences between the groups.

Figure 6.

Effect of the cigarette smoke extract (CSE) on the airway epithelial AJC gene expression and barrier integrity *in vitro*. **A.** Viability of differentiated 16HBE14o⁻ cells apically stimulated with 1%, 2%, or 4% CSE for 6 days as described in Materials & Methods compared to unstimulated cells; n=3 in each group; N.S., non-significant difference ($p>0.05$) for all 3 comparisons. **B.** Expression of AJC genes (CLDN1, CLDN8, CGN, CDH1), PTEN pathway genes (PTEN, FOXO3), and oxidative-stress related genes (CYP1A1, CYP1B1, NQO1) in 16HBE14o⁻ cells stimulated with 2% CSE for 6 days as described in Materials & Methods compared to unstimulated cells determined by TaqMan PCR; y-axis - normalized expression levels; p values indicate statistical differences between the groups; n=3 in each group; **C.** Transepithelial resistance of differentiated 16HBE14o⁻ cell monolayers (ordinate; expressed as % of initial level) was measured during the 6-day period of stimulation with 2% CSE, as described in Materials & Methods compared to control (unstimulated) cells; p values indicate significant differences between the groups; n=3 in each group.

Table 1
Changes in the AJC Gene Expression in the SAE of Healthy Smokers and Smokers with COPD¹

Gene category		Gene	Change vs N ³		p value (vs NS)	
Gene name	symbol ²	S ⁴	COPD ⁵	S	COPD	
Transmembrane tight junction (TJ) genes						
Claudin 1	CLDN1			<0.001	6.4 × 10 ⁻⁰⁵	
Claudin 2	CLDN2			N.D. ⁶	N.D.	
Claudin 3	CLDN3			<0.009	0.023	
Claudin 4	CLDN4			N.S. ⁷	0.022	
Claudin 5	CLDN5			N.D.	N.D.	
Claudin 6	CLDN6			N.D.	N.D.	
Claudin 7	CLDN7			N.S.	<0.007	
Claudin 8	CLDN8			2.7 × 10 ⁻⁰⁷	2.9 × 10 ⁻⁰⁸	
Claudin 9/HCTP4-binding protein	CLDN9			<0.002	6.0 × 10 ⁻⁰⁴	
Claudin 10	CLDN10			4.7 × 10 ⁻¹¹	2.2 × 10 ⁻¹¹	
Claudin 11	CLDN11			N.D.	N.D.	
Claudin 12	CLDN12			N.S.	N.S.	
Claudin 14	CLDN14			N.D.	N.D.	
Claudin 15	CLDN15			N.S.	N.S.	
Claudin 16	CLDN16			N.S.	<0.003	
Claudin 17	CLDN17			N.D.	N.D.	
Claudin 18	CLDN18			N.S.	N.S.	
Claudin 19	CLDN19			N.S.	N.S.	
Claudin 20	CLDN20			N.D.	N.D.	
Occludin	OCLN			4.4 × 10 ⁻⁵	3.9 × 10 ⁻⁰⁵	
F11 receptor	F11R			<0.03	N.S.	
Junctional adhesion molecule 2	JAM2			N.D.	N.D.	
Junctional adhesion molecule 3	JAM3			N.S.	<0.02	
WNK lysine deficient protein kinase 4	WNK4			4.7 × 10 ⁻¹¹	1.4 × 10 ⁻⁰⁹	
Coxsackie virus and adenovirus receptor	CXADR			<0.005	5.8 × 10 ⁻⁰⁵	

Gene name	Gene category	Gene	Change vs N ³		p value (vs NS)	
			S ⁴	COPD ⁵	S	COPD
Cytoplasmic TJ genes						
Tight junction protein 1 (zona occludens 1)		TJP1			<0.02	5.5 × 10 ⁻⁰⁵
Tight junction protein 2 (zona occludens 2)		TJP2			N.S.	N.S.
Tight junction protein 3 (zona occludens 3)		TJP3			<0.005	N.S.
Membrane protein, palmitoylated 1, 55kDa		MPP1			<0.03	N.S.
Membrane protein, palmitoylated 7 (MAGUK p55 subfamily member 7)		MPP7			<0.04	5.2 × 10 ⁻⁰⁴
Symplekin		SYMPK			<0.003	N.S.
Membrane associated guanylate kinase, WW and PDZ domain containing 1		MAG11			8.7 × 10 ⁻⁰⁶	4.5 × 10 ⁻⁰⁶
Membrane associated guanylate kinase, WW and PDZ domain containing 2		MAG12			1.8 × 10 ⁻⁰⁹	2.8 × 10 ⁻⁰⁷
Membrane associated guanylate kinase, WW and PDZ domain containing 3		MAG13			8.3 × 10 ⁻⁰⁵	0.022
Multiple PDZ domain protein		MPDZ			<0.002	5.2 × 10 ⁻⁰⁴
Cold shock domain protein A		CSDA			<0.002	N.S.
Cingulin		CGN			2.7 × 10 ⁻⁰⁷	7.6 × 10 ⁻⁰⁵
Polarity complex genes						
Par-3 partitioning defective 3 homolog (C. elegans)		PARD3			<0.004	N.S.
Par-6 partitioning defective 6 homolog beta (C. elegans)		PARD6B			<0.005	5.0 × 10 ⁻⁰⁴
Protein kinase C, zeta		PRKCZ			<0.003	N.S.
Protein kinase C, iota		PRKCI			<0.02	<0.02
Serine/threonine kinase 11 (Peutz-Jeghers syndrome)		STK11			N.S.	N.S.
Scribbled homolog (Drosophila)		SCRIB			N.S.	N.S.
Lethal giant larvae homolog 1 (Drosophila)		LLGL1			N.S.	N.S.
Discs, large homolog 7 (Drosophila)		DLG7			N.S.	N.S.
Discs, large homolog 1 (Drosophila)		DLG1			<0.03	<0.04
Crumbs homolog 3 (Drosophila)		CRB3			N.S.	N.S.
InaD-like (Drosophila)		INADL			N.S.	<0.004
Membrane protein, palmitoylated 5 (MAGUK p55 subfamily member 5)		MPP5			8.3 × 10 ⁻⁰⁵	<0.006
Adherens junctions genes						
Cadherin 1, type 1, E-cadherin (epithelial)		CDH1			<0.03	<0.01
Cadherin 2, type 1, N-cadherin (neuronal)		CDH2			0.002	1.0 × 10 ⁻⁰⁵

Gene name	Gene category	Gene symbol ²	Change vs N ³		p value (vs NS)	
			S ⁴	COPD ⁵	S	COPD
Cadherin 3, type 1, P-cadherin (placental)		CDH3			4.7 × 10 ⁻⁰⁴	N.S.
Catenin (cadherin-associated protein), alpha 1, 102kDa		CTNNA1			N.S.	N.S.
Catenin (cadherin-associated protein), alpha-like 1		CTNNAL1			1.7 × 10 ⁻⁰⁶	N.S.
Catenin (cadherin-associated protein), beta 1, 88kDa		CTNNB1			<0.003	<0.005
Catenin, beta like 1 ; catenin, beta like 1		CTNNB1			N.S.	N.S.
Catenin, beta interacting protein 1		CTNNBIP1			<0.006	N.S.
Catenin (cadherin-associated protein), delta 1		CTNND1			<0.03	N.S.
Protein tyrosine phosphatase, receptor type, M		PTPRM			4.7 × 10 ⁻⁰⁵	N.S.
Myeloid/lymphoid or mixed-lineage leukemia; translocated to, 4		MLLT4			2.4 × 10 ⁻⁰⁵	7.0 × 10 ⁻⁰⁵
Poliovirus receptor-related 1		PVRL1			N.D.	N.D.
Poliovirus receptor-related 2		PVRL2			N.S.	N.S.
Poliovirus receptor-related 3		PVRL3			8.7 × 10 ⁻⁰⁸	<0.001
Apical polarity-regulating and other AJC-related genes						
Angiomotin		AMOT			N.D.	N.D.
Numb homolog (Drosophila)		NUMB			N.S.	N.S.
Hepatocyte nuclear factor 4, alpha		HNF4A			N.D.	N.D.
Phosphatase and tensin homolog (mutated in multiple advanced cancers 1)		PTEN			5.1 × 10 ⁻⁰⁴	2.8 × 10 ⁻⁰⁴
Forkhead box A1		FOXA1			3.0 × 10 ⁻⁰⁶	3.3 × 10 ⁻⁰⁵
Forkhead box A2		FOXA2			4.7 × 10 ⁻⁰⁴	7.6 × 10 ⁻⁰⁵

¹ Color codes for significantly differentially expressed genes (vs healthy nonsmokers); red = significantly up-regulated genes (vs healthy nonsmokers); blue = significantly down-regulated genes (vs healthy nonsmokers)

² For genes with more than 1 probe set, the probe set with the best P call and/or (for significant genes) the lowest p value (vs healthy nonsmokers) is included

³ NS = healthy nonsmokers

⁴ S = healthy smokers

⁵ COPD = COPD smokers

⁶ N.D. = not detected (P call <20% in both study groups)

⁷ N.S. = not significant