Published in final edited form as:

For *Ecol Manage*. 2013 November 15; 308: . doi:10.1016/j.foreco.2013.07.052.

Is *Markhamia lutea*'s abundance determined by animal foraging?

Colin A. Chapman^{a,*}, Tyler R. Bonnell^b, Raja Sengupta^b, Tony L. Goldberg^c, and Jessica M. Rothman^d

^aMcGill School of Environment and Department of Anthropology, McGill University, Montreal, Quebec, Canada H3A 2T7 and Wildlife Conservation Society, Bronx, New York, USA

^bDepartment of Geography, McGill University, Montreal, Quebec, Canada

^cDepartment of Pathobiological Sciences, University of Wisconsin-Madison, Madison Wisconsin, United States of America

^dDepartment of Anthropology, Hunter College of the City University of New York, and New York Consortium in Evolutionary Primatology, 695 Park Avenue, New York, NY 10065 USA

Abstract

Understanding the determinants of tropical forest tree richness and spatial distribution is a central goal of forest ecology; however, the role of herbivorous mammals has received little attention. Here we explore the potential for red colobus monkeys (*Procolobus rufomitratus*) to influence the abundance of *Markhamia lutea* trees in a tropical forest by feeding extensively on the tree's flowers, such that this tree population is not able to regularly set fruit. Using 14 years of data from Kibale National Park, Uganda, we quantify *M. lutea* flower and fruit production. Similarly, using 21 years of data, we quantify temporal changes in the abundance of stems in size classes from 1 m tall and above. Our analyses demonstrate that *M. lutea* is rarely able to produce fruit and that this corresponds to a general decline in its abundance across all size classes. Moreover, using 7 years of feeding records, we demonstrate that red colobus feed on *M. lutea*, consuming large amounts of leaf and flower buds whenever they were available, suggesting that this behavior limits fruit production. Therefore, we suggest that red colobus are presently important for structuring the distribution and abundance of *M. lutea* in Kibale. This dynamic raises the intriguing question of how a large *M. lutea* population was able to originally establish. There is no evidence of a change in red colobus population size; however, if this old-growth forest is in a non-equilibrium state, *M. lutea* may have become established when red colobus ate a different diet.

Keywords

Tree Mortality; Seed Dispersal; Herbivory; Kibale National Park; Non-Equilibrium State

1. Introduction

Tropical rain forests are renowned for their extraordinary tree species richness (Connell, 1978; Phillips *et al.*, 1994; Wright, 2002), with many forests containing more than 100 tree

© 2013 Elsevier B.V. All rights reserved.

*Corresponding Author McGill School of Environment and Department of Anthropology, 855 Sherbrooke St W., McGill University, Montreal, Quebec, Canada H3A 2T7 Colin.Chapman@McGill.ca Phone 514-378-5596 .

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

species per ha and as many as 283 species per ha (Phillips *et al.*, 1994). Understanding the factors that explain this richness and determine the causes of the spatial distribution of trees has been a central goal of forest ecology for decades (Connell, 1971; Grubb, 1977; Hubbell, 1979; Grubb, 1996; Terborgh *et al.*, 2002; Lawes *et al.*, 2008; Condit *et al.*, 2012; Dalling *et al.*, 2012). Variation in the abiotic environment has been shown by some studies to be important in predicting tree diversity and distribution (Hubbell *et al.*, 1999; Harms *et al.*, 2001; Valencia *et al.*, 2004; John *et al.*, 2007). Topography in particular is often implicated (Grubb, 1996; Valencia *et al.*, 2004), suggesting that soil moisture or nutrient availability play an important role in influencing tree distribution. For example, on Barro Colorado Island, Panama 52 of 171 shrub and tree species show distributional biases with respect to swamps and 44 species show distributional biases with respect to slopes (Harms *et al.*, 2001; Dalling *et al.*, 2012). Similarly, John *et al.* (2007) demonstrated that 104 of 258 tree species on BCI had biased distributions based on soil chemistry.

Tree diversity patterns and distributional biases can be affected by the actions of animals through seed dispersal (Wright, 2002; Balcomb and Chapman, 2003; Condit *et al.*, 2012) and herbivory (Janzen, 1970; Zanne and Chapman, 2005; Zanne *et al.*, 2005; McCall and Irwin, 2006). Insects and small rodents are typically the focus of herbivory studies and there is usually an emphasis on the seed and seedling life history stages (Howe, 1990; Terborgh and Wright, 1994; Sagers and Coley, 1995; Wright and Duber, 2001; McCall and Irwin, 2006). By contrast, the importance of non-rodent mammalian herbivores and their impact on other tree life stages has received comparatively little attention, with the exception of very large scale impacts. Perhaps the best known large scale example of animals influencing tree distribution involves elephants (*Loxodonta africana*), which play an important role in maintaining open wooded grasslands (Laws, 1970; Dublin *et al.*, 1990) and converting forest to grassland (Stuart *et al.*, 1985). Animals that modify the environment and change, maintain, and/or create new habitats in ways that affect resource availability for other species are called “ecosystem engineers” (Jones *et al.*, 1994; Jones *et al.*, 1997; Power, 1997; Wilby, 2002; Crain and Bertness, 2006; Wright and Jones, 2006; Chapman *et al.*, 2013).

Here, we investigate how herbivore could influence tree distribution patterns and explore the potential for red colobus monkeys (*Procolobus rufomitratus*) to systematically influence the abundance of *Markhamia lutea* (Bignoniaceae) trees by feeding on the tree’s flowers to the extent that the population is not able to regularly set fruit. Struhsaker (1978) documented that red colobus frequently eat all or the majority of the flowers of *M. lutea* and hypothesized that this limited fruit set. He quantified red colobus feeding over 130 months and document that *M. lutea* constituted 8.66% of the overall red colobus diet, with floral buds consumed on 59 occasions, flowers on 34 occasions, and fruit on no occasions (Struhsaker, 2010). To further evaluate this potential we first describe the number of occasions that individual trees monitored over 176 months were able to successfully flower, and did not have their developing flowers eaten by red colobus. Second, we quantified feeding of a single group of red colobus on *M. lutea* flowers over 81 months. Third, we quantify fruit set of individual trees in the area used by a red colobus focal group over almost 15 years. Finally, we describe the recruitment and stand structure of *M. lutea* on permanent plots over 24 years and examine whether the number of trees, their cumulative size, and the number of individuals recruiting into the smaller size classes decline over time.

2. Materials and Methods

2.1 Study site

This study was conducted in Kibale National Park, a mid-altitude, moist-evergreen forest in western Uganda (0°13 – 0°41 N and 30°19 – 30°32 E)(Chapman and Lambert, 2000). The area has been protected since it became a forest reserve in 1932, eventually being designated

a national park in 1993. During the time it was a forest reserve, several areas of the forest were logged (Chapman *et al.*, 2000; Chapman and Lambert, 2000), but the K-30 area (282 ha) was not commercially harvested, though some stems were removed selectively before 1970 (Struhsaker, 1997). Rainfall is heterogeneous across the park (Stampone *et al.*, 2011), but the area of the study receives an average of 1696 mm annually (1990-2011; Chapman and Chapman unpublished data collected at Makerere University Biological Field Station).

2.2 Flower Production

From 1998 to the present, we conducted phenological monitoring of 10 individuals of 36 tree species including *M. lutea*. Here, we compile phenological data from June 1998 to March 2013 ($n = 176$ months). We only monitored trees greater than the smallest DBH known to be capable of reproduction. *M. lutea* is a common edge species and also occurs in the forest in tree fall gaps (Hamilton, 1991). The flowers of *M. lutea* are bright yellow, occur in showy terminal clusters, and each trumpet shaped flower is up to 6 cm long, with 5 frilly lobes (Hamilton, 1991); thus it is very unlikely that the observer missed recording an incident when a tree had mature flowers. The red colobus often eat developing flowers. In these cases we did not score the tree as flowering since flowers never got the chance to develop to the stage where they opened and could receive pollen.

2.3 Consumption

We observed a single social group of red colobus for 88 months (May 2006 to March 2013). From May 2006 to May 2010 we conducted 30-minute scan samples. By June 2010 we had learned to easily recognize individuals and we conducted 15-minute scan samples. In total, we collected 52,700 scan samples, which represents approximately 4140 hours of observations. During these scans we recorded the individual's identity, age and sex class, behavior (e.g., feeding, traveling, being social [i.e., grooming, playing]), and the species and plant part being consumed if the animal was feeding. Data were collected by CAC and five trained field assistants who have worked with the project for up to 24 years. We conducted an intensive training period to standardize all data collection techniques and minimize interobserver error prior to beginning data collection.

2.4 Fruit Set

The presence of fruit was evaluated from June 1998 to March 2013 ($n = 176$ months) using the same 10 *M. lutea* individuals for which flowering was monitored. The fruits of *M. lutea* are 75 cm long, thin, brown capsules, that hang in clusters and split open to release seeds approximately 2.5 cm long with transparent wings (Hamilton, 1991). The conspicuous nature of the *M. lutea* fruit makes it unlikely that any fruiting events were overlooked.

2.5 Stand structure

We used data from permanent plots to test the hypotheses that the number of trees, their cumulative size, and the number of individuals recruiting into the smaller size classes were declining over time. In December 1989 we established 11 permanent tree plots (200×10 m; total area = 2.2 ha) in the relatively undisturbed section of the forest. These plots were resurveyed in May 2000, September-November 2006 (Chapman *et al.*, 2010a), and January 2013. In these plots, each tree with a DBH > 10 cm was identified to species-level, marked with an aluminum tag, and measured. When plots were resurveyed, all tagged trees were located again and measured. In addition, all new trees recruiting into the > 10 cm DBH size class were located, identified, tagged, and measured. Changes in the distribution of tree size, across years, were tested using an Anderson-Darling k-sample test.

In 1991, we identified and measured the DBH of all plants that were greater than 1.2 m in height and less than 10 cm DBH. This sampling was done in 4 m wide $\times$ 200 m long plots that were 2 m from trails and within the larger vegetation plots ($n=26$ transects). We resampled the same areas using the same methods approximately 20 years later (between July 2011 and Jan 2012). We used paired t-tests to compare stems in different size classes up to 6 cm DBH between times (stems greater than 6 cm DBH and <10 cm were rare and thus were not considered). In total, 131 *M. lutea* individuals were identified and measured in 1991 and 95 in 2011. An analysis using only the plots in the old-growth forest reveals the same trends (i.e., significant or not) as the analysis which uses plots in all areas; here we only report results for all areas since the sample is more robust

3. Results

3.1 Flower Production

All of the 10 individual *M. lutea* trees that were monitored over the 176 months produced fruit or flowered at least once, with the exception of the largest tree which did not fruit or flower (88 cm DBH). Thus, ninety percent of the trees flowered over the 176 months. However, mature flowers were only recorded in 10 of the 176 months and in only one month did two individuals have flowers, thus most of the monitored individuals flowered only once in 14.7 years.

3.2 Consumption

The red colobus regularly eat young leaves, flowers, and leaf petioles. In months that the red colobus ate from *M. lutea* trees they spent 1.26% of their feeding time eating from this species. Over a home range that averaged 45 ha (range 23-66 ha; Snaith and Chapman, 2008), they ate leaf and flower buds in 62 of the 80 months that observations were made. *M. lutea* flower consumption represents 30.7 % of all flower consumption. They were only observed to eat mature flowers on one occasion. Furthermore, in every month that one of the ten monitored trees had flowers, the red colobus ate the flowers or buds.

3.3 Fruit Set

Only on 3 occasions did one of the ten trees that were monitored for 176 months bear ripe fruit. Fruiting occurred in April 2010, April 2011, and August 2011; each event was a different tree. On four occasions trees produced unripe fruit, but these fruit did not develop to ripeness. Assuming that a tree could potentially fruit for 1 month a year, the 10 monitored trees each had at least 14 chances to fruit, for a cumulative of 140 potential events. Observed fruiting was therefore only 2.1% of potential fruiting. Among individual trees, 1 tree did not fruit or flower, 4 trees produced reproductive part on 1 occasion, 2 produced them on 3 occasions, 1 produced them on 4 occasions, and 1 produced them on 5 occasions.

3.4 Stand structure

Both the number of *M. lutea* trees and their cumulative size (DBH) declined over time in the old-growth forest (Fig. 1a, b). An examination of the stand structure in progressively larger 10 cm size bins in the four time periods (1989, 1999, 2006, and 2013), reveals a progressive decline in the number of trees in each size class, with the exception of the 50 to 59.9 size class where sample sizes were very low (Fig. 2). The distribution of tree sizes was not significantly different over the 24 years (Anderson-Darling k-sample test $P=0.77$), suggesting an even loss of trees across size classes.

By sampling subplots within the larger vegetation plots for stems from 1m in height to 10 cm DBH, we were able to locate 131 stems in 1991 and 95 in 2013 in all forestry

compartments. The classes (<1 cm DBH, <2 cm DBH.... to <6 cm DBH) were fewer in 2011 than in 1991 ($P < 0.001$ in all cases).

4. Discussion

Using data spanning 14 years on *M. lutea* flower and fruit production and spanning 21 years on the abundance of stems in size classes from 1 m tall to the largest trees, we documented that this species is rarely able to produce fruit in Kibale National Park, Uganda, which corresponds with a general decline in this species abundance. Moreover, we show that red colobus feed in these trees and often concentrate their feeding on immature flowers and buds when they are available. Thus, this herbivore is likely playing an important role in structuring the distribution pattern of *M. lutea*. Although we cannot rule out other factors affecting flower production (e.g., temperature or soil chemistry), observation by Struhsaker in the early 1970 came to similar conclusions (Struhsaker, 1978). Furthermore, both Tom Struhsaker and Colin Chapman who together have conducted research in Kibale for over 43 years have frequently seen *M. lutea* trees fruiting and flowering in the farmlands neighboring the national park where red colobus are absent, suggesting that the presence of red colobus in the national park is limiting flower production and thereby tree recruitment.

This raises the intriguing question: How was a large *M. lutea* population able to establish in the National Park to begin with? Over the last few hundred years the forest outside of what is now Kibale National Park was converted from forest to agricultural and grazing lands. With this conversion, the red colobus in the forests being converted likely moved into what is now the national park. This would have resulted in an artificially high population density in the park and could have led to the overexploitation of *M. lutea*'s floral buds and flowers. However, by the 1950s the majority of the forest that could be cleared was felled (Chapman and Lambert, 2000), so the population within the park has had approximately 60 years to adjust to an appropriate carrying capacity. Census data (groups per km²) and detailed group counts suggest that the red colobus population has been stable since the 1970s (Chapman *et al.*, 2010b; Gogarten *et al.*, Submitted). Population genetic analyses of red colobus also suggest long-term population stability (Allen *et al.*, 2012). Thus, recent demographic expansion of red colobus does not explain current trends of low *M. lutea* fruit set success. Rather, we suggest that changes in red colobus foraging behaviour, independent of demographic changes, may be impacting the *M. lutea* population that became established long ago. We have previously documented a gradual change in forest composition in the old-growth area of Kibale (Chapman *et al.*, 2010a). Species that frequently recruit into areas of disturbance (e.g., *Celtis durandii* a favored red colobus food) have a more strongly negative annualized rate of population change than trees recruiting into the understory or canopy gaps, and these species are declining in their average cumulative diameter at breast height, but not as rapidly as *M. lutea* which recruit along edges and canopy gaps (Chapman *et al.*, 2010a). These changes suggest that the forest is currently in a non-equilibrium state. If these changes have led to the decline in the abundance of preferred foods, the red colobus may have shifted to eating foods of lower preference, and one such food could be *M. lutea* flowers.

Other primate species have been described to feed extensively on certain species' flowers and may similarly be impacting their distribution and abundance. For example, spider monkeys (*Ateles geoffroyi*) limit the fruit set of *Symphonia globulifera* by eating its flowers (Riba-Hernandez and Stoner, 2005). For three months *S. globulifera* flowers represented between 86 to 100% of a study groups monthly foraging time, resulting in the complete detachment of flowers in 80% of their feeding episodes. In the subsequent fruiting season, none of these trees set fruit. In contrast 70% of trees outside of the home range of the spider monkey group fruited. Similarly, giraffes (*Giraffa camelopardalis*) consumed approximately

85% of *Acacia nigrescens* flowers within the height range they can reach, which significantly reduced flower set (Fleming *et al.*, 2006). While insect florivory has been described in a number of studies (McCall and Irwin, 2006), the impact of mammals has received little attention (but see Marell *et al.*, 2009; Pearson *et al.*, 2012). The mammal examples that do exist and the data presented here suggest that more attention should be placed on understanding the long-term effects of mammalian foraging patterns on the population structure of tropical trees.

The distribution of *M. lutea* trees over its range could be influenced by the abundance of red colobus. The five recognized species of red colobus (Grubb, 1990) have a very patchy distribution and are absent from large areas of equatorial Africa (e.g., Gabon, mainland Equatorial Guinea, and much of Cameroon) (Kingdon, 1971; Struhsaker, 1975; Oates, 1994; Struhsaker, 2010). Similarly, the Uganda red colobus (*Procolobus rufomitratus tephrosceles*) occurs in a few isolated forests along the western side of Tanzania and in Kibale (Struhsaker, 2005). This patchy distribution of an important herbivore offers a means to test the hypothesis that mammalian herbivores can be playing significant roles in structuring forest plant communities.

Acknowledgments

Funding for the research in Kibale was provided by the Canada Research Chairs Program, Natural Science and Engineering Research Council of Canada, Fonds Québécois de la Recherche sur la Nature et les Technologies, National Geographic and by NIH grant TW009237 as part of the joint NIH-NSF Ecology of Infectious Disease program and the UK Economic and Social Research Council. Tyler Bonnell was supported by an FQRNT Fellowship. Thanks to Richard Wrangham for initiating the phenology monitoring with CC in 1989. Permission to conduct this research was given by the National Council for Science and Technology and the Uganda Wildlife Authority. We thank Lauren Chapman, Tom Struhsaker, and Amy Zanne for helpful comments on this research.

References

- Allen JM, Miyamoto MM, Akwari J, Margarini K, Unvari-Martin J, Carter T, Chapman CA. Population genetics of habitat sensitive red colobus suggest long-term stability of Kibale National Park. *Ecology and Evolution*. 2012; 2:2829–2842. [PubMed: 23170217]
- Balcomb SR, Chapman CA. Bridging the gap: Influence of seed deposition on seedling recruitment in a primate-tree interaction. *Ecological Monographs*. 2003; 73:625–642.
- Chapman CA, Balcomb SR, Gillespie TR, Skorupa JP, Struhsaker TT. Long-term effects of logging on African primate communities: a 28-year comparison from Kibale National Park, Uganda. *Conser. Biol.* 2000; 14:207–217.
- Chapman CA, Bonnell TR, Gogarten JF, Lambert JE, Omeja PA, Twinomugisha D, Wasserman MD, Rothman JM. Primates as ecosystem engineers. *International Journal of Primatology*. 2013; 34:1–14.
- Chapman CA, Chapman LJ, Jacob AL, Rothman JM, Omeja PA, Reyna-Hurtado R, Hartter J, Lawes MJ. Tropical tree community shifts: Implications for wildlife conservation. *Biological Conservation*. 2010a; 143:366–374.
- Chapman CA, Lambert JE. Habitat alteration and the conservation of African primates: Case study of Kibale National Park, Uganda. *American Journal of Primatology*. 2000; 50:169–185. [PubMed: 10711532]
- Chapman CA, Struhsaker TT, Skorupa JP, Snaith TV, Rothman JM. Understanding long-term primate community dynamics: Implications of forest change. *Ecological Applications*. 2010b; 20:179–191. [PubMed: 20349839]
- Condit R, Chisholm RA, Hubbell SP. Thirty years of forest census at Barro Colorado and the importance of immigration in maintaining diversity. *PLoS ONE*. 2012; 7:e49826. [PubMed: 23226222]
- Connell, JH. On the role of natural enemies in preventing competitive exclusion in some marine animals and in rain forest trees. In: den Boer, PJ.; Gradwell, GR., editors. *Dynamics of populations*.

- Centre for Agricultural Publishing and Documentation; Wageningen, The Netherlands: 1971. p. 298–313.
- Connell JH. Diversity in tropical rain forests and coral. *Science*. 1978;1302–1309. 1999. [PubMed: 17840770]
- Crain CM, Bertness MD. Ecosystem engineering across environmental gradients: implications for conservation and management. *BioScience*. 2006; 56:211–218.
- Dalling JW, Schnitzer SA, Baldeck C, Harms KE, John R, Mangan SA, Lobo E, Yavitt JB, Hubbell SP. Resource-based habitat associations in a neotropical liana community. *Journal of Ecology*. 2012; 100:1174–1182.
- Dublin HT, Sinclair ARE, McGlade J. Elephants and fire as causes of multiple stable states in the Serengeti Mara woodlands. *Journal of Animal Ecology*. 1990; 59:1147–1164.
- Fleming PA, Hofmeyr SD, Nicolson SW, duToit JT. Are giraffes pollinators or flower predators of *Acacia nigrescens* in Kruger National Park, South Africa? *Journal of Tropical Ecology*. 2006; 22:247–253.
- Gogarten, JF.; Jacob, AL.; Ghai, RR.; Rothman, JM.; Twinomugisha, D.; Wasserman, MD.; Chapman, CA. Causes and consequences of changing group sizes in a primate community over 15+ years: Implications of non-equilibrium states. Submitted
- Grubb, P. Primate geography in the Afro-tropical forest biome. In: Peters, G.; Hutterer, R., editors. *Vertebrates in the tropics*. Museum Alexander Koenig; Bonn: 1990. p. 187–214.
- Grubb PJ. The maintenance of species-richness in plant communities: The importance of regeneration niche. *Biology Review*. 1977; 52:107–145.
- Grubb, PJ. Rainforest dynamics: the need for new paradigms. In: Edwards, DS., editor. *Tropical rainforest research – current issues*. Kluwer; Dordrecht: 1996. p. 215–233.
- Hamilton, AC. A field guide to Uganda forest trees. Makerere University Printery; Kampala: 1991.
- Harms KE, Condit R, Hubbell SP, Foster RB. Habitat associations of trees and shrubs in a 50-ha neotropical forest plot. *Journal of Ecology*. 2001; 89:947–959.
- Howe HF. Survival and growth of juvenile *Virola surinamensis* in Panama: effects of herbivory and canopy closure. *Journal of Tropical Ecology*. 1990; 6:259–280.
- Hubbell SP. Tree dispersion, abundance, and diversity in a tropical dry forest. *Science*. 1979; 203:1299–1309. [PubMed: 17780463]
- Hubbell SP, Foster RB, O'Brien ST, Harms KE, Condit R, Wechsler B, Wright SJ, Loode SL. Light-gaps, disturbances, recruitment limitation, and tree diversity in a Neotropical forest. *Science*. 1999; 283:554–557. [PubMed: 9915706]
- Janzen DH. Herbivores and the number of tree species in tropical forests. *American Naturalist*. 1970; 104:501–528.
- John R, Dalling JW, Harms KE, Yavitt JB, Stallard RF, Mirabello M, Hubbell SP, Valencia R, Navarrete H, Vallejo M, Foster RB. Soil nutrients influence spatial distributions of tropical tree species. *Proceedings of the National Academy of Sciences of the United States of America*. 2007; 104:864–869. [PubMed: 17215353]
- Jones CG, Lawton JH, Shachak M. Organisms as ecosystem engineers. *Oikos*. 1994; 69:373–386.
- Jones CG, Lawton JH, Shachak M. Ecosystem engineering by organisms: why semantics matters. *Trends in Ecology and Evolution*. 1997; 12:275. [PubMed: 21238068]
- Kingdon, J. East African mammals: An atlas of evolution in Africa. Vol. I. Academic Press; London: 1971.
- Lawes MJ, Griffiths ME, Midgley JJ, Boudreau S, Eeley HAC, Chapman CA. Tree spacing and area of competitive influence do not scale with tree size in an African rainforest. *Journal of Vegetation Science*. 2008; 19:729–738.
- Laws RM. Elephants as agents of habitat and landscape change in East Africa. *Oikos*. 1970; 21:1–15.
- Marell A, Archaux F, Korboulewsky N. Floral herbivory of the wood anemone (*Anemone nemorosa* L.) by roe deer (*Capreolus capreolus* L.). *Plant Species Biology*. 2009; 24:209–214.
- McCall A, Irwin RE. Florivory: the intersection of pollination and herbivory. *Ecology Letters*. 2006; 9:1351–1365. [PubMed: 17118009]

- Oates, JF. The natural history of African colobines. In: Davies, AG.; Oates, JF., editors. *Colobine Monkeys: their ecology, behaviour and evolution*. Cambridge University Press; Cambridge: 1994.
- Pearson DE, Potter T, Maron JL. Biotic resistance: exclusion of native rodent consumers releases populations of weaker invader. *Journal of Ecology*. 2012; 100:1383–1390.
- Phillips OL, Hall P, Gentry AH, Sawyer SA, Vázquez R. Dynamics and species richness of tropical rain forests. *Proceedings of the National Academy of Sciences of the United States of America*. 1994; 91:2905–2809. [PubMed: 8159677]
- Power M. Ecosystem engineering by organisms: why semantics matters, reply. *Trends in Ecology and Evolution*. 1997; 12:275–276. [PubMed: 21238069]
- Riba-Hernandez P, Stoner KE. Massive destruction of *Symphonia globulifera* (Clusiaceae) flowers by Central American spider monkeys (*Ateles geoffroyi*). *Biotropica*. 2005; 37:274–278.
- Sagers CL, Coley PD. Benefits and costs of defense in a neotropical shrub. *Ecology*. 1995; 76:1835–1843.
- Snaith TV, Chapman CA. Red colobus monkeys display alternative behavioural responses to the costs of scramble competition. *Behavioural Ecology*. 2008; 19:1289–1296.
- Stampone M, Hartter J, Chapman CA, Ryan SJ. Trends and variability in localized precipitation around Kibale National Park, Western Uganda, Africa. *Research Journal of Environmental and Earth Sciences*. 2011; 3:14–23.
- Struhsaker, TT. *The Red Colobus Monkey*. University of Chicago Press; Chicago: 1975.
- Struhsaker, TT. Interrelations of red colobus monkeys and rain-forest trees in the Kibale Forest, Uganda. In: Montgomery, GG., editor. *The ecology of arboreal folivore*. Smithsonian Institution Press; Washington: 1978. p. 397–422.
- Struhsaker, TT. *Ecology of an African rain forest: logging in Kibale and the conflict between conservation and exploitation*. University of Florida Press; Gainesville: 1997.
- Struhsaker TT. Conservation of red colobus and their habitats. *International Journal of Primatology*. 2005; 26:525–538.
- Struhsaker, TT. *The red colobus monkey: Variation in demography, behavior, and ecology of endangered species*. Oxford University Press; Oxford: 2010.
- Stuart NOE, Hatton JC, Spencer DHN. The effect of long-term exclusion of large herbivores on vegetation in Murchison Falls National Park, Uganda. *Biological Conservation*. 1985; 22:229–245.
- Terborgh, J.; Pitman, M.; Silman, H.; Schichter, P.; Nunez, V. Maintenance of tree diversity in tropical forests. In: Levey, D.; Silva, W.; Galetti, M., editors. *Seed dispersal and frugivory: ecology, evolution and conservation*. CABI Publishing; Wallingford: 2002. p. 1–18.
- Terborgh J, Wright SJ. Effects of mammalian herbivores on plant recruitment in two neotropical forests. *Ecology*. 1994; 75:1829–1833.
- Valencia R, Foster RB, Munoz GV, Condit R, Svenning J-C, Hernandez C, Romoleroux K, Losos E, Magard E, Balslev H. Tree species distributions and local habitat variation in the Amazon: large forest plot in eastern Ecuador. *Journal of Ecology*. 2004; 92:214–229.
- Wilby A. Ecosystem engineering: a trivialized concept? *Trends in Ecology and Evolution*. 2002; 17:307.
- Wright JP, Jones CG. The concept of organisms as ecosystem engineers ten years on: progress, limitations, and challenges. *BioScience*. 2006; 56:203–209.
- Wright JS, Duber HC. Poachers and forest fragmentation alter seed dispersal, seed survival, and seedling recruitment in the palm *Attalea butyraceae*, with implications for tropical tree diversity. *Biotropica*. 2001; 33:583–595.
- Wright SJ. Plant diversity in tropical forests: a review of mechanisms of species coexistence. *Oecologia*. 2002; 130:1–14.
- Zanne AE, Chapman CA. Diversity of woody species in forest, treefall gaps, and edge in Kibale National Park, Uganda. *Plant Ecology*. 2005; 178:121–139.
- Zanne AE, Chapman CA, Kitajima K. Evolutionary and ecological correlates of early seedling morphology in East African trees and shrubs. *American Journal Of Botany*. 2005; 92:972–978. [PubMed: 21652480]

HIGHLIGHTS

Determinants of tropical forest tree richness and spatial distribution

Red colobus monkeys influence the abundance of *Markhamia lutea*

14 years of data on *M. lutea* flower and fruit production

21 years of data on stand structure

red colobus are important for structuring the distribution of *M. lutea* in Kibale

Fig. 1.
The number and cumulative DBH (cm) of *Markhamia lutea* trees found in permanent plots during 4 time periods.

Fig. 2.
Number of *Markhamia lutea* trees found in four DBH size classes in Kibale National Park, Uganda at four time periods.