Published in final edited form as:

Psychol Med. 2010 June ; 40(6): . doi:10.1017/S0033291709991218.

Early Response to Antidepressant Treatment in Bulimia Nervosa

Robyn Sysko, Ph.D.^{1,2}, Nanshi Sha, M.S.³, Yuanjia Wang, Ph.D.^{3,4}, Naihua Duan, Ph.D.^{1,3,4}, and B. Timothy Walsh, M.D.^{1,2}

¹Department of Psychiatry, College of Physicians and Surgeons of Columbia University, New York, NY

²Division of Clinical Therapeutics, New York State Psychiatric Institute, New York, NY

³Department of Biostatistics, Mailman School of Public Health, Columbia University, New York, NY

⁴Division of Biostatistics and Data Coordination, New York State Psychiatric Institute, New York, NY

Abstract

Background—Bulimia Nervosa (BN) is a serious psychiatric disorder characterized by frequent episodes of binge eating and inappropriate compensatory behavior. Numerous trials have found that antidepressant medications are efficacious for the treatment of BN. Early response to antidepressant treatment, in the first few weeks after medication is initiated, may provide clinically useful information about an individual's likelihood of ultimately benefitting or not responding to such treatment. The purpose of this study was to examine the relationship between initial and later response to fluoxetine, the only antidepressant medication approved by the FDA for the treatment of BN, with the goal of developing guidelines to aid clinicians in deciding when to alter the course of treatment.

Methods—Data from the two largest medication trials conducted in BN (n=785) were used. Receiver operating characteristic (ROC) curves were constructed to assess whether symptom change during the first several weeks of treatment was associated with eventual non-response to fluoxetine at the end of the trial.

Results—Eventual non-responders to fluoxetine could be reliably identified by the third week of treatment.

Conclusions—Patients with BN who will fail to report a 60% or greater decrease in the frequency of binge eating or vomiting at week three are unlikely to respond to fluoxetine. As no reliable relationships between pre-treatment characteristics and eventual response to pharmacotherapy have been identified for BN, early response is one of the only available indicators to guide clinical management.

Correspondence: Dr. Robyn Sysko, Columbia Center for Eating Disorders, New York State Psychiatric Institute, 1051 Riverside Drive, Unit 98, New York, NY, 10032, phone: 212-543-4296, fax: 212-543-5607, syskor@childpsych.columbia.edu.

Presented in part at the 14th Annual Meeting of the Eating Disorder Research Society, September 25–27, 2008, Montreal, Canada, and the 42nd Annual Meeting of the Association for Behavioral and Cognitive Therapies, November 13–16, 2008, Orlando, Florida.

Declaration of Interest: This work was funded, in part, by the Rahr Eating Disorder Research Project Fund, and Dr. Sysko is supported, in part, by 5T32MH015144 from the National Institutes of Health. In the last two years, Dr. Walsh reported receiving research support from Abbott Laboratories and AstraZeneca and Dr. Duan received research support from Pfizer Pharmaceuticals. Dr. Sysko reports holding stock in Pfizer Pharmaceuticals. Dr. Wang, and Mr. Sha report no biomedical financial interests or potential conflicts of interest in the last two years.

Introduction

Bulimia Nervosa (BN) is a serious psychiatric disorder characterized by frequent episodes of binge eating and inappropriate compensatory behavior. Numerous trials have found that antidepressant medications, including both tricyclic antidepressants (TCAs) and selective serotonin reuptake inhibitors (SSRIs), produce consistently superior reductions in bulimic symptoms than those achieved with placebo (Shapiro *et al.* 2007). While TCAs and SSRIs appear to have roughly equal efficacy, SSRI antidepressants have fewer side effects and are better tolerated (Zhu & Walsh, 2002). Fluoxetine, an SSRI, is the only antidepressant currently approved by the FDA for the treatment of BN. Two large multi-site studies, sponsored by Eli Lilly & Company, evaluated fluoxetine for the treatment of BN, and found that fluoxetine at 60 mg/d produced significantly greater reductions in bulimic symptoms when compared to fluoxetine 20 mg/d and placebo (Fluoxetine Bulimia Nervosa Collaborative Study Group, 1992; Goldstein *et al.* 1995).

In general, it has proven difficult to identify reliable pre-treatment predictors of response to psychiatric treatments. However, in recent years, an increasing number of studies have examined early or rapid response to treatment as an indicator of eventual response. Early response refers to the change in symptoms during the first several weeks of treatment and rapid response or “sudden gains,” denotes a quick and clinically-meaningful change in symptoms. Across psychiatric disorders such as mood and anxiety disorders, psychosis, and eating disorders, early or rapid response can be useful in predicting treatment outcome (e.g., Grilo & Masheb, 2007; Kinon *et al.* 2008; le Grange *et al.* 2008; Masheb & Grilo, 2007; Pollack *et al.* 2008; Szegedi *et al.* 2009; Tao *et al.* 2009). In addition, for patients with eating disorders, early and rapid response are robust predictors of short- and long-term response to cognitive-behavioral therapy (CBT; Agras *et al.* 2000; Fairburn *et al.* 2004; Grilo & Masheb, 2007; Kaplan *et al.* 2009; Wilson *et al.* 1999; Wilson *et al.* 2002). However, data on response to antidepressant medications are more limited. An analysis of rapid response among patients with binge eating disorder found that non-response to medication (fluoxetine or placebo) in the first four weeks of treatment was associated with later non-response (Grilo *et al.* 2006). Grilo and colleagues (2006) also observed that patients who did not demonstrate an early rapid response to CBT continued to improve throughout treatment while initial non-responders to medication experienced limited additional benefit from fluoxetine or placebo. Another study suggested that patients with BN who will not respond to antidepressant medication can be reliably identified early in treatment (Walsh *et al.* 2006). Specifically, 80–85% of patients would be correctly identified as eventual non-responders to desipramine by the second week of treatment. However, this study had several important limitations, including the relatively small sample of patients with BN ($n=77$), and the now infrequent use of desipramine to treat BN.

Decreases in binge eating and purging may be evident among patients with BN in the first few weeks after antidepressant treatment is initiated. However, there are no guidelines for clinicians to indicate when the response should be declared inadequate and the treatment terminated (Sadock & Sadock, 2007), and recommendations for prescribing antidepressants to patients with BN often omit information about the time course for discontinuing these medications (Becker *et al.* 2008; Walsh, 2008). The ability to recognize non-response rapidly would substantially enhance clinical management. Alternative psychological and pharmacological treatments could be implemented, reducing the delay to symptom relief and patient withdrawal from treatment because of frustration over lack of response. Thus, the purpose of the current study was to examine the response to antidepressant medication in the largest treatment trials conducted for BN (Fluoxetine Bulimia Nervosa Collaborative Study Group, 1992; Goldstein *et al.* 1995) to determine whether clinically useful guidelines can be

developed to predict eventual non-response to fluoxetine on the basis of symptomatic response during the first several weeks of treatment.

Method

Description of studies

Data for the current study were provided by Eli Lilly and Company from two previously published studies of fluoxetine (Fluoxetine Bulimia Nervosa Collaborative Study Group, 1992; Goldstein *et al.* 1995) for the treatment of DSM-III R BN. The first study (Fluoxetine Bulimia Nervosa Collaborative Study Group, 1992) randomly assigned patients with BN to either fluoxetine (20 mg/d or 60 mg/d) or placebo at 13 sites in the US and Canada. A total of 387 patients participated in the study (n=129 placebo, n=129 20 mg fluoxetine, n=129 60 mg fluoxetine). The second study (Goldstein *et al.* 1995) enrolled patients at 15 sites across the United States, and examined fluoxetine at a 60 mg/d dosage in comparison to placebo. Two hundred and ninety-six patients were randomly assigned to receive fluoxetine and 102 patients to receive placebo for 16 weeks. Thus, the sample for this study included a total of 785 patients: 231 receiving placebo and 554 receiving fluoxetine (n=129 fluoxetine 20 mg; n=425 fluoxetine 60 mg). As the 1992 trial was only eight weeks in duration, our primary analyses were limited to the first eight weeks of antidepressant treatment.

The inclusion and exclusion criteria for the two studies were very similar. Patients in both studies were at least 18 years of age and met DSM-III R criteria for BN (American Psychiatric Association, 1987). Although twice weekly episodes of binge eating over the prior three months are required by DSM-III-R and DSM-IV (American Psychiatric Association, 2000) for a diagnosis of BN, these studies included patients with more severe symptoms. Specifically, individuals needed to report at least three episodes of binge eating weekly for the prior six months. Mean ages and body weights between the studies were very similar (1992 study: 27.2 yrs and 60.6 kg; 1995 study: 26.7 yrs and 58.0 kg). The 1992 study included only women and the 1995 study included 15 (1.9%) men. Both studies assessed binge eating and vomiting frequencies using a self-report weekly measure of binge eating and vomiting.

As the data provided by Eli Lilly and Company were completely anonymous, this project did not constitute human subjects research according to the federal definition [45CFR46.102(f)], and Institutional Review Board review was not required for this study.

Statistical Analyses

Receiver operating characteristic (ROC) curves were constructed to assess whether symptom change during the first several weeks of treatment predicted eventual non-response to fluoxetine at the end of the trial. ROC curves provide a graphical display of the accuracy of a measure (e.g., number of binge episodes in the first four weeks of treatment) in correctly predicting an outcome (e.g., non-response to treatment at weeks seven and eight). All possible values for “hits” (sensitivity), and “false alarms” (1-specificity) are plotted for various thresholds of the measure, and the ROC curve can be examined to optimize the tradeoff between sensitivity and specificity (McFall & Treat, 1999). The ROC curves were used to assess the association between the change in symptoms (binge eating and vomiting) at weeks one through four and the primary binary outcome measure (responder or non-responder).

The primary ROC analyses used the entire patient sample, pooled across treatment arms, to predict non-response defined as the failure to achieve at least a 75% reduction in symptoms at weeks seven and eight. Non-response was chosen as the primary outcome because patients who fail to respond and require a change in treatment are the more clinically

important group to identify. The criterion of a 75% reduction in symptoms has been used as an outcome in other medications studies of BN (Fluoxetine Bulimia Nervosa Collaborative Study Group, 1992; Goldstein *et al.* 1995; Pope *et al.* 1983; Walsh *et al.* 1987) and allowed for comparisons with our earlier study of early response to desipramine (Walsh *et al.* 2006). Further, while abstinence is an important goal for a full course of treatment for BN, a 75% reduction is a clinically significant change in the first seven or eight weeks of treatment. In the previous analysis of early response (Walsh *et al.* 2006), similar results were obtained for ROC analyses restricted to individuals receiving medication and analyses including both medication and placebo. Thus, we used the entire sample for our primary analyses and examined differences between treatment groups (20 mg fluoxetine, 60 mg fluoxetine, placebo) in secondary analyses. Area under the curve (AUC) was calculated for each ROC curve. The AUC measures the overall accuracy of the prediction, or how well the prediction separates the group being tested into those with and without the outcome in question (response vs. non-response). As a general guideline, an AUC = 0.5 provides no discrimination (e.g., chance), AUC's between 0.7 and 0.8 are considered acceptable, AUC's between 0.8 and 0.90 are considered excellent, and AUC's > 0.9 are considered outstanding (Hosmer & Lemeshow, 2000).

We conducted a variety of secondary analyses. First, ROC curves were constructed using alternative outcomes, including the prediction of other levels of response commonly employed in studies of BN, including the failure to achieve at least a 50% decline or a 100% decline (full abstinence) from baseline. Second, separate ROC curves were derived for patients assigned to 20 and 60 mg of fluoxetine and placebo, and the data from the 1992 and 1995 studies. Third, we applied Obuchowski's method (Obuchowski, 2006) that generalizes the usual dichotomous ROC analysis to examine changes in the number of episodes of binge eating and vomiting as continuous outcome measures. Fourth, we analyzed the frequency of participants' binge eating and vomiting during weeks seven and eight using a Poisson regression with the following predictors: vomiting frequency at week four, ethnicity, baseline BMI, etc. We allowed for possible over-dispersion and estimated scale parameters using Pearson's chi-square statistic. Finally, we examined ROC curves predicting the failure to achieve a 75% decline from baseline among the subset of patients who received a longer course of treatment (outcome = average of weeks fifteen and sixteen).

Results

Among the 785 patients randomized to medication or placebo, the average age of the patients was 28.0 ± 7.8 years (range 17.0–63.0) and mean body mass index (kg/m²) was 22.3 ± 3.7 (range 15.9–45.7).

Missing Data

A total of four patients (1992 study, n=1 placebo group; 1995 study, n=3 60 mg fluoxetine) reported no binge eating episodes at baseline, 67 (n=66 1992 study, n=26 placebo, n=20 fluoxetine 20 mg, n=20 fluoxetine 60 mg; 1995 study, n=1 placebo group) reported no vomiting at baseline, and one patient (1992 study, placebo) reported neither binge eating or vomiting. The Fluoxetine Bulimia Nervosa Collaborative Study Group (1992) reported that 67 of 387 patients enrolled in the study did not use self-induced vomiting as their primary compensatory behavior. These individuals reported abuse of laxatives or diuretics, fasting, strict dieting, or rigorous exercise, with many patients indicating the use of more than one method to prevent weight gain. Thus, the number of patients who did not report vomiting episodes at baseline in the 1992 study may be explained by the use of other forms of inappropriate compensatory behavior. It is not entirely clear why other patients (n=6) reported no binge eating or vomiting at baseline; however, this may be related to reactivity, including the timing of the baseline assessment, which occurred post placebo wash-out, or

the time-frame for the measurement, which was only the prior week. The 72 individuals without binge eating or vomiting at baseline were excluded from the ROC analyses described below, resulting in an initial sample of 713 patients. Among these 713 patients, the average number of binge eating episodes was 11.5 ± 11.2 (range 1.0–150.0) and the average vomiting episodes per week was 13.6 ± 16.5 (range 1.0–225.0).

Primary Analysis: Predicting Non-Response at Weeks Seven and Eight

ROC curves predicted non-response at weeks seven and eight using the percentage change in binge eating or vomiting during the first four weeks of the study. Missing data are excluded from the fitting of ROC curves. At weeks seven and eight, a total of 557 and 510 patients had data on binge eating frequency and vomiting, respectively. A total of 375 patients (67.3%; n=108 placebo; n= 74 fluoxetine 20 mg; n=193 fluoxetine 60 mg) failed to show a 75% or greater decrease in binge eating at weeks seven and eight, and 348 (68.2%; n=100 placebo; n= 56 fluoxetine 20 mg; n=192 fluoxetine 60 mg) were classified as non-responders for vomiting. Similar results were obtained when the ROC curves were fitted separately by treatment arms (placebo, fluoxetine 20 mg, or fluoxetine 60 mg), and by study (1992 or 1995). Thus, data from the combined sample are presented in Figures 1A and 1B.

In an ROC analysis, AUC indicates the accuracy of the prediction (Table 1). Starting with the week three data, the AUCs were in the excellent range (0.808 for binge eating, 0.815 for vomiting). Only the ROC curves constructed from data at week four had greater AUCs (0.828 for binge eating, 0.819 for vomiting), but these AUCs were still in the excellent range (Hosmer & Lemeshow, 2000), suggesting that the accuracy of the prediction from response at the fourth week of treatment was not substantially better. A large proportion of the patients who would fail to respond to medication by weeks seven and eight could be correctly classified by week three. If medication was discontinued for patients who failed to demonstrate a reduction in binge eating of approximately 60% at Week 3, our analyses indicated that 78% of patients who would have failed to respond to medication would be correctly identified (sensitivity), and 27.5% who would have responded to medication at weeks seven and eight would be misclassified as non-responders (1-specificity). A similar pattern was observed for vomiting, with a cut point of a decrease in vomiting of approximately 60% at week three, 79% of the eventual non-responders would be correctly classified as failing to respond to fluoxetine and 26% of eventual responders would be misclassified as non-responders.

Secondary Analyses

Secondary ROC analyses predicting 50% and 100% reduction in binge eating and vomiting at Week 8 found similar results to the analyses using 75% response. Specifically, the AUCs for week three were in the acceptable range when predicting a 100% reduction in binge eating (0.778), and in the excellent range when predicting a 50% reduction in binge eating (0.819) or vomiting (0.819), or a 100% response for vomiting (0.824). Exploratory analyses using Obuchowski's method (Obuchowski, 2006) for a continuous outcome for binge eating and vomiting did not produce more accurate predictions than the dichotomous indicator of response or non-response based on a 75% reduction in bulimic symptoms. The Poisson model, which included additional predictors, produced an ROC curve with an AUC slightly better than the AUCs for the percent change in binge eating or vomiting at the first or second week of treatment, and similar to AUCs observed for change in binge eating or vomiting at the third or fourth weeks of treatment. A subset of patients had data on binge eating (n=327) and vomiting (n=304) at weeks fifteen and sixteen. The ROC curves for the longer-term response identified week three as the most accurate in predicting non-response (AUCs = 0.763 for binge eating, 0.757 for vomiting). A cut point of approximately 60% at week three would identify 69% and 68% of the eventual non-responders for binge eating and vomiting,

respectively, and 28% and 30% of eventual responders would be misclassified as non-responders.

Discussion

The purpose of the current study was to evaluate whether clinically useful guidelines could be developed to predict eventual non-response to fluoxetine on the basis of response during the first several weeks of treatment using the largest extant sample of patients with BN (Fluoxetine Bulimia Nervosa Collaborative Study Group, 1992; Goldstein *et al.* 1995). The results suggest that a majority of patients with BN who will not respond to fluoxetine can be identified early in treatment, specifically by the third week. At week three, if fluoxetine was discontinued for patients who demonstrated less than a 60% reduction in binge eating, approximately 73% would be correctly classified as eventual non-responders to fluoxetine and 23% of patients who would eventually respond to fluoxetine would be misclassified as non-responders. For patients with less than a 60% decrease in vomiting at week three, approximately 72% would be correctly classified as non-responders and 22% would be misclassified.

These data, in combination with the findings of our previous research (Walsh *et al.* 2006), suggest that eventual antidepressant treatment response in BN can be reasonably predicted from change in symptoms during the first several weeks of treatment. Further, these results are consistent with studies of cognitive behavioral therapy (CBT), which has been extensively studied in randomized controlled trials of adults with BN (National Institute for Clinical Excellence, 2004). Studies by Wilson, Fairburn, and colleagues (Wilson *et al.* 1999; Wilson *et al.* 2002) suggest that response to CBT in BN begins in the first few weeks after the initiation of treatment, and early response to CBT is also predictive of longer-term outcome (Fairburn *et al.*, 2004). However, while early response is a reliably observed predictor of response to psychological and pharmacological treatments for patients with BN, the mechanisms of this phenomenon are not known.

The results of the current study also suggest applications for the clinical treatment of BN. Some patients with BN may be reluctant to take a medication (Wilson & Fairburn, 2007), but might be willing to consider a brief trial of fluoxetine to evaluate whether they are likely to eventually respond to antidepressant treatment. The ability to make rapid adjustments to the course of treatment if a patient does not experience a 50% reduction in symptoms by the third week might improve treatment efficacy and compliance. Although approximately 20% of non-responders are misclassified at week 3 using the 50% criterion, clinicians might still consider switching to another medication treatment or CBT at week 3 to evaluate whether a different intervention might provide greater symptom relief.

There are several limitations to the current study, including the use of retrospective analyses, and the use of criterion for response was based on data from a self-report daily diary and not an interview. In addition, as the analyses examined non-response only during acute treatment and not over a follow-up period, the findings may not generalize over longer periods of time or following the discontinuation of fluoxetine. Finally, this study focused on patients reporting vomiting as their primary means of compensation, and the results may be different for individuals with BN using other types of purging behaviors (e.g., laxatives). Strengths of the current study include the sample size, the consistency of the findings regardless of the cutpoint or endpoint (seven-eight weeks vs. fifteen-sixteen weeks) used for eventual response, and the examination of response to fluoxetine, the only medication currently approved by the FDA for the treatment of BN.

In summary, this study focused on early response to fluoxetine among patients with BN. By the third week of treatment, the majority of patients who will eventually fail to respond to fluoxetine can be identified with reasonable accuracy, and the medication can be discontinued for individuals who have not experienced a 60% or greater reduction in bulimic symptoms. Since no reliable relationships between pre-treatment characteristics and eventual response to pharmacotherapy have been identified, early response is one of the only available indicators to guide clinical management.

Acknowledgments

The authors would like to acknowledge the assistance of Eli Lilly and Company, and Dr. Charles M. Beasley Jr., in particular, for generously providing the data presented here.

References

- American Psychiatric Association. Diagnostic and Statistical Manual of Mental Disorders. 3. American Psychiatric Association Press; Washington, DC: 1987. revision
- American Psychiatric Association. Diagnostic and Statistical Manual of Mental Disorders. 4. American Psychiatric Association Press; Washington, DC: 2000. text revision
- Agras WS, Crow SJ, Halmi KA, Mitchell JE, Wilson GT, Kraemer HC. Outcome predictors for the cognitive behavior treatment of bulimia nervosa: data from a multisite study. *American Journal of Psychiatry*. 2000; 157:1302–1308. [PubMed: 10910795]
- Becker, AE.; Mickley, DW.; Derenne, JL.; Klibanski, A. Eating disorders: Evaluation and management. In: Stern, TA.; Rosenbaum, JF.; Fava, M.; Biederman, J.; Rauch, SL., editors. *Massachusetts General Hospital: Comprehensive Clinical Psychiatry*. Mosby, Inc; Philadelphia, PA: 2008. p. 499-518.
- Fairburn CG, Agras WS, Walsh BT, Wilson GT, Stice E. Prediction of outcome in bulimia nervosa by early change in treatment. *American Journal of Psychiatry*. 2004; 161:2322–2324. [PubMed: 15569910]
- Fluoxetine Bulimia Nervosa Collaborative Study Group. Fluoxetine in the treatment of bulimia nervosa: A multicenter, placebo-controlled, double-blind trial. *Archives of General Psychiatry*. 1992; 49:139–147. [PubMed: 1550466]
- Goldstein DJ, Wilson MG, Thompson VL, Potvin JH, Rampey AH. The Fluoxetine Bulimia Nervosa Research Group. Long-term fluoxetine treatment of bulimia nervosa. *British Journal of Psychiatry*. 1995; 166:660–666. [PubMed: 7620754]
- Grilo CM, Masheb RM. Rapid response predicts binge eating and weight loss in binge eating disorder: findings from a controlled trial of orlistat with guided self-help cognitive behavioral therapy. *Behaviour Research and Therapy*. 2007; 45:2537–2550. [PubMed: 17659254]
- Grilo CM, Masheb RM, Wilson GT. Rapid response to treatment for binge eating disorder. *Journal of Consulting and Clinical Psychology*. 2006; 74:602–613. [PubMed: 16822116]
- Hosmer, DW.; Lemeshow, S. *Applied logistic regression*. 2. Wiley; New York: 2000.
- Kaplan AS, Walsh BT, Olmsted M, Attia E, Carter JC, Devlin MJ, Pike KM, Woodside B, Rockert W, Roberto CA, Parides M. The slippery slope: prediction of successful weight maintenance in anorexia nervosa. *Psychological Medicine*. 2009; 10:1–9.
- Kinon BJ, Chen L, Ascher-Svanum H, Stauffer VL, Kollack-Walker S, Sniadecki JL, Kane JM. Predicting response to atypical antipsychotics based on early response in the treatment of schizophrenia. *Schizophrenia Research*. 2008; 102:230–240. [PubMed: 18423985]
- le Grange D, Doyle P, Crosby RD, Chen E. Early response to treatment in adolescent bulimia nervosa. *International Journal of Eating Disorders*. 2008; 41:755–757. [PubMed: 18570193]
- Masheb RM, Grilo CM. Rapid response predicts treatment outcomes in binge eating disorder: implications for stepped care. *Journal of Consulting and Clinical Psychology*. 2007; 75:639–644. [PubMed: 17663617]
- McFall RM, Treat TA. Quantifying the information value of clinical assessments with signal detection theory. *Annual Review of Psychology*. 1999; 50:215.

- National Institute for Clinical Excellence. Eating disorders: Core interventions in the treatment and management of eating disorders in primary and secondary care. National Institute for Clinical Excellence; London: 2004.
- Obuchowski NA. An ROC-type measure of diagnostic accuracy when the gold standard is continuous-scale. *Statistical Medicine*. 2006; 25:481–493.
- Pollack MH, Kornstein SG, Spann ME, Crits-Christoph P, Raskin J, Russell JM. Early improvement during duloxetine treatment of generalized anxiety disorder predicts response and remission at endpoint. *Journal of Psychiatric Research*. 2008; 42:1176–1184. [PubMed: 18348888]
- Pope HG, Hudson JI, Jonas JM, Yurgelun-Todd D. Bulimia treated with imipramine: A placebo-controlled, double-blind study. *American Journal of Psychiatry*. 1983; 140:554–558. [PubMed: 6342421]
- Sadock, BJ.; Alcott Sadock, V. Kaplan & Sadock's Synopsis of Psychiatry: Behavioral Sciences/Clinical Psychiatry. 10. Lippincott Williams & Wilkins; Philadelphia, PA: 2007.
- Shapiro JR, Berkman ND, Brownley KA, Sedway JA, Lohr KN, Bulik CM. Bulimia nervosa treatment: A systematic review of randomized controlled trials. *International Journal of Eating Disorders*. 2007; 40:321–336. [PubMed: 17370288]
- Szegedi A, Jansen WT, van Willigenburg AP, van der Meulen E, Stassen HH, Thase ME. Early improvement in the first 2 weeks as a predictor of treatment outcome in patients with major depressive disorder: a meta-analysis including 6562 patients. *Journal of Clinical Psychiatry*. 2009; 70:344–53. [PubMed: 19254516]
- Tao R, Emslie G, Mayes T, Nakonezny P, Kennard B, Hughes C. Early prediction of acute antidepressant treatment response and remission in pediatric major depressive disorder. *Journal of the American Academy of Child and Adolescent Psychiatry*. 2009; 48:71–78. [PubMed: 19057412]
- Walsh, BT. Eating disorders. In: Tasman, A.; Kay, J.; Lieberman, JA.; First, MB.; Maj, M., editors. *Psychiatry*. 3. John Wiley & Sons, Ltd; West Sussex, England: 2008. p. 1609-1625.
- Walsh BT, Sysko R, Parides MK. Early response to medication among women with bulimia nervosa. *International Journal of Eating Disorders*. 2006; 39:72–75. [PubMed: 16254873]
- Walsh BT, Wilson GT, Loeb KL, Devlin MJ, Pike KM, Roose SP, Fleiss J, Waternaux C. Medication and psychotherapy in the treatment of bulimia nervosa. *American Journal of Psychiatry*. 1987; 154:523–31. [PubMed: 9090340]
- Wilson, GT.; Fairburn, CG. Treatments for Eating Disorders. In: Nathan, PE.; Gorman, JM., editors. *Treatments That Work*. 3. Oxford University Press; New York: 2007. p. 579-610.
- Wilson GT, Fairburn CG, Agras WS, Walsh BT, Kraemer H. Cognitive-behavioral therapy for bulimia nervosa: Time course and mechanisms of change. *Journal of Consulting and Clinical Psychology*. 2002; 70:267–274. [PubMed: 11952185]
- Wilson GT, Loeb KL, Walsh BT, Labouvie E, Petkova E, Liu X, Waternaux C. Psychological versus pharmacological treatments of bulimia nervosa: Predictors and processes of change. *Journal of Consulting and Clinical Psychology*. 1999; 67:451–459. [PubMed: 10450615]
- Zhu AJ, Walsh BT. Pharmacologic treatment of eating disorders. *Canadian Journal of Psychiatry*. 2002; 47:227–234.

Figure 1.

Figures 1A and 1B. ROC Curves predicting the failure to achieve a 75% or greater reduction in binge eating (1A) or vomiting (1B) by weeks 7 & 8 based on the change in symptoms from baseline during the first four weeks of treatment.

Table 1

Areas Under the Curve for Binge Eating and Vomiting at Each Week of Treatment

Binge Eating		
Test Variable	AUC	SE
% Change in Binge Eating from Baseline to Week 1	0.647	0.025
% Change in Binge Eating from Baseline to Week 2	0.756	0.023
% Change in Binge Eating from Baseline to Week 3	0.808	0.020
% Change in Binge Eating from Baseline to Week 4	0.828	0.020
Vomiting		
Test Variable	AUC	SE
% Change in Vomiting from Baseline to Week 1	0.695	0.026
% Change in Vomiting from Baseline to Week 2	0.739	0.025
% Change in Vomiting from Baseline to Week 3	0.815	0.021
% Change in Vomiting from Baseline to Week 4	0.819	0.021

AUC=Area Under the Curve; SE= Standard Error