Current FDA-Related Drug Information

Approvals, Submission, and Important Labeling Changes for US Marketed Pharmaceuticals

Danial E. Baker, PharmD, FASHP, FASCP*

This monthly feature will help readers keep current on new drugs, new indications, dosage forms, and safety-related changes in labeling or use. Efforts have been made to ensure the accuracy of this information; however, if there are any questions, please let me know at bakerdan@wsu.edu.

Table 1. New drugs approved by the US Food and Drug Administration (FDA): November 20, 2012 through December 20, 2012

Cabozantinib – <i>Cometriq</i> (Exelixis)	
<i>Comparative agents:</i>	Vandetanib
<i>Indication:</i>	Treatment of patients with progressive, metastatic medullary thyroid cancer
<i>Mechanism of action:</i>	Kinase inhibitor
<i>Common adverse effects:</i>	Diarrhea, stomatitis, palmar-plantar erythrodysesthesia syndrome, decreased weight, decreased appetite, nausea, fatigue, oral pain, hair color changes, dysgeusia, hypertension, abdominal pain, and constipation
<i>Dosage form & strength:</i>	Capsules; 20 mg, 80 mg
<i>Product labeling:</i>	http://www.accessdata.fda.gov/drugsatfda_docs/label/2012/203756lbl.pdf
Influenza virus vaccine – <i>Flucelvax</i> (Novartis)	
<i>Comparative agents:</i>	Influenza vaccines
<i>Indication:</i>	Prevention of influenza
<i>Mechanism of action:</i>	Immunostimulation
<i>Common adverse effects:</i>	Local and systemic reactions in adults were injection site pain, injection site erythema, headache, fatigue, myalgia, and malaise.
<i>Dosage form & strength:</i>	Injection; 0.5 mL
<i>Product labeling:</i>	http://www.fda.gov/downloads/BiologicsBloodVaccines/Vaccines/ApprovedProducts/UCM329134.pdf
Pasireotide diaspertate – <i>Signifor</i> (Novartis)	
<i>Comparative agents:</i>	Somatostatin analogs
<i>Indication:</i>	Treatment of adult patients with Cushing's disease for whom pituitary surgery is not an option or has not been curative
<i>Mechanism of action:</i>	Binds to human somatostatin receptors resulting in inhibition of ACTH secretion, which leads to decreased cortisol secretion
<i>Common adverse effects:</i>	Diarrhea, nausea, hyperglycemia, cholelithiasis, headache, abdominal pain, fatigue, and diabetes mellitus
<i>Dosage form & strength:</i>	Injection; 0.3 mg/mL, 0.6 mg/mL, 0.9 mg/mL single-dose ampules
<i>Product labeling:</i>	http://www.accessdata.fda.gov/drugsatfda_docs/label/2012/200677lbl.pdf

(continued)

*Director, Drug Information Center, College of Pharmacy, Washington State University Spokane, PO Box 1495, Spokane, WA 99210-1495.

Table 1. New drugs approved by the US Food and Drug Administration (FDA): November 20, 2012 through December 20, 2012 (CONT.)

Ponatinib – Iclusig (ARIAD Pharmaceuticals)	
<i>Comparative agents:</i>	Imatinib, lapatinib
<i>Indication:</i>	Treatment of adult patients with chronic phase, accelerated phase, or blast phase chronic myeloid leukemia (CML) that is resistant or intolerant to prior tyrosine kinase inhibitor therapy or Philadelphia chromosome positive acute lymphoblastic leukemia (Ph+ALL) that is resistant or intolerant to prior tyrosine kinase inhibitor therapy
<i>Mechanism of action:</i>	Inhibited the in vitro tyrosine kinase activity of ABL and T315I mutant AB plus other mechanisms; reduced the size of tumors expressing native or T315I mutant BCR-ABL
<i>Common adverse effects:</i>	Hypertension, rash, abdominal pain, fatigue, headache, dry skin, constipation, arthralgia, nausea, pyrexia, thrombocytopenia, anemia, neutropenia, lymphopenia, and leukopenia
<i>Dosage form & strength:</i>	Tablet; 15 mg and 45 mg
<i>Product labeling:</i>	http://www.accessdata.fda.gov/drugsatfda_docs/label/2012/203469lbl.pdf
Raxibacumab – (GlaxoSmithKline)	
<i>Comparative agents:</i>	Ciprofloxacin
<i>Indication:</i>	Treatment of inhalational anthrax
<i>Mechanism of action:</i>	Neutralizing antibody; recombinant, human, IgG1 λ monoclonal antibody
<i>Common adverse effects:</i>	Headache, infections, nausea, pain in extremity, pruritus
<i>Dosage form & strength:</i>	Injection; 50 mg/mL to 1700 mg
<i>Product labeling:</i>	http://www.accessdata.fda.gov/drugsatfda_docs/label/2012/125349s000lbl.pdf

Table 2. New dosage forms and indications approved by the FDA: November 20, 2012 through December 20, 2012

Generic name	Brand name (Company)	Indication and comments
New indications		
Abiraterone acetate	<i>Zytiga</i> (Janssen Biotech)	Treatment of patients with metastatic castration-resistant prostate cancer in combination with prednisone
Eltrombopag	<i>Promacta</i> (GlaxoSmith Kline)	Treatment of thrombocytopenia in patients with chronic hepatitis C to allow the initiation and maintenance of interferon-based therapy
Unoprostone isopropyl	<i>Rescula</i> (Sucampo Pharma)	Treatment of open-angle glaucoma or ocular hypertension

Table 3. Agents pending FDA approval: November 20, 2012 through December 20, 2012

Generic name	Brand name (Company)	Indication and comments
Recommended for approval by an FDA advisory panel		
Telavancin	<i>Vibativ</i> (Theravance)	Mixed vote for its use in the treatment of hospital-acquired pneumonia. Advisory panel voted against (9-6) for its use in the treatment of hospital-acquired pneumonia, but supported its efficacy (13-2) for treating nosocomial pneumonia when alternative anti-infectives are not available. It should be considered useful in the treatment of nosocomial pneumonia patients with methicillin-resistant <i>Staphylococcus aureus</i> who have creatinine clearance above 30 mL/min.
Agents scheduled for review by an FDA advisory panel		
Rintatolimod	<i>Ampligen</i> (Hemispherx Biopharma)	Treatment of patients with chronic fatigue syndrome

Table 4. New drug / biologics license applications filed by manufacturer: November 20, 2012 through December 20, 2012

Generic name Brand name (Company)	Comparative agents	Indication	Mechanism of action	Common adverse effects	Dosage form or route
Canagliflozin / metformin (Janssen Research & Development)	None	Type 2 diabetes	Sodium-glucose transport protein (SGLT2) inhibitor; inhibits glucose reabsorption in the kidney	Headache, nausea	Oral
Cariprazine (Forest Laboratories)	Atypical antipsychotics	Treatment of bipolar I disorder and schizophrenia	D2 and D3 receptor antagonist	Restlessness, tremors, dyspepsia, vomiting	Oral

Table 5. Supplemental applications filed by manufacturer: November 20, 2012 through December 20, 2012

Generic name	Brand name (Company)	Comments
Erlotinib	<i>Tarceva</i> (Astellas Pharma)	First-line treatment of patients with locally advanced or metastatic non-small-cell lung cancer whose tumors have epidermal growth factor receptor (EGFR) activating mutations
Tacrolimus	(Astellas Pharma)	Extended-release capsule formulation for the prophylaxis or organ rejection in adult kidney transplant recipients and adult male liver transplant recipients.
Ustekinumab	<i>Stelara</i> (Janssen Biotech)	Treatment of adult patients with active psoriatic arthritis

Table 6. Significant labeling changes or “Dear Health Professional” letters related to safety^a

Generic name <i>Brand name</i> (Company)	Warning
Ondansetron <i>Zofran</i> (GlaxoSmithKline)	Removal of the 32 mg single intravenous dose product because of serious cardiac risks (eg, QT interval prolongation) http://www.fda.gov/Safety/MedWatch/SafetyInformation/SafetyAlertsforHumanMedicalProducts/ucm330772.htm
Sodium oxybate <i>Xyrem</i> (Jazz Pharmaceuticals)	Use with alcohol or central nervous system depressant drugs can impair consciousness and may lead to severe respiratory depression. http://www.fda.gov/Safety/MedWatch/SafetyInformation/SafetyAlertsforHumanMedicalProducts/ucm332430.htm?source=govdelivery
Telaprevir <i>Incivek</i> (Vertex Pharmaceuticals)	Case reports of serious skin reactions, some fatal, when telaprevir is used with peginterferon alfa and ribavirin to treat hepatitis C. If the patient develops a rash with systemic symptoms or a progressive severe rash, all 3 drugs should be discontinued immediately and the patient should receive urgent medical care. http://www.fda.gov/Safety/MedWatch/SafetyInformation/SafetyAlertsforHumanMedicalProducts/ucm332860.htm

^aPractitioners are encouraged to check the FDA’s MEDWATCH Web site (<http://www.fda.gov/medwatch/safety.htm>) for updated information.