Published in final edited form as:

Breast Cancer Res Treat. 2013 June ; 139(3): . doi:10.1007/s10549-013-2602-2.

Metalloproteinase-disintegrin ADAM12 is associated with a breast tumor-initiating cell phenotype

Hui Li,

Department of Biochemistry and Molecular Biophysics, Kansas State University, 141 Chalmers Hall, Manhattan, KS 66506, USA

Sara Duhachek-Muggy,

Department of Biochemistry and Molecular Biophysics, Kansas State University, 141 Chalmers Hall, Manhattan, KS 66506, USA

Suzanne Dubnicka, and

Department of Statistics, Kansas State University, Manhattan, KS 66506, USA

Anna Zolkiewska

Department of Biochemistry and Molecular Biophysics, Kansas State University, 141 Chalmers Hall, Manhattan, KS 66506, USA

Abstract

Members of the ADAM family of proteases have been associated with mammary tumorigenesis. Gene profiling of human breast tumors identified several intrinsic subtypes of breast cancer, which differ in terms of their basic biology, response to chemotherapy/radiation, preferential sites of metastasis, and overall patient survival. Whether or not the expression of individual ADAM proteases is linked to a particular subtype of breast cancer and whether the functions of these ADAMs are relevant to the cancer subtype have not been investigated. We analyzed several transcriptomic datasets and found that *ADAM12L* is specifically up-regulated in claudin-low tumors. These tumors are poorly differentiated, exhibit aggressive characteristics, have molecular signatures of epithelial-to-mesenchymal transition (EMT), and are rich in markers of breast tumor-initiating cells (BTICs). Consistently, we find that *ADAM12L*, but not the alternative splice variant *ADAM12S*, is a part of stromal, mammosphere, and EMT gene signatures, which are all associated with BTICs. In patients with estrogen receptor-negative tumors, high expression of *ADAM12L*, but not *ADAM12S*, is predictive of resistance to neoadjuvant chemotherapy. Using MCF10DCIS.com breast cancer cells, which express the endogenous ADAM12L and efficiently form mammospheres when plated at the density of single cell per well, we show that ADAM12L plays an important role in supporting mammosphere growth. We postulate that ADAM12L may serve as a novel marker and/or a novel therapeutic target in BTICs.

Keywords

Metalloprotease; Disintegrin; Claudin-low tumors; Tumor-initiating cells; Epithelial-to-mesenchymal transition; Mammospheres

© Springer Science+Business Media New York 2013

A. Zolkiewska zolkiea@ksu.edu.

Conflict of interest The authors declare that they have no conflict of interest.

Electronic supplementary material The online version of this article (doi:10.1007/s10549-013-2602-2) contains supplementary material, which is available to authorized users.

Introduction

Members of the ADAM (a disintegrin and metalloproteinase) family are frequently up-regulated in breast tumors [1]. It has been postulated that they may serve as attractive new biomarkers and/or therapeutic targets in breast cancer [2]. However, very little is currently known about patterns of ADAM expression in individual molecular subtypes of breast cancer. Based on gene expression profiling, breast cancers can be classified into several intrinsic subtypes: luminal A and B, HER2-enriched, basal-like, claudin-low, and normal-like [3]. Each of these subtypes is characterized by a unique gene signature, basic biology, response to endocrine and chemotherapy regimens, preferential sites of metastasis, and overall patient survival [4–7]. To consider any ADAM-based approaches, it is important to establish whether the expression of individual ADAM proteases is linked to a particular subtype of breast cancer and whether the functions of these ADAMs contribute to subtype-specific phenotypes.

ADAM12 is one of the most prominent breast cancer-associated *ADAM* genes. *ADAM12* mRNA is alternatively spliced in human [8]. *ADAM12L* (transcript variant 1, exons 1–18, 20–24) encodes the long, transmembrane protein isoform. *ADAM12S* (transcript variant 2, exons 1–19) gives rise to the short, secreted protein isoform. While *ADAM12L* levels in breast tumors have been consistently found higher than those in normal mammary tissue [9–11], the status of *ADAM12S* is less clear. *ADAM12S* was reported to be up-regulated in breast cancer based on qRT-PCR [12], but these results are at odds with DNA microarray profiling of ductal and lobular carcinomas versus normal ductal and lobular cells [11]. Several other studies observed increased expression of ADAM12 in breast tumors at the mRNA or protein level, but the reagents (PCR primers or antibodies) did not distinguish between ADAM12L and ADAM12S splice variants/protein isoforms [13–15].

Due to the differences in membrane association, the biological functions of ADAM12L and ADAM12S may be different as well. Analysis of the expression levels of these two variants in individual breast cancer subtypes may provide important clues about the role of ADAM12 in mammary oncogenesis. In this work, we analyzed a number of transcriptomic datasets and found that *ADAM12L* is specifically up-regulated in claudin-low tumors. These tumors are poorly differentiated, exhibit aggressive characteristics, have molecular signatures of EMT, and are enriched in features of mammary stem cells [16] and BTICs [17–21]. Consistently, *ADAM12L*, but not *ADAM12S*, is a part of the stromal, mammosphere, and EMT gene signatures, which are all associated with BTICs. In patients with estrogen receptor (ER)-negative tumors, high expression of *ADAM12L* is predictive of resistance to neoadjuvant chemotherapy. Importantly, we found that ADAM12L supports tumorsphere growth of MCF10DCIS.com breast cancer cells, suggesting that ADAM12L function may actively contribute to the BTICs phenotype.

Methods

Gene expression datasets

Gene expression datasets were retrieved from Gene Expression Omnibus (GEO, <http://www.ncbi.nlm.nih.gov/geo/>). GSE series numbers and patient characteristic are provided in Table 1. The list of gene probe sets is provided in Online Resource Table S1.

Statistical analyses

ADAM expression levels in breast cancer subtypes were analyzed using the one-way analysis of variance (ANOVA) with Bonferroni post-test analysis for multiple pairwise comparisons. ANOVA, unpaired *t* test, and paired *t* test were performed using GraphPad Prism, version 5.0 (GraphPad Software, Inc.). To identify associations between variables

and chemotherapy response, univariate and multivariate logistic regression analyses were performed using the R software [22]. *ADAM12L* and *ADAM12S* gene expression levels were treated as continuous variables; all other explanatory variables were binary. Only the complete cases were used in the multivariate models.

Cells and cell treatment

HMLE and HMLE-Twist-ER cells (a gift from Dr. Robert A. Weinberg, Whitehead Institute for Biomedical Research, Cambridge, MA) were maintained as described [23]. To induce EMT, HMLE cells were treated with 2.5 ng/ml of TGF β 1 (R&D Systems), and HMLE-Twist-ER cells were incubated with 20 nM of 4-hydroxy-tamoxifen (Sigma) for 12–20 days. Control cells were treated with vehicle only. MCF10DCIS.com cells (Asterand) were grown in DMEM/F12 (1:1) with 5 % horse serum and 29 mM NaHCO₃. For *ADAM12* knockdown, MCF10DCIS.com cells were incubated with MISSION™ Lentiviral shADAM12 Transduction Particles (Sigma, clone ID TRCN0000047037) or with MISSION™ Non-Target shRNA Control Transduction Particles (Sigma, SHC002V), according to the manufacturer's instructions. shADAM12 clone TRCN0000047037 was selected from five different shADAM12 clones, because preliminary analysis established that it produced the most efficient *ADAM12* knockdown. After 1 day, media containing lentiviral particles were replaced with fresh media, and after additional 24 h, stably transduced cells were selected with 3 μ g/ml of puromycin for 7 days. Stable overexpression of the wild-type ADAM12L and the G479E mutant in MCF10DCIS.com cells was performed by retroviral transduction, as described [24].

qRT-PCR

qRT-PCR was performed as described [25]. The following primers were used: h*ADAM12L* (F), 5'-AGCCACACCAGGATAGAGAC-3'; h*ADAM12L* (R), 5'-CGCCTTGAGTGACACTACAG-3'; h-*ACTIN* (F), 5'-TTGCCGACAGGATGCAGAA-3'; and h-*ACTIN* (R), 5'-GCCGATCCACACGGAGTACT-3'.

Tumorsphere growth

Adherent cells were detached using 0.25 % trypsin and 1 mM EDTA, counted, and suspended in complete mammary epithelial basal medium (Lonza), supplemented with 20 ng/ml EGF, 20 ng/ml bFGF, 1 $\times$ B27, 4 μ g/mL heparin, and 1 % penicillin/streptomycin. After serial dilution to a concentration of 10 cells/ml, 100 μ l of cell suspension was plated in 96-well ultralow attachment plate (Corning). One day after plating, each well was examined using a light microscope to determine the number of cells present; wells containing more than one cell were excluded from further analysis. Cells were allowed to grow undisturbed for 10 days and then each well was examined again to record the number of spheres formed.

Immunoblotting

Immunoblotting was performed using anti-ADAM12 (Ab #3394), as described [24].

Results

ADAM12L is elevated in claudin-low tumors

We examined the expression levels of all catalytically active, not testis-specific, ADAM metalloproteases [26] in different molecular subtypes of breast tumors from the human tumor database at the University of North Carolina (the UNC337 dataset, Ref. [18]). Pairwise comparison of claudin-low tumors versus basal-like, HER2-enriched, luminal A, luminal B, and normal-like tumors showed that *ADAM12L* was the only *ADAM* that was

significantly elevated in claudin-low tumors (Fig. 1). *ADAM12S* was not represented on Agilent platforms used for gene profiling in that study.

To directly compare *ADAM12L* and *ADAM12S* levels in different subtypes of breast cancer, we used three other datasets, EMC286, EMC192, and MSK82 [27–29], profiled on Affymetrix platforms. Tumor classification into basal-like, claudin-low, HER2-enriched, combined luminal, and normal-like subtypes was adopted after the study of Harrell et al. [6]. While *ADAM12L* was significantly up-regulated in claudin-low tumors, no differences in the expression level of *ADAM12S* were observed between different tumor subtypes (Fig. 2).

Claudin-low signatures are enriched in residual tumors remaining after endocrine therapy for ER-positive disease [20]. We analyzed *ADAM12L* and *ADAM12S* levels before and after letrozole treatment in two different datasets [20,30]. In both datasets, *ADAM12L*, but not *ADAM12S*, was significantly increased in tumors after letrozole therapy (Fig. 3).

ADAM12L is associated with stromal/mesenchymal and mammosphere gene signatures

Claudin-low subtype is characterized by high expression of stromal/mesenchymal genes, a high abundance of CD44 and a low amount of CD24, and is associated with the mammosphere gene signature [17, 18]. Therefore, we analyzed *ADAM12L* and *ADAM12S* levels in the relevant, publicly available gene expression studies. In a study where breast tumor epithelium and stroma were isolated using laser capture microdissection [31], *ADAM12L* expression was significantly higher in stromal than in epithelial cells (Fig. 4a). Strikingly, *ADAM12S* showed an opposite pattern of expression, and it was significantly enriched in epithelial cells (Fig. 4a). When primary breast tumor cells were sorted by flow cytometry into CD44⁺/CD24^{−/low} cells and “other” populations [20], *ADAM12L* was enriched in the CD44⁺/CD24^{−/low} population. Although the difference in *ADAM12L* expression did not reach statistical significance, it was clearly more pronounced than the difference in *ADAM12S* expression (Fig. 4b). Finally, *ADAM12L*, but not *ADAM12S*, was significantly higher in cancer mammospheres grown in suspension than in primary bulk tumors [20] (Fig. 4c). These observations are consistent with the previously reported up-regulation of *ADAM12L* in mammospheres formed by normal mammary epithelial cells [32].

ADAM12L is induced during epithelial-to-mesenchymal transition (EMT)

Analysis of the results of several gene expression studies indicates that *ADAM12L* is significantly up-regulated during EMT. First, in human mammary epithelial (HMLE) cells induced to undergo EMT by overexpressing TGF- β 1, transcription factors Twist, Gsc, or Snail [33], or by knocking down expression of E-cadherin [34], *ADAM12L* was significantly up-regulated during EMT (Fig. 4d). Second, *ADAM12L* expression was higher in spontaneously arising mesenchymal subpopulations of cells (MSPs) than in CD24^{hi} epithelial subpopulation of HMLE cells [35] (Fig. 4e). Third, in Ras-transformed HMLE cells treated in culture with either paclitaxel, a commonly used breast cancer chemotherapeutic drug, or salinomycin, a compound that selectively targets cells after EMT [36], *ADAM12L* was significantly lower in salinomycin-treated cells (Fig. 4f). In contrast, *ADAM12S* was not associated with EMT in any of the datasets analyzed in Fig. 4d–f.

ADAM12L predicts resistance to chemotherapy in ER-negative breast tumors

EMT and mammosphere gene signatures are important characteristics of BTICs [37–40]. BTICs are largely resistant to chemotherapy [41–43]. Consistently, a stromal metagene, a group of highly co-expressed 50 genes with a pattern of expression similar to that of EMT and mammospheres, was reported to predict resistance to neoadjuvant chemotherapy in ER-negative breast tumors [44]. While *ADAM12* was listed as one of the 50 genes of the

stromal metagene, the relative contribution of *ADAM12L* and *ADAM12S* variants was not addressed. Here, we asked whether there is a difference in the predictive value of *ADAM12L* and *ADAM12S* to chemotherapy and whether *ADAM12L* is an independent predictor of chemoresistance.

Among patients with ER-negative tumors undergoing preoperative anthracycline-based chemotherapy (the EORTC10994 dataset, [Ref. 44]), *ADAM12L* was significantly lower in cases with pathologic complete response (pCR, $n = 28$) than in patients with near pCR ($n = 37$) (Fig. 5a). In contrast, there was no difference in the level of *ADAM12S* expression between these two groups. In another study of the response to anthracycline/taxane-based neoadjuvant therapy (the MAQCII/MDACC dataset, [refs. 45, 46]), *ADAM12L* was significantly lower in ER-negative patients with pCR ($n = 40$) compared to patients with residual disease ($n = 50$) (Fig. 5b). *ADAM12S* again did not significantly differ between the two groups.

To determine if *ADAM12L* or *ADAM12S* provides predictive information for resistance of ER-negative tumors to chemotherapy in the two datasets, univariate and multivariate logistic regression analyses were performed with the nodal status and T stage (for the EORTC10994 dataset), and the nodal status, T stage, patient age, and tumor grade (for the MAQCII/MDACC dataset) as categorical variables, and with *ADAM12L* and *ADAM12S* gene expression levels as continuous variables. The information on patient age and tumor grade was not available for the EORTC10994 dataset.

ADAM12L was the only significant variable in univariate analysis in both datasets (Table 2). *ADAM12L* was also a predictive factor of pCR in the multivariate analysis of the EORTC10994 dataset [odds ratio (OR) 0.37; 95 % confidence interval (CI) 0.16–0.77; $P = 0.0129$], with the nodal status, T stage, and *ADAM12S* included in the model. In multivariate analysis of the MAQCII/MDACC dataset, *ADAM12L* was borderline significantly associated with pCR (OR 0.76, CI 0.54–1.04, $P = 0.0936$). Similar borderline significance of *ADAM12L* was achieved when age and grade were omitted from the model to match the number of variables with the EORTC10994 dataset (results not shown). *ADAM12S* was not predictive of pCR in any model (Table 2).

ADAM12L in normal and neoplastic mammary cells

To begin to characterize possible roles of ADAM12L protein in BTICs, we investigated the relationship between ADAM12L, EMT, and sphere growth. First, we validated the results of gene expression profiling by qRT-PCR. HMLE cells were induced to undergo EMT with TGF β , and EMT was confirmed by the loss of expression of E-cadherin. *ADAM12L* level increased by ~fivefold and ~tenfold after 12 or 20 days of treatment with TGF β (Fig. 6a). Similarly, when HMLE cells with tamoxifen-inducible expression of TWIST1, HMLE-TWIST-ER were induced to undergo EMT by 12- or 20-day treatment with 4-hydroxy-tamoxifen, *ADAM12L* was up-regulated by ~twofold and ~threefold, respectively (Fig. 6b). For mammosphere experiments, we chose MCF10DCIS.com cells, an ER-negative, CD44-positive breast cancer cell line that forms spheres with high efficiencies when plated in ultralow attachment plates in serum-free media supplemented with growth factors [47]. Comparing to the monolayer conditions, the level of *ADAM12L* was ~fivefold higher in MCF10DCIS.com cells grown as spheres (Fig. 6c). Furthermore, among all *ADAM* mRNA encoding catalytically active, transmembrane, non-testis-specific *ADAMs* (i.e., *ADAM8*, *9*, *10*, *12*, *15*, *17*, *19*, *28*, and *33*), the only other *ADAM* that was up-regulated in spheres was *ADAM28* (result not shown). We calculated the mRNA copy numbers for *ADAM12L* and *ADAM28* in MCF10DCIS.com spheres and found that the abundance of *ADAM12L* was at least two orders of magnitude higher than that of *ADAM28* (result not shown), suggesting that *ADAM12* is the main *ADAM* up-regulated in cells grown as spheres.

An important question is whether ADAM12L plays an active role in the process of EMT or sphere growth. By up-regulating or down-regulating *ADAM12L* in mammary MCF10A or HMLE cells, we did not observe a significant effect on the time course of EMT induced by limiting doses of TGF (results not shown). In contrast, ADAM12L appeared to support sphere formation in [MCF10DCIS.com](#) cells. To ensure that spheres contain the progeny of single cells truly capable of proliferation in suspension (Fig. 7a), and not simply clusters of originally plated cells, [MCF10DCIS.com](#) cells were seeded at the density of 1 cell/well. Down-regulation of *ADAM12* by shRNA significantly decreased the sphere formation efficiency (Fig. 7b). In addition, cells with down-regulated *ADAM12L* level were efficiently excluded from spheres (Fig. 7c), further suggesting that ADAM12L protein may have an important function during sphere formation.

The shRNA construct used in the experiments shown in Fig. 7 was more potent in down-regulating *ADAM12L* than several other shRNA tested, but it targeted both the *ADAM12L* and *ADAM12S* variants. As another approach to specifically limit the function of the ADAM12L variant in [MCF10DCIS.com](#) cells, we overexpressed the dominant-negative mutant of ADAM12L, G479E [48]. While overexpression of the wild-type (WT) ADAM12L did not significantly increase the efficiency of sphere formation, the number of spheres formed by cells overexpressing the G479E mutant was significantly lower than the number of spheres formed by cells with overexpression of the WT ADAM12L (Fig. 8a). The level of ADAM12L in spheres was further compared to the level of ADAM12L in cells grown as monolayers by Western blotting (Fig. 8b). We observed a weak signal representing the endogenous ADAM12L and strong signals for the WT and G479E proteins. The WT protein existed as the full-length, nascent form of ~120 kDa and the processed, active form of ~90 kDa. As expected, the G479E mutant was not processed and lacked the active form [48]. Importantly, the ratio of the amount of G479E mutant protein in spheres versus monolayers was significantly lower than the ratio of the WT ADAM12L protein (nascent and processed forms combined) (Fig. 8c). This result indicated that the G479E mutant might have been selectively excluded from the spheres, further suggesting an important role of the endogenous, active ADAM12L in sphere formation.

Discussion

Our analysis of a number of transcriptomic datasets for primary breast tumors demonstrates that *ADAM12L* and *ADAM12S* transcripts show distinct patterns of expression in breast cancer. This finding is remarkable, because the two corresponding protein isoforms, ADAM12L and ADAM12S, have different biochemical properties and unique intracellular localizations. Since *ADAM12L*, and not *ADAM12S*, is up-regulated in claudin-low tumors that are enriched for BTIC features [20], and since it is a part of the EMT and mammosphere gene signatures, both of which are also linked to BTICs [20, 21, 42], we postulate that any potential function of ADAM12 in BTICs requires membrane anchoring present in ADAM12L, and not in ADAM12S. While the mechanisms responsible for the differential regulation of *ADAM12L* and *ADAM12S* in breast cancer are not known, the most likely scenario involves post-transcriptional up-regulation of *ADAM12L* via specific elements in its unique 3'-UTR, a hypothesis that is currently being tested in our laboratory.

Our analysis also reveals that among different molecular subtypes of breast cancer, *ADAM12L* is specifically up-regulated in the claudin-low subtype. Interestingly, recent profiling of histologically benign tissue adjacent to breast tumors identified two distinct subtypes of tumor microenvironment, named "Active" and "Inactive" [49]. The "Active" cancer-adjacent tissue showed features of EMT and expressed claudin-low-related genes, including *ADAM12L*. These claudin-low gene expression features were more common in the extratumoral than intratumoral locations and were independent of whether the tumor

itself was claudin-low or not. Furthermore, claudin-low features in normal breast tissues from non-diseased women have been shown to be associated with high risk of developing breast cancer [50]. *ADAM12L* was also significantly increased in this healthy claudin-low group, reinforcing the inherent link between *ADAM12L* and the claudin-low phenotype.

ADAM12 was reported earlier to be a part of stromal gene signatures [44, 51, 52], and here we show that this conclusion applies selectively to the *ADAM12L* variant. While it is accepted that the stromal gene cluster represents reactive tumor stroma, the cellular origin of the stromal gene signature is less clear. Since the pattern of gene expression within the stromal cluster is similar to the EMT and mammosphere signatures, it has been argued that EMT of tumor epithelial cells could account for increased stromal gene expression [44]. The EMT program directly contributes to the resistance of chemotherapeutic agents [36, 53–56]. Consistently, we found *ADAM12L* to be an independent predictor of chemoresistance to neoadjuvant anthracycline-based therapy in ER-negative breast tumors from the EORTC10994 dataset. In the MAQCII/MDACC dataset, the association between the level of *ADAM12L* expression in ER-negative tumors and resistance to anthracycline/taxane-based neoadjuvant therapy was borderline significant. Since the pathological response of claudin-low tumors to anthracyclines and taxanes are similar [57], we believe that a somewhat lesser power of *ADAM12L* to predict response to chemotherapy in the MAQCII/MDACC datasets may be related to a different method of tissue harvesting [fine needle aspiration (FNA) versus core biopsy in the EORTC10994 dataset]. As FNA samples typically contain little stromal component [58, 59], contribution of stromal/mesenchymal genes to gene signatures in the MAQCII/MDACC dataset must have been naturally underestimated.

The defining characteristics of human breast cancer subtypes, including the claudin-low subtype, are conserved among human cancer cell lines and among mouse models of the disease [17, 18]. Because *ADAM12L* is up-regulated in human claudin-low tumors, we believe that claudin-low cell lines and mouse models are best suited to study the role of *ADAM12L* in mammary tumorigenesis. Of note, the role of *ADAM12* in breast cancer was previously addressed using luminal MCF7 and T47D cell lines [12, 60] or TgMMTV-PyMT mice [13, 61], which are models of the luminal tumors [17, 18]. Application of claudin-low cell lines or mouse models in the future may be necessary to obtain new insights into the biological role of *ADAM12L* in the breast cancer disease.

In conclusion, up-regulation of *ADAM12L* in claudinlow tumors, during EMT, and in mammospheres, as well as poor chemotherapy response of tumors with high expression of *ADAM12L*, all link *ADAM12L* to a BTIC phenotype. Whether or not *ADAM12L* may serve as a marker for BTICs, or at least a certain subpopulation of BTICs, is currently being investigated in our laboratory. In particular, we are performing orthotopic cell transplantation assays to directly compare the tumorigenic potential of cells with high versus low expression of the endogenous *ADAM12L*. Sphere formation assay, while representing a simple and frequently used tool for evaluation of stem-like properties of cells and a surrogate assay for tumorigenicity, has important limitations. As it has been demonstrated previously, not all spheres are necessarily derived from stem-like cells. In fact, for primary mouse mammary epithelial cells, the proportion of spheres containing the regenerative stem cells can be as low as 15–33 % [62]. Therefore, sphere formation assays alone may not provide an accurate measure of the frequency of BTICs, and limiting dilution transplantation assays are needed to directly evaluate the tumorigenic potential of *ADAM12L*-expressing cells.

Finally, the fact that down-regulation of *ADAM12L* compromised sphere formation of MCF10DCIS.com cells suggests that *ADAM12L* protein may play an active role in sphere

growth and/or may increase survival of cells deprived of attachment to the matrix. Molecular mechanisms by which ADAM12L would enhance sphere growth are currently not known, but they may involve modulation of critical signaling pathways such as Notch, EGFR, or TGF [24, 63, 64]. If an active role of ADAM12L in maintaining the BTICs phenotype is confirmed, ADAM12L may be considered as a candidate for a therapeutic target in these cells.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work was supported by NIH grants R15CA151065 and R01CA172222, and Innovative Research Award from Terry C. Johnson Center for Basic Cancer Research at KSU. This is contribution 13-289-J from Kansas Agricultural Experiment Station.

References

1. Murphy G. The ADAMs: signalling scissors in the tumour microenvironment. *Nat Rev Cancer*. 2008; 8:929–941. doi:10.1038/nrc2459. [PubMed: 19005493]
2. Duffy MJ, Mullooly M, O'Donovan N, Sukor S, Crown J, Pierce A, McGowan PM. The ADAMs family of proteases: new biomarkers and therapeutic targets for cancer? *Clin Proteomics*. 2011; 8:9. doi:10.1186/1559-0275-8-9. [PubMed: 21906355]
3. Perou CM, Borresen-Dale AL. Systems biology and genomics of breast cancer. *Cold Spring Harbor Persp Biol*. 2011; 3(2):1–17. doi:10.1101/cshperspect.a003293.
4. Comprehensive molecular portraits of human breast tumours. *Nature*. 2012; 490:61–70. doi:10.1038/nature11412. [PubMed: 23000897]
5. Prat A, Ellis MJ, Perou CM. Practical implications of gene-expression-based assays for breast oncologists. *Nat Rev Clin Oncol*. 2012; 9:48–57. doi:10.1038/nrclinonc.2011.178. [PubMed: 22143140]
6. Harrell JC, Prat A, Parker JS, Fan C, He X, Carey L, Anders C, Ewend M, Perou CM. Genomic analysis identifies unique signatures predictive of brain, lung, and liver relapse. *Breast Cancer Res Treat*. 2012; 132:523–535. doi:10.1007/s10549-011-1619-7. [PubMed: 21671017]
7. Heiser LM, Sadanandam A, Kuo WL, Benz SC, Goldstein TC, Ng S, Gibb WJ, Wang NJ, Ziyad S, Tong F, Bayani N, Hu Z, Billig JJ, Dueregger A, Lewis S, Jakkula L, Korkola JE, Durinck S, Pepin F, Guan Y, Purdom E, Neuvial P, Bengtsson H, Wood KW, Smith PG, Vassilev LT, Hennessy BT, Greshock J, Bachman KE, Hardwicke MA, Park JW, Marton LJ, Wolf DM, Collisson EA, Neve RM, Mills GB, Speed TP, Feiler HS, Wooster RF, Haussler D, Stuart JM, Gray JW, Spellman PT. Sub-type and pathway specific responses to anticancer compounds in breast cancer. *Proc Natl Acad Sci USA*. 2012; 109:2724–2729. doi:10.1073/pnas.1018854108. [PubMed: 22003129]
8. Kveiborg M, Albrechtsen R, Couchman JR, Wewer UM. Cellular roles of ADAM12 in health and disease. *Int J Biochem Cell Biol*. 2008; 40:1685–1702. doi:10.1016/j.biocel.2008.01.025. [PubMed: 18342566]
9. Iba K, Albrechtsen R, Gilpin BJ, Loechel F, Wewer UM. Cysteine-rich domain of human ADAM 12 (meltrin alpha) supports tumor cell adhesion. *Am J Pathol*. 1999; 154:1489–1501. pii:S0002-9440(10)65403-X. [PubMed: 10329602]
10. Bertucci F, Finetti P, Cervera N, Charafe-Jauffret E, Mamessier E, Adelaide J, Debono S, Houvenaeghel G, Maraninchi D, Viens P, Charpin C, Jacquemier J, Birnbaum D. Gene expression profiling shows medullary breast cancer is a subgroup of basal breast cancers. *Cancer Res*. 2006; 66(9):4636–4644. doi:10.1158/0008-5472.CAN-06-0031. [PubMed: 16651414]
11. Turashvili G, Bouchal J, Baumforth K, Wei W, Dziechciarkova M, Ehrmann J, Klein J, Fridman E, Skarda J, Srovnal J, Hajduch M, Murray P, Kolar Z. Novel markers for differentiation of lobular and ductal invasive breast carcinomas by laser microdissection and microarray analysis. *BMC Cancer*. 2007; 7:55. doi:10.1186/1471-2407-7-55. [PubMed: 17389037]

12. Roy R, Rodig S, Bielenberg D, Zurakowski D, Moses MA. ADAM12 transmembrane and secreted isoforms promote breast tumor growth: a distinct role for ADAM12-S protein in tumor metastasis. *J Biol Chem*. 2011; 286:20758–20768. doi:10.1074/jbc.M110.216036. [PubMed: 21493715]
13. Kveiborg M, Fröhlich C, Albrechtsen R, Tischler V, Dietrich N, Holck P, Kronqvist P, Rank F, Mercurio AM, Wewer UM. A role for ADAM12 in breast tumor progression and stromal cell apoptosis. *Cancer Res*. 2005; 65:4754–4761. doi:10.1158/0008-5472.CAN-05-0262. [PubMed: 15930294]
14. Lendeckel U, Kohl J, Arndt M, Carl-McGrath S, Donat H, Rocken C. Increased expression of ADAM family members in human breast cancer and breast cancer cell lines. *J Cancer Res Clin Oncol*. 2005; 131:41–48. doi:10.1007/s00432-004-0619-y. [PubMed: 15565459]
15. Mitsui Y, Mochizuki S, Kodama T, Shimoda M, Ohtsuka T, Shiomi T, Chijiwa M, Ikeda T, Kitajima M, Okada Y. ADAM28 is overexpressed in human breast carcinomas: implications for carcinoma cell proliferation through cleavage of insulin-like growth factor binding protein-3. *Cancer Res*. 2006; 66(20):9913–9920. doi:10.1158/0008-5472.CAN-06-0377. [PubMed: 17047053]
16. Lim E, Vaillant F, Wu D, Forrest NC, Pal B, Hart AH, Asselin-Labat ML, Gyorki DE, Ward T, Partanen A, Feleppa F, Huschtscha LI, Thorne HJ, Fox SB, Yan M, French JD, Brown MA, Smyth GK, Visvader JE, Lindeman GJ. Aberrant luminal progenitors as the candidate target population for basal tumor development in BRCA1 mutation carriers. *Nat Med*. 2009; 15:907–913. doi:10.1038/nm.2000. [PubMed: 19648928]
17. Herschkowitz JI, Simin K, Weigman VJ, Mikaelian I, Usary J, Hu Z, Rasmussen KE, Jones LP, Assefnia S, Chandrasekharan S, Backlund MG, Yin Y, Khramtsov AI, Bastein R, Quackenbush J, Glazer RI, Brown PH, Green JE, Kopelovich L, Furth PA, Palazzo JP, Olopade OI, Bernard PS, Churchill GA, Van Dyke T, Perou CM. Identification of conserved gene expression features between murine mammary carcinoma models and human breast tumors. *Genome Biol*. 2007; 8:R76. doi:10.1186/gb-2007-8-5-r76. [PubMed: 17493263]
18. Prat A, Parker JS, Karginova O, Fan C, Livasy C, Herschkowitz JI, He X, Perou CM. Phenotypic and molecular characterization of the claudin-low intrinsic subtype of breast cancer. *Breast Cancer Res*. 2010; 12:R68. doi:10.1186/bcr2635. [PubMed: 20813035]
19. Hennessy BT, Gonzalez-Angulo AM, Stenke-Hale K, Gilcrease MZ, Krishnamurthy S, Lee JS, Fridlyand J, Sahin A, Agarwal R, Joy C, Liu W, Stivers D, Baggerly K, Carey M, Lluch A, Monteaudo C, He X, Weigman V, Fan C, Palazzo J, Hortobagyi GN, Nolden LK, Wang NJ, Valero V, Gray JW, Perou CM, Mills GB. Characterization of a naturally occurring breast cancer subset enriched in epithelial-to-mesenchymal transition and stem cell characteristics. *Cancer Res*. 2009; 69:4116–4124. doi:10.1158/0008-5472.CAN-08-3441. [PubMed: 19435916]
20. Creighton CJ, Li X, Landis M, Dixon JM, Neumeister VM, Sjolund A, Rimm DL, Wong H, Rodriguez A, Herschkowitz JI, Fan C, Zhang X, He X, Pavlick A, Gutierrez MC, Renshaw L, Larionov AA, Faratian D, Hilsenbeck SG, Perou CM, Lewis MT, Rosen JM, Chang JC. Residual breast cancers after conventional therapy display mesenchymal as well as tumor-initiating features. *Proc Natl Acad Sci USA*. 2009; 106:13820–13825. doi:10.1073/pnas.0905718106. [PubMed: 19666588]
21. Creighton CJ, Chang JC, Rosen JM. Epithelial-mesenchymal transition (EMT) in tumor-initiating cells and its clinical implications in breast cancer. *J Mammary Gland Biol Neoplasia*. 2010; 15:253–260. doi:10.1007/s10911-010-9173-1. [PubMed: 20354771]
22. Ihaka R, Gentleman R. R: a language for data analysis and graphics. *J Comput Graph Stat*. 1996; 5:299–314.
23. Mani SA, Guo W, Liao MJ, Eaton EN, Ayyanan A, Zhou AY, Brooks M, Reinhard F, Zhang CC, Shipitsin M, Campbell LL, Polyak K, Briskin C, Yang J, Weinberg RA. The epithelial-mesenchymal transition generates cells with properties of stem cells. *Cell*. 2008; 133:704–715. doi:10.1016/j.cell.2008.03.027. [PubMed: 18485877]
24. Li H, Duhachek-Muggy S, Qi Y, Hong Y, Behbod F, Zolkiewska A. An essential role of metalloprotease-disintegrin ADAM12 in triple-negative breast cancer. *Breast Cancer Res Treat*. 2012; 135:759–769. doi:10.1007/s10549-012-2220-4. [PubMed: 22926263]

25. Li H, Solomon E, Duhachek Muggy S, Sun D, Zolkiewska A. Metalloprotease-disintegrin ADAM12 expression is regulated by Notch signaling via microRNA-29. *J Biol Chem*. 2011; 286:21500–21510. doi:10.1074/jbc.M110.207951. [PubMed: 21518768]
26. Edwards DR, Handsley MM, Pennington CJ. The ADAM metalloproteinases. *Mol Aspects Med*. 2008; 29:258–289. doi:10.1016/j.mam.2008.08.001. [PubMed: 18762209]
27. Wang Y, Klijn JG, Zhang Y, Sieuwerts AM, Look MP, Yang F, Talantov D, Timmermans M, Meijer-van Gelder ME, Yu J, Jatke T, Berns EM, Atkins D, Foekens JA. Gene-expression profiles to predict distant metastasis of lymph-node-negative primary breast cancer. *Lancet*. 2005; 365:671–679. doi:10.1016/S0140-6736(05)17947-1. [PubMed: 15721472]
28. Bos PD, Zhang XH, Nadal C, Shu W, Gomis RR, Nguyen DX, Minn AJ, van de Vijver MJ, Gerald WL, Foekens JA, Massague J. Genes that mediate breast cancer metastasis to the brain. *Nature*. 2009; 459:1005–1009. doi:10.1038/nature08021. [PubMed: 19421193]
29. Minn AJ, Gupta GP, Siegel PM, Bos PD, Shu W, Giri DD, Viale A, Olshen AB, Gerald WL, Massague J. Genes that mediate breast cancer metastasis to lung. *Nature*. 2005; 436:518–524. doi:10.1038/nature03799. [PubMed: 16049480]
30. Miller WR, Larionov AA, Renshaw L, Anderson TJ, White S, Murray J, Murray E, Hampton G, Walker JR, Ho S, Krause A, Evans DB, Dixon JM. Changes in breast cancer transcriptional profiles after treatment with the aromatase inhibitor, letrozole. *Pharmacogenet Genomics*. 2007; 17:813–826. doi:10.1097/FPC.0b013e32820b853a. [PubMed: 17885619]
31. Boersma BJ, Reimers M, Yi M, Ludwig JA, Luke BT, Stephens RM, Yfantis HG, Lee DH, Weinstein JN, Ambs S. A stromal gene signature associated with inflammatory breast cancer. *Int J Cancer*. 2008; 122:1324–1332. doi:10.1002/ijc.23237. [PubMed: 17999412]
32. Dontu G, Abdallah WM, Foley JM, Jackson KW, Clarke MF, Kawamura MJ, Wicha MS. In vitro propagation and transcriptional profiling of human mammary stem/progenitor cells. *Genes Dev*. 2003; 17:1253–1270. doi:10.1101/gad.106180317/10/1253. [PubMed: 12756227]
33. Taube JH, Herschkowitz JI, Komurov K, Zhou AY, Gupta S, Yang J, Hartwell K, Onder TT, Gupta PB, Evans KW, Hollier BG, Ram PT, Lander ES, Rosen JM, Weinberg RA, Mani SA. Core epithelial-to-mesenchymal transition interactome gene-expression signature is associated with claudin-low and metaplastic breast cancer subtypes. *Proc Natl Acad Sci USA*. 2010; 107:15449–15454. doi:10.1073/pnas.1004900107. [PubMed: 20713713]
34. Onder TT, Gupta PB, Mani SA, Yang J, Lander ES, Weinberg RA. Loss of E-cadherin promotes metastasis via multiple downstream transcriptional pathways. *Cancer Res*. 2008; 68:3645–3654. doi:10.1158/0008-5472.CAN-07-2938. [PubMed: 18483246]
35. Scheel C, Eaton EN, Li SH, Chaffer CL, Reinhardt F, Kah KJ, Bell G, Guo W, Rubin J, Richardson AL, Weinberg RA. Paracrine and autocrine signals induce and maintain mesenchymal and stem cell states in the breast. *Cell*. 2011; 145:926–940. doi:10.1016/j.cell.2011.04.029. [PubMed: 21663795]
36. Gupta PB, Onder TT, Jiang G, Tao K, Kuperwasser C, Weinberg RA, Lander ES. Identification of selective inhibitors of cancer stem cells by high-throughput screening. *Cell*. 2009; 138:645–659. doi:10.1016/j.cell.2009.06.034. [PubMed: 19682730]
37. Visvader JE, Lindeman GJ. Cancer stem cells: current status and evolving complexities. *Cell Stem Cell*. 2012; 10:717–728. doi:10.1016/j.stem.2012.05.007. [PubMed: 22704512]
38. Baccelli I, Trumpp A. The evolving concept of cancer and metastasis stem cells. *J Cell Biol*. 2012; 198:281–293. doi:10.1083/jcb.201202014. [PubMed: 22869594]
39. Charafe-Jauffret E, Ginestier C, Birnbaum D. Breast cancer stem cells: tools and models to rely on. *BMC Cancer*. 2009; 9:202. doi:10.1186/1471-2407-9-202. [PubMed: 19555472]
40. Velasco-Velazquez MA, Homsí N, De La Fuente M, Pestell RG. Breast cancer stem cells. *Int J Bioch Cell Biol*. 2012; 44(4):573–577. doi:10.1016/j.biocel.2011.12.020.
41. Li X, Lewis MT, Huang J, Gutierrez C, Osborne CK, Wu MF, Hilsenbeck SG, Pavlick A, Zhang X, Chamness GC, Wong H, Rosen J, Chang JC. Intrinsic resistance of tumorigenic breast cancer cells to chemotherapy. *J Nat Cancer Inst*. 2008; 100:672–679. doi:10.1093/jnci/djn123. [PubMed: 18445819]
42. Dave B, Mittal V, Tan NM, Chang JC. Epithelial-mesenchymal transition, cancer stem cells and treatment resistance. *Breast Cancer Res*. 2012; 14:202. doi:10.1186/bcr2938. [PubMed: 22264257]

43. Lacerda L, Pusztai L, Woodward WA. The role of tumor initiating cells in drug resistance of breast cancer: implications for future therapeutic approaches. *Drug Resist Updat*. 2010; 13:99–108. doi: 10.1016/j.drug.2010.08.001. [PubMed: 20739212]
44. Farmer P, Bonnefoi H, Anderle P, Cameron D, Wirapati P, Becette V, Andre S, Piccart M, Campone M, Brain E, Macgrogan G, Petit T, Jassem J, Bibeau F, Blot E, Bogaerts J, Aguet M, Bergh J, Iggo R, Delorenzi M. A stroma-related gene signature predicts resistance to neoadjuvant chemotherapy in breast cancer. *Nat Med*. 2009; 15:68–74. doi:10.1038/nm.1908. [PubMed: 19122658]
45. Ayers M, Symmans WF, Stec J, Damokosh AI, Clark E, Hess K, Lecoche M, Metivier J, Booser D, Ibrahim N, Valero V, Royce M, Arun B, Whitman G, Ross J, Sneige N, Hortobagyi GN, Pusztai L. Gene expression profiles predict complete pathologic response to neoadjuvant paclitaxel and fluorouracil, doxorubicin, and cyclophosphamide chemotherapy in breast cancer. *J Clin Oncol*. 2004; 22:2284–2293. doi:10.1200/JCO.2004.05.166. [PubMed: 15136595]
46. Popovici V, Chen W, Gallas BG, Hatzis C, Shi W, Samuelson FW, Nikolsky Y, Tsyganova M, Ishkin A, Nikolskaya T, Hess KR, Valero V, Booser D, Delorenzi M, Hortobagyi GN, Shi L, Symmans WF, Pusztai L. Effect of training-sample size and classification difficulty on the accuracy of genomic predictors. *Breast Cancer Res*. 2010; 12:R5. doi:10.1186/bcr2468. [PubMed: 20064235]
47. Valdez KE, Fan F, Smith W, Allred DC, Medina D, Behbod F. Human primary ductal carcinoma in situ (DCIS) subtype-specific pathology is preserved in a mouse intraductal (MIND) xenograft model. *J Pathol*. 2011; 225:565–573. doi:10.1002/path.2969. [PubMed: 22025213]
48. Dyczynska E, Syta E, Sun D, Zolkiewska A. Breast cancer-associated mutations in metalloprotease disintegrin ADAM12 interfere with the intracellular trafficking and processing of the protein. *Int J Cancer*. 2008; 122:2634–2640. doi:10.1002/ijc.23405. [PubMed: 18241035]
49. Roman-Perez E, Casbas-Hernandez P, Pirone JR, Rein J, Carey LA, Lubet RA, Mani SA, Amos KD, Troester MA. Gene expression in extratumoral microenvironment predicts clinical outcome in breast cancer patients. *Breast Cancer Res*. 2012; 14:R51. doi:10.1186/bcr3152. [PubMed: 22429463]
50. Haakensen VD, Lingjaerde OC, Luders T, Riis M, Prat A, Troester MA, Holmen MM, Frantzen JO, Romundstad L, Navjord D, Bukholm IK, Johannesen TB, Perou CM, Ursin G, Kristensen VN, Borresen-Dale AL, Helland A. Gene expression profiles of breast biopsies from healthy women identify a group with claudin-low features. *BMC Med Genomics*. 2011; 4:77. doi: 10.1186/1755-8794-4-77. [PubMed: 22044755]
51. Farmer P, Bonnefoi H, Becette V, Tubiana-Hulin M, Fumoleau P, Larsimont D, Macgrogan G, Bergh J, Cameron D, Goldstein D, Duss S, Nicoulaz AL, Brisken C, Fiche M, Delorenzi M, Iggo R. Identification of molecular apocrine breast tumours by microarray analysis. *Oncogene*. 2005; 24(29):4660–4671. doi:10.1038/sj.onc.1208561. [PubMed: 15897907]
52. Desmedt C, Haibe-Kains B, Wirapati P, Buyse M, Larsimont D, Bontempi G, Delorenzi M, Piccart M, Sotiriou C. Biological processes associated with breast cancer clinical outcome depend on the molecular subtypes. *Clin Cancer Res*. 2008; 14:5158–5165. doi: 10.1158/1078-0432.CCR-07-4756. [PubMed: 18698033]
53. Hollier BG, Evans K, Mani SA. The epithelial-to-mesenchymal transition and cancer stem cells: a coalition against cancer therapies. *J Mammary Gland Biol Neoplasia*. 2009; 14:29–43. doi: 10.1007/s10911-009-9110-3. [PubMed: 19242781]
54. Cheng GZ, Chan J, Wang Q, Zhang W, Sun CD, Wang LH. Twist transcriptionally up-regulates AKT2 in breast cancer cells leading to increased migration, invasion, and resistance to paclitaxel. *Cancer Res*. 2007; 67:1979–1987. doi:10.1158/0008-5472.CAN-06-1479. [PubMed: 17332325]
55. Kajita M, McClinic KN, Wade PA. Aberrant expression of the transcription factors snail and slug alters the response to genotoxic stress. *Mol Cell Biol*. 2004; 24:7559–7566. doi:10.1128/MCB.24.17.7559-7566.2004. [PubMed: 15314165]
56. Yu M, Bardia A, Wittner BS, Stott SL, Smas ME, Ting DT, Isakoff SJ, Ciciliano JC, Wells MN, Shah AM, Concannon KF, Donaldson MC, Sequist LV, Brachtel E, Sgroi D, Baselga J, Ramaswamy S, Toner M, Haber DA, Maheswaran S. Circulating breast tumor cells exhibit dynamic changes in epithelial and mesenchymal composition. *Science*. 2013; 339:580–584. doi: 10.1126/science.1228522. [PubMed: 23372014]

57. Martin M, Romero A, Cheang MC, Lopez Garcia, Asenjo JA, Garcia-Saenz JA, Oliva B, Roman JM, He X, Casado A, de la Torre J, Furio V, Puente J, Caldes T, Vidart JA, Lopez-Tarruella S, Diaz-Rubio E, Perou CM. Genomic predictors of response to doxorubicin versus docetaxel in primary breast cancer. *Breast Cancer Res Treat.* 2011; 128:127–136. doi:10.1007/s10549-011-1461-y. [PubMed: 21465170]
58. Gonzalez-Angulo AM, Iwamoto T, Liu S, Chen H, Do KA, Hortobagyi GN, Mills GB, Meric-Bernstam F, Symmans WF, Pusztai L. Gene expression, molecular class changes, and pathway analysis after neoadjuvant systemic therapy for breast cancer. *Clinical Cancer Res.* 2012; 18:1109–1119. doi:10.1158/1078-0432.CCR-11-2762. [PubMed: 22235097]
59. Symmans WF, Ayers M, Clark EA, Stec J, Hess KR, Sneige N, Buchholz TA, Krishnamurthy S, Ibrahim NK, Buzdar AU, Theriault RL, Rosales MF, Thomas ES, Gwyn KM, Green MC, Syed AR, Hortobagyi GN, Pusztai L. Total RNA yield and microarray gene expression profiles from fine-needle aspiration biopsy and core-needle biopsy samples of breast carcinoma. *Cancer.* 2003; 97:2960–2971. doi:10.1002/cncr.11435. [PubMed: 12784330]
60. Roy R, Moses MA. ADAM12 induces estrogen-independence in breast cancer cells. *Breast Cancer Res Treat.* 2012; 131(3):731–741. doi:10.1007/s10549-011-1431-4. [PubMed: 21387162]
61. Frohlich C, Nehammer C, Albrechtsen R, Kronqvist P, Kveiborg M, Sehara-Fujisawa A, Mercurio AM, Wewer UM. ADAM12 produced by tumor cells rather than stromal cells accelerates breast tumor progression. *Mol Cancer Res.* 2011; 9:1449–1461. doi:10.1158/1541-7786.MCR-11-0100. [PubMed: 21875931]
62. Moraes RC, Zhang X, Harrington N, Fung JY, Wu M-F, Hilsenbeck SG, Allred DC, Lewis MT. Constitutive activation of smoothened (SMO) in mammary glands of transgenic mice leads to increased proliferation, altered differentiation and ductal dysplasia. *Dev.* 2007; 134:1231–1242. doi:10.1242/dev.02797.
63. Dyczynska E, Sun D, Yi H, Sehara-Fujisawa A, Blobel CP, Zolkiewska A. Proteolytic processing of delta-like 1 by ADAM proteases. *J Biol Chem.* 2007; 282:436–444. doi:10.1074/jbc.M605451200. [PubMed: 17107962]
64. Atfi A, Dumont E, Colland F, Bonnier D, L'Helgoualc'h A, Prunier C, Ferrand N, Clement B, Wewer UM, Theret N. The disintegrin and metalloproteinase ADAM12 contributes to TGF-beta signaling through interaction with the type II receptor. *J Cell Biol.* 2007; 178:201–208. doi: 10.1083/jcb.200612046. [PubMed: 17620406]

Fig. 1.

Expression levels of catalytically active, transmembrane ADAMs vary across the intrinsic subtypes of breast cancer. Gene expression data for the UNC337 cohort profiled on Agilent microarrays were retrieved from GEO:GSE18229. Box-and-whisker plots of log₂-transformed, median-centered values are shown (whiskers: 5–95 percentile). Statistical significance of the differences between groups was determined using one-way ANOVA with Bonferroni post-test analysis. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$, **** $P < 0.0001$. *B* basal, *CL* claudin-low, *H* HER2-enriched, *LA* luminal A, *LB* luminal B, *NL* normal-like

Fig. 2.

ADAM12L, but not *ADAM12S*, is up-regulated in claudin-low breast tumors. Gene expression data for the EMC286, EMC192, and MSK82 cohorts profiled on Affymetrix microarrays were retrieved from GEO:GSE2034, GSE12276, and GSE2603, respectively. The intrinsic subtypes of breast cancer were adopted after [6]. Box-and-whisker plots of log₂-transformed, median-centered, and z-transformed values are shown (whiskers: 5–95 percentile). Statistical significance of the differences between groups was determined using one-way ANOVA with Bonferroni post-test analysis. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$, **** $P < 0.0001$. *B* basal, *CL* claudin-low, *H* HER2-enriched, *L* luminal, *NL* normal-like

Fig. 3.

ADAM12L, but not *ADAM12S*, is up-regulated after endocrine therapy. Gene expression data for patients with ER-positive breast tumors, before and after letrozole treatment, were retrieved from GEO:GSE10281 (a) and GSE5462 (b) datasets, respectively. Values were log2-transformed and centered on pre-treatment expression levels. Statistical significance of the differences in pre-treatment and post-treatment *ADAM12L* and *ADAM12S* levels were analyzed by paired *t* test. Red and blue lines indicate the mean expression values before and after treatment, respectively

Fig. 4.

ADAM12L, but not *ADAM12S*, is associated with stromal, mammosphere, and EMT gene signatures. **a–f** *ADAM12L* and *ADAM12S* gene expression data were retrieved from GEO using the indicated GSE accession numbers. Log2-transformed, median-centered values are shown. **a** *ADAM12L* and *ADAM12S* expression in paired samples of breast tumor stroma and epithelium isolated using laser capture microdissection. *P* values for paired *t* test are shown. Red and blue lines indicate the mean expression values of *ADAM12L* and *ADAM12S* in the stroma and the epithelium, respectively. **b** *ADAM12L* and *ADAM12S* expression in CD44⁺/CD24^{-low} sorted populations of primary breast cancer cells versus the remaining "other" flow-sorted fractions. **c** *ADAM12L* and *ADAM12S* expression in primary breast cancer cells grown as spheres versus bulk tumors. In **b** and **c**, box-and-whisker plots are shown, and *P* values were calculated by unpaired *t* test. In **a–c**, the numbers (*n*) of primary tumors are indicated. **d** Fold change in *ADAM12L* and *ADAM12S* expression levels in HMLE cells undergoing EMT. *P* value was calculated by one-way ANOVA. **e** *ADAM12L* and *ADAM12S* expression in mesenchymal subpopulations (MSPs) and in pure epithelial fractions (CD24^{hi}) of HMLE cells. **f** *ADAM12L* and *ADAM12S* expression in transformed HMLER cells treated with 10 nM paclitaxel or 1 μ M salinomycin for 1 week. In **e** and **f**, *P* values for unpaired *t* test are shown.

Fig. 5.

Pre-treatment expression levels of *ADAM12L*, but not *ADAM12S*, in ER-negative breast tumors are higher in patients with poor response to neoadjuvant chemotherapy. Gene expression data for patients receiving anthracycline-based chemotherapy (**a**) or anthracycline plus taxane-based chemotherapy (**b**) were retrieved from GEO:GSE4779 and GSE20194, respectively. Log2-transformed, median-centered values are shown. **a** pCR, pathologic complete response, npCR, near pCR. **b** pCR versus RD (residual disease)

Fig. 6.

Validation of the up-regulation of *ADAM12L* mRNA during EMT and in non-adherent spheres by qRT-PCR. **a** HMLE cells were incubated for 0, 12, and 20 days with TGF β . **b** HMLE-Twist-ER cells were incubated for 0, 12, and 20 days with 4-hydroxy-tamoxifen (4-OH-Tam) to induce Twist expression. *Insets* in **a** and **b** show decrease in E-cadherin expression after 12 days of treatment, a hallmark of EMT. **c** MCF10DCIS.com cells were cultured as attached monolayers or seeded at 100 cells/well in ultralow attachment 96-well plates and incubated for 10 days in serum-free medium with EGF and bFGF. *ADAM12L* mRNA was measured by qRT-PCR and normalized to β -ACTIN; the values at day 0 (**a**, **b**) or in monolayers (**c**) were set as one. Statistical significance of *ADAM12L* up-regulation was evaluated with one-sample *t* test. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$

Fig. 7.

Down-regulation of *ADAM12* in [MCF10DCIS.com](#) cells inhibits sphere formation. [MCF10DCIS.com](#) cells stably transduced with shADAM12 or shControl vectors were seeded at 1 cell/well in ultralow attachment 96-well plates and incubated for 10 days in serum-free medium with EGF and bFGF. **a** Phase-contrast image of a typical sphere at day 10. **b** Sphere formation efficiency was determined in three independent experiments, shown in *red*, *green*, and *blue*, respectively. *P* value was determined using paired *t* test. **c** The endogenous *ADAM12L* mRNA levels in cells grown as monolayers or spheres were determined by qRT-PCR and normalized to β -ACTIN; the values in shControl cells were set as one. Statistical significance of *ADAM12L* up-regulation was evaluated with one-sample *t* test. **P* < 0.05, *ns* non-significant

Fig. 8.

The effect of ADAM12L overexpression in [MCF10DCIS.com](#) cells on sphere formation. [MCF10DCIS.com](#) cells stably transduced with the wild-type ADAM12L (WT), a dominant-negative G479E mutant ADAM12L [48], or empty vector (EV) were seeded to form spheres at 1 cell/well, as in Fig. 6. **a** Sphere formation efficiency was determined in six experiments, each shown in *different colors*. *P* value was determined using paired *t* test; *ns* non-significant. **b** ADAM12L protein level in monolayers (M) and in spheres (S) was determined by Western blotting; β -actin is a gel-loading control. *Arrowhead* indicates the nascent full-length ADAM12L, *arrow* shows the mature processed form of the wild-type ADAM12L. **c** Quantification of the band intensities from panel **b**. *P* value was determined using unpaired *t* test

Table 1

Patient characteristics

Dataset name	UNC337	EMC286	EMC192	MSK82	GSE10281	GSE5462	GSE5847	GSE7513	GSE7515	EORTC 10994	MAQCII/MDACC
GEO Accession Number	GSE18229	GSE2034	GSE12276	GSE2603	GSE10281	GSE5462	GSE5847	GSE7513	GSE7515	GSE4779	GSE20194
Number of patients	337	286	192	82	18	58	48	15	15	102	229
Age (years)											
Median	55	52	50	54.5	Unknown	Unknown	57	Unknown	Unknown	Unknown	51
Range	24–89	26–83	24–82	30–86	Unknown	Unknown	Unknown	Unknown	Unknown	Unknown	26–79
T stage											
T0/1	62	146	50	12	0	0	4	0	0	2	24
T2	137	131	107	58	0	0	30	0	0	63	131
T3/4	71	8	26	12	0	0	14	0	0	34	74
Unknown	67	1	9	0	18	58	0	15	15	3	0
Tumor grade											
Grade 1	25	44	37	21	0	0	0	0	0	0	13
Grade 2	82	5	2	0	0	0	0	0	0	0	93
Grade 3	147	19	13	5	0	0	0	0	0	0	123
Grade 4	0	92	54	15	0	0	0	0	0	0	0
Unknown	83	126	86	41	18	58	48	15	15	102	0
Lymph node status											
Positive	138	0	144	54	0	0	0	0	0	62	162
Negative	134	286	48	28	0	0	0	0	0	37	67
Unknown	65	0	0	0	18	58	48	15	15	3	0
ER status											
Positive	153	209	116	46	18	58	21	0	0	37	139
Negative	113	77	73	36	0	0	26	0	0	65	90
Unknown	71	0	3	0	0	0	1	15	15	0	0
PR status											
Positive	108	166	92	35	0	0	0	0	0	25	104
Negative	138	107	90	47	0	0	0	0	0	75	125
Unknown	91	13	10	0	18	58	48	15	15	2	0
HER2 status											

Dataset name	UNC337	EMC286	EMC192	MSK82	GSE10281	GSE5462	GSE5847	GSE7513	GSE7515	EORTC 10994	MAQCII/MDACC
Positive	60	46	132	17	0	0	5	0	0	0	40
Negative	214	207	38	59	0	0	38	0	0	0	189
Unknown	63	33	22	6	18	58	5	15	15	102	0
Endocrine therapy	Unknown	0	36	10	18	58	0	Unknown	Unknown	0	0
Chemotherapy	Unknown	0	46	25	0	0	19	Unknown	Unknown	102	248
Reference	18	27	28	29	20	30	31	20	20	44	45,46

Table 2

Predictive factors of pathologic complete response in ER-negative tumors

Variable	<u>Univariate logistic regression</u>		<u>Multivariate logistic regression</u>	
	OR (95 % CI)	P value	OR (95 % CI)	P value
EORTC 10994 (GSE4779)				
Node (N1–N2 vs N0)	0.77 (0.27–2.18)	0.6210	0.78 (0.25–2.45)	0.6727
T stage (T3 vs T1–T2)	0.88 (0.31–2.45)	0.8000	0.76 (0.24–2.34)	0.6382
<i>ADAM12L</i>	0.36 (0.16–0.72)	0.0065	0.37 (0.16–0.77)	0.0129
<i>ADAM12S</i>	3.78 (0.45–36.60)	0.2320	3.54 (0.36–39.73)	0.2858
MAQCII/MDACC (GSE20194)				
Age (>median vs median)	0.51 (0.22–1.18)	0.1210	0.49 (0.19–1.20)	0.1209
Node (N1–N3 vs N0)	2.72 (0.93–9.16)	0.0804	3.38 (1.04–13.22)	0.0560
T stage (T3–T4 vs T1–T2)	0.52 (0.14–1.78)	0.3040	0.35 (0.08–1.37)	0.1413
Grade (3 vs 2)	2.25 (0.69–8.78)	0.2010	2.26 (0.62–9.76)	0.2375
<i>ADAM12L</i>	0.75 (0.55–0.99)	0.0497	0.76 (0.54–1.04)	0.0936
<i>ADAM12S</i>	0.83 (0.55–1.24)	0.3670	0.97 (0.62–1.51)	0.8961

OR odds ratio for having pathologic complete response versus residual disease, CI confidence interval