Published in final edited form as:

Womens Health Issues. 2013 ; 23(2): . doi:10.1016/j.whi.2012.12.005.

Human Papillomavirus Vaccine Coverage Among Females Aged 11 to 17 in Texas Counties: An Application of Multilevel, Small Area Estimation

Jan M. Eberth, PhD^{a,*}, Md Monir Hossain, PhD^b, Jasmin A. Tiro, PhD^c, Xingyou Zhang, PhD^d, James B. Holt, PhD^d, and Sally W. Vernon, PhD^e

^aDivision of Quantitative Sciences, Department of Biostatistics, MD Anderson Cancer Center, Houston, Texas

^bDivision of Epidemiology and Biostatistics, Cincinnati Children's Hospital Medical Center, Cincinnati, Ohio

^cDepartment of Clinical Sciences, University of Texas Southwestern Medical Center, Dallas, Texas

^dDivision of Adult & Community Health, Centers for Disease Control & Prevention, Atlanta, Georgia

^eDivision of Health Promotion & Behavioral Sciences, University of Texas School of Public Health, Houston, Texas

Abstract

Background—Local data are often used to plan and evaluate public health interventions and policy. With increasingly fewer public resources to collect sufficient data to support direct estimation of local outcomes, methods for deriving small area estimates are vital. The purpose of this study is to describe the county-level geographic distribution of human papillomavirus (HPV) vaccine coverage among adolescent females in Texas using multilevel small area estimation.

Methods—Multilevel (individual, county, public health region) random-intercept logit models were fit to HPV vaccination data (1 dose Gardasil) from the 2008 Behavioral Risk Factor Surveillance System. Using the parameter estimates from the final model, we simulated 10,000 data sets for each regression coefficient from the normal distribution and applied them to the logit model to estimate HPV vaccine coverage in each county.

Results—County-level coverage estimates ranged from 7% to 29%, compared with the state average of 18% (95% confidence interval [CI], 13.59–21.88). Many Southwestern border and metropolitan counties exhibited high coverage estimates. Low coverage estimates were noted in the Panhandle, Southeastern border region, and Northeast. Significant correlations were observed between HPV vaccination and Hispanic ethnicity, county poverty, and public health region poverty.

Conclusion—Harnessing the flexibility of multilevel small area models to estimate HPV vaccine coverage at the county level, we have provided data that may inform the development of health education programs/policies, the provision of health services, and the planning of new

research studies. Additionally, we have provided a framework for modeling other health outcomes at the county level using national survey data.

Introduction

The U.S. Food and Drug Administration approved the use of Gardasil (HPV4), a vaccine against four prevalent strains of human papillomavirus (HPV) for females aged 9 to 26 in June 2006 (U.S. Food and Drug Administration, 2006). Despite reports of high acceptability and willingness to vaccinate against HPV (Zimet, Liddon, Rosenthal, Lazcano-Ponce, & Allen, 2006), early research showed that uptake was suboptimal (Gottlieb et al., 2009; Jain, Stokley, & Yankey, 2008; Kahn et al., 2008; Rosenthal et al., 2008). Initiation of the HPV4 series (3 doses given over 6 months) grew over time, increasing to 48.7% of U.S. females aged 13 to 17 in 2010 (32% completed the vaccine series); however, initiation rates varied considerably by geographic region (range across 50 states, 28.8%–73.0%; Dorell, Stokley, Yankey, Liang, & Markowitz, 2011). In an analysis of 2008 Behavioral Risk Factor Surveillance System (BRFSS) data, Pruitt and Schootman (2010) found that HPV vaccine coverage differed significantly across states and counties, a finding explained in part by state- and county-level poverty. Although studies indicate the presence of geographic variation in HPV vaccine coverage across states and selected regions, and have found that group-level covariates such as poverty (Pruitt & Schootman, 2010) and education (Chao, Velicer, Slezak, & Jacobsen, 2010) are associated with HPV vaccination, local level estimates are not uniformly available in the literature.

Local data on HPV vaccine coverage are important for several reasons, including the planning of health promotion programs, identifying target populations for services (e.g., free vaccine clinics), and evaluating the impact of policies and programs designed to improve vaccine uptake. National health surveys provide much of the data used to conduct population-based public health research in the United States. Whereas these surveys provide estimates at the national, state, and sometimes county levels, their ability to provide local estimates is limited owing to costs, sample size constraints (i.e., 50 respondents per geographic unit recommended for direct estimation; Siegel, Frazier, Mariolis, Brackbill, & Smith, 1993), and privacy protections. Thus, researchers often rely on small area estimation (SAE), a set of indirect estimation techniques for producing small area estimates using data collected for larger areas and census tabulations for the small area of interest. This study used multilevel, SAE to describe the county-level geographic distribution of HPV vaccine coverage (1 dose HPV4) among females aged 11 to 17 years living in Texas. The methodological framework we discuss herein can be used by other researchers to analyze their own health outcome(s) of interest using a variety of national and state-based population surveys.

Analysis

Overview

In multilevel small area models (a type of synthetic estimation), model coefficients are linked with demographic- and geographic-specific population data (hereafter referred to as the auxiliary dataset) to predict the outcome for each small area or population subgroup. We applied multilevel SAE to predict county-level coverage of HPV vaccination. The 2008 BRFSS was used to develop a three-level (individual, county, public health region [PHR]), random-intercept logit model. Several factors informed our choice of covariates and model-building strategy. First, covariates included in the final model (both individual and group levels) were required to be available in both the individual and auxiliary datasets to permit estimation of small area means. Second, all individual- and group-level covariates considered for inclusion had known (Chao, Velicer, Slezak, & Jacobsen, 2009; Chao et al.,

2010; Dempsey, Cohn, Dalton, & Ruffin, 2010; Dorell, Stokley, Yankey, Cohn, & Markowitz, 2010; Gottlieb et al., 2009; Neubrand, Breitkopf, Rupp, Breitkopf, & Rosenthal, 2009; Pruitt & Schootman, 2010; Rondy, van Lier, van de Kassteele, Rust, & de Melker, 2010) or hypothesized associations with HPV vaccination in adolescents. Finally, because Texas counties are nested within PHRs and vaccine coverage data in Texas is often presented for PHRs, we included PHR-level random effects in the model to ensure the correct selection of group-level covariates and parameter estimates. A summary of these methods is shown in Figure 1.

SAS Version 9.1 (SAS Institute, Cary, NC) was used for data management and MLwiN Version 2.20 was used to implement multilevel models. Our study was granted exempt status by our Institutional Review Board.

Data Collection and Management

We used the 2008 Texas BRFSS (U.S. Centers for Disease Control and Prevention, 2008b) to obtain the dependent variable and individual-level covariates of interest. County- and PHR-level data were acquired from multiple sources including the Association of Religious Data Archives, U.S. Department of Agriculture Office of Management and Budget, Texas Cancer Registry, Texas Department of Health and Human Services, and U.S. Census Small Area Health Insurance Estimates Program. County-specific population estimates, stratified by gender, age, and race/ethnicity (the auxiliary dataset), were obtained from the Vintage 2008 Bridged-Race Postcensal Population Estimates file available from the Health Center for Health Statistics (Centers for Disease Control and Prevention, 2009).

The BRFSS is a population-based, random digit-dial survey conducted annually by the states, through a cooperative agreement with the U.S. Centers for Disease Control and Prevention. In addition to the core questions asked by all states in 2008, Texas adopted the optional HPV modules. In Texas, the proportion of contacted persons who completed the interview was 75% (Centers for Disease Control and Prevention, 2008a). Information on the survey design and sampling strategy are available elsewhere (Centers for Disease Control and Prevention, 2008a, 2008b).

When children resided in the home of an adult respondent (referred to as the parent/caregiver), he or she answered questions about one randomly selected child. Our sample consisted of 574 parents/caregivers, all of whom 1) had a county of residence listed in the public use dataset, 2) had a randomly selected 11- to 17-year-old daughter for whom survey data were collected, and 3) reported their level of educational attainment. The daughter's age and race/ethnicity, as well as the parent/caregiver's educational attainment, are known predictors of HPV vaccine uptake (Chao et al., 2009, 2010; Conroy et al., 2009; Dempsey et al., 2010; Dorell et al., 2010; Gottlieb et al., 2009; Neubrand et al., 2009; Pruitt & Schootman, 2010; Wong et al., 2011); thus, any person with missing data on these items was excluded. Additionally, participants who did not know their daughter's HPV vaccination status were excluded. These criteria reduced the sample size from 623 to 574.

In selecting group-level covariates for examination, we made an a priori decision to include county-level poverty because of its known positive association with HPV vaccine initiation (Pruitt & Schootman, 2010). Because the majority of covariates of interest had not been previously studied, we implemented an exploratory approach to select additional covariates for examination. We were specifically interested in testing the effects of access, religious adherence, sexually transmitted disease and cervical cancer prevalence, and uptake of other childhood and adolescent vaccines. When there were multiple measures representing a specific concept (e.g., access to care could be measured by density of physicians or *Vaccines For Children* providers in the county, urban/rural classifications, or designation as a Health

Professional Shortage Area), we conducted principal components analysis to extract one representative covariate for each concept. The covariate with the strongest correlation to the first principal component was then examined in our models. Principal components analysis was not done for PHR-level covariates because of limited variability ($n = 11$). This approach resulted in examination of the group-level covariates listed in Table 1.

Although sampling weights are provided by the BRFSS, they are not efficient for use in small area models because the weights are constructed at the state level. The problem is compounded for multilevel models, which traditionally have not made use of weights associated with complex survey data (Carle, 2009). As suggested by Carle (2009) and Rabe-Hesketh and Skrondal (2006), we rescaled the level 1 sampling weights (i.e., CHILDDWT) to produce better point estimates and reduce the risk of biased estimates when the cluster size is small. We also included weights at levels 2 and 3 as proposed by Goldstein (2010). Technical documentation for the scaling and creation of these weights is provided in Appendix A.

Statistical Analysis

The outcome was initiation of HPV vaccination (1 dose HPV4, yes or no) among females aged 11 to 17. A series of increasingly complex models were fit to the data, starting with the empty model and adding covariates sequentially by level (Table 1). Level 1 covariates were added to the empty model in step 2 and were retained for inclusion regardless of statistical significance in steps 3 through 5. At steps 3 and 4, group-level covariates were added to the model simultaneously. Covariates were removed from the model sequentially, starting with the least significant, until only covariates with $p < .1$ remained. Using a more conservative 0.25 p -value, as suggested by Hosmer and Lemeshow (2000), resulted in the same selection of covariates. Because auxiliary data were only stratified by age and race/ethnicity, parental education was removed from the final model (step 5). A post-estimation adjustment for education at the county-level was incorporated as suggested by Congdon (2009; equation in Appendix A). First-order marginal quasi-likelihood, followed by second-order penalized quasi-likelihood (uses marginal quasi-likelihood estimates as starting values; provides robust estimates when sample sizes are small), was used in all models. Interaction terms were not tested owing to the small sample size.

We applied the parameter estimates from the final model to simulate 10,000 probabilities of HPV vaccine uptake for each race- and age-specific subgroup each county. The probabilities were adjusted for the educational mix of each county. We then calculated the mean value, standard deviation, and 95% confidence interval (CI) of these simulated subgroup probabilities. The variance associated with the PHR-level random intercept was very small and thus ignored in the simulation. The county random intercept was only estimated for sampled counties in the 2008 Texas BRFSS. Finally, we multiplied the race- and age-specific adjusted probabilities above by the corresponding county subpopulation to generate summary estimates of HPV vaccine coverage for each county.

To test the accuracy of our multilevel model, we aggregated our county-level estimates up the state level and compared the resulting model-based estimate with the direct estimate of HPV vaccine coverage at the state level. We did the same validation checks with our age- and race-specific county estimates. These analyses allow us to examine whether our small area estimates (aggregated up to the state level) produce the expected results at the state level, where direct estimation of HPV vaccine coverage is possible.

Results

Sample Characteristics

Our sample consisted of 574 parents/caregivers residing in 99 Texas counties (39% of counties). The adolescent females were primarily White (54%) or Hispanic (33%), with educated parents/caregivers (34% had a bachelor's degree or higher and 27% had some college education), and a mean age of 14 years. The large majority of respondents (88%) indicated they were parents (biologic, step, or adoptive) of the adolescent female. Many of the participants came from large metropolitan counties including Bexar ($n = 55$), El Paso ($n = 44$), Harris ($n = 40$), Lubbock ($n = 31$), Tarrant ($n = 31$), Travis ($n = 36$), and Val Verde ($n = 30$), which is not surprising given their greater probability of being selected as a survey participant.

Model Results

In all models, there was small variability at the county- and PHR-level (Table 2). At the county level, we evaluated the area level variance using the median odds ratio (MOR) (not able to be estimated at the PHR-level). Because the MOR was not far from 1.00 in the final model, we concluded that there were minimal differences between counties in the individual probability of obtaining HPV vaccination.

Hispanic ethnicity was the only significant level 1 covariate. Compared with Whites, Hispanic females aged 11 to 17 were significantly more likely to be vaccinated against HPV (odds ratio [OR], 1.77; 95% CI, 1.28–2.46). The removal of parental education from the final model impacted the strength and significance of the covariate (OR, 1.34; 95% CI, 0.95–1.91), suggesting a possible suppression effect. We suspect the likely reason for this effect is educational distribution across racial/ethnic groups. In our sample, Whites, Blacks, and other races had positively skewed distributions (i.e., toward more education), whereas Hispanics had a negative skew. In contrast with several studies finding a positive effect of age on HPV vaccine uptake (Chao et al., 2010; Conroy et al., 2009; Dempsey et al., 2010; Gottlieb et al., 2009; Pruitt & Schootman, 2010), we found a null effect that likely reflects how we defined the variable (i.e., continuous vs. categorical).

We found a significant, positive association between county-level poverty and HPV vaccination. The odds of being vaccinated increased by about 9% for every 1% increase in county-level poverty. A more meaningful interpretation, based on the distribution of poverty in Texas (range across counties, 6.2%–36.8%), would be the change in odds for every 5% or 10% change in county-level poverty. For these values, we would expect to see the odds of vaccination increase by 54% and 139%, respectively. Our model also showed a significant, negative association between PHR-level poverty and HPV vaccination. Thus, for a 5% or 10% change in PHR-level poverty (vs. a 1% change, as shown in Table 2), the odds of vaccination decreased by 35% and 58%, respectively.

Small Area Estimates

In 2008, 17.7% (95% CI, 13.59–21.88) of Texas females aged 11 to 17 were vaccinated against HPV (based on direct, weighted estimation of HPV vaccine initiation data from the 2008 Texas BRFSS). That same year, the national average for females aged 11 to 17 was 30.4%. County-level model-based small area estimates ranged from 6.8% to 29.0% (Figure 2; Appendix B). No estimate is provided for Loving county because there were no representative females. The state-level indirect estimate was 15.5% (95% CI, 14.1–16.9), which lies within the confidence interval of its direct estimate of HPV vaccine coverage (95% CI, 13.59–21.88). Our race/ethnicity- and age-specific indirect estimates, scaled up to the state level, were also within the confidence intervals of their respective direct estimates,

with the exception of the indirect estimate for HPV vaccine coverage among 13-year-old girls in Texas (Table 3).

Discussion

Researchers have used model-based SAE to examine a variety of health behaviors and outcomes including asthma, obesity, smoking, mammography utilization, and health insurance coverage (Congdon, 2010; Goodman, 2010; Li, Land, Zi, Keithly, & Kelsey, 2009; Mendez-Luck, Yu, Meng, Jhawar, & Wallace, 2007; Schneider, Lapane, Clark, & Rakowski, 2009; Yu, Meng, Mendez-Luck, Jhawar, & Wallace, 2007). Using BRFSS data, Jia, Muennig, and Borawski (2004) found that regression models provided better precision than traditional synthetic and spatial smoothing methods in SAE. Multilevel modeling is a particularly useful SAE approach because it simultaneously examines individual and place-based factors, and borrows strength from the global model to make small area inferences. The increasing availability of software products that can be used to perform multilevel SAE using weighted survey data (e.g., MLwiN, STATA's GLAMM, and Mplus), makes it possible for researchers without formal training in statistics to conduct small area research.

Examining health behaviors and outcomes at the local level facilitates policy making, program development and evaluation, and hypothesis generation. At the policy end, legislators need information about their jurisdictions to make informed choices about bills related to health education programs and mandates. For example, Mendez-Luck and co-workers (2007) provided asthma prevalence data at the Assembly district level in California to assist in advocacy and policy-making activities. With data on disease rates over space and time, healthcare administrators and researchers can also develop and evaluate health promotion and outreach programs, targeted and tailored to the communities with the greatest needs. Finally, SAE provides a foundation for hypothesis generation. Knowing which areas are most affected by a particular disease, or conversely, which have the best health behaviors, can lead to formulating and testing hypotheses about the causal mechanisms involved.

SAE is ideal for studying HPV vaccination for several reasons, including the following: 1) The Texas ImmTrac Vaccine Registry is a voluntary "opt-in" program, which limits the generalizability of the data, and 2) the data are not actively collected by schools or public health departments since it is not required for entry into Texas public schools. Our findings indicate wide variability in HPV vaccine coverage across counties (7%–29%). Many of the Southwest border counties exhibited high coverage rates, as did many large, metropolitan counties (e.g., Dallas, Brazos). Low coverage rates were common in the Panhandle (i.e., northern-most counties), Southeast border region, and the Northeast. Our multilevel model also showed some counterintuitive findings. Although county-level poverty was associated with increased odds of vaccination, PHR-level poverty was associated with decreased odds. Our findings paralleled those reported by Pruitt and Schootman (2010), in which HPV vaccination was positively associated with county-level poverty and negatively associated with state-level poverty. This suggests that county-level poverty may be more indicative of better financed public health programs and/or is strongly related to the racial/ethnic makeup of the county. However, not all counties were represented in our sample, which may have resulted in a biased measure of association for county-level poverty. The ability to interpret our findings on PHR-level poverty is limited by two issues: 1) PHR-level poverty represents a greater scale of geographic aggregation (i.e., sets of geographically contiguous counties, typically anchored by at least one metropolitan area), resulting in limited data variability and generalizability, and 2) the sample of respondents within each PHR varied widely, with some PHRs having respondents drawn from multiple counties and others having respondents from only one or two counties.

SAE has some inherent limitations. First, not all covariates of interest can be included. At level 1, only those covariates that are present in both the individual-level and auxiliary dataset can be included. It is possible, however, to make post-model adjustments (as we did for education) at the expense of model simplicity. Second, having no or few respondents at higher levels can limit estimation of random effects. In our study, we handled this by assigning the mean of simulated random effects to all non-sampled counties. Alternatively, one can ignore the random effects altogether (as we did for PHRs). Finally, there is often no gold standard for comparison of small area estimates. Without a reference, one cannot easily assess the accuracy of small area estimates. Several methods have been proposed to deal with this issue, including cross-validation using part of the sample, a different dataset, or data from another year (as long as the outcome is stable over time), as well as aggregation of one's estimates to a higher geographic level, where direct estimation and thus comparison and calibration (Yu et al., 2007) are possible.

Two lines of future research are proposed: Examination of how alternate modeling strategies might affect our estimates, and estimation of HPV vaccine coverage in other population groups. Alternative modeling strategies include 1) use of different sampling weights, 2) specifying other hierarchical structures, 3) removing random effects, 4) adding a spatial component to incorporate neighborhood information, and 5) evaluating the potential of other software programs (e.g., STATA's GLAMM procedure) to perform similar analyses with greater ease. Estimation of vaccine coverage among other population groups, such as males aged 9 to 26 and females aged 18 to 26, is also warranted.

Acknowledgments

This work was supported by National Cancer Institute preand post-doctoral fellowships (R25-CA057712, R25-CA57730) and a National Institutes of Health Cancer Center Support Grant (CA016672). The authors wish to acknowledge the excellent editorial contributions of Dr. Sandi Pruitt (UT Southwestern), Dr. Graham Moon (University of Southampton), and the Doctoral Behavioral Science Seminar at the University of Texas, School of Public Health at Houston. We also thank Dr. Christina Dorell, from the Centers for Disease Control & Prevention, whose suggestions were integral to the planning of this study. The findings and conclusions in this report are those of the authors and do not necessarily represent the official position of the National Cancer Institute, the National Institutes of Health, or the Centers for Disease Control and Prevention. Gardasil is a registered trademark of Merck & Co., Inc.

Appendix A: Technical Documentation of Methods

[Equation 1: General]

$$\begin{aligned} \text{logit}(p_{ijk}) = & \gamma_{0j} + \gamma_{0k} + \beta_1 \text{Level 1 Variables}_{ijk} \\ & + \beta_{2,jk} \text{Level 2 Variables}_{jk} \\ & + \beta_{3,k} \text{Level 3 Variables}_k \end{aligned}$$

[Equation 1 – Applied Preliminary Model]

$$\begin{aligned} \text{logit}(p_{ijk}) = & \beta_{0j} + \beta_{0k} \\ & + \beta_1 \text{Age}_{ijk} + \beta_2 <\text{HS}_{ijk} + \beta_3 \text{HS graduate}_{ijk} \\ & + \beta_4 \text{Some college}_{ijk} + \beta_5 \text{Bachelors plus}_{ijk} \\ & + \beta_6 \text{Whit}_{ijk} + \beta_7 \text{Black}_{ijk} + \beta_8 \text{Hispanic}_{ijk} \\ & + \beta_9 \text{Other races}_{ijk} + \beta_{10,jk} \text{Country Poverty}_{jk} \\ & + \beta_{11,k} \text{PHR Poverty}_k \end{aligned}$$

where:

$\beta_{0j} = \beta_0 + \mu_j$ (country – level random – intercept).
 $\beta_{0k} = \beta_0 + \mu_k$ (PHR – level random – intercept).
 β_1 = Regression coefficient for age.
 β_1 = Regression coefficient for < High school education (reference)
 β_3 = Regression coefficient for HS graduate.
 β_4 = Regression coefficient for some college.
 β_5 = Regression coefficient for Bachelors plus.
 β_6 = Regression coefficient for White (reference).
 β_7 = Regression coefficient for Black.
 β_8 = Regression coefficient for Hispanic.
 β_9 = Regression coefficient for other races.
 β_{10} = Regression coefficient for country poverty.
 β_{11} = Regression coefficient for PHR poverty.
 i (subjects) = 1, ..., m_{jk}
 j (county) = 1, ..., n_k
 k (PHR) = 1, ..., 11.
 μ = area level variance.

[Equation 2: Applied final model]

$$\begin{aligned}
 \text{logit}(p_{ijk}) = & \beta_{0j} + \beta_{0k} \\
 & + \beta_1 \text{Age}_{ijk} + \beta_6 \text{White}_{ijk} + \beta_7 \text{Black}_{ijk} \\
 & + \beta_8 \text{Hispanic}_{ijk} + \beta_9 \text{Other races}_{ijk} \\
 & + \beta_{10,jk} \text{Country Poverty}_{ijk} \\
 & + \beta_{11,k} \text{PHR Poverty}_k
 \end{aligned}$$

where:

$\beta_{0j} = \beta_0 + \mu_j$ (county – level random – intercept)
 $\beta_{0k} = \beta_0 + \mu_k$ (PHR – level random – intercept)
 * μ_k will be set to 0 in the simulations and calculation of small area means.
 $\beta_{0j} = \beta_0 + \mu_j$ (county – level random – intercept)
 $\beta_{0k} = \beta_0 + \mu_k$ (PHR – level random – intercept)
 β_1 = Regression coefficient for age.
 β_6 = Regression coefficient for White (reference)
 β_7 = Regression coefficient for Black.
 β_8 = Regression coefficient for Hispanic.
 β_9 = Regression coefficient for other races.
 β_{10} = Regression coefficient for county poverty.

β_{11} = Regression coefficient for PHR poverty.

i (subjects) $1, \dots, m_{jk}$

j (county) $= 1, \dots, n_k$

k (PHR) $= 1, \dots, 11$.

μ = area level variance.

[Equation 3: Weighted regression coefficients for education]

$$L_j = \sum \pi|e_j \times [\exp(\beta_x \text{Education}_{ijk}) / 1 + \exp(\beta_x \text{Education}_{ijk})]$$

where:

$x = 1 - 4$.

i (subjects) $1, \dots, m_{jk}$

j (county) $= 1, \dots, n_k$

k (PHR) $= 1, \dots, 11$.

$\pi|e$ = the proportion of persons in county j with educational attainment e

[Equation 4: Simulated mean random effect (level 2 only)]

$$\begin{aligned} x_{ij} &\sim \text{normal}(\mu_i, \sigma_i^2) \text{ (parameters from MLwiN)} \\ \bar{x}_i &= \frac{1}{10,000} \sum x_{ij} \text{ (for sampled countries)} \\ \bar{x} &= \frac{1}{99} \sum \bar{x}_i \text{ (for non-sampled countries)} \end{aligned}$$

where:

$i = 1, 2, \dots, 99$ (number of sampled counties, and thus estimated random effects).

$j = 1, 2, \dots, 10,000$ (number of simulations).

[Equation 5: Simulated means and 95% confidence intervals, adjusted for county education]

Step 1: Simulations

$$\begin{aligned} \beta_0 &\sim \text{normal}(-1.839, 0.835)^* \\ \beta_1 &\sim \text{normal}(0.025, 0.047) \\ \beta_7 &\sim \text{normal}(-0.075, 0.556) \\ \beta_8 &\sim \text{normal}(0.299, 0.177) \\ \beta_9 &\sim \text{normal}(0.254, 0.539) \\ \beta_{10} &\sim \text{normal}(0.087, 0.034) \\ \beta_{11} &\sim \text{normal}(-0.085, 0.032) \end{aligned}$$

$$\begin{aligned} x_{ij} &\sim \text{normal}(\mu_i, \sigma_i^2) \\ r_{ij} &\sim \text{normal}(\bar{x}, \sigma_i^2) \end{aligned}$$

* Values taken from the model (i.e., regression coefficients and standard errors).

Step 2: Calculate adjusted means

Race- and age-specific county means:

$$\bar{p}_{RAj} = \frac{1}{n} \sum \left(\frac{\exp^{\beta_0 + (x_{ij} \text{ or } r_{ij}) + \beta_1 \text{ age} + \text{etc} \dots}}{1 + \exp^{\beta_0 + (x_{ij} \text{ or } r_{ij}) + \beta_1 \text{ age} + \text{etc} \dots}} * L_j \right)$$

Age-specific county means:

$$\bar{p}_{Aj} = \frac{\sum (N_{RAj} * p_{RAj})}{N_{Aj}}$$

Race-specific county means:

$$\bar{p}_{Rj} = \frac{\sum (N_{RAj} * p_{RAj})}{N_{Rj}}$$

County means:

$$\bar{p}_j = \frac{\sum (N_{RAj} * p_{RAj})}{N_j}$$

Step 3: Calculate standard deviation

$$\begin{aligned} S_{RAj} &= \sqrt{\frac{1}{n-1} \sum (p_{RAj} - \bar{p}_{RAj})^2} \\ S_{Aj} &= \sqrt{\frac{1}{n-1} \sum (p_{Aj} - \bar{p}_{Aj})^2} \\ S_{Rj} &= \sqrt{\frac{1}{n-1} \sum (p_{Rj} - \bar{p}_{Rj})^2} \\ S_j &= \sqrt{\frac{1}{n-1} \sum (p_j - \bar{p}_j)^2} \end{aligned}$$

Step 4: Calculate 95% confidence interval

$$95\% CI = \bar{p}_{RAj} \pm t_{(1-\alpha/2; n-1)} \left(\frac{S}{\sqrt{n}} \right)$$

where:

β_0 = Intercept.

β_1 = Regression coefficient for age.

β_7 = Regression coefficient for race (Black vs. White)

β_8 = Regression coefficient for race (Hispanic vs. White)

β_9 = Regression coefficient for race (other races vs. White)

β_{10} = Regression coefficient for county poverty.

β_{11} = Regression coefficient for PHR poverty.

L = weighted regression coefficient for education.

RA = race and age specific.

A = age specific.

R = race specific.

j = county specific.

$n = 10,000$ (number of simulations)

$t = 1.96$ (from Student's t distribution)

N = population count

$$P_{RAj} = \frac{\exp^{\beta_0 + \beta_1 \text{ age} + \text{etc} \dots}}{1 + \exp^{\beta_0 + \beta_1 \text{ age} + \text{etc} \dots}} * L_j$$

p = probability of vaccination.

s = standard deviation of simulated mean.

μ = area level variance.

$\bar{X}$ = simulated mean random effect for nonsampled counties (see equation 4).

[Equation 6: Calculation of median odds ratio]

Step 1: Calculate median odds ratio

$$\text{Median Odds Ratio (MOR)} = \exp^{0.95 * \sqrt{\mu_j}}$$

where:

μ = area level variance.

j = county-specific.

[Equation 7: Level 2 weight calculation]

$$w'_k = W_k K / \sum W_k$$

$$\text{where } W_k = \frac{\sum_i w_{ik}}{n_k},$$

n_k = number of respondents in a county (or PHR)

w_{ik} = raw CHLDWT values.

$K = 254$ (11 for PHR)

* We extended this formula to the level 3 (PHR) weights.

Appendix B

Appendix B

County-Level Estimates of Human Papilloma Virus Vaccine Coverage among Females Aged 11 to 17: Texas, 2008

County name	Mean (%)	Lower 95% CI (%)	Upper 95% CI (%)
Anderson	18.00	11.84	25.08
Andrews	13.87	8.48	20.18
Angelina	16.29	10.59	22.76
Aransas	9.92	5.11	16.10
Archer	9.01	4.32	15.61
Armstrong	10.17	4.87	17.57
Atascosa	16.44	9.83	23.81
Austin	11.94	7.42	17.24
Bailey	15.71	9.51	22.83
Bandera	12.46	7.06	19.29
Bastrop	13.56	8.78	19.15
Baylor	15.59	9.23	23.21
Bee	16.64	9.08	25.17
Bell	14.18	9.44	19.51
Bexar	16.54	10.33	23.67
Blanco	11.87	6.86	17.95
Borden	10.24	4.82	18.13
Bosque	13.91	8.16	20.92
Bowie	16.56	10.25	23.79
Brazoria	11.78	7.62	16.82
Brazos	24.96	17.97	32.09
Brewster	11.27	5.97	17.78
Briscoe	13.93	8.49	20.28
Brooks	19.63	9.43	30.88
Brown	14.98	9.08	22.14
Burleson	14.60	9.37	20.79
Burnet	13.28	7.90	19.71
Caldwell	15.20	9.91	21.18
Calhoun	14.42	9.05	20.52
Callahan	12.94	7.06	20.23
Cameron	20.61	10.34	31.72
Camp	15.10	9.94	20.86
Carson	9.37	4.68	15.88
Cass	15.32	9.06	22.62
Castro	18.06	11.38	25.49
Chambers	10.84	6.47	16.28
Cherokee	19.08	12.79	26.22
Childress	21.34	14.55	28.52

County name	Mean (%)	Lower 95% CI (%)	Upper 95% CI (%)
Clay	10.32	5.08	17.31
Cochran	19.62	12.80	26.98
Coke	14.59	8.76	21.60
Coleman	18.34	11.00	26.52
Collin	10.80	6.76	15.98
Collingsworth	18.20	11.51	25.91
Colorado	17.30	11.61	23.50
Comal	9.60	5.50	14.98
Comanche	15.96	9.73	23.04
Concho	22.95	15.24	31.12
Cooke	13.54	8.16	20.00
Coryell	14.67	9.45	20.87
Cottle	18.42	12.09	25.34
Crane	11.79	6.95	17.51
Crockett	14.18	8.21	21.33
Crosby	20.90	13.52	28.78
Culberson	15.21	8.01	23.50
Dallam	12.55	7.59	18.34
Dallas	18.95	13.59	24.85
Dawson	20.19	13.23	27.51
Deaf-Smith	17.68	10.51	25.49
Delta	16.43	9.19	25.13
Denton	10.63	6.66	15.78
DeWitt	16.63	10.96	23.01
Dickens	22.77	14.93	30.99
Dimmit	28.99	19.13	38.48
Donley	15.41	8.55	23.77
Duval	15.06	6.66	24.83
Eastland	16.40	9.50	24.38
Ector	15.70	9.80	22.50
Edwards	19.67	12.57	27.41
El-Paso	12.27	7.82	17.71
Ellis	17.23	8.64	26.86
Erath	19.73	12.79	27.57
Falls	21.68	15.64	27.95
Fannin	16.35	9.73	23.93
Fayette	12.07	7.31	17.90
Fisher	13.67	8.53	19.93
Floyd	19.32	12.49	26.82
Foard	15.23	8.83	22.85
Fort-Bend	11.18	7.61	15.34
Franklin	13.82	8.21	20.85

County name	Mean (%)	Lower 95% CI (%)	Upper 95% CI (%)
Freestone	13.90	8.70	20.08
Frio	25.20	15.37	34.89
Gaines	16.06	10.15	22.65
Galveston	13.38	8.76	18.80
Garza	18.96	12.46	26.19
Gillespie	10.03	5.50	15.85
Glasscock	10.56	5.80	16.77
Goliad	13.41	7.93	19.85
Gonzales	16.50	10.72	22.87
Gray	13.26	7.81	19.85
Grayson	13.92	8.17	20.96
Gregg	13.19	8.33	18.79
Grimes	14.32	9.31	20.10
Guadalupe	10.99	6.57	16.11
Hale	17.50	11.06	24.67
Hall	20.89	14.36	27.97
Hamilton	13.27	7.07	21.31
Hansford	11.89	6.86	17.88
Hardeman	15.51	9.61	22.57
Hardin	10.79	5.40	17.81
Harris	16.45	11.41	21.95
Harrison	13.28	8.16	19.44
Hartley	10.10	5.23	16.68
Haskell	19.07	12.33	26.58
Hays	15.44	9.99	21.75
Hemphill	9.48	5.03	15.44
Henderson	13.71	7.89	20.69
Hidalgo	21.30	10.54	32.87
Hill	15.60	9.79	22.24
Hockley	16.24	10.30	22.99
Hood	12.38	6.81	19.31
Hopkins	13.86	8.19	20.66
Houston	20.39	13.68	27.74
Howard	19.88	13.34	26.88
Hudspeth	19.97	11.20	29.42
Hunt	15.24	9.40	22.04
Hutchinson	11.38	6.64	17.24
Irion	10.90	6.49	16.37
Jack	13.02	7.24	20.32
Jackson	12.29	7.65	18.01
Jasper	15.73	9.33	23.17
Jeff-Davis	8.21	4.08	14.07

County name	Mean (%)	Lower 95% CI (%)	Upper 95% CI (%)
Jefferson	14.06	9.31	19.54
Jim-Hogg	12.65	5.31	21.98
Jim-Wells	11.71	5.47	19.74
Johnson	11.76	6.92	17.74
Jones	19.85	12.89	27.32
Karnes	23.42	15.94	30.90
Kaufman	11.89	7.37	17.34
Kendall	9.20	4.97	14.76
Kenedy	8.99	3.42	17.27
Kent	10.67	4.77	18.98
Kerr	12.72	7.57	19.01
Kimble	16.37	10.25	23.24
King	12.20	5.63	21.43
Kinney	19.43	11.96	27.55
Kleberg	13.93	7.05	22.39
Knox	18.13	12.12	24.97
La-Salle	15.09	8.93	22.33
Lamar	19.19	12.32	26.53
Lamb	14.81	9.16	21.63
Lampasas	25.47	15.58	35.35
Lavaca	11.45	6.62	17.51
Lee	13.02	8.20	18.52
Leon	14.05	8.46	20.66
Liberty	15.62	9.87	22.31
Limestone	16.80	11.24	23.04
Lipscomb	12.17	7.20	18.16
Live-Oak	9.10	4.47	14.90
Llano	13.25	7.42	20.44
Loving	N/A	N/A	N/A
Lubbock	14.83	9.46	20.93
Lynn	17.26	11.17	24.25
Madison	20.78	13.92	28.14
Marion	19.37	13.61	25.78
Martin	6.82	2.98	12.52
Mason	19.19	13.08	25.82
Matagorda	18.00	11.25	25.70
Maverick	15.44	9.78	22.16
McCulloch	14.12	8.37	21.01
McLennan	18.08	12.51	23.98
McMullen	23.60	13.46	33.92
Medina	15.30	9.43	22.06
Menard	19.88	12.53	27.84

County name	Mean (%)	Lower 95% CI (%)	Upper 95% CI (%)
Midland	11.89	7.32	17.39
Milam	15.63	10.35	21.73
Mills	17.18	10.46	24.95
Mitchell	22.24	15.01	29.76
Montague	12.42	6.52	19.80
Montgomery	11.39	6.79	17.22
Moore	13.08	7.60	19.56
Morris	15.16	9.53	21.72
Motley	16.49	10.15	24.01
Nacogdoches	18.56	12.61	24.98
Navarro	18.77	13.00	25.00
Newton	16.49	10.10	23.58
Nolan	17.94	11.79	24.85
Nueces	9.53	4.70	15.90
Ochiltree	12.04	7.02	17.94
Oldham	12.67	6.85	19.87
Orange	11.04	5.85	17.86
Palo-Pinto	14.99	9.11	21.84
Panola	12.45	7.31	18.89
Parker	12.47	6.90	19.68
Parmer	13.72	7.92	20.82
Pecos	19.71	12.30	27.68
Polk	12.35	7.38	18.54
Potter	19.98	13.84	26.61
Presidio	13.69	6.10	22.97
Rains	13.64	7.52	21.31
Randall	9.32	5.06	15.22
Reagan	10.89	6.07	17.03
Real	16.56	10.00	24.20
Red-River	15.75	9.67	22.78
Reeves	25.30	15.78	34.76
Refugio	7.86	3.89	12.97
Roberts	8.29	3.57	15.45
Robertson	18.58	12.82	24.85
Rockwall	10.20	5.96	15.84
Runnels	16.31	10.21	22.96
Rusk	14.06	8.84	20.64
Sabine	12.07	6.19	19.57
San-Augustine	18.34	11.32	25.94
San-Jacinto	14.90	8.78	22.10
San-Patricio	8.99	4.52	14.76
San- Saba	20.19	13.19	27.61

County name	Mean (%)	Lower 95% CI (%)	Upper 95% CI (%)
Schleicher	14.75	8.90	21.40
Scurry	16.53	10.65	23.10
Shackelford	11.06	5.52	18.26
Shelby	18.44	12.27	25.30
Sherman	13.58	8.25	19.82
Smith	12.75	8.15	18.43
Somervell	11.94	6.93	18.33
Starr	22.17	10.95	34.06
Stephens	15.68	9.41	22.84
Sterling	12.42	6.87	19.17
Stonewall	14.49	8.35	21.88
Sutton	12.37	7.30	18.35
Swisher	17.56	11.35	24.52
Tarrant	14.59	9.85	20.01
Taylor	14.91	9.45	21.19
Terrell	19.82	12.74	27.37
Terry	21.57	14.44	29.00
Throckmorton	12.85	6.41	21.35
Titus	14.20	9.03	20.20
Tom-Green	14.99	9.68	21.21
Travis	15.93	10.77	21.74
Trinity	16.06	9.79	23.62
Tyler	15.21	8.38	23.15
Upshur	12.82	7.07	19.83
Upton	14.65	8.88	21.20
Uvalde	21.84	13.39	30.41
Val-Verde	18.91	10.80	27.96
Van-Zandt	13.56	7.35	21.11
Victoria	14.47	9.08	20.65
Walker	22.09	15.67	28.81
Waller	15.88	10.59	21.54
Ward	15.96	10.24	22.37
Washington	13.55	8.61	19.38
Webb	16.74	7.39	27.43
Wharton	16.57	11.19	22.50
Wheeler	11.41	6.58	17.46
Wichita	13.04	8.06	19.12
Wilbarger	14.38	9.13	20.51
Willacy	18.01	8.57	28.84
Williamson	9.59	5.83	14.53
Wilson	10.96	6.42	16.43
Winkler	14.59	8.97	21.16

County name	Mean (%)	Lower 95% CI (%)	Upper 95% CI (%)
Wise	11.81	6.78	18.19
Wood	13.02	7.16	20.28
Yoakum	12.61	7.27	18.88
Young	13.50	7.51	20.60
Zapata	14.68	6.34	24.50
Zavala	28.98	18.34	39.25

N/A, not applicable.

Author Descriptions

Jan M. Eberth, PhD, is an Assistant Professor of Epidemiology at the University of South Carolina, Arnold School of Public Health. Her research focuses on disparities in access, utilization, and quality of cancer prevention and treatment services.

Md Monir Hossain, PhD, is an Assistant Professor in the Department of Pediatrics at Cincinnati Children's Hospital Medical Center. Dr. Hossain works to develop innovative statistical methods for cluster detection and small area health analyses.

Jasmin A. Tiro, PhD, is an Assistant Professor in the Department of Clinical Sciences at the University of Texas Southwestern Medical Center. Her research centers on evaluating HPV vaccination behavior and cancer prevention delivery in clinical settings.

Xingyou Zhang, PhD, is a geographer and statistician in the Division of Adult and Community Health at the Centers for Disease Control and Prevention. Dr. Zhang's research focuses on the integration of geographic information and technology in epidemiologic studies.

James B. Holt, PhD, is a geographer in the Division of Adult and Community Health at the Centers for Disease Control and Prevention, and Team Leader of the Analytic Methods Branch. He utilizes geospatial data and methods for chronic disease prevention research.

Sally W. Vernon, PhD, is Professor of Epidemiology and Behavioral Sciences at The University of Texas Health Science Center at Houston School of Public Health. She is recognized for her work in the area of behavioral research to promote cancer screening.

References

- Carle AC. Fitting multilevel models in complex survey data with design weights: Recommendations. *BMC Medical Research Methodology*. 2009; 9:49. [PubMed: 19602263]
- Chao C, Velicer C, Slezak JM, Jacobsen SJ. Correlates for completion of 3-dose regimen of HPV vaccine in female members of a managed care organization. *Mayo Clinic Proceedings*. 2009; 84:864–870. [PubMed: 19797775]
- Chao C, Velicer C, Slezak JM, Jacobsen SJ. Correlates for human papillomavirus vaccination of adolescent girls and young women in a managed care organization. *American Journal of Epidemiology*. 2010; 171:357–367. [PubMed: 20047978]
- Congdon P. A multilevel model for cardiovascular disease prevalence in the US and its application to micro area prevalence estimates. *International Journal of Health Geographics*. 2009; 8:6. [PubMed: 19183458]
- Congdon P. A multilevel model for comorbid outcomes: Obesity and diabetes in the US. *International Journal of Environmental Research and Public Health*. 2010; 7:333–352. [PubMed: 20616977]

- Conroy K, Rosenthal SL, Zimet GD, Yan J, Bernstein DI, Glynn S, et al. Human papillomavirus vaccine uptake, predictors of vaccination, and self-reported barriers to vaccination. *Journal of Women's Health*. 2009; 18:1679–1686.
- Dempsey A, Cohn L, Dalton V, Ruffin M. Patient and clinic factors associated with adolescent human papillomavirus vaccine utilization within a university-based health system. *Vaccine*. 2010; 28:989–995. [PubMed: 19925899]
- Dorell C, Stokley S, Yankey D, Cohn A, Markowitz L. National, state, and local area vaccination coverage among adolescents aged 13–17 years: United States, 2009. *Morbidity & Mortality Weekly Report*. 2010; 59:1018–1023. [PubMed: 20724968]
- Dorell C, Stokley S, Yankey D, Liang JL, Markowitz L. National and state vaccination coverage among adolescents aged 13 through 17 years: United States, 2010. *Morbidity and Mortality Weekly Report*. 2011; 60:1117–1123. [PubMed: 21866084]
- Goldstein, H. Multilevel statistical models. 4th ed. John Wiley & Sons, Ltd; West Sussex, UK: 2010.
- Goodman MS. Comparison of small-area analysis techniques for estimating prevalence by race. *Preventing Chronic Disease*. 2010; 7:A33. [PubMed: 20158961]
- Gottlieb SL, Brewer NT, Sternberg MR, Smith JS, Ziarnowski K, Liddon N, et al. Human papillomavirus vaccine initiation in an area with elevated rates of cervical cancer. *Journal of Adolescent Health*. 2009; 45:430–437. [PubMed: 19837348]
- Hosmer, DW.; Lemeshow, S. Applied logistic regression. 2nd ed. John Wiley & Sons, Inc; Hoboken, NJ: 2000.
- Jain N, Stokley S, Yankey D. Vaccination coverage among adolescents aged 13–17 years: United States, 2007. *Morbidity & Mortality Weekly Report*. 2008; 57:1100–1103. [PubMed: 18846032]
- Jia H, Muennig P, Borawski E. Comparison of small-area analysis techniques for estimating county-level outcomes. *American Journal of Preventive Medicine*. 2004; 26:453–460. [PubMed: 15165663]
- Kahn JA, Rosenthal SL, Jin Y, Huang B, Namakydoust A, Zimet GD. Rates of human papillomavirus vaccination, attitudes about vaccination, and human papillomavirus prevalence in young women. *Obstetrics and Gynecology*. 2008; 111:1103–1110. [PubMed: 18448742]
- Li W, Land T, Zi Z, Keithly L, Kelsey JL. Small-area estimation and prioritizing communities for tobacco control efforts in Massachusetts. *American Journal of Public Health*. 2009; 99:470. [PubMed: 19150913]
- Mendez-Luck CA, Yu H, Meng YY, Jhavar M, Wallace SP. Estimating health conditions for small areas: Asthma symptom prevalence for state legislative districts. *Health Services Research*. 2007; 42:2389–2409. [PubMed: 17995549]
- Neubrand TPL, Breitkopf CR, Rupp R, Breitkopf D, Rosenthal SL. Factors associated with completion of the human papillomavirus vaccine series. *Clinical Pediatrics*. 2009; 48:966–969. [PubMed: 19483128]
- Pruitt SL, Schootman M. Geographic disparity, area poverty, and human papillomavirus vaccination. *American Journal of Preventive Medicine*. 2010; 38:525–533. [PubMed: 20409501]
- Rabe-Hesketh S, Skrondal A. Multilevel modelling of complex survey data. *Journal of the Royal Statistical Society: Series A (Statistics in Society)*. 2006; 169:805–827.
- Rondy M, van Lier A, van de Kasstele J, Rust L, de Melker H. Determinants for HPV vaccine uptake in the Netherlands: A multilevel study. *Vaccine*. 2010; 28:2070–2075. [PubMed: 20045095]
- Rosenthal SL, Rupp R, Zimet GD, Meza HM, Loza ML, Short MB, et al. Uptake of HPV vaccine: Demographics, sexual history and values, parenting style, and vaccine attitudes. *Journal of Adolescent Health*. 2008; 43:239–245. [PubMed: 18710678]
- Schneider KL, Lapane KL, Clark MA, Rakowski W. Using small-area estimation to describe county-level disparities in mammography. *Preventing Chronic Disease*. 2009; 6:A125. [PubMed: 19755001]
- Siegel PZ, Frazier EL, Mariolis P, Brackbill RM, Smith C. Behavioral Risk Factor Surveillance, 1991: Monitoring progress toward the nation's year 2000 health objectives. *Morbidity and Mortality Weekly Report Surveillance Summaries*. 1993; 42(SS-4):1–21.

- U.S. Centers for Disease Control and Prevention. [Retrieved September 2, 2011] Behavioral Risk Factor Surveillance System: 2008 summary data quality report. 2008a. from ftp://ftp.cdc.gov/pub/Data/Brfss/2008_Summary_Data_Quality_Report.pdf
- U. S. Centers for Disease Control and Prevention. [Retrieved January 20, 2011] BRFSS Annual Survey Data: Survey data and documentation. 2008b. from http://www.cdc.gov/brfss/technical_infodata/surveydata/2008.htm
- U.S. Centers for Disease Control and Prevention. [Retrieved January 20, 2011] U.S. Census populations with bridged race categories. 2009. from http://www.cdc.gov/nchs/nvss/bridged_race.htm
- U.S. Food and Drug Administration. [Retrieved January 20, 2011] FDA licenses new vaccine for prevention of cervical cancer and other diseases in females caused by human papillomavirus. 2006. from <http://www.fda.gov/NewsEvents/Newsroom/PressAnnouncements/2006/ucm108666.htm>
- Wong CA, Berkowitz Z, Dorell CG, Price RA, Lee J, Saraiya M. Human papillomavirus vaccine uptake among 9- to 17-year-old girls. *Cancer*. 2011; 117:5612–5620. [PubMed: 21692069]
- Yu H, Meng YY, Mendez-Luck CA, Jhavar M, Wallace SP. Small-area estimation of health insurance coverage for California legislative districts. *American Journal of Public Health*. 2007; 97:731–737. [PubMed: 17329663]
- Zimet GD, Liddon N, Rosenthal SL, Lazcano-Ponce E, Allen B. Chapter 24: Psychosocial aspects of vaccine acceptability. *Vaccine*. 2006; 24(S3):201–209.

Figure 1.

Summary of data collection, management, and analysis strategy. PCA, principal component analysis; PHR, public health region. ^aPreliminary model includes age, race/ethnicity, parental education, county and PHR poverty. ^bTechnical documentation (i.e., equations/Equ.) can be found in Appendix A. ^cFinal model includes all the variables from the preliminary model except for parental education. Although parental education is an important predictor of HPV vaccination, it could not be included in the final model because the corresponding variable was not present in the Auxiliary Dataset. A post-adjustment method, described by Congdon (25), was used as an alternative to inclusion of parental education in the final model.

Figure 2.
County-level estimates of HPV vaccine coverage among females aged 11 to 17 in Texas.

Table 1

Model-Building Strategy for Determining the Final Model of HPV Vaccine Initiation among Texas Females Aged 11 to 17

Step	Model Type*	Covariates Examined	Covariates Retained
1	Empty model	N/A	N/A
2	Covariates at level 1	Child's age	Child's age
		Child's race/ethnicity	Child's race/ethnicity
		Parental education	Parental education
3	Covariates at levels 1 and 2	Child's age	Child's age
		Child's race/ethnicity	Child's race/ethnicity
		Parental education	Parental education
		County poverty (%)	County poverty (%)
		County chlamydia rate	
		County cervical cancer incidence rate	
		County polio vaccination (%) among 7th graders	
		County measles vaccination (2nd dose, %) among kindergarteners	
		County rurality index (metropolitan, micropolitan, noncore)	
		County uninsurance (%)	
		County evangelical religious adherence rate	
4	Covariates at all levels: Preliminary model	Child's age	Child's age
		Child's race/ethnicity	Child's race/ethnicity
		Parental education	Parental education
		County poverty (%)	County poverty (%)
		PHR poverty (%)	PHR poverty (%)
		PHR chlamydia rate	
		PHR cervical cancer incidence rate	
		PHR polio vaccination (%) among 7th graders	
		PHR measles vaccination (2nd dose, %) among kindergarteners	
		PHR uninsurance (%)	
5	Covariates at all levels: Final model	N/A	Child's age
			Child's race/ethnicity
			County poverty (%)
			PHR poverty (%)

Abbreviations: HPV4, Gardasil; N/A, not applicable; PHR, public health region.

* All models contain both county- and PHR-level random intercepts.

Table 2

Random-Intercept Logit Models of HPV Vaccine Initiation among Texas Females Aged 11 to 17

Parameters	Model 1, OR (95% CI)	Model 2, OR (95% CI)	Model 3, OR (95% CI)	Model 4, OR (95% CI)	Model 5, OR (95% CI)
Fixed effects					
Intercept	0.26 (0.21–0.32)	0.10 (0.02–0.50)	0.05 (0.01–0.30)	0.09 (0.01–0.72)	0.16 (0.03–0.82)
Child's age	N/A	1.03 (0.93–1.14)	1.03 (0.93–0.14)	1.03 (0.92–1.14)	1.03 (0.94–1.12)
White	N/A	Ref	Ref	Ref	Ref
Black	N/A	1.04 (0.33–3.23)	1.06 (0.35–3.22)	1.02 (0.34–3.06)	0.93 (0.31–2.76)
Hispanic	N/A	2.00 (1.48–2.70)	1.70 (1.17–2.47)	1.77 (1.28–2.46)	1.34 (0.95–1.91)
Other races	N/A	1.28 (0.51–3.20)	1.25 (0.46–3.40)	1.24 (0.42–3.63)	1.29 (0.45–3.71)
<HS education	N/A	Ref	Ref	Ref	N/A
HS graduate	N/A	0.78 (0.37–1.66)	0.82 (0.39–1.73)	0.85 (0.40–1.77)	N/A
Some college	N/A	1.80 (0.88–3.66)	1.94 (0.94–4.02)	1.97 (0.95–4.10)	N/A
Bachelor's	N/A	1.72 (0.94–3.16)	1.84 (0.98–3.48)	1.84 (0.97–3.47)	N/A
County poverty*	N/A	N/A	1.05 (0.99–1.11)	1.09 (1.02–1.18)	1.09 (1.02–1.17)
PHR poverty	N/A	N/A	N/A	0.92 (0.86–0.99)	0.92 (0.86–0.98)

Abbreviations: CI, confidence interval; HPV4, Gardasil; HS, high school; OR, odds ratio; PHR, public health region; SE, standard error.

* County- and PHR-level poverty were measured continuously.

Table 3

Comparison of Indirect and Direct Estimation of Human Papilloma Virus (HPV) Vaccine Coverage at the State Level

HPV Vaccine Coverage in Texas	Indirect Estimate (95% CI)	Direct Estimate (95% CI)
Texas	15.51 (14.14–16.93)	17.74 (13.59–21.88)
Texas, racial/ethnic categories		
White	13.55 (11.93–15.32)	13.69 (9.18–18.82)
Black	14.45 (11.46–17.72)	21.70 (5.97–37.42)
Hispanic	17.61 (15.00–20.31)	22.42 (14.03–30.80)
Other races	16.27 (13.19–19.60)	23.75 (0–54.58)
Texas, age categories (yrs)		
11	14.70 (11.41–28.50)	11.48 (3.55–19.41)
12	14.94 (11.63–18.73)	12.20 (5.95–18.46)
13	15.25 (11.88–19.18)	27.81 (16.71–38.92)
14	15.55 (12.13–19.37)	14.53 (4.84–24.22)
15	15.78 (12.24–19.64)	27.69 (12.80–42.59)
16	16.04 (12.55–19.85)	16.68 (7.89–25.46)
17	16.32 (12.82–20.28)	15.95 (6.37–25.53)