Published in final edited form as:

Biochemistry. 2013 July 23; 52(29): . doi:10.1021/bi400681d.

Dependence of the product chain-length on detergents for long-chain *E*-polyprenyl diphosphate synthases

Jian-Jung Pan, Gurusankar Ramamoorthy, and C. Dale Poulter*

Department of Chemistry, University of Utah, 315 South 1400 East, Salt Lake City, Utah 84112

Abstract

Long-chain *E*-polyprenyl diphosphate synthases (*E*-PDS) catalyze repetitive addition of isopentenyl diphosphate (IPP) to the growing prenyl chain of an allylic diphosphate. The polyprenyl diphosphate products are required for the biosynthesis of ubiquinones and menaquinones required for electron transport during oxidative phosphorylation to generate ATP. *In vitro*, the long-chain PDSs require addition of phospholipids or detergents to the assay buffer to enhance product release and maintain efficient turnover. During preliminary assays of product chain-length with anionic, zwitterionic, and non-ionic detergents, we discovered considerable variability. Examination of a series of non-ionic PEG detergents with several long-chain *E*-PDSs from different organisms revealed that *in vitro* incubations with nonaethylene glycol monododecyl ether or Triton X-100 typically gave chain lengths that corresponded to those of the isoprenoid moieties in respiratory quinones synthesized *in vivo*. In contrast incubations in buffer with *n*-butanol, CHAPS, DMSO, *n*-octyl- β -glucopyranoside, or β -cyclodextrin or in buffer without detergent typically proceeded more slowly and gave a broad range of chain lengths.

Polyprenyl diphosphate synthases (PDS) catalyze 1'-4 chain elongation by the sequential addition of the growing isoprenoid chain in an allylic diphosphate to isopentenyl diphosphate (IPP) to generate linear allylic diphosphates with two or more isoprene units. The enzymes are selective for the ultimate chain-length of their products and the stereochemistry of the double bond, *E* or *Z*, introduced during each addition to IPP. The *E*-PDSs generally catalyze the formation of products with shorter chains, C₁₀₋₅₀. *Z*-PDSs typically give compounds with eleven (C₅₅) or more isoprene units including natural rubber, although a few instances of short-chain *Z*-PDSs have been reported.¹⁻² The linear polyprenyl diphosphates are essential substrates for the biosynthesis of all but the simplest isoprenoid compounds found in Nature. Farnesyl diphosphate (FPP, C₁₅) is required for synthesis of sterols, sesquiterpenes, and farnesylated proteins, while geranylgeranyl diphosphate (GGPP, C₂₀) is the precursor for carotenoids, diterpenes, and geranylgeranylated proteins. These metabolites perform numerous essential functions in cells and are a rich source for potential human therapeutics. The longer chain *E*-allylic diphosphates (typically C₃₀ – C₅₀) are required for the biosynthesis of the isoprenoid side chain of ubiquinones (UQ) and menaquinones (MQ) in different species, such as UQ-6 in *Saccharomyces cerevisiae*, UQ-8 in *E. coli* and UQ-10 in humans. The respiratory quinones are intermediates in the electron transport in the cellular respiratory metabolism to generate ATP. *Z*-PDSs synthesize linear C₅₅ – C₁₀₀₊ isoprenoid molecules that serve as a lipid carrier during peptidoglycan biosynthesis of bacterial cell walls and glycoprotein biosynthesis in eukaryotes.

*Corresponding author: Tel.: (801) 581-6685. poulter@chem.utah.edu..

Supporting Information Available

TLCs 1-4 with *n*-butanol, SDS, DMSO and OGP; Michaelis-Menten plots for GIs 53711383 and 19551716; TLCs with different allylic substrates, DMAPP and FPP (adapted from Ref. 9). This material is available free of charge via the Internet at <http://pubs.acs.org>.

Purified long-chain PDSs require phospholipids or detergents for efficient turnover, presumably due to the slow release of the hydrophobic products. Their stimulatory effect appears to be enzyme-dependent.³ For instance, Triton X-100 is a good activator for the *L. plantarum*, *M. luteus* and *Bacillus subtilis* undecaprenyl diphosphate (UPPS) synthases, while deoxycholate is only effective for the *L. plantarum* enzyme. The stimulatory effect of phospholipids also varies with enzyme and the type of phospholipids. In addition to stimulating activity, phospholipids and detergents can also affect the product profile. For example, the chain-length of major product for *M. luteus* UPPS shifted from C₅₀ to C₅₅ when Triton X-100 was replaced by a phospholipid extract. Without a phospholipid or detergent in the assay buffer, *E. coli* UPPS slowly produced a series of products with chain lengths up to C₇₅.⁴ A microsomal preparation of dehydrololichyl diphosphate synthase, a C₉₀ and C₉₅ Z-PDS from rat liver, gave shorter C₅₅-C₈₀ products with increasing concentrations of different detergents.⁵

Only a few studies have been reported for the effects of detergent and phospholipids on the activity and product distribution of long-chain *E*-PDSs. The activity of solanesyl diphosphate (SPP, C₄₅) synthase is stimulated by Tween-80, bovine serum albumin (BSA) and bacitracin. The *M. luteus* enzyme was also stimulated by a high molecular-mass fraction (HMF) from the bacterium. HMF is thought to bind to the C₄₅-PP products and presumably promotes turnover by providing a hydrophobic reservoir for the products.⁶ In related studies with *E. coli* octaprenyl diphosphate (OPP, C₄₀) synthase, CHAPS and Triton X-100 were stimulatory, while Tween-80, deoxycholic acid and *n*-octyl glucoside were not.⁷ BSA and HMF were also activators. The effects of these reagents on product distribution were not reported, except that in the absence of Triton X-100 over-elongated products (C₄₅-C₆₀) were found. Similar results were observed for *Paracoccus denitrificans* decaprenyl diphosphate (DPP, C₅₀) synthase, which gave a C₆₅ product in the absence of Triton X-100,⁸ and a *Cryptosporidium parvum* nonspecific PDS, where the chain lengths of the products increased from C₂₀-C₂₅ in the presence of Tween-20 to C₄₀ in its absence.⁹ We now report an investigation of the effect of several different detergents on the activities and product distributions as part of a study of the functions of long-chain *E*-PDSs.

Materials and methods

Reagents

All the chemicals were purchased from Sigma-Aldrich. [4-¹⁴C]IPP was purchased from PerkinElmer. DMAPP, GPP and FPP were synthesized according to that of Davisson et al.¹⁰ Potato acid phosphatase was purchased from Roche.

Enzymes

All the enzymes were provided by the Enzyme Function Initiative (EFI) Protein Core at the Albert Einstein College of Medicine. The detailed procedures of the enzyme expression and purification are published elsewhere.¹¹

Product analysis

Incubations were carried out at 30 °C for 2 h in 40 µL of a mixture of 35 mM HEPES buffer, pH 7.6, containing 5 mM β-mercaptoethanol, 10 mM MgCl₂, and 200 µM [¹⁴C]IPP and 50 µM allylic substrate (DMAPP, GPP or FPP), unless otherwise specified. The reaction was quenched with 0.5 M EDTA, pH 8.0, extracted with 3× 80 µL of water-saturated *n*-butanol. The butanol extract was transferred to an Eppendorf tube; solvent was removed on a SpeedVacTM; and 200 µL of 50 mM sodium acetate buffer, pH 4.7, containing 2 units of potato acid phosphatase (2 units/mg), 0.1 % (v/v) Triton X-100, and 20 % (v/v) *n*-propanol were added. The mixture was incubated overnight at 37 °C and the isoprenoid alcohols were

extracted with $3 \times 200 \mu\text{L}$ of hexanes or methyl *t*-butyl ether (MTBE). The volume was concentrated on the SpeedVacTM and resuspended in $15 \mu\text{L}$ hexanes or MTBE. Samples were spotted on RP-C18 TLC plates and developed with 19:1 (v/v) acetone:water. The developed TLC plates were visualized by phosphorimaging (Molecular Dynamics storage phosphor screen) using a Typhoon 8600 Variable Mode Imager (GE Healthcare).

Kinetic studies

Initial velocities were measured by the acid-lability assay¹² for incubations of $400 \mu\text{L}$ samples of 35 mM HEPES buffer, pH 7.6, containing 5 mM β -mercaptoethanol, 10 mM MgCl_2 , varied concentrations of [^{14}C]IPP, FPP, and detergents at 37°C for 10 min. The reaction was initiated by addition of enzyme, samples were removed periodically, and immediately quenched with $\text{MeOH:HCl}=3:1$, followed by incubation at 37°C for 10 min. The sample was extracted with $400 \mu\text{L}$ of ligroin (b.p. $95\text{--}110^\circ\text{C}$), and the radioactivity in a $200 \mu\text{L}$ portion was quantified by liquid scintillation spectrometry on a Tri-Carb 2910TR liquid scintillation analyzer (PerkinElmer). The data were fit to the Michaelis-Menten equation using GraFit 5.0 (Erithacus Software).

Results and discussion

As part of a study to determine the function of isoprenoid chain elongation prenyltransferases identified in this work by their GI (geneInfo number) designations, we examined a diverse group of putative chain elongation enzymes within a larger cluster of proteins in the isoprenoid synthase superfamily.¹¹ One of the challenges in the study was the selection of a buffer system that gave maximal activity and a set of products whose chain lengths most closely approximated those formed *in vivo*. Since examination of previous reports gave no clear consensus for which set of reaction conditions were optimal, the chain-length selectivity of several *E*-PDSs for a series of reagents that could alter the solubility of the hydrophobic long-chain isoprenyl diphosphate products was studied.

Effect of buffer composition on product distributions

Product distributions used for the PDSs were studied in HEPES buffer supplemented with the following reagents: *n*-butanol, dimethyl sulfoxide (DMSO), sodium dodecyl sulfate (SDS), CHAPS, β -cyclodextrin (CD), 3,6,9-heptakis-(2,6-di-*O*-methyl)- β -cyclodextrin (MCD), *n*-octyl- β -glucopyranoside (OGP), Triton X-100 (TX), pentaethylene glycol monododecyl ether (PEG12E), hexaethylene glycol monododecyl ether (HEG12E), octaethylene glycol monododecyl ether (OEG12E), nonaethylene glycol monododecyl ether (NEG12E) and octaethylene glycol tetradecyl ether (OEG14E). Incubations with enzyme GI 19551716 in buffer containing 5% – 25% (v/v) *n*-butanol gave little turnover, resulting in trace amounts of products, perhaps due to denaturation of the enzymes (Supplementary data, TLC1). Similar results were obtained with several enzymes in the presence of 0.5 mM SDS (Supplementary data, TLC2). In the presence of DMSO or OGP, the PDSs were less selective and gave longer chain lengths than for incubations in 0.1 % (1.6 mM) TX (see SI, TLCs1,3,4). Product distributions in buffer containing CHAPS were dependent on detergent concentration, and are compared with 0.1% TX in Figures 1A and 1B. The predominant product for enzyme GI 19551716 is a C_{45} allylic diphosphate in buffer with TX, accompanied by smaller amounts of C_{40} and C_{50} products. The C_{45} diphosphate also predominates in the presence of CHAPS; however, the amount of C_{50} material increases slightly at 2.5 mM and the amount of C_{40} product increases in 5 mM relative to TX (Figure 1A). As the concentration of TX was reduced from 0.1% to 0.01%, the product distribution shifted to longer chain lengths, as seen for GI 19551716, GI 53711383, and GI 67866738, where the profiles are similar at 0.1% and 0.05% but shift substantially at 0.01% (Figure

1B). Product distributions in 2 mM CHAPS are similar to those for 0.01% TX. The optimal concentration of TX for selective production of products appears to be 0.05-0.1%.

Considerable variation was seen in chain lengths for different PDSs in buffers containing CD or MCD relative to 0.1% TX. At 2 mM MCD (Figure 2A), the product distributions for GI 19551716 (C₄₅), GI 53711383 (C₅₅), and GI 67866738 (C₅₀) were similar to those for 0.1% Triton X-100. Buffer with 1.6 mM CD gave a slight increase in higher chain-length products for GI 19551716. A more substantial increase in chain-length (C₅₀ to C₅₅) was seen for GI 67866738, while GI 53711383 gave a substantial shift to higher chain lengths with C₆₅ allylic diphosphate as the major product along with substantial amounts of C₆₀–C₇₅ chains. The profiles for GI 19551716 were similar for buffers containing 0.2 mM CD and MCD (Figure 2B), while for GI 53711383, longer chain products were produced less selectively. GI 16131077, which appears to be inherently less selective in 0.1% TX buffer than the other three enzymes listed in Figure 2 (compare lanes 2), gave an even broader distribution of products in 0.2 mM CD and MCD buffers.

The PEG-based non-ionic detergent TX consistently gave the narrowest distribution of chain lengths. However, TX is a mixture consisting of a 4-(1,1,3,3-tetramethylbutyl)phenyl moiety attached to PEG units of different lengths. This heterogeneity sometimes complicates the isolation and purification of the products. We compared 2 mM concentrations of a series of dodecyl ether PEG detergents containing PEG₅ (PEG12E), PEG₆ (HEG12E), PEG₈ (OEG12E), and PEG₉ (NEG12E) units and a PEG₈ tetradecyl ether (OEG14E) with 0.1% TX to examine the effect of the non-ionic detergents on the length of the hydrocarbon chain (Figure 3). While all of the alkoxy PEG detergents gave narrow product distributions, the profiles for NEG12E most closely matched those of TX and, in one case (see GI 67866738 in Figure 3B), was more selective.

The long-chain *E*-PDSs typically prefer DMAPP or FPP as the initial allylic substrate and the IPP:allylic substrate ratio can affect product distributions.¹³⁻¹⁶ One might expect that the longest chain product would depend strongly on the initial IPP:allylic diphosphate ratio, for example IPP:DMAPP = 4:1 would give mostly the C₂₅ diphosphate. However, long-chain PDSs selectively elongate by binding the intermediates tighter than the starting substrates, and terminate when the growing chain fills the elongation cavity.^{4,7} Over elongation is seen under forcing conditions, presumably from a change in conformation of the enzyme induced by the isoprenoid chain to create a longer binding channel.¹¹ Figure 4 shows the product distributions for four enzymes and different ratios of DMAPP and FPP. The product distributions were similar in 0.1% TX buffer for DMAPP and FPP at 4:1 and 1:2 ratios of IPP:allylic substrate. In buffer without TX, the reactions were slower, less selective, and somewhat over-elongated, even when the IPP:allylic ratio was 1:2.

Additional comparisons are shown in Figure 5. Over elongation was seen for the allylic substrates FPP and geranyl diphosphate (C₁₀, GPP) in the absence of TX, but both gave a narrower range of chain lengths than DMAPP. This observation suggests that the longer allylic substrates, FPP and GPP, help promote chain-length selectivity by competing better than DMAPP with longer chain-length products for binding to the enzymes. Interestingly, the product distributions at 20:1 ratios of IPP:FPP or DMAPP in 24 hour incubations (Figure 6) were similar to those determined for 2 h incubations with 4:1 IPP and allylic substrates. The product distributions for two enzymes, GI 29376566 and GI 58578715, were similar with, or without, TX in the buffer. It is possible that these enzymes are normally associated with another protein *in vivo*, that facilitates product release.⁶ In addition, it is also important to note the influence that a His₆ tag can exert on the distributions of products. Those shown in Figures 3A,B,D and 5A,B for GI 67866738, a C₅₀ elongation enzyme, are similar for C-terminal His₆ tagged, and N-terminal His₆ tagged versions of the protein. In contrast, N-

terminal His₆ tagged GI 39934115 are C₅₀ elongation enzymes, while the C-terminal His₆ tagged protein gives approximately equal amounts of C₄₅ and C₅₀ products (Figures 3C, 5A and 5B). A much larger difference is seen for GI 19551716, where the C-terminal tagged proteins synthesize C₄₅ products, while the N-terminally tagged enzyme give a broad distribution of products between C₄₅ and C₇₀ (Figures 3C, 6A, and 6B). Thus, it is advisable to work with tagless or both N- and C-terminal tag versions of the recombinant proteins when determining function.

Effects of detergents on kinetic parameters

Steady-state kinetic constants were determined for PDSs GI 53711383 and GI 19551716 in buffer containing 0.1% TX. For GI 53711383, $k_{\text{cat}} = 0.03 \text{ s}^{-1}$, $K_{\text{M}}^{\text{IPP}} = 11 \text{ }\mu\text{M}$, and $K_{\text{M}}^{\text{FPP}} = 0.05 \text{ }\mu\text{M}$ and for GI 19551716, $k_{\text{cat}} = 0.01 \text{ s}^{-1}$, $K_{\text{M}}^{\text{IPP}} = 13 \text{ }\mu\text{M}$, and $K_{\text{M}}^{\text{FPP}} = 0.28 \text{ }\mu\text{M}$. Relative rates were then measured for both enzymes in buffers containing saturating concentrations of IPP (50 μM) and FPP (5 μM) and no detergent or 2 mM OGP, 2 mM NEG12E, or 0.1% TX. For GI 53711383, ~10-fold stimulations of the rate for chain elongation were seen for NEG12E and TX (Figure 7A), while the rate in buffer containing OGP is similar to that in the absence of a detergent. The enzyme gave a tight distribution consisting of C₅₀ and C₅₅ products in TX and NEG12E buffers, and a broad distribution between C₃₀ and C₆₅ in OGP or without detergent (Figure 8A). Detergents had a much smaller effect on the rate of chain elongation for GI 19551716 (Figure 7B) with a slight increase for NEG12E and slight decreases for OGP and TX. Tight product distributions with a maximum at C₄₅ were seen for NEG12E and TX. A slight increase in the amount of C₅₀ diphosphate was seen with OGP (Figure 8B). A broader distribution between C₄₅ and C₆₀ was seen without detergent. Previously, small 3-fold rate enhancements were reported for the *E. coli* C₄₀ synthase in TX.⁷ The rate enhancement was not large enough to switch the rate-determining step from the product release to the chemical steps,⁷ and the authors suggested that another mechanism for enhancing product release/sequestration might operate *in vivo*.

Conclusions

As chain-length increases, the polyisoprenoid diphosphate products of chain elongation reactions become increasingly hydrophobic. *In vivo*, the longer chain-length compounds can affiliate with membrane, bind to and transported by proteins, and converted to downstream metabolites. All of these scenarios remove the products from the vicinity of the enzyme and promote turnover. *In vitro*, the hydrophobic products accumulate and can bind to the enzyme, thereby slowing the chain elongation process by product inhibition and serve as substrates for further prenyltransfers to produce “over-elongated” products. Inclusion of detergents in the assay buffer typically promotes turnover and selectivity.

We and others found that Triton X-100 (TX) consistently stimulates activity and enhances selectivity.^{13,16,17} In addition, we found that the chain lengths of polyisoprenoid diphosphates generated by *E*-PDSs in TX buffer are similar to those found in metabolites containing the polyisoprenoid chains synthesized *in vivo*. As seen in Table 1, the principal products we detected for nine long chain *E*-PDSs in the presence of TX closely matched the chain lengths of the polyisoprenoid chains found in menaquinone and ubiquinones isolated from the corresponding organisms. TX may facilitate the rate and selectivity of chain elongation by solubilizing the hydrophobic products and competing with polyisoprenoid products for binding to the active site of the chain elongation enzymes. A crystal structure of decaprenyl diphosphate synthase shows two molecules of TX in the active site tunnel.¹⁸ Detergent binding is consistent with the observation that activity is reduced to levels seen in its absence while selectivity is retained at high concentrations (> 1%, 16 mM).

Finally, Triton X-100 is a good detergent for *in vitro* kinetic and product studies of *E*-PDSs. However, TX is a mixture of compounds where the hydrophobic (tetramethylbutyl)phenyl moiety is attached to a series PEG units of different lengths consisting of an average 9.5 ethylene glycol monomers. In some applications, the mixture of detergent molecules complicates the isolation and purification of products from the enzyme catalyzed reactions. As a result, we were interested in finding a monomeric non-ionic detergent that would minimize this problem and examined a series of *n*-alkyl PEG ethers. Preliminary experiments indicated that the lengths of the hydrocarbon and PEG regions of the detergents were important. Of the PEGs studied, we found that NEG12E consistently gave polyprenyl diphosphates with the narrowest product distributions and the chain lengths corresponded to those synthesized *in vivo* and to the chain lengths found for TX. Clearly care should be taken in the selection of detergents for *in vitro* studies of the long-chain elongation enzymes, and NEG12E and TX are good choices for a wide variety of *E*-PDSs.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This project was supported by the Enzyme Function Initiative, NIH grant GM093342.

Funding support

This work was supported by the Enzyme Function Initiative, NIH grant GM093342.

Abbreviations

PDS	polyprenyl diphosphate synthases
DMAPP	dimethylallyl diphosphate
IPP	isopentenyl diphosphate
ATP	adenosine triphosphate
PEG	polyethylene glycol
DMSO	dimethyl sulfoxide
GPP	geranyl diphosphate
FPP	farnesyl diphosphate
GGPP	geranylgeranyl diphosphate
OPP	octaprenyl diphosphate
SPP	solanesyl diphosphate
DPP	decaprenyl diphosphate
UPP	undecaprenyl diphosphate
UQ	ubiquinone
MQ	menaquinone
BSA	bovine serum albumin
HMF	high molecular-mass fraction
CHAPS	3-[(3-cholamidopropyl)dimethylamino]-1-propanesulfonate

MTBE	methyl <i>t</i> -butyl ether
TLC	thin-layer chromatography
HEPES	4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid
SDS	sodium dodecyl sulfate
CD	β -cyclodextrin
MCD	3,6,9-heptakis-(2,6-di-O-methyl)- β -cyclodextrin
OGP	<i>n</i> -octyl- β -glucopyranoside
TX	Triton X-100
PEG12E	pentaethylene glycol monododecyl ether
HEG12E	hexaethylene glycol monododecyl ether
OEG12E	octaethylene glycol monododecyl ether
NEG12E	nonaethylene glycol monododecyl ether
OEG14E	octaethylene glycol tetradecyl ether

References

- Schulbach MC, Brennan PJ, Crick DC. Identification of a short (C₁₅) chain *Z*-isoprenyl diphosphate synthase and a homologous long (C₅₀) chain isoprenyl diphosphate synthase in *Mycobacterium tuberculosis*. *J. Biol. Chem.* 2000; 275:22876–22881. [PubMed: 10816587]
- Akhtar TA, Matsuba Y, Schauvinhold I, Yu G, Lees HA, Klein SE, Pichersky E. The tomato *cis*-prenyltransferase gene family. *Plant J.* 2013; 73:640–652. [PubMed: 23134568]
- Allen CM. Purification and characterization of undecaprenylpyrophosphate synthetase. *Method Enzymol.* 1985; 110:281–299.
- Pan J-J, Chiou S-T, Liang P-H. Product distribution and pre-steady state kinetic analysis of *Escherichia coli* undecaprenyl pyrophosphate synthase reaction. *Biochemistry.* 2000; 39:10936–10942. [PubMed: 10978182]
- Matsuoka S, Sagami H, Kurisaki A, Ogura K. Variable product specificity of microsomal dehydrololichyl diphosphate synthase from rat liver. *J. Biol. Chem.* 1991; 266:3464–3468. [PubMed: 1995611]
- Ohnuma, S.-i.; Koyama, T.; Ogura, K. Purification of solanesyl-diphosphate synthase from *Micrococcus luteus*: a new class of prenyltransferase. *J. Biol. Chem.* 1991; 266:23706–23713. [PubMed: 1748647]
- Pan J-J, Kuo T-H, Chen Y-K, Yang L-W, Liang P-H. Insight into the activation mechanism of *Escherichia coli* octaprenyl pyrophosphate synthase derived from pre-steady kinetic analysis. *Biochim. Biophys. Acta.* 2002; 1594:64–73. [PubMed: 11825609]
- Takahashi S, Nishino T, Koyama T. Isolation and expression of *Paracoccus denitrificans* decaprenyl diphosphate synthase gene for production of ubiquinone-10 in *Escherichia coli*. *Biochem. Eng. J.* 2003; 16:183–190.
- Artz JD, Dunford JE, Arrowood MJ, Dong A, Chruszcz M, Kavanagh K, Minor W, Russell RGG, Ebetino FH, Oppermann U, Hui R. Targeting a uniquely nonspecific prenyl synthase with bisphosphonates to combat cryptosporidiosis. *Chem. Biol.* 2008; 15:1296–1306. [PubMed: 19101474]
- Davisson VJ, Woodside AB, Neal TR, Stremmler KE, Muehlbacher M, Poulter CD. Phosphorylation of isoprenoid alcohols. *J. Org. Chem.* 1986; 51:4768–4779.
- Wallrapp FH, Pan J-J, Ramamoorthy G, Almonacid DE, Hillerich BS, Seidel R, Patskovsky Y, Babbitt PC, Almo SC, Jacobson MP, Poulter CD. Prediction of function for the polyprenyl

- transferase subgroup in the isoprenoid synthase superfamily. *Proc Natl. Acad. Sci. U.S.A.* 2013; 110:E1196–1202. [PubMed: 23493556]
12. Laskovics FM, Krafcik JM, Poulter CD. Prenyltransferase. Kinetic studies of the 1'-4 coupling reaction with avian liver enzyme. *J. Biol. Chem.* 1979; 254:9458–9463. [PubMed: 489545]
 13. Ferella M, Montalvetti A, Rohloff P, Miranda K, Fang J, Reina S, Kawamukai M, Búa J, Nilsson D, Pravia C, Katzin A, Cassera MB, Åslund L, Andersson B, Docampo R, Bontempi EJ. A solanesyl-diphosphate synthase localizes in glycosomes of *Trypanosoma cruzi*. *J. Biol. Chem.* 2006; 281:39339–39348. [PubMed: 17062572]
 14. Hsieh F-L, Chang T-H, Ko T-P, Wang AJ-H. Structure and mechanism of an *Arabidopsis* medium/long-chain-length prenyl pyrophosphate synthase. *Plant physiol.* 2011; 155:1079–1090. [PubMed: 21220764]
 15. Hirooka K, Izumi Y, Nakazawa Y, Fukusaki E, Kobayashi A. Functional analysis of two solanesyl diphosphate synthases from *Arabidopsis thaliana*. *Biosci. Biotechnol. Biochem.* 2005; 69:592–601. [PubMed: 15784989]
 16. Zahiri HS, Noghabi KA, Shin YC. Biochemical characterization of the decaprenyl diphosphate synthase of *Rhodobacter sphaeroides* for coenzyme Q10 production. *Appl. Microbiol. Biotechnol.* 2006; 73:796–806. [PubMed: 16896603]
 17. Jones MO, Perez-Fons L, Robertson FP, Bramley PM, Fraser PD. Functional characterization of long-chain prenyl diphosphate synthases from tomato. *Biochem. J.* 2013; 449:729–740. [PubMed: 23126257]
 18. Chang S-Y, Ko T-P, Liang P-H, Wang AH-J. Catalytic mechanism revealed by the crystal structure of undecaprenyl pyrophosphate synthase in complex with sulfate, magnesium, and Triton. *J. Biol. Chem.* 2003; 278:29298–29307. [PubMed: 12756244]
 19. Asai K, Fujisaki S, Nishimura Y, Nishino T, Okada K, Nakagawa T, Kawamukai M, Matsuda H. The identification of *Escherichia coli* ispB (CEL) gene encoding the octaprenyl diphosphate synthase. *Biochem. Biophys. Res. Commun.* 1994; 202:340–345. [PubMed: 8037730]
 20. Collins MD, Pirouz T, Goodfellow M, Minnikin DE. Distribution of menaquinones in actinomycetes and corynebacteria. *J. Gen. Microbiol.* 1977; 100:221–230. [PubMed: 894261]
 21. Barton, LL. Structural and functional relationships in prokaryotes. Springer; New York: 2005. Electron Transport and Coupled Phosphorylation; p. p41321
 22. Kim SB, Goodfellow M. *Streptomyces avermitilis* sp. nov., nom. rev., a taxonomic home for the avermectin-producing streptomycetes. *Int. J. Syst. Evol. Microbiol.* 2002; 52:2011–2014. [PubMed: 12508861]
 23. Ramana VV, Chakravarthy SK, Raj PS, Kumar BV, Shobha E, Ramaprasad EVV, Sasikala C, Ramana CV. Descriptions of *Rhodopseudomonas parapalustris* sp. nov., *Rhodopseudomonas harwoodiae* sp. nov. and *Rhodopseudomonas pseudopalustris* sp. nov., and emended description of *Rhodopseudomonas palustris*. *Int. J. Syst. Evol. Microbiol.* 2012; 62:1790–1798. [PubMed: 21986724]
 24. Lambert MA, Moss CW. Cellular fatty acid compositions and isoprenoid quinone contents of 23 *Legionella* species. *J. Clin. Microbiol.* 1989; 27:465–573. [PubMed: 2715320]
 25. Ramotar K, Conly JM, Chubb H, Louie TJ. Production of menaquinones by intestinal anaerobes. *J. Inf. Diseases.* 1984; 150:213–218. [PubMed: 6470528]
 26. Zhang D, Shrestha B, Li BZ, Tan T. Ubiquinone-10 production using *Argobacterium tunefaciens* gene in *Escherichia coli* by coexpression system. *Mol. Biotechnol.* 2007; 35:1–14. [PubMed: 17401144]
 27. Liu X, Wu H, Ye J, Yuan Q, Zhang H. Cloning and characterization of the ddsA gene encoding decaprenyl diphosphate synthase from *Rhodobacter capsulatus* B10. *Can. J. Microbiol.* 2006; 52:1141–1147. [PubMed: 17473883]

Figure 1.

Product distributions for GI 19551716, GI 67866738 and GI 53711383 for 2 h incubations in buffer containing TX or CHAPS. (A) Production distributions for GI 19551716 in buffer containing 0.1% TX (lane 1), 2.5 mM CHAPS (lane 2), 5.0 mM CHAPS (lane 3). (B) Production distributions for GI 19551716, GI 67866738 and GI 53711383 in buffer containing 0.1% TX (lane 1), 0.05 % TX (lane 2), 0.01 % TX (lane 3), 2.0 mM CHAPS (lane 4). ^aC-terminal His₆ tag.

Figure 2.

Product distributions of GI 16131077, GI 19551716, GI 53711383, and GI 67866738 for 2 h incubations in TX (0.1%, lane 2), CD (1.6 mM, lane 1; 0.2 mM, lane 4), MCD (2 mM, lane 3; 0.2 mM, lane 5). ^aC-terminal His₆ tag. ^bN-terminal His₆ tag.

Figure 3.

Product distributions for GI 15640461, GI 16131077, GI 19551716, GI 39934115, GI 53711383, and GI 67866738 for 2 h incubations. Part A; lane 1, 0.1% TX; lane 2, 2 mM PEG12E; Part B; lane 1, 0.1% TX; lane 2, 2 mM NEG12E; Part C; lane 1, 0.1% TX; lane 3, 2mM HEG12E; lane 4, 2 mM OEG12E; Part D; lane 1, 0.1% TX; lane 2; 2 mM OEG14E. ^aC-terminal His₆ tag. ^bN-terminal His₆ tag.

Figure 4.

Product distributions for GI 1955176, GI 53711383, GI 67866738, and GI 77862362 for 2 h incubations with different ratios of IPP:DMAPP and IPP:FPP. Lane 1, IPP:DMAPP=4:1, 0.1 % TX; Lane 2, IPP:DMAPP=1:2, 0.1 % TX; Lane 3, IPP:DMAPP=1:2, no TX; Lane 4, IPP:FPP=4:1, 0.1 % TX; Lane 5, IPP:FPP=1:2, 0.1 % TX; Lane 6, IPP:FPP=1:2, no TX. ^aC-terminal His₆ tag. ^bN-terminal His₆ tag.

Figure 5.

Product distributions for GI 29831407, GI 39934115, and 67866738 for 2 h incubations in the presence and absence of TX. Part A, IPP:allylic diphosphate = 4:1. Lane 1, FPP, 0.1% TX; Lane 2, FPP, no TX; Lane 3, DMAPP, no TX. Part B, IPP:allylic diphosphate = 4:1. Lane 1, GPP, 0.1% TX; Lane 2, GPP, no TX; Lane 3, DMAPP, no TX.. ^a. C-terminal His₆ tag. ^b. N-terminal His₆ tag.

Figure 6.

Product distributions for 24 h incubations in the presence and absence of TX where IPP:FPP = 20:1. Part A, GI 19551716, GI 29376566, GI 29831407, GI 39934115, 53711383, and GI 67866738; Part B, GI 15640461, GI 16131077, GI 19551716, GI 52842862, and GI 58578715. Lane 1, no TX; Lane 2, 0.1 % TX. ^aC-terminal His₆ tag. ^bN-terminal His₆ tag.

Figure 7.

Relative chain elongation rates for GI 53711383 (Part A) and GI 19551716 (Part B). Buffers contained 50 μ M IPP, 5 μ M FPP, and no detergent, 2 mM OGP, 2 mM NEG12E, or 0.1 % TX. ^aC-terminal His₆ tag.

Figure 8. Product distributions for GI 53711383 (Part A) and GI 19551716 (Part B). Buffers contained 50 μ M IPP, 5 μ M FPP, and none, 2 mM OGP, 2 mM NEG12E, or 0.1 % TX. ^aC-terminal His₆ tag.

Table 1The comparison of *in vitro* and *in vivo* chain-lengths for *E*-PDSs.

GI	Organism	<i>In vitro</i>	<i>In vivo</i> ^a	Refs.
67866738	decaprenyl diphosphate synthase [Agrobacterium tumefaciens]	C ₅₀	C ₅₀ (UQ-10)	19
19551716	geranylgeranyl pyrophosphate synthase [Corynebacterium glutamicum ATCC 13032]	C ₄₅	C ₄₅ (MK-9)	20
53711383	octaprenyl-diphosphate synthase [Bacteroides fragilis YCH46]	C ₅₀ /C ₅₅	C ₅₀ /C ₅₅ (MK-10/11)	21
39934115	farnesyltranstransferase [Rhodopseudomonas palustris CGA009]	C ₅₀	C ₅₀ (UQ-10)	22
29831407	polyprenyl diphosphate synthase [Streptomyces avermitilis MA-4680]	C ₄₅	C ₄₅ (MK-9)	23
29376566	heptaprenyl diphosphate synthase, component II, putative [Enterococcus faecalis V583]	C ₄₅	C ₄₅ (DMK-9)	24
16131077	octaprenyl diphosphate synthase [Escherichia coli str. K-12 substr. MG1655]	C ₄₅ -C ₅₅	C ₄₀ (UQ-8)	25
77862362	decaprenyl diphosphate synthase [Rhodobacter capsulatus]	C ₅₀ -C ₆₀	C ₅₀ (UQ-10)	26
52842862	octaprenyl diphosphate synthase IspB [Legionella pneumophila subsp. pneumophila str. Philadelphia 1]	C ₆₀ -C ₇₀	C ₅₅ -C ₆₅ (UQ-11/12/13)	27
58578715	octaprenyl-diphosphate synthase [Ehrlichia ruminantium str. Welgevonden]	C ₅₅ -C ₆₅	NK	-
15640461	octaprenyl-diphosphate synthase [Vibrio cholerae O1 biovar El Tor str. N16961]	C ₄₅ -C ₆₀	NK	-

^aWith TX.^bUQ: ubiquinone. MK: menaquinone. DMK: demethylmenaquinone. NK: Not known.