Received:
18 September 2013

Revised:
11 October 2013

Accepted:
22 October 2013

doi: 10.1259/bjr.20130599

Cite this article as:

O'Connor KM, Barest G, Moritani T, Sakai O, Mian A. "Dazed and diffused": making sense of diffusion abnormalities in neurologic pathologies. *Br J Radiol* 2013;86:20130599.

REVIEW ARTICLE

"Dazed and diffused": making sense of diffusion abnormalities in neurologic pathologies

¹K M O'CONNOR, MD, ¹G BAREST, MD, ²T MORITANI, MD, PhD, ¹O SAKAI, MD, PhD and ¹A MIAN, MD

¹Department of Radiology, Boston Medical Center, Boston University School of Medicine, Boston, MA, USA

²Department of Radiology, University of Iowa Hospital, Iowa City, IA, USA

Address correspondence to: Dr Asim Mian

E-mail: asim.mian@bmc.org

ABSTRACT

To review diffusion abnormalities seen in diffusion-weighted MRI in neurological pathologies. We examine the clinical significance of the abnormalities in a broad spectrum of neurological diseases and highlight our current understanding of their causes. Diffusion abnormalities seen on diffusion-weighted MRI can play an important role in the diagnosis and follow-up of a broad spectrum of neurological diseases. A thorough understanding of the appearance and significance of these abnormalities is critical in patient management.

Although the best known use of diffusion-weighted imaging (DWI) in MRI of the brain is detection of acute vaso-occlusive infarct, diffusion abnormalities occur in a wide spectrum of neurological pathologies. These diffusion abnormalities provide information that can directly affect the management of a patient, whether identifying pathology that would otherwise go unrecognised, narrowing a broad differential diagnosis, identifying conditions that require emergent intervention, evaluating response to treatment or providing prognostic information about a patient's expected clinical course. In this review, we explain the significance of diffusion abnormalities in a variety of neurological diseases and highlight our current understanding of the causes of these diffusion abnormalities.

ISCHAEMIC

Hypoxic-ischaemic injury

Hypoxic-ischaemic injury to the brain is a result of hypoxia with or without associated ischaemia. In adults, hypoxic-ischaemic injury most commonly occurs in the setting of cardiac arrest, whereas dehydration, perinatal anoxia and child abuse are the most common causes in children [1,2].

In general, the brain is more resilient to hypoxic insults than to insults that involve both hypoxia and ischaemia, and the distribution of diffusion abnormalities can suggest the underlying cause [3]. For instance, with carbon monoxide poisoning, where oxygen delivery to the brain is impaired but blood flow is preserved, diffusion abnormalities primarily involve the globus pallidus bilaterally [3,4]. By contrast, with cardiac arrest, where both oxygen delivery and blood flow to

the brain are impaired, involvement of the cortex, watershed zones, basal ganglia, thalamus and hippocampus is more typical [1–3]. Whatever the cause, the diffusion abnormality is usually bilateral and symmetric [2].

Diffusion abnormalities caused by hypoxic-ischaemic injury can be seen within 1 hour of the acute insult, hours before CT or other MR sequences [1,2,5]. The abnormalities evolve before peaking approximately 2.0–4.5 days after the initial insult [6]. These diffusion abnormalities are thought to be caused, at least in part, by the glutamate excitotoxic pathway where reduced energy-dependent glutamate reuptake within the synaptic clefts results in cytotoxic oedema (Figure 1). The glutamate receptors are concentrated in the dendrites of the cortical and deep grey matter, and this may explain the distribution of diffusion abnormalities in hypoxic-ischaemic injury [1,2].

INFECTIOUS, BACTERIAL

Cerebral abscess

Cerebral abscesses most commonly occur in the setting of sinusitis, otitis/mastoiditis and head trauma [7]. The most commonly presenting symptom is headache, whereas fever is found in only approximately 50% of patients [7]. Although cerebral abscesses can occur anywhere in the brain, they are most likely to occur in the frontal and temporal lobes [7]. Cerebral abscesses are also typically centred at the grey–white junction, a fact that has been attributed to the marked tortuosity of subcortical vessels that may serve as a trap for infectious organisms [8].

Figure 1. Relationship between glutamate and diffusion abnormality. Normally, glutamate concentration in the synaptic cleft is carefully regulated with excess glutamate (yellow circles) removed by adjacent glial cells and by the pre-synaptic neuron itself. However, when pathology leads to excess glutamate in the synaptic cleft, the glutamate can bind excessively to the non-N-methyl-D-aspartate receptors on the post-synaptic cell. This allows excess Na^+ (red circles) to enter the post-synaptic cell and results in cytotoxic oedema with diffusion abnormality.

DWI is useful for distinguishing cerebral abscesses, which can be rapidly fatal, from other less acute causes of peripherally enhancing lesions such as primary and metastatic malignancies [9] (Figure 2). In particular, diffusion restriction within a lesion favours cerebral abscess over necrotic tumour, which is attributed to the increased viscosity and cellularity of pus [9–11]. One notable exception, however, is brain metastases from some adenocarcinomas, particularly small-cell adenocarcinomas, which can also demonstrate restricted diffusion, for reasons that are not well understood [9,12].

Epidural and subdural empyema

Epidural and subdural empyemas are uncommon intracranial infections that usually arise as a complication of sinusitis or mastoiditis [13]. With epidural empyemas, symptoms of headache, fever and meningismus are often mild, and their onset may delay the development of empyema by days to weeks, which may be attributable to the protective nature of the dura [13,14]. By contrast, these same symptoms are fulminant in the case of subdural empyemas and rapidly progress to severe neurological impairment, septic shock and often seizure [13]. Given the association of empyemas with sinusitis, both epidural and subdural empyemas are commonly found adjacent to the frontal sinuses [13] (Figure 3). In contrast to subdural empyemas, which commonly result in cerebral oedema with mass effect, the brain parenchyma adjacent to epidural empyemas is less likely to demonstrate signal abnormalities, probably because the dura is interposed between the empyema and the brain [14].

Figure 2. Images from a 56-year-old male with a cerebral abscess. (a) Axial post-contrast T_1 weighted image demonstrates a lobulated lesion within the right parietal lobe with thick peripheral enhancement (arrow). (b, c) Axial diffusion-weighted imaging ($b=1000$) (b) and apparent diffusion coefficient map (c) images demonstrate restricted diffusion within the lesion (arrows) consistent with a cerebral abscess. Surrounding vasogenic oedema is also noted (arrowhead).

(a)

(b)

(c)

Figure 3. Images from a 13-year-old male with sinusitis complicated by epidural empyema. (a) Sagittal post-contrast T_1 weighted image demonstrates marked mucosal enhancement of the fluid-filled paranasal sinuses consistent with acute sinusitis (arrowhead). An extra-axial peripherally enhancing fluid collection along the frontal convexity consistent with an epidural empyema is noted (arrow). (b, c) Axial diffusion-weighted imaging ($b=1000$) (b) and apparent diffusion coefficient map (c) images demonstrate restricted diffusion within the epidural empyema (arrows).

Although diffusion restriction is characteristic of subdural empyemas, epidural empyemas demonstrate restricted diffusion less consistently [14]. This may be related to the more insidious presentation of patients with epidural empyemas compared with those with subdural empyemas, as chronic pus may restrict diffusion less than fresh pus [14]. Nevertheless, a recent study demonstrated that normalisation of apparent diffusion coefficient (ADC) values within an empyema correlated with clinical improvement of the patient, whereas persistently decreased ADC values correlated with ongoing sepsis [15].

INFECTIOUS, VIRAL

Herpes encephalitis

Herpes encephalitis is the most common cause of sporadic viral encephalitis in the USA [16]. Presenting symptoms include acute onset of fever, headache and seizure [16]. Early diagnosis is essential given that the mortality rate is >70% in untreated patients and that early treatment with acyclovir is one of the few factors that improve prognosis [16,17].

Diffusion-weighted abnormalities in herpes encephalitis include bilateral, asymmetrical regions of diffusion-weighted hyperintensity involving the medial temporal lobes and inferior frontal lobes, with relative sparing of the basal ganglia and white matter [18] (Figure 4). Although, typically, there is corresponding T_2 hyperintensity, several studies have suggested that the diffusion-weighted signal changes are more sensitive than those on conventional MR sequences [19,20]. Moreover, DWI may be more reliable than even polymerase chain reaction testing for the early diagnosis of herpes encephalitis [21].

Diffusion restriction is common in herpes encephalitis [19,20,22], but lesions that demonstrate T_2 shine-through have also been described [20,22]. Diffusion restriction may result from the glutamate excitotoxic pathway, as the concentration of glutamate in the cerebrospinal fluid increases significantly in the setting of herpes encephalitis [23].

INFECTIOUS, PRION

Creutzfeldt-Jakob Disease

Creutzfeldt-Jakob Disease (CJD) is a prion-mediated rapidly progressive dementia that classically presents with ataxia, memory impairment and cognitive decline [24]. The disease takes on different forms, depending on the origin of the abnormal prion protein, including spontaneous, familial and variant CJD. Spontaneous CJD is by far the most common of these forms, accounting for 85% of the cases [24].

Although electroencephalography and cerebrospinal fluid biomarkers have been used to diagnose CJD, MRI with diffusion weighting has been proven to be the most sensitive and specific test for diagnosing CJD [25]. Diffusion restriction involving the bilateral caudate, putamen, pulvinar nuclei of the thalami, and frontotemporoparietal cortex with relative sparing of the globus pallidus and occipital cortex is the earliest imaging change [25–28] (Figure 5). Abnormal T_2 /fluid-attenuated inversion recovery (FLAIR) hyperintensity in the same distribution can also be seen, but these changes typically occur later in the disease and can be subtle compared with the DWI changes [25–27].

Figure 4. Images from a 69-year-old male with herpes encephalitis. (a, b) Axial diffusion-weighted imaging ($b=1000$) (a) and apparent diffusion coefficient (ADC) map (b) images demonstrate bilateral but asymmetric increase in diffusion-weighted signal, right more than left, within the medial temporal lobes (arrows) without evidence of restriction on the ADC map.

(a)

(b)

Moreover, T_2 /FLAIR hyperintensity in the absence of restricted diffusion is suggestive of rapidly progressive dementias other than CJD [25].

Several theories have been proposed to explain why CJD causes diffusion restriction. It is known that the spongiform degeneration caused by CJD results in intraneuronal microvacuoles, and these microvacuoles may result in restricted diffusion [28]. Some have proposed that the prion proteins themselves serve to restrict diffusion by binding water molecules and thereby restricting their diffusion [30]. Others have noted that prion deposition correlates with abnormalities in local glutamate receptor expression, suggesting that the glutamate excitotoxic pathway may also contribute to diffusion restriction [31].

DEVELOPMENTAL/CONGENITAL

Epidermoid cyst

Epidermoid cysts are uncommon developmental/congenital lesions that account for 0.2–1.8% of intracranial neoplasms [32]. The two most common locations for epidermoid cysts are the cerebellopontine angle (40–60%) and fourth ventricle (5–18%) [32].

Epidermoid cysts demonstrate diffusion-weighted hyperintensity, a feature that helps distinguish them from arachnoid cysts. The exact cause for the diffusion-weighted hyperintensity, however, is disputed. Some authors suggest that the hyperintensity represents T_2 shine-through only [32], whereas others suggest that the

Figure 5. Images from a 51-year-old male with Creutzfeldt-Jakob disease. (a, b) Axial diffusion-weighted imaging ($b=1000$) (a) and apparent diffusion coefficient map (b) images demonstrate regions of restricted diffusion involving the caudate nuclei and putamina bilaterally (arrows) as well as the pulvinar of the left thalamus (arrowheads). Reproduced with permission from Springer Science+Business Media and originally published in [29].

(a)

(b)

keratinaceous material within the epidermoid cyst restricts diffusion [33].

INFLAMMATORY/DEVELOPMENTAL

Cholesteatoma

Cholesteatomas are benign but erosive lesions consisting of a keratin-filled sac lined with stratified squamous epithelium [34]. Cholesteatomas are most commonly acquired rather than being congenital and, when acquired, typically arise from the pars flaccida of the tympanic membrane with extension into Prussak's space [34]. Symptoms result from erosion of adjacent structures, including the middle ear ossicles, inner ear and facial nerve [35]. Treatment is generally surgical, although residual or recurrent disease is more frequent, so second-look surgeries within 6–18 months of initial surgery are commonly performed [35].

Cholesteatomas demonstrate increased diffusion-weighted signal for reasons that are not entirely clear (Figure 6). Some authors have proposed that the hyperintensity results from a combination of T_2 shine-through and diffusion restriction [36,37], whereas others have suggested that the hyperintensity reflects T_2 shine-through only [38]. In either case, diffusion-weighted hyperintensity is a useful feature for both the initial detection of cholesteatoma and the detection of residual or recurrent disease in post-surgical patients [35–38]. Post-surgical granulation tissue, in particular, can be distinguished in part from a recurrent or residual cholesteatoma by its lack of diffusion-weighted hyperintensity [35,36,39]. Indeed, newer DWI techniques have shown promise for identifying cholesteatomas <5 mm and may obviate the need for second-look surgeries altogether [35,39,40].

NEOPLASTIC, TYPICALLY BENIGN

Meningioma

Meningiomas are the most common extra-axial neoplasm accounting for 13–26% of primary intracranial neoplasms [41]. Most commonly, meningiomas arise sporadically in patients in their fourth to sixth decades and are discovered incidentally [42].

DWI plays no role in the diagnosis of typical meningiomas and indeed can be somewhat variable [43]. DWI may, however, be useful to differentiate typical meningiomas from atypical and malignant meningiomas that account for approximately 9.6% of all meningiomas and are much more likely to recur [43]. Making the distinction with conventional MRI alone can be challenging because worrisome features such as heterogeneity, perilesional oedema, irregular surface and invasion of adjacent structures can also be seen in typical meningiomas [44]. However, several recent studies have found that atypical or malignant meningiomas can be reliably distinguished from typical meningiomas by their lower ADC values, a fact attributed in part to their increased cellularity [43,45,46].

NEOPLASTIC, MALIGNANT

Glioblastoma

Glioblastoma is an aggressive World Health Organization Grade IV malignancy with a survival rate of less than 1 year [47]. Patients are typically older than 40 years and present with symptoms of headache, memory loss, motor or language deficits, personality change or seizure [47].

Figure 6. Images from a 58-year-old male with recurrent cholesteatoma. (a) Axial post-contrast T_1 weighted image demonstrates a non-enhancing intermediate signal lesion within the left middle ear, adjacent to the site of prior surgery (arrow). (b, c) Axial diffusion-weighted imaging ($b=1000$) (b) and apparent diffusion coefficient map images demonstrate restricted diffusion within the lesion (arrows).

Imaging of glioblastoma typically demonstrates an infiltrating, heterogeneously enhancing lesion with mass effect within the supratentorial white matter that may cross the midline [47]. Central necrosis is a hallmark feature of glioblastoma, attributed to its rapid growth and subsequent tendency to outstrip its blood supply [47]. Owing to these areas of necrosis, up to 88% of glioblastomas demonstrate T_2 shine-through as their predominant feature on diffusion imaging [48]. Although the more cellular components of glioblastomas do correspond to areas with decreased ADC values, these decreases may not be readily visible on qualitative evaluation of the diffusion images.

Diffusion characteristics of glioblastoma are potentially useful in several clinical scenarios. As previously noted, necrotic glioblastomas can be readily differentiated from cerebral abscesses by their lack of diffusion restriction [48]. Also, quantitative analyses of ADC values have shown that ADC values in high-grade gliomas such as glioblastoma are significantly lower than ADC values in low-grade gliomas, a distinction with clear prognostic implications [49,50]. Studies have also suggested that ADC values may help distinguish glioblastoma from lymphoma, as the ADC values of lymphoma tend to be lower than those of glioblastoma [50–52]. More recently, diffusion restriction has been considered in the evaluation of recurring glioblastoma in patients treated with bevacizumab. Interpretation is challenging, however, since new areas of restricted diffusion can represent either tumour recurrence [53] or hypoxia, presumably, secondary to the anti-angiogenic properties of bevacizumab [54].

Lymphoma

Primary central nervous system (CNS) lymphoma accounts for 1–5% of all primary malignancies of the CNS, with diffuse large B-cell non-Hodgkin lymphomas representing >95% of primary CNS lymphomas [55,56]. The mean age at onset in immunocompetent patients is 50–60 years, whereas in immunocompromised patients, the mean age is in the thirties [56,57].

Primary CNS lymphomas in the immunocompetent patient have typical imaging features. Lesions typically demonstrate restricted diffusion on MRI, a finding attributed to the high cellularity of the lymphomas [55,58] (Figure 7). In addition, lesions are solitary in approximately 60–80% of cases [55,56], demonstrate homogeneous enhancement in up to 90% of cases and rarely demonstrate necrosis, haemorrhage or calcification [55,56,59]. Lesions are typically distributed in a periventricular location and may involve the corpus callosum [55,56]. Imaging of primary CNS lymphoma in the immunocompromised patient demonstrates similar findings, although multiplicity of lesions, peripheral enhancement, necrosis and haemorrhage are more often seen [55,59].

Diffusion characteristics of primary CNS lymphoma have several potential clinical applications. First, as described above, quantitative analysis of ADC values may help differentiate primary CNS lymphomas from gliomas. Additionally, a recent study found that the ADC values of untreated primary CNS lymphoma lesions may help distinguish patients who may require whole-brain radiation and prolonged chemotherapy from patients who can simply be followed after a standard chemotherapy regimen [60]. ADC values have also shown promise in distinguishing primary CNS

Figure 7. Images from a 43-year-old male with HIV and central nervous system lymphoma (courtesy of Barest G, Boston, MA). (a) Axial non-contrast head CT demonstrates a large slightly hyperattenuating lesion (arrow) centred in the left frontal lobe with mass effect on the left lateral ventricle. (b, c) Axial diffusion-weighted imaging ($b=1000$) (b) and apparent diffusion coefficient map (c) images demonstrate restricted diffusion within the lesion (arrows).

lymphoma from toxoplasmosis in immunocompromised patients. In particular, a study demonstrated that the ADC values in toxoplasmosis abscesses are significantly higher than those in lymphoma lesions [58].

NEOPLASTIC, MALIGNANT

Metastases

Approximately 10% of cancer patients develop brain metastases [61]. Approximately 80% of metastatic lesions are supratentorial, 15% are infratentorial and 5% occur in the brainstem [62]. Notably, metastatic disease to the brain presents as a solitary lesion approximately half the time [63].

Diffusion-weighted signal is extremely variable in metastatic lesions and therefore generally not useful in their characterisation. For instance, a recent study of 51 patients with metastatic brain lesions found that 18 lesions were hypointense on diffusion-weighted sequences, 17 were hyperintense and 16 were isointense [63]. However, several studies have suggested that ADC values may be useful to distinguish solitary metastases from primary brain malignancies [64–66]. Specifically, although peritumoral oedema is common in both metastases and primary brain malignancies, the ADC values in the peritumoral oedema of metastases have been shown to be higher than those of primary brain malignancies. It is believed that this difference is owing to the more infiltrative nature of primary malignancies, which results in an “infiltrative oedema” as opposed to simple vasogenic oedema [67].

DEMYELINATING

Multiple sclerosis

Multiple sclerosis is a demyelinating disease of young adults that typically presents with a combination of neurological symptoms, the onset of which can be sudden or insidious [68]. In up to 90% of patients with multiple sclerosis, symptoms will remit and relapse early in the disease, although in later years symptoms become slowly progressive in most of these patients [69].

Acute multiple sclerosis plaques typically demonstrate increased diffusion-weighted signal with increased ADC values [37,70,71]. As the plaques age, the ADC values decrease but remain elevated compared with normal white matter [72–74]. In acute cases, the increased ADC values have been attributed to loss of myelin and increased extracellular water in the setting of active inflammation [74]. In chronic cases, after the acute inflammatory process has resolved, extracellular water decreases but still remains increased relative to normal white matter [74].

Rare multiple sclerosis plaques demonstrate restricted diffusion. A recent study of 13 such plaques demonstrated that the majority of the lesions were symptomatic, enhancing and periventricular in location, and all of the lesions ultimately demonstrated increased ADC values on follow-up [75]. Although it is not clear why some multiple sclerosis plaques demonstrate restricted diffusion, it has been observed that glutamate is increased in the cerebrospinal fluid of patients with acute multiple sclerosis and that macrophages and microglia produce high levels of glutamate near areas of axonal damage [76,77]. It has therefore been proposed that this excess glutamate may lead to excitotoxic injury to

oligodendrocytes, myelin sheaths and axons, resulting in cytotoxic plaques [23].

TRAUMATIC

Diffuse axonal injury

Diffuse axonal injury is the result of traumatic acceleration and deceleration injuries that result in “shearing” of brain tissue with different densities or rigidities [78,79]. Patients present with a sudden loss of consciousness that begins at the moment of impact [79].

Imaging findings of diffuse axonal injury are often much less impressive than would be expected, given the patient’s poor neurological status. Initial non-contrast head CT, for instance, may be normal [79]. On MRI, multiple, small, round lesions may be seen at the grey–white matter junction, in the upper dorsolateral brainstem, in the deep grey matter and in the corpus callosum on T_2 /FLAIR, gradient echo and diffusion-weighted sequences [78,80]. Interestingly, no single sequence is likely to demonstrate that all of the lesions as multiple distinct processes lead to the lesions [78].

Lesions that are apparent on diffusion-weighted sequences typically demonstrate restricted diffusion [23,78] (Figure 8). The mechanism for this remains incompletely understood, although a recent theory postulates that diffusion restriction is a result of leakage of glutamate from damaged axons with subsequent cytotoxic injury to adjacent glial cells [23]. Occasionally, lesions demonstrate increased diffusion-weighted signal with normal-to-increased ADC values, possibly representing T_2 shine-through from post-traumatic vasogenic oedema [78,79]. Interestingly, the best imaging-based predictor of neurologic disability may be the total volume of diffusion-weighted signal abnormality seen on MRI, whether restricted or not [78].

TOXIC AND METABOLIC

Heroin toxicity

Heroin abuse can result in a spongiform leukoencephalopathy, particularly when impure heroine fumes are inhaled. Symptoms are typically non-specific, and a mortality rate of 23% has been reported [81].

MRI demonstrates confluent, bilateral, symmetric diffusion-weighted and T_2 /FLAIR hyperintensities in a typical distribution involving the posterior cerebral and cerebellar white matter, cerebellar peduncles, brainstem, splenium of the corpus callosum and posterior limbs of the internal capsules [82,83]. Notably, the diffusion-weighted abnormalities may be detected before T_2 /FLAIR abnormalities [84].

Diffusion restriction is variable in heroin-induced leukoencephalopathy [82–86]. When present, diffusion restriction has been attributed to vacuolisation within preserved myelin sheaths, leading to the entrapment of water molecules [30,84,86]. At least one study has proposed that the coalescing of smaller vacuoles into larger vacuoles as the disease progresses may increase diffusibility of water molecules over time and thereby account for the apparent variability in diffusion restriction [85].

Figure 8. Images from a 6-year-old male with diffuse axonal injury after being struck by a car. (a, b) Axial diffusion-weighted imaging ($b=1000$) (a) and apparent diffusion coefficient map (b) images demonstrate a representative punctate lesion in the splenium of the corpus callosum with restricted diffusion (arrows). Additional lesions were seen in the body of the corpus callosum and at the grey-white matter junction in the frontal lobe (not shown).

(a)

(b)

Methotrexate toxicity

Intrathecal and intravenous administration of methotrexate is used to treat CNS leukaemia because of its ability to cross the blood-brain barrier, and 3–10% of treated patients develop methotrexate-induced leukoencephalopathy [87]. Typical symptoms mimic those of ischaemic infarct [87] but resolve within 1 or 2 days, often spontaneously [88]. Symptom onset may occur as early 2 weeks or as late as 2 years after treatment [87].

DWI of methotrexate-induced leukoencephalopathy demonstrates confluent areas of restricted diffusion involving the periventricular white matter [89] and occasionally the cerebellar

Figure 9. Images from a 50-year-old female with methotrexate toxicity. (a, b) Axial diffusion-weighted imaging ($b=1000$) (a) and apparent diffusion coefficient map (b) images demonstrate regions of restricted diffusion involving the corona radiata (arrows). No corresponding signal abnormality was detected on T_2 weighted images (c). Reproduced with permission from Springer Science+Business Media and originally published in [29].

(a)

(b)

(c)

Figure 10. Images from a 38-year-old female with a history of alcoholism, malnutrition and hepatic failure presenting with osmotic myelinolysis (courtesy of Barest G, Boston, MA). (a, b) Axial diffusion-weighted imaging ($b=1000$) (a) and apparent diffusion coefficient map (b) images demonstrate a region of restricted diffusion centred in the right pons (arrows) with sparing of the right corticospinal tract (arrowheads). (c) Axial fluid-attenuated inversion recovery image from follow-up MRI performed 2 weeks later demonstrates the typical “owl-eye” appearance of the pons seen in osmotic myelinolysis.

white matter and cerebral cortex [87] (Figure 9). T_2 images may be normal initially but subsequently demonstrate hyperintensity [87]. Interestingly, the diffusion restriction is typically transient with diffusion signal abnormalities partially or completely resolving within days of symptom resolution [88–91]. This has led several authors to speculate that the underlying cause of diffusion restriction in methotrexate-induced leukoencephalopathy is reversible cytotoxic oedema within myelin sheaths but with no underlying structural damage [88–91], a process that is probably mediated by glutamate [23,92].

Osmotic myelinolysis

Patients with osmotic myelinolysis may present with confusion, quadriplegia, pseudobulbar palsy or locked-in syndrome [93]. Osmotic myelinolysis classically occurs in the setting of rapidly corrected severe hyponatremia of several days duration [94]. Conditions associated with osmotic myelinolysis include alcoholism, alcohol withdrawal, malnutrition, liver transplantation, cirrhosis and severe burns [95]. Prognosis is quite variable, with approximately one-third of patients recovering completely, one-third having mild residual neurologic deficits and one-third remaining severely neurologically impaired [96].

Osmotic myelinolysis initially demonstrates symmetrical regions of restricted diffusion that may involve the central pons, cerebellum, basal ganglia, lateral geniculate bodies, external and extreme capsule, hippocampus or cerebral cortex [97,98]. When the pons is involved, it demonstrates a characteristically symmetric region of restricted diffusion involving the central pons that spares both the periphery of the pons and the corticospinal tracts [98] (Figure 10). This appearance has been likened to “owl eyes”, and helps distinguish osmotic myelinolysis from a pontine infarct, which is typically wedge-shaped and does not spare the periphery of the pons.

Diffusion restriction in the affected areas can be seen within 24 h of the symptom onset [98]. In some studies, diffusion abnormalities were seen days to weeks before abnormalities appeared on conventional MRI sequences [98]. Over the course of weeks, the diffusion abnormalities may completely resolve or demonstrate T_2 shine-through [95,98,99]. Resolution of diffusion abnormalities

may predict a better prognosis for the patient [99], although this is controversial [96].

The cause of diffusion abnormalities in osmotic myelinolysis is incompletely understood. One hypothesis is that early diffusion restriction may represent intramyelinic cytotoxic oedema that subsequently evolves into vasogenic oedema as myelin is destroyed [95].

OTHER

Posterior reversible encephalopathy syndrome

Patients with posterior reversible encephalopathy syndrome typically present with a combination of headache, seizure, visual changes and altered mental status [100]. Although the diagnosis is commonly considered in pre- or post-partum women with hypertension, it can occur in other situations, including post-transplant toxicity to cyclosporine A [100]. The underlying cause is thought to be dysfunctional autoregulation in the posterior circulation, resulting in increased permeability of the capillaries, leakage of fluid into the interstitium and subsequent vasogenic oedema [100].

Imaging typically demonstrates normal to slightly increased diffusion-weighted signals with markedly increased ADC values consistent with vasogenic oedema [98,99]. These changes in signal are typically bilateral and involve the cortical and sub-cortical regions of the occipital and posterior parietal lobes, but they can also involve the frontal lobes, temporal lobes, cerebellum, brainstem and bilateral basal ganglia [100,101]. Mild mass effect and similarly distributed parenchymal enhancement may also be seen [100]. The diffusion abnormality normally resolves once the underlying cause is controlled; however, if left untreated the syndrome can progress to cytotoxic oedema with massive infarction and death [100].

CONCLUSION

The appearance and distribution of diffusion abnormalities on diffusion-weighted MR images can play an important role in the diagnosis and follow-up of a broad spectrum of neurological diseases. A thorough understanding of these abnormalities is critical to patient management.

REFERENCES

- Arbelaez A, Castillo M, Mukherji SK. Diffusion-weighted MR imaging of global cerebral anoxia. *AJNR Am J Neuroradiol* 1999;20:999–1007.
- Gutierrez LG, Rovira A, Portela LA, Leite Cda C, Lucato LT. CT and MR in non-neonatal hypoxic-ischemic encephalopathy: radiological findings with pathophysiological correlations. *Neuroradiology* 2010;52:949–76.
- Singhal AB, Topcuoglu MA, Koroshetz WJ. Diffusion MRI in three types of anoxic encephalopathy. *J Neurol Sci* 2002;196:37–40.
- Prockop LD, Naidu KA. Brain CT and MRI findings after carbon monoxide toxicity. *J Neuroimaging* 1999;9:175–81.
- Wijdicks EF, Campeau NG, Miller GM. MR imaging in comatose survivors of cardiac resuscitation. *AJNR Am J Neuroradiol* 2001;22:1561–5.
- Wijman CA, Mlynash M, Caulfield AF, Hsia AW, Eyngorn I, Bammer R, et al. Prognostic value of brain diffusion-weighted imaging after cardiac arrest. *Ann Neurol* 2009;65:394–402. doi: 10.1002/ana.21632
- Nathoo N, Nadvi SS, Narotam PK, van Dellen JR. Brain abscess: management and outcome analysis of a computed tomography era experience with 973 patients. *World Neurosurg* 2011;75:716–26; discussion 612–717. doi: 10.1016/j.wneu.2010.11.043
- Nonaka H, Akima M, Hatori T, Nagayama T, Zhang Z, Ihara F. The microvasculature of the cerebral white matter: arteries of the subcortical white matter. *J Neuropathol Exp Neurol* 2003;62:154–61.

9. Hartmann M, Jansen O, Heiland S, Sommer C, Munkel K, Sartor K. Restricted diffusion within ring enhancement is not pathognomonic for brain abscess. *AJNR Am J Neuroradiol* 2001;22:1738–42.
10. Desprechins B, Stadnik T, Koerts G, Shabana W, Breucq C, Osteaux M. Use of diffusion-weighted MR imaging in differential diagnosis between intracerebral necrotic tumors and cerebral abscesses. *AJNR Am J Neuroradiol* 1999;20:1252–7.
11. Noguchi K, Watanabe N, Nagayoshi T, Kanazawa T, Toyoshima S, Shimizu M, et al. Role of diffusion-weighted echo-planar MRI in distinguishing between brain brain abscess and tumour: a preliminary report. *Neuroradiology* 1999;41:171–4.
12. Hayashida Y, Hirai T, Morishita S, Kitajima M, Murakami R, Korogi Y, et al. Diffusion-weighted imaging of metastatic brain tumors: comparison with histologic type and tumor cellularity. *AJNR Am J Neuroradiol* 2006;27:1419–25.
13. Osborn MK, Steinberg JP. Subdural empyema and other suppurative complications of paranasal sinusitis. *Lancet Infect Dis* 2007;7:62–7. doi: [10.1016/S1473-3099\(06\)70688-0](https://doi.org/10.1016/S1473-3099(06)70688-0)
14. Tsuchiya K, Osawa A, Katase S, Fujikawa A, Hachiya J, Aoki S. Diffusion-weighted MRI of subdural and epidural empyemas. *Neuroradiology* 2003;45:220–3. doi: [10.1007/s00234-003-0949-5](https://doi.org/10.1007/s00234-003-0949-5)
15. Fanning NF, Laffan EE, Shroff MM. Serial diffusion-weighted MRI correlates with clinical course and treatment response in children with intracranial pus collections. *Pediatr Radiol* 2006;36:26–37. doi: [10.1007/s00247-005-0019-8](https://doi.org/10.1007/s00247-005-0019-8)
16. Whitley RJ. Viral encephalitis. *N Engl J Med* 1990;323:242–50. doi: [10.1056/NEJM199007263230406](https://doi.org/10.1056/NEJM199007263230406)
17. Raschilas F, Wolff M, Delatour F, Chaffaut C, De Broucker T, Chevret S, et al. Outcome of and prognostic factors for herpes simplex encephalitis in adult patients: results of a multicenter study. *Clin Infect Dis* 2002;35:254–60. doi: [10.1086/341405](https://doi.org/10.1086/341405)
18. Kapur N, Barker S, Burrows EH, Ellison D, Brice J, Illis LS, et al. Herpes simplex encephalitis: long term magnetic resonance imaging and neuropsychological profile. *J Neurol Neurosurg Psychiatry* 1994;57:1334–42.
19. Kuker W, Nagele T, Schmidt F, Heckl S, Herrlinger U. Diffusion-weighted MRI in herpes simplex encephalitis: a report of three cases. *Neuroradiology* 2004;46:122–5. doi: [10.1007/s00234-003-1145-3](https://doi.org/10.1007/s00234-003-1145-3)
20. Heiner L, Demaerel P. Diffusion-weighted MR imaging findings in a patient with herpes simplex encephalitis. *Eur J Radiol* 2003;45:195–8.
21. Akylidz BN, Gumus H, Kumandas S, Coskun A, Karakukucu M, Yikmaz A. Diffusion-weighted magnetic resonance is better than polymerase chain reaction for early diagnosis of herpes simplex encephalitis: a case report. *Pediatr Emerg Care* 2008;24:377–9. doi: [10.1097/PEC.0b013e318177a712](https://doi.org/10.1097/PEC.0b013e318177a712)
22. Sener RN. Herpes simplex encephalitis: diffusion MR imaging findings. *Comput Med Imaging Graph* 2001;25:391–7.
23. Moritani T, Smoker WR, Sato Y, Numaguchi Y, Westesson PL. Diffusion-weighted imaging of acute excitotoxic brain injury. *AJNR Am J Neuroradiol* 2005;26:216–28.
24. Appleby BS, Appleby KK, Crain BJ, Onyike CU, Wallin MT, Rabins PV. Characteristics of established and proposed sporadic Creutzfeldt-Jakob disease variants. *Arch Neurol* 2009;66:208–15. doi: [10.1001/archneurol.2008.533](https://doi.org/10.1001/archneurol.2008.533)
25. Vitali P, Maccagnano E, Caverzasi E, Henry RG, Haman A, Torres-Chae C, et al. Diffusion-weighted MRI hyperintensity patterns differentiate CJD from other rapid dementias. *Neurology* 2011;76:1711–19. doi: [10.1212/WNL.0b013e31821a4439](https://doi.org/10.1212/WNL.0b013e31821a4439)
26. Ukisu R, Kushihashi T, Kitanosono T, Fujisawa H, Takenaka H, Ohgiya Y, et al. Serial diffusion-weighted MRI of Creutzfeldt-Jakob disease. *AJR Am J Roentgenol* 2005;184:560–6. doi: [10.2214/ajr.184.2.01840560](https://doi.org/10.2214/ajr.184.2.01840560)
27. Murata T, Shiga Y, Higano S, Takahashi S, Mugikura S. Conspicuity and evolution of lesions in Creutzfeldt-Jakob disease at diffusion-weighted imaging. *AJNR Am J Neuroradiol* 2002;23:1164–72.
28. Mittal S, Farmer P, Kalina P, Kingsley PB, Halperin J. Correlation of diffusion-weighted magnetic resonance imaging with neuropathology in Creutzfeldt-Jakob disease. *Arch Neurol* 2002;59:128–34.
29. Moritani T, Ekholm S, Westesson P-LA. Diffusion-weighted MR imaging of the brain. 2nd edn. Berlin, Germany: Springer; 2009.
30. Geschwind MD, Potter CA, Sattavat M, Garcia PA, Rosen HJ, Miller BL, et al. Correlating DWI MRI with pathologic and other features of Jakob-Creutzfeldt disease. *Alzheimer Dis Assoc Disord* 2009;23:82–7.
31. Ferrer I, Puig B. GluR2/3, NMDAepsilon1 and GABAA receptors in Creutzfeldt-Jakob disease. *Acta Neuropathol* 2003;106:311–18. doi: [10.1007/s00401-003-0732-z](https://doi.org/10.1007/s00401-003-0732-z)
32. Hakyemez B, Aksoy U, Yildiz H, Ergin N. Intracranial epidermoid cysts: diffusion-weighted, FLAIR and conventional MR findings. *Eur J Radiol* 2005;54:214–20. doi: [10.1016/j.ejrad.2004.06.018](https://doi.org/10.1016/j.ejrad.2004.06.018)
33. Annet L, Duprez T, Grandin C, Dooms G, Collard A, Cosnard G. Apparent diffusion coefficient measurements within intracranial epidermoid cysts in six patients. *Neuroradiology* 2002;44:326–8.
34. Som PM, Curtin HD. Head and neck imaging. 4th edn. St Louis, MO: Mosby; 2003.
35. Schwartz KM, Lane JJ, Bolster BD Jr, Neff BA. The utility of diffusion-weighted imaging for cholesteatoma evaluation. *AJNR Am J Neuroradiol* 2011;32:430–6. doi: [10.3174/ajnr.A2129](https://doi.org/10.3174/ajnr.A2129)
36. Aikele P, Kittner T, Offergeld C, Kaftan H, Huttenbrink KB, Laniado M. Diffusion-weighted MR imaging of cholesteatoma in pediatric and adult patients who have undergone middle ear surgery. *AJR Am J Roentgenol* 2003;181:261–5. doi: [10.2214/ajr.181.1.1810261](https://doi.org/10.2214/ajr.181.1.1810261)
37. Schaefer PW, Grant PE, Gonzalez RG. Diffusion-weighted MR imaging of the brain. *Radiology* 2000;217:331–45. doi: [10.1148/radiology.217.2.r00nv24331](https://doi.org/10.1148/radiology.217.2.r00nv24331)
38. Vercruysse JP, De Foer B, Pouillon M, Somers T, Casselman J, Offeciers E. The value of diffusion-weighted MR imaging in the diagnosis of primary acquired and residual cholesteatoma: a surgical verified study of 100 patients. *Eur Radiol* 2006;16:1461–7. doi: [10.1007/s00330-006-0160-2](https://doi.org/10.1007/s00330-006-0160-2)
39. Sharifian H, Taheri E, Borghei P, Shakiba M, Jalali AH, Roshanfekar M, et al. Diagnostic accuracy of non-echo-planar diffusion-weighted MRI versus other MRI sequences in cholesteatoma. *J Med Imaging Radiat Oncol* 2012;56:398–408. doi: [10.1111/j.1754-9485.2012.02377.x](https://doi.org/10.1111/j.1754-9485.2012.02377.x)
40. Dremmen MH, Hofman PA, Hof JR, Stokroos RJ, Postma AA. The diagnostic accuracy of non-echo-planar diffusion-weighted imaging in the detection of residual and/or recurrent cholesteatoma of the temporal bone. *AJNR Am J Neuroradiol* 2012;33:439–4. doi: [10.3174/ajnr.A2824](https://doi.org/10.3174/ajnr.A2824)
41. Kleihues P, Cavenee WK, International Agency for Research on Cancer. Pathology and genetics of tumours of the nervous system. Lyon, France: IARC Press; 2000: 314 p.
42. Rockhill J, Mrugala M, Chamberlain MC. Intracranial meningiomas: an overview of diagnosis and treatment. *Neurosurg Focus* 2007;23:E1. doi: [10.3171/FOC-07/10/E1](https://doi.org/10.3171/FOC-07/10/E1)
43. Hakyemez B, Yildirim N, Gokalp G, Erdogan C, Parlak M. The contribution of diffusion-weighted MR imaging to distinguishing typical from atypical

- meningiomas. *Neuroradiology* 2006;48:513–20. doi: [10.1007/s00234-006-0094-z](https://doi.org/10.1007/s00234-006-0094-z)
44. Sanverdi SE, Ozgen B, Oguz KK, Mut M, Dolgun A, Soylemezoglu F, et al. Is diffusion-weighted imaging useful in grading and differentiating histopathological subtypes of meningiomas? *Eur J Radiol* 2012;81:2389–95. doi: [10.1016/j.ejrad.2011.06.031](https://doi.org/10.1016/j.ejrad.2011.06.031)
 45. Yin B, Liu L, Zhang BY, Li YX, Li Y, Geng DY. Correlating apparent diffusion coefficients with histopathologic findings on meningiomas. *Eur J Radiol* 2012;81:4050–6. doi: [10.1016/j.ejrad.2012.06.002](https://doi.org/10.1016/j.ejrad.2012.06.002)
 46. Nagar VA, Ye JR, Ng WH, Chan YH, Hui F, Lee CK, et al. Diffusion-weighted MR imaging: diagnosing atypical or malignant meningiomas and detecting tumor dedifferentiation. *AJNR Am J Neuroradiol* 2008;29:1147–52. doi: [10.3174/ajnr.A0996](https://doi.org/10.3174/ajnr.A0996)
 47. Dean BL, Drayer BP, Bird CR, Flom RA, Hodkar JA, Coons SW, et al. Gliomas: classification with MR imaging. *Radiology* 1990;174:411–15. doi: [10.1148/radiology.174.2.2153310](https://doi.org/10.1148/radiology.174.2.2153310)
 48. Hakymez B, Erdogan C, Yildirim N, Parlak M. Glioblastoma multiforme with atypical diffusion-weighted MR findings. *Br J Radiol* 2005;78:989–92. doi: [10.1259/bjr/12830378](https://doi.org/10.1259/bjr/12830378)
 49. Kono K, Inoue Y, Nakayama K, Shakudo M, Morino M, Ohata K, et al. The role of diffusion-weighted imaging in patients with brain tumors. *AJNR Am J Neuroradiol* 2001;22:1081–8.
 50. Kitis O, Altay H, Calli C, Yuntun N, Akalin T, Yurtseven T. Minimum apparent diffusion coefficients in the evaluation of brain tumors. *Eur J Radiol* 2005;55:393–400. doi: [10.1016/j.ejrad.2005.02.004](https://doi.org/10.1016/j.ejrad.2005.02.004)
 51. Stadnik TW, Chaskis C, Michotte A, Shabana WM, van Rompaey K, Luybaert R, et al. Diffusion-weighted MR imaging of intracerebral masses: comparison with conventional MR imaging and histologic findings. *AJNR Am J Neuroradiol* 2001;22:969–76.
 52. Guo AC, Cummings TJ, Dash RC, Provenzale JM. Lymphomas and high-grade astrocytomas: comparison of water diffusibility and histologic characteristics. *Radiology* 2002;224:177–83. doi: [10.1148/radiol.2241010637](https://doi.org/10.1148/radiol.2241010637)
 53. Gerstner ER, Frosch MP, Batchelor TT. Diffusion magnetic resonance imaging detects pathologically confirmed, non-enhancing tumor progression in a patient with recurrent glioblastoma receiving bevacizumab. *J Clin Oncol* 2010;28:e91–93.
 54. Rieger J, Bahr O, Muller K, Franz K, Steinbach J, Hattingen E. Bevacizumab-induced diffusion-restricted lesions in malignant glioma patients. *J Neurooncol* 2010;99:49–56. doi: [10.1007/s11060-009-0098-8](https://doi.org/10.1007/s11060-009-0098-8)
 55. Haldorsen IS, Espeland A, Larsson EM. Central nervous system lymphoma: characteristic findings on traditional and advanced imaging. *AJNR Am J Neuroradiol* 2011;32:984–92. doi: [10.3174/ajnr.A2171](https://doi.org/10.3174/ajnr.A2171)
 56. Yap KK, Sutherland T, Liew E, Tartaglia CJ, Pang M, Trost N. Magnetic resonance features of primary central nervous system lymphoma in the immunocompetent patient: a pictorial essay. *J Med Imaging Radiat Oncol* 2012;56:179–86.
 57. Kongsiriwattanukul S, Suankratay C. Central nervous system infections in HIV-infected patients hospitalized at King Chulalongkorn Memorial Hospital. *J Med Assoc Thai* 2011;94:551–8.
 58. Camacho DL, Smith JK, Castillo M. Differentiation of toxoplasmosis and lymphoma in AIDS patients by using apparent diffusion coefficients. *AJNR Am J Neuroradiol* 2003;24:633–7.
 59. Berger JR. Mass lesions of the brain in AIDS: the dilemmas of distinguishing toxoplasmosis from primary CNS lymphoma. *AJNR Am J Neuroradiol* 2003;24:554–5.
 60. Barajas RF Jr, Rubenstein JL, Chang JS, Hwang J, Cha S. Diffusion-weighted MR imaging derived apparent diffusion coefficient is predictive of clinical outcome in primary central nervous system lymphoma. *AJNR Am J Neuroradiol* 2010;31:60–6. doi: [10.3174/ajnr.A1750](https://doi.org/10.3174/ajnr.A1750)
 61. Kamar FG, Posner JB. Brain metastases. *Semin Neurol* 2010;30:217–35. doi: [10.1055/s-0030-1255225](https://doi.org/10.1055/s-0030-1255225)
 62. Christiaans MH, Kelder JC, Arnoldus EP, Tijssen CC. Prediction of intracranial metastases in cancer patients with headache. *Cancer* 2002;94:2063–8.
 63. Gaudino S, Di Lella GM, Russo R, Lo Russo VS, Piludu F, Quaglio FR, et al. Magnetic resonance imaging of solitary brain metastases: main findings of nonmorphological sequences. *Radiol Med* 2012;117:1225–41.
 64. Chiang IC, Kuo YT, Lu CY, Yeung KW, Lin WC, Sheu FO, et al. Distinction between high-grade gliomas and solitary metastases using peritumoral 3-T magnetic resonance spectroscopy, diffusion, and perfusion imagings. *Neuroradiology* 2004;46:619–27.
 65. Krabbe K, Gideon P, Wagn P, Hansen U, Thomsen C, Madsen F. MR diffusion imaging of human intracranial tumours. *Neuroradiology* 1997;39:483–9.
 66. Rollin N, Guyotat J, Streichenberger N, Honnorat J, Tran Minh VA, Cotton F. Clinical relevance of diffusion and perfusion magnetic resonance imaging in assessing intra-axial brain tumors. *Neuroradiology* 2006;48:150–9. doi: [10.1007/s00234-005-0030-7](https://doi.org/10.1007/s00234-005-0030-7)
 67. Lee EJ, terBrugge K, Mikulis D, Choi DS, Bae JM, Lee SK, et al. Diagnostic value of peritumoral minimum apparent diffusion coefficient for differentiation of glioblastoma multiforme from solitary metastatic lesions. *AJR Am J Roentgenol* 2011;196:71–6. doi: [10.2214/AJR.10.4752](https://doi.org/10.2214/AJR.10.4752)
 68. Calabresi PA. Diagnosis and management of multiple sclerosis. *Am Fam Physician* 2004;70:1935–44.
 69. Lublin FD, Reingold SC. Defining the clinical course of multiple sclerosis: results of an international survey. National Multiple Sclerosis Society (USA) Advisory Committee on Clinical Trials of New Agents in Multiple Sclerosis. *Neurology* 1996;46:907–11.
 70. Horsfield MA, Larsson HB, Jones DK, Gass A. Diffusion magnetic resonance imaging in multiple sclerosis. *J Neurol Neurosurg Psychiatry* 1998;64(Suppl 1):S80–84.
 71. Castriota-Scanderbeg A, Sabatini U, Fasano F, Floris R, Fraracci L, Mario MD, et al. Diffusion of water in large demyelinating lesions: a follow-up study. *Neuroradiology* 2002;44:764–7. doi: [10.1007/s00234-002-0806-y](https://doi.org/10.1007/s00234-002-0806-y)
 72. Christiansen P, Gideon P, Thomsen C, Stubgaard M, Henriksen O, Larsson HB. Increased water self-diffusion in chronic plaques and in apparently normal white matter in patients with multiple sclerosis. *Acta Neurol Scand* 1993;87:195–9.
 73. Larsson HB, Thomsen C, Frederiksen J, Stubgaard M, Henriksen O. In vivo magnetic resonance diffusion measurement in the brain of patients with multiple sclerosis. *Magn Reson Imaging* 1992;10:7–12.
 74. Tievsky AL, Ptak T, Farkas J. Investigation of apparent diffusion coefficient and diffusion tensor anisotropy in acute and chronic multiple sclerosis lesions. *AJNR Am J Neuroradiol* 1999;20:1491–9.
 75. Balashov KE, Lindzen E. Acute demyelinating lesions with restricted diffusion in multiple sclerosis. *Mult Scler* 2012;18:1745–53. doi: [10.1177/1352458512445407](https://doi.org/10.1177/1352458512445407)
 76. Stover JF, Pleines UE, Morganti-Kossmann MC, Kossmann T, Lowitzsch K, Kempfski OS. Neurotransmitters in cerebrospinal fluid reflect pathological activity. *Eur J Clin Invest* 1997;27:1038–43.
 77. Werner P, Pitt D, Raine CS. Multiple sclerosis: altered glutamate homeostasis in lesions correlates with oligodendrocyte and axonal damage. *Ann Neurol* 2001;50:169–80.

78. Schaefer PW, Huisman TA, Sorensen AG, Gonzalez RG, Schwamm LH. Diffusion-weighted MR imaging in closed head injury: high correlation with initial glasgow coma scale score and score on modified Rankin scale at discharge. *Radiology* 2004;233: 58–66. doi: [10.1148/radiol.2323031173](https://doi.org/10.1148/radiol.2323031173)
79. Kinoshita T, Moritani T, Hiwatashi A, Wang HZ, Shrier DA, Numaguchi Y, et al. Conspicuity of diffuse axonal injury lesions on diffusion-weighted MR imaging. *Eur J Radiol* 2005;56:5–11. doi: [10.1016/j.ejrad.2005.04.001](https://doi.org/10.1016/j.ejrad.2005.04.001)
80. Liu AY, Maldjian JA, Bagley LJ, Sinson GP, Grossman RI. Traumatic brain injury: diffusion-weighted MR imaging findings. *AJNR Am J Neuroradiol* 1999;20:1636–41.
81. Wolters EC, van Wijngaarden GK, Stam FC, Rengelink H, Lousberg RJ, Schipper ME, et al. Leukoencephalopathy after inhaling “heroin” pyrolysate. *Lancet* 1982;2:1233–7.
82. Offiah C, Hall E. Heroin-induced leukoencephalopathy: characterization using MRI, diffusion-weighted imaging, and MR spectroscopy. *Clin Radiol* 2008;63:146–52. doi: [10.1016/j.crad.2007.07.021](https://doi.org/10.1016/j.crad.2007.07.021)
83. Jee RC, Tsao WL, Shyu WC, Yen PS, Hsu YH, Liu SH. Heroin vapor inhalation-induced spongiform leukoencephalopathy. *J Formos Med Assoc* 2009;108:518–22. doi: [10.1016/S0929-6646\(09\)60101-7](https://doi.org/10.1016/S0929-6646(09)60101-7)
84. Bega DS, McDaniel LM, Jhaveri MD, Lee VH. Diffusion weighted imaging in heroin-associated spongiform leukoencephalopathy. *Neurocrit Care* 2009;10:352–4. doi: [10.1007/s12028-008-9172-x](https://doi.org/10.1007/s12028-008-9172-x)
85. Bartlett E, Mikulis DJ. Chasing “chasing the dragon” with MRI: leukoencephalopathy in drug abuse. *Br J Radiol* 2005;78:997–1004. doi: [10.1259/bjr/61535842](https://doi.org/10.1259/bjr/61535842)
86. Chen CY, Lee KW, Lee CC, Chin SC, Chung HW, Zimmerman RA. Heroin-induced spongiform leukoencephalopathy: value of diffusion MR imaging. *J Comput Assist Tomogr* 2000;24:735–7.
87. Rollins N, Winick N, Bash R, Booth T. Acute methotrexate neurotoxicity: findings on diffusion-weighted imaging and correlation with clinical outcome. *AJNR Am J Neuroradiol* 2004;25: 1688–95.
88. Salkade PR, Lim TA. Methotrexate-induced acute toxic leukoencephalopathy. *J Cancer Res Ther* 2012;8:292–6. doi: [10.4103/0973-1482.98993](https://doi.org/10.4103/0973-1482.98993)
89. Baehring JM, Fulbright RK. Delayed leukoencephalopathy with stroke-like presentation in chemotherapy recipients. *J Neurosurg Psychiatry* 2008;79:535–9. doi: [10.1136/jnnp.2007.123737](https://doi.org/10.1136/jnnp.2007.123737)
90. Haykin ME, Gorman M, van Hoff J, Fulbright RK, Baehring JM. Diffusion-weighted MRI correlates of subacute methotrexate-related neurotoxicity. *J Neurooncol* 2006;76:153–7.
91. Balin J, Parmar H, Kujawski L. Conventional and diffusion-weighted MRI findings of methotrexate related sub-acute neurotoxicity. *J Neurol Sci* 2008;269:169–71. doi: [10.1016/j.jns.2007.12.012](https://doi.org/10.1016/j.jns.2007.12.012)
92. Weller M, Marini AM, Finiels-Marlier F, Martin B, Paul SM. MK-801 and memantine protect cultured neurons from glutamate toxicity induced by glutamate carboxypeptidase-mediated cleavage of methotrexate. *Eur J Pharmacol* 1993;248: 303–12.
93. Wright DG, Laurenro R, Victor M. Pontine and extrapontine myelinolysis. *Brain* 1979; 102:361–85.
94. Sterns RH, Cappuccino JD, Silver SM, Cohen EP. Neurologic sequelae after treatment of severe hyponatremia: a multicenter perspective. *J Am Soc Nephrol* 1994;4:1522–30.
95. Forster A, Nolte I, Wenz H, Al-Zghloul M, Kerl HU, Brockmann C, et al. Value of diffusion-weighted imaging in central pontine and extrapontine myelinolysis. *Neuroradiology* 2013;55:49–56. doi: [10.1007/s00234-012-1083-z](https://doi.org/10.1007/s00234-012-1083-z)
96. Menger H, Jorg J. Outcome of central pontine and extrapontine myelinolysis (n = 44). *J Neurol* 1999;246:700–5.
97. Gocht A, Colmant HJ. Central pontine and extrapontine myelinolysis: a report of 58 cases. *Clin Neuropathol* 1987;6: 262–70.
98. Ruzek KA, Campeau NG, Miller GM. Early diagnosis of central pontine myelinolysis with diffusion-weighted imaging. *AJNR Am J Neuroradiol* 2004;25: 210–13.
99. Dervisoglu E, Yegenaga I, Anik Y, Sengul E, Turgut T. Diffusion magnetic resonance imaging may provide prognostic information in osmotic demyelination syndrome: report of a case. *Acta Radiol* 2006;47: 208–12.
100. Covarrubias DJ, Luetmer PH, Campeau NG. Posterior reversible encephalopathy syndrome: prognostic utility of quantitative diffusion-weighted MR images. *AJNR Am J Neuroradiol* 2002;23: 1038–48.
101. Donmez FY, Basaran C, Kayahan Ulu EM, Yildirim M, Coskun M. MRI features of posterior reversible encephalopathy syndrome in 33 patients. *J Neuroimaging* 2010;20:22–8. doi: [10.1111/j.1552-6569.2008.00306.x](https://doi.org/10.1111/j.1552-6569.2008.00306.x)