Published in final edited form as:

Cerebellum. 2012 June ; 11(2): . doi:10.1007/s12311-011-0347-1.

Parasagittal Zones in the Cerebellar Cortex Differ in Excitability, Information Processing, and Synaptic Plasticity

Timothy J. Ebner, Xinming Wang, Wangcai Gao, Samuel W. Cramer, and Gang Chen

Department of Neuroscience, University of Minnesota, Lions Research Building, Room 421, 2001 Sixth Street SE, Minneapolis, MN 55455, USA

Timothy J. Ebner: ebner001@umn.edu

Abstract

At the molecular and circuitry levels, the cerebellum exhibits a striking parasagittal zonation as exemplified by the spatial distribution of molecules expressed on Purkinje cells and the topography of the afferent and efferent projections. The physiology and function of the zonation is less clear. Activity-dependent optical imaging has proven a useful tool to examine the physiological properties of the parasagittal zonation in the intact animal. Recent findings show that zebrin II-positive and zebrin II-negative zones differ markedly in their responses to parallel fiber inputs. These findings suggest that cerebellar cortical excitability, information processing, and synaptic plasticity depend on the intrinsic properties of different parasagittal zones.

Keywords

Parasagittal zones; Cerebellar cortex; Parallel fibers; Purkinje cells; Zebrin II

Introduction

Purkinje cells (PCs), the sole output neurons of the cerebellar cortex, express a multitude of molecules in either a zebrin II-positive or zebrin II-negative parasagittal banding pattern [1, 2]. Both the climbing fiber projection from the inferior olive and the PC projection to the cerebellar nuclei are organized in matching parasagittal zones [3]. Mossy fibers also terminate in longitudinal zones. Climbing and mossy fiber afferents show spatial correspondence with the molecular markers [1]. However, the functional implications of this parasagittal architecture remain elusive [1]. While cerebellar afferents generate parasagittally oriented responses in the cerebellar cortex [1] and synaptic plasticity appears to vary among zones [4], a greater understanding is needed of the physiological properties of the parasagittal zones and how any differences relate to the underlying molecular and cellular architecture. Studies using flavoprotein optical imaging are helping to shed light on the physiological characteristics of the parasagittal zones.

Parasagittal Organization of Molecular Layer Inhibition

Flavoprotein imaging revealed that low-frequency stimulation (10 Hz) of parallel fibers (PFs) evokes not only the expected “beam” of increased fluorescence along the activated

PFs but also parasagittal bands of decreased fluorescence [5]. Where the bands cross the beam, there is a relative decrease in fluorescence. Blocking GABA_A receptors abolishes the bands of decreased fluorescence [5, 6]. Furthermore, PF stimulation evokes a decrease in simple spike firing in PCs located within an inhibitory band but not in PCs located between the bands. Also, intracellular Ca²⁺ decreases within the bands. Together, these findings demonstrate that the bands of decreased fluorescence are due to GABAergic molecular layer inhibition that generates both “off-beam” inhibition lateral to the beam and local “on-beam” inhibition [5].

The inhibitory bands are structural, extending across a folium and align with zebrin II-positive bands [5]. The inhibitory bands are evoked by peripheral stimulation (e.g., stimulation of the ipsilateral vibrissal pad) and modulate the responses to peripheral stimulation, suggesting a role in controlling the spatial patterning of cerebellar cortical activity. Although the signaling mechanism that generates the parasagittal inhibition is not known, newer findings suggest a key role for excitatory amino acid transporter type 4 (EAAT4) [7]. Importantly, EAAT4 is expressed parasagittally on PCs in zebrin II-positive bands [1]. The inhibitory bands may have implications for cerebellar disease. In a transgenic mouse model of spinocerebellar ataxia type 8, the inhibitory banding pattern is lost as is the normal responsiveness to GABA_A antagonists [6]. The reduction in molecular layer inhibition likely disrupts cerebellar cortical function and contributes to the striking motor phenotype in SCA8 mice.

Parasagittal Organization of Long-latency Patches

Flavoprotein imaging uncovered another functional parasagittal compartmentalization. In addition to the beam-like response at a short latency, high-frequency PF stimulation evokes patches of increased fluorescence along the beam at 20–25 s latencies [8]. Consistent with the dependency on high-frequency PF stimulation, these long-latency patches are abolished by type 1 metabotropic glutamate receptor (mGluR₁) antagonists. Ionotropic glutamate receptor antagonists have no effect. The long-latency patches are evoked in zebrin II-positive parasagittal bands. The long-latency patches reflect the release of Ca²⁺ from intracellular stores as blockers of phospholipase C β (PLC β) and ryanodine receptors completely suppress the long-latency patches. The long-latency patches exhibit a robust, mGluR₁-dependent long-term potentiation to high-frequency PF stimulation demonstrating that synaptic plasticity differs among the zones [8, 9].

The finding that the long-latency patches are mediated by mGluR₁ receptors and the release of internal Ca²⁺ suggests a possible mechanism. Both mGluR₁ receptors and the downstream signaling cascade in PCs are organized parasagittally [10]. The PLC β 3 isoform is found on PCs in zebrin II-positive bands while the PLC β 4 isoform and the splice variant, mGluR_{1b}, are found on PCs in complementary zebrin II-negative bands [10]. The mGluR_{1b} isoform has less potency in coupling to the PLC downstream signaling pathway than does the mGluR_{1a} isoform and these differences in potency may be involved in the expression and unique properties of the long-latency patches [8].

In summary, parasagittal zones respond differentially to PF stimulation, including excitability and synaptic plasticity that likely underlies distinct functions. The next challenge is to understand the implications for cerebellar function.

Acknowledgments

We wish to thank Lijuan Zhou and Michael McPhee for technical support and Kris Bettin for manuscript preparation. This study was supported in part by NIH grants NS048944, NS18338, and NS062158 and a grant from the Minnesota Medical Foundation.

References

1. Apps R, Hawkes R. Cerebellar cortical organization: a one-map hypothesis. *Nat Rev Neurosci.* 2009; 10:670–81. [PubMed: 19693030]
2. Hawkes R, Herrup K. Aldolase C/zebrin II and the regionalization of the cerebellum. *J Mol Neurosci.* 1995; 6:147–58. [PubMed: 8672398]
3. Voogd, J.; Bigare, F. Topographical distribution of olivary and corticonuclear fibers in the cerebellum. In: Courville, J.; DeMontigny, C.; Lamarre, Y., editors. *The inferior olivary nucleus.* New York: Raven; 1980. p. 207-34.
4. Wadiche JI, Jahr CE. Patterned expression of Purkinje cell glutamate transporters controls synaptic plasticity. *Nat Neurosci.* 2005; 8:1329–34. [PubMed: 16136036]
5. Gao W, Chen G, Reinert KC, Ebner TJ. Cerebellar cortical molecular layer inhibition is organized in parasagittal zones. *J Neurosci.* 2006; 26:8377–87. [PubMed: 16899733]
6. Moseley ML, Zu T, Ikeda Y, et al. Bidirectional expression of CUG and CAG expansion transcripts and intranuclear polyglutamine inclusions in spinocerebellar ataxia type 8. *Nat Genet.* 2006; 38:758–69. [PubMed: 16804541]
7. Reinert KC, Gao W, Chen G, Wang X, Peng Y, Ebner TJ. Cellular and metabolic origins of flavoprotein autofluorescence in the cerebellar cortex *in vivo*. *Cerebellum.* 2011; 10:585–99. [PubMed: 21503591]
8. Wang X, Chen G, Gao W, Ebner TJ. Parasagittally aligned, mGluR1-dependent patches are evoked at long latencies by parallel fiber stimulation in the mouse cerebellar cortex *in vivo*. *J Neurophysiol.* 2011; 105:1732–46. [PubMed: 21289138]
9. Wang X, Chen G, Gao W, Ebner TJ. Long-term potentiation of the responses to parallel fiber stimulation in mouse cerebellar cortex *in vivo*. *Neurosci.* 2009; 162:713–22.
10. Hartmann J, Konnerth A. Mechanisms of metabotropic glutamate receptor-mediated synaptic signaling in cerebellar Purkinje cells. *Acta Physiol.* 2009; 195:79–90.